Title:

Unique composition and hyperstability of the ventral disc, *Giardia*'s complex microtubule

organelle

Authors:

Nosala, C., Hagen, K.D., Chase, T.M., Jones, K., Loudermilk, R., Nguyen, K., and S.C. Dawson

Address:

Department of Microbiology and Molecular Genetics

One Shields Avenue

University of California, Davis

Davis, CA 95616

530-752-3633

scdawson@ucdavis.edu

Abstract

Giardia is a common protistan parasite that causes diarrheal disease worldwide. Motile

trophozoites colonize the small intestine, attaching to the villi with the ventral disc, a unique

and complex microtubule (MT) organelle. Attachment to the host epithelium allows *Giardia* to

resist peristalsis during infection of the host gastrointestinal tract. Despite our emerging view of

the complexity of ventral disc architecture, we are still in the very preliminary stages of

understanding how specific structural elements contribute to disc stability or generate forces

for attachment. The ventral disc is a large, dome-shaped, spiral MT array decorated with

microribbon-crossbridge protein complexes (MR-CB) that extend upward into the cytoplasm.

Using a new high salt method for disc biochemical fractionation followed by shotgun proteomic

analyses and validation by GFP-tagging, we identified 54 new disc-associated proteins (DAPs).

Of 87 DAPs confirmed to date, 54 localize only to the disc, and the remainder localize to

additional structures such as the flagella, basal bodies, or median body. Almost one third of

DAPs lack any homology to proteins in other eukaryotes and another one third simply contain

ankyrin repeat domains. Many DAPs localize to specific structural regions of the disc, including

the ventral groove region and disc margin. We also demonstrate that the disc is a stable

structure lacking canonical MT dynamic instability and show that the disc structure and

composition remain intact after detergent extraction in up to 2M potassium chloride. Future

genetic, biochemical, and functional analyses of DAPs will be central toward understanding disc

architecture, assembly and dynamics.

Introduction

Microtubules (MTs) in protists assemble into cytoskeletal arrays and organelles that adopt shapes, functions, or regulatory mechanisms not present in model eukaryotic systems (Dawson and Paredez, 2013). The ubiquity and diversity of unique cytoskeletal organelles in microbial eukaryotes underscores the fact that cytoskeletal variation is the norm rather than the exception (Russell et al., 2017). Protistan cytoskeletal organelles are often composed of conserved MAPs as well as proteins that lack homology to proteins in other eukaryotes (Hagen et al., 2011; Hu et al., 2006; Preisner et al., 2016), and thus may be an untapped reservoir of non-canonical MT-binding proteins governing MT assembly, nucleation, or dynamics (Hampl et al., 2009).

One such unique microtubule organelle is present in *Giardia,* a widespread protistan parasite that causes diarrheal disease in humans and livestock (Einarsson et al., 2016). Commonly ingested from contaminated water sources, *Giardia* cysts excyst into flagellated trophozoites and proliferate in the proximal small intestine (Nosala and Dawson, 2015). There they attach to the intestinal microvilli using the ventral disc (Elmendorf et al., 2003; Nosala and Dawson, 2015), a microtubule (MT) organelle (Crossley and Holberton, 1983a; Crossley and Holberton, 1985; Feely et al., 1982; Friend, 1966; Holberton, 1973; Holberton, 1981) 8 μm in diameter with parallel, uniformly spaced MTs that spiral one and a quarter turns to form a dome. An overlap zone occurs between the upper and lower portions, and the ventral groove region extends over the ventral flagella (Figure 1A, B).

The overall architecture of the ventral disc was first described by Cheissin over 50 years

ago (Cheissin, 1964), and abundant MT-associated proteins termed 'giardins' were isolated

from the disc two decades later (Crossley and Holberton, 1983a). Early work described the

overall structural elements of associated with the ventral disc MT spiral array (Friend, 1966),

and the first 3D high-resolution structure of the ventral disc was obtained recently using cryo-

electron tomography (cryo-ET) (Schwartz et al., 2012). Cryo-ET of whole isolated ventral discs

defined an elaborate cytoskeletal architecture with dense protein complexes coating nearly

every protofilament of the microtubule spiral array (Figure 1A). In sum, the disc contains more

than 1.2 mm of tubulin forming roughly one hundred MTs that vary in length from 2 to 18 μm

(Brown et al., 2016). Associated with the entire length of the MT spiral are unique repetitive

substructural elements – the trilaminar microribbons – that extend 150-400 nm dorsally into

the cytoplasm and are believed to stabilize the overall disc structure (Holberton, 1973;

Holberton, 1981) (see Figure 1C, D). Regularly spaced "crossbridge" structures link adjacent

microribbons (Holberton, 1973) and repeat every 16 nm (Figure 1D), corresponding to the

distance of two alpha/beta tubulin dimers. Repeating every 8 nm are sidearms and paddles that

are spaced at the distance of a single alpha/beta tubulin dimer. Other repetitive MT-associated

substructures include three MT-associated proteins (gMAPs 1-3) and three MT inner proteins

(gMIPS 5, 7 and 8). An associated structure at the disc periphery, the lateral crest (Figure 1C),

forms a seal with surfaces in early attachment (Feely, 1990; Feely et al., 1982).

Despite our new understanding of the complexity of ventral disc architecture (Brown et

al., 2016), we still know very little about the composition of the disc or the contribution of

specific structural elements to disc stability and the generation of attachment forces. In a

comprehensive proteomic analysis of detergent-extracted, isolated ventral discs, we previously

identified over twenty new disc-associated proteins (now referred to as DAPs, rather than

giardins) that specifically localize to regions of the ventral disc or lateral crest (Hagen et al.,

2011). The complexity of the disc architecture as revealed by cryo-ET implied that there could

be disc proteins that were still undiscovered.

Here we have identified 54 additional disc-associated proteins (DAPs) using a new

method of disc biochemical fractionation in high salt followed by shotgun proteomic analyses.

Localization of these candidate DAPs to the ventral disc was confirmed by GFP tagging. We also

demonstrate that the complex disc is a "hyperstable" structure in that drugs that normally

affect MT dynamic instability have no effect on disc microtubules (Dawson et al., 2007) and no

turnover of disc-associated proteins (DAPs) has been observed (Dawson et al., 2007; Hagen et

al., 2011). The overall ventral disc structure remains intact after detergent extraction in up to

2M potassium chloride. Further dissection of the mechanisms underlying the disc's unusual

stability will first require thorough knowledge of the functional roles of unique disc-associated

proteins.

Materials and Methods

Giardia **culture conditions**

Giardia strains were cultured in sterile 16 ml screw-capped disposable tubes (BD Falcon)

containing 12 ml modified TYI-S-33 medium supplemented with bovine bile and 5% adult and

5% fetal bovine serum [56]. Cultures were incubated upright at 37°C without shaking and

reached confluence at ~48 hr. Puromycin (50 μg/ml) was added when required to maintain

episomal plasmids in GFP strains.

C-terminal GFP tagging of candidate disc-associated proteins

Giardia GFP strains were created using Gateway cloning as previously described (Hagen et al.,

2011). To add C-terminal GFP tags to candidate DAPs, PCR forward primers were designed to

bind 200 bp upstream of the gene to include the *Giardia* native promoter. Full-length genes

lacking stop codons were amplified from *Giardia lamblia* strain WBC6 genomic DNA using either

PfuTurbo Hotstart PCR Master Mix or Easy-A High Fidelity PCR Master Mix (Agilent) and were

cloned into the ThermoFisher Scientific Gateway entry vectors pENTR/D-TOPO (blunt-end,

directional TOPO cloning) or pCR8/GW/TOPO (for efficient TOPO TA cloning), respectively.

Inserts were sequenced to confirm gene identity and correct orientation. To generate DAP-GFP

fusions, entry vectors were recombined with the *E. coli*/*Giardia* Gateway destination vector

pcGFP1F.pac (Dawson and House, 2010) via LR recombination using LR Clonase II Plus

(ThermoFisher Scientific). Clones were screened by AscI digest and Qiagen's Plasmid Plus Midi

Kit was used to prepare bulk plasmid DNA. To create *Giardia* strains expressing GFP-tagged

candidate DAPs, 20 μg plasmid DNA was introduced into 10^7 *G. lamblia* strain WBC6 (ATCC

50803) trophozoites by electroporation using a Bio-Rad GenePulserXL as previously described

(Dawson et al., 2007). Transfectants were initially selected with 10 μg /ml puromycin; the

antibiotic concentration was increased to 50 μg /ml once cultures reached 50% confluence

(typically 7 to 10 days). Cultures were maintained with selection for at least two weeks prior to

the preparation of frozen stocks.

Biochemical fractionation of the Giardia cytoskeleton

Detergent extraction of *Giardia's* microtubule cytoskeleton was done as previously described

(Hagen et al., 2011). Medium was decanted from one confluent culture tube and trophozoites

were washed *in situ* three times with 5 mls of warmed 1X HEPES Buffered Saline (HBS). To

demembranate cytoskeletons, cells were iced for 15 minutes in the last 1X HBS wash, pelleted

by centrifugation (1000 × g, 5 min), and resuspended in 1 ml of 1X PHEM (60 mM PIPES, 25 mM

HEPES, 10 mM EGTA, 1 mM MgCl2, pH 7.4) containing 1% Triton X-100 and 1M KCl. 1X HALT

protease inhibitor cocktail (Roche) was added to prevent proteolysis. This solution was

transferred to an Eppendorf tube and vortexed continuously for 30 min at medium speed.

Ventral disc cytoskeletons were then pelleted as above and washed twice in 1X PHEM lacking

both Triton X-100 and KCl. Sufficient extraction of cytoskeletons was confirmed by wet mount

using DIC microscopy.

Cytoskeletons were fractionated as previously described (Crossley and Holberton, 1983a;

Crossley and Holberton, 1983b; Crossley and Holberton, 1985) and Figure 2A). Following

detergent extraction, an aliquot of demembranated cytoskeletons in PHEM was retained as

'P1.' Cytoskeletons were then pelleted as described above, washed, and resuspended in CB

buffer (10mM Tris, 1mM EDTA, pH 7.7) for 48 hours to dissolve the crossbridges. Leftover

cytoskeletal complexes were pelleted at 1000 x g for 5 min and the supernatant was retained as

'S2.' Complexes were next washed and resuspended in MR buffer (10mM HEPES, 5mM EDTA,

pH 8.7) for 48 hours to dissolve microribbons. The remaining tubulin complexes were pelleted,

and the supernatant was retained as 'S3.' The pelleted tubulin complexes were resuspended in

1X PHEM and retained as 'P3.'

SDS-PAGE and Western Blotting

A total of 20 µg of total cell lysate in RIPA buffer was run on SDS/PAGE using Precise Tris·HEPES

protein gels (ThermoFisher), 4–20%. Gels were transferred onto 0.22 micron nitrocellulose and

blocked for 1 h at room temperature in 5% (wt/ vol) milk in PBS with 0.05% Tween. Antibodies

were diluted in 5% (wt/vol) milk in PBS with 0.05% Tween as follows: anti-HA 1:5,000, (Sigma)

1:5,000, goat antimouse IgG HRP (Bio-Rad) 1:10000. ECL substrate (Millipore) was used for

detection and development. For SDS-PAGE of fractionated proteins, 20 ul of total undiluted

fraction samples were run as for western blots. Gels were stained in True Blue for 1 hour then

rinsed with MilliQ water and incubated with MilliQ water overnight at room temperature to

help destain.

Proteomic analyses of fractions and mass spectrometry

All MS/MS samples were analyzed using X! Tandem (The GPM, thegpm.org; version X! Tandem

Alanine (2017.2.1.4)). X! Tandem was used to search the uniprotgiardiaintestinalis_Craprev

database (14528 entries) assuming the digestion enzyme trypsin and with a fragment ion mass

tolerance of 20 PPM and a parent ion tolerance of 20 PPM. X! Tandem variable modifications

used were Glu->pyro-Glu of the N-terminus, ammonia-loss of the N-terminus, gln->pyro-Glu of

the N-terminus, deamidated of asparagine and glutamine, oxidation of methionine and

tryptophan and dioxidation of methionine and tryptophan. Scaffold (version Scaffold_4.8.4,

Proteome Software Inc., Portland, OR) was used to validate MS/MS based peptide and protein

identifications. Peptide identifications were accepted if they exceeded specific database search

engine thresholds. Protein identifications were accepted if they contained at least 5 identified

peptides. Proteins that contained similar peptides and could not be differentiated based on

MS/MS analysis alone were grouped to satisfy the principles of parsimony. Proteins sharing

significant peptide evidence were grouped into clusters.

Live imaging of GFP-tagged DAP strains

Strains were thawed from frozen stocks and cultured for 24 to 48 hours prior to live imaging.

Trophozoites were iced 15 min, harvested by centrifugation at 1000 x g, resuspended in warm

medium, and incubated in 96-well black glass bottom imaging plates (Cellvis, Mountain View,

CA) for up to 2 hours at 37C in a nitrogen-enriched atmosphere to promote attachment. The

medium was replaced with 1X HBS, and the trophozoites were incubated under the same

conditions for 30 min. Additional warm HBS washes were performed as needed during imaging

to remove detached cells. Imaging was performed using differential interference contrast (DIC)

and epifluorescence with a Leica DMI6000B wide-field inverted fluorescence microscope

(PlanApo 100X, NA 1.40 oil immersion objective). Optical sections were acquired at 0.2-µm

intervals using Metamorph (MDS Technologies). Images were processed using Fiji, and two-

dimensional maximum intensity projections were created from the three-dimensional data sets

if required.

Structured Illumination Microscopy

3D stacks were collected at 0.125 um intervals on the Nikon N-SIM Structured Illumination

Super-resolution Microscope with a 100x/NA 1.49 objective, 100 EX V-R diffraction grating, and

an Andor iXon3 DU-897E EMCCD. Images were recollected and reconstructed in the "2D-SIM"

mode (no out of focus light removal; reconstruction used three diffraction grating angles each

with three translations).

Transmission Electron Microscopy

Cytoskeleton preps of wild type and knockdown cells were prepared as above and applied to

400 mesh formvar/carbon coated glow-discharged grids. Negative staining was performed by

applying 1% phosphotungstic acid, pH 5.4, and grids were dried by blotting without washes.

For thin sections, *Giardia* trophozoites that were pelleted or attached to ACLAR discs (Electron

Microscopy Sciences) were fixed for 10 minutes in 4% paraformaldehyde and secondarily fixed

for 1 hour in 1% osmium tetroxide. Cells were washed three times with cold ddH2O to remove

fixative, dehydrated through ascending concentrations of ethanol (30%, 50%), and incubated

for 1 hour in 2% uranyl acetate in 50% ethanol. Dehydration continued through 70% and 3 x

95% ethanol, and was completed with three changes in 100% ethanol for a minimum of 10

minutes each change. Cell were embedded in 1:1 epoxy resin:acetone overnight at room

temperature. The next day the resin was removed and replaced with 100% resin twice for two

hours each. The ACLAR discs were placed at the bottom of a flat bottom beam capsule with the

cells facing up and the capsule was filled with fresh resin. The blocks were polymerized at 70C

overnight. The blocks were trimmed, and thin sections were cut with a Leica UCT

ultramicrotome (Leica Ultracut UCT, Leica, Vienna, Austria) and stained with 2% uranyl acetate

in 70% ethanol and lead citrate before viewing in the Talos L120C electron microscope

(FEI/TheromoScientific Company, Hillsboro, OR., U.S.A. made in Eindhoven, The Netherlands) at

100KV. Images were acquired using the fully integrated Ceta CMOS camera.

Results

Biochemical fractionation readily extracts disc and axoneme proteins for shotgun proteomic

analysis and mass spectrometry.

Using cryo-ET of detergent extracted ventral discs, Brown et al. (Brown et al., 2016) recently

defined specific regional variations in the ventral disc architecture, such as the ventral groove,

disc margin and overlap zone areas. To find additional proteins that comprise these structurally

distinct regions of the disc, we developed a novel biochemical fractionation protocol to disrupt

the disc and used proteomic analysis and GFP tagging to identify candidate proteins.

Detergent extraction of *Giardia* trophozoites with Triton X-100 removes membrane and cytosol

from the microtubule cytoskeleton and yields intact ventral discs that are stable in PHEM buffer

for weeks (Crossley and Holberton, 1983a; Crossley and Holberton, 1983b; Crossley and

Holberton, 1985; Holberton and Ward, 1981). Here we extracted trophozoites with both 1%

Triton X-100 and 1M KCl (Figure 2A, P1 fraction). Microtubules and other disc proteins (Figure

2B, C) were retained following the high salt extraction, confirming the overall stability of the

disc and cytoskeleton. Negative staining of the P1 fraction (Figure 2D) showed the loss of the

nuclei, and retention of the disc, flagella, and the funis. Subsequent extractions with TRIS-EDTA and HEPES buffers were used to destabilize and disrupt the disc, resulting in pellet or supernatant fractions (S2, S3, P2, and P3). Using a strain with fluorescently tagged delta-giardin (Figure 2C, D), we determined that these additional treatments caused significant changes to the ventral disc ultrastructure (S3 fraction), including a relaxation of the normally domed and closed ventral disc spiral, and separation of microtubule/microribbon pairs from crossbridges (Figure 2E, F). Flagellar components are removed from the disc fractions in the procedure (Figure 2E, F). SDS-PAGE of each fraction resolved proteins enriched in each pellet (P1, P3) and supernatant (S2, S3) fraction.

Proteomics and mass spectrometry of ventral disc fractions

Following mass spectrometry of each fraction (P1, P3 and S2, S3) and compared the overlap between proteins in various fractions. One hundred and five proteins were identified with at least five hits in any fraction after mass spectrometry (Table 1). Highly abundant known disc-associated proteins were identified in every fraction sampled (tubulin, MBP, delta giardin, etc). Other proteins had significant overlap between all fractions, yet sixty-two proteins were found to be exclusively enriched in S2 as compared to S3.

A revised inventory of disc-associated proteins

In a prior proteomic analysis of detergent-extracted, isolated ventral discs, we identified over

twenty new DAPs that specifically localize to regions of the ventral disc or lateral crest (Hagen

et al., 2011). To determine the cellular localization of candidate DAPs identified here, we again

used C-terminal GFP tagging of candidate DAPs under the control of native promoters (Figure 3,

Figure 4 and Supplemental Figures 1-4). The disc-specific localization of new 54 DAPs was

confirmed with this method, bringing the total number of verified DAPs to 87 (Table 1). Many

of the newly identified DAPs also localize to one or several other cytoskeletal structures (Figure

3A), such as the membrane bound or cytoplasmic regions of the eight axonemes or the eight

basal bodies (31 total DAPs), the median body (23 total DAPs), or the lateral crest structure

surrounding the disc (4 total DAPs). Forty-two of the 87 known DAPs only to specific regions of

the ventral disc, and not to other cytoskeletal structure (Figure 3).

DAP compositional variation in distinct regions of the ventral disc

To classify the 54 new DAPs, we divided them into categories based on their GFP localization to

six distinct regions of the disc: the disc body (body), the disc margin (DM), the overlap zone

(OZ), the ventral groove (VG), the dense bands (DB) and the lateral crest (LC) (Figure 4 and

Table 1)*.* Some DAPs localized to more than one disc region, and some also localized to other

cytoskeletal structures. Overall, the majority of DAPs localize to more than one region of the

disc (Figure 4 and Table 1). Thrity-six DAPs localize to only one of the six disc regions: six DAPS

to the disc body only; three to overlap zone; twenty-two to the disc margin; four to the dense

bands, and only one DAP is associated specifically with the ventral groove. Most DAPs localize

to the disc body, overlap zone, or disc margin regions, which highlights the overall structural

complexity of these areas of the disc relative to other regions such as the dense bands that

have five DAPS.

The majority of DAPs lack known MT-binding motifs

Of the total confirmed DAPs, only two are related to known MT-binding proteins (Figure 5).

Specifically, one DAP has a CAP-GLY domain protein (DAP5374) (Weisbrich et al., 2007), and one

kinesin, kinesin-6a (DAP102455), localizes to the ventral disc in the disc margin region. Twenty-

six of the eight-seven DAPs lack homology to known proteins in other eukaryotes. An

additional twenty-nine DAPs simply contain ankyrin repeat domains (Figure 3C and Figure 4).   A

handful of the identified DAPs are members of conserved protein families, including: three

members of the striated fiber (SF)–assemblins (beta-giardin, delta-giardin, and SALP-1 (Palm et

al., 2003)); four annexin family members (e.g., alpha-giardins (Bauer et al., 1999; Peattie, 1990;

Weiland et al., 2005; Weiland et al., 2003)), and thirteen NEK kinases (Figure 5 and Table 1).

Nine of the thirteen NEKs lack conserved catalytic residues or are pseudokinases which may,

however, still retain kinase activity (Hagen et al., 2011). Other DAPs include those with a WD40

repeat domain (DAP15218), a DUF1126 domain (DAP41512), a Mlf1p domain (DAP16424), an

ENTH domain (DAP3256), or SHIPPO repeat domains (DAP103164 and DAP9148).

Ventral disc composition and integrity is insensitive to microtubule drugs and high salt

extraction.

To evaluate the effect of MT stabilizing or depolymerizing drugs such as Taxol or nocodazole on disc structure, we treated cells with each of these drugs separately and quantified the area and overall structure of the disc body, overlap zone and central bare area (Nosala et al., 2018). Treatment with either MT stabilizing or depolymerizing drugs had no effect on disc area or conformation (Figure 6A). In contrast, other cytoskeletal elements such as the flagella or median body are dynamic and sensitive to these MT drugs (Figure 6B).

We also determined that the detergent extraction of ventral disc cytoskeletons with up to 2M KCl had little effect on the recovery of disc-associated proteins as seen by SDS-PAGE (Figure 6C). Furthermore, the GFP-tagged disc protein DAP16343 (or MBP) remains associated with the disc even after extraction with 1M KCl, and both alpha tubulin and delta-giardin, a component of the microribbons, remain associated with the disc following salt extraction (Figure 6D).

Discussion

Paradigms of microtubule function, dynamics, and nucleation have classically been defined by studies of the mitotic spindle and cilium in model systems. Yet for nearly 300 years, diverse and unique interphase cytoskeletal organelles have also been described in microbial eukaryotes, or protists (Dawson and Paredez, 2013). In these organisms, microtubules are often arranged into higher-order assemblies that have forms and functions that are quite different from those of model organisms (Chaaban and Brouhard, 2017). The complex architecture and understudied functional abilities of non-canonical microtubule organelles like the ventral disc challenge our conceptions of the capabilities of cytoskeletal polymers.

The ventral disc is critical for parasite attachment to the host epithelium, and *Giardia*

trophozoites are believed to attach to surfaces via hydrodynamic suction-based forces that

result in a pressure differential underneath the disc relative to the outside medium (Woessner

and Dawson, 2012). Overall conformational changes of regions of the disc may create or

modulate these attachment forces in the absence of canonical MT dynamics (Woessner and

Dawson, 2012). Defining the molecular mechanisms that underlie disc functions in attachment

requires a more thorough understanding of the novel proteins and complexes that decorate

almost all protofilaments of the disc MTs, and vary in composition and conformation along the

length of the microtubules (Brown et al., 2016).

A dramatically expanded inventory of disc-associated proteins

Given the relatively small number of proteins known to regulate microtubule dynamics and

assembly, how do diverse eukaryotic cells assemble and regulate elaborate microtubule

organelles like the ventral disc? Diverse protistan cytoskeletal structures are often composed of

proteins that lack homology to proteins in other eukaryotes (Hagen et al., 2011; Hu et al., 2006;

Preisner et al., 2016), and thus may be an untapped reservoir of non-canonical MT-binding

proteins governing MT assembly, nucleation, or dynamics (Hampl et al., 2009). Some DAPs may

represent novel MT binding proteins with unique functions, such as facilitating the curved MT

array of the disc. Only one DAP has a bonafide MT binding motif – DAP5374, a CAP-Gly protein

(Weisbrich et al., 2007). One other MT-associated protein– kinesin-6a (DAP102455) – localizes

to the ventral disc, in the disc margin region. Fifty-four of the total 87 DAPs localize exclusively

to the disc, and the remainder also localize to other microtubule structures such as the flagella,

basal bodies, or median body which may indicate their more general MT-binding properties.

Thus, while DAPs are primarily uncharacterized with respect to their microtubule binding or biochemical properties, many still may directly influence ventral disc MT nucleation, MT + end binding, or MT stability. DAPs may also contribute to novel aspects of the disc structure, likely generating and stabilizing the curved spiral array of microtubules (Brown et al., 2016; Schwartz et al., 2012).

Using a new method for disc biochemical fractionation followed by shotgun proteomic analyses, we identified and confirmed 54 new disc-associated proteins (DAPs), adding to the 33 previously described DAPs (Nosala et al., 2018). Many of the 87 known DAPs are likely structural components of the disc architectural elements (e.g., microribbons, crossbridges, disc margin). Overall, the ventral disc is composed of novel, non-homologous proteins and ankyrin repeat proteins (Figure 5), which are known to mediate protein-protein interactions, protein folding, and protein stability (Li et al., 2006). Eighty-five DAPS lack homology to known MT-binding proteins, and more than a third are novel, lacking homology to proteins outside of *Giardia* species – including those of other closely-related diplomonads like *Spironucleus* (Andersson et al., 2007). Among this group may be proteins involved in regulating the assembly or disassembly of the intricate disc architecture.

Close to 30 DAPs simply contain ankyrin repeat domains. Ankyrins can also interact directly with tubulin *in vitro* (Bennett and Davis, 1981; Davis and Bennett, 1984). The 30 disc-associated ankyrin repeat proteins may contribute to disc assembly or architecture, as ankyrin repeat proteins have been associated with the microtubule cytoskeleton in diverse eukaryotes. For

example, in the apicomplexan protist *Toxoplasma gondii*, the MT based apical complex acts as

an invasion machine that is necessary for infection of host cells and parasite replication, and is

constructed from canonical tubulins, non-canonical tubulin polymers, and novel proteins (Hu et

al., 2006; Morrissette, 2015). Like the ventral disc, the conoid is largely composed of ankyrin-

repeat containing proteins that are integral to conoid structure and functioning (Long et al.,

2017). Ankyrin proteins stabilize subsets of microtubules in human erythrocytes, muscles, and

neurons [42]. Thus, the adaptability of ankyrin repeat domains may facilitate the stability of

microtubule organelles in *Giardia* and other eukaryotes.

Giardia has an expanded repertoire of over 70 NEK kinases (Manning et al., 2011), and

13 NEK kinases are associated with the disc. NEKs in the disc could act as structural or

regulatory proteins involved in disc biogenesis or disassembly during cell division. Five NEK

kinase DAPs also have ankyrin repeat domains. Nine of the twelve disc-associated NEKs lack

conserved catalytic residues or are pseudokinases, yet may still retain kinase activity (Hagen et

al., 2011).

Regional variation in disc composition is correlated to similar variations in disc architecture

The strikingly distinct localization patterns of the 54 disc and median body associated DAPs that

delineate the disc (Figure 4 and Supplemental 1-4) are consistent with the structural variations

in the disc defined by cryo-ET (Brown et al., 2016). Due to regional variations in the disc

architecture, a single microtubule can be coated with different DAPs in different regions,

beginning at the "dense band" nucleation zone and terminating at the disc margin (Brown et

al., 2016). The majority of disc microtubules terminate with their plus ends either at the disc

margin or in the overlap zone, with a small subset observed to terminate within the disc body

itself (Brown et al., 2016).

Of the 54 disc and median-body localizing DAPs, six localize specifically to the disc body,

with 22 DAPs the disc margin and two the lateral crest, three to the overlap zone, four to the

dense bands, and one to the ventral groove region (Figure 4). The function and mechanism by

which regional variation in DAP localization is generated is unknown, yet the localization of

DAPs to these specific regions (e.g., disc margin, dense bands, etc.) implies unique molecular

properties associated with these regions.

Do DAPs confer "hyperstability" to the ventral disc?

Emerging microbial eukaryotic model systems like *Giardia* offer a wealth of unexplored

cytoskeletal organelles with unique properties and functions (Russell et al., 2017). The

insensitivity of the disc to MT drugs and its integrity following extraction in high salt, or

"hyperstability," could be explained by the high number of DAPs and protein complexes coating

the disc microtubules (Brown et al., 2016). The extensive microribbon and crossbridge

complexes may also contribute to disc stability, as Holberton and Ward previously observed

that detergent-extracted discs unwind and lose cohesion as crossbridge connections dissociate

over time (Holberton and Ward, 1981). This was particularly evident in the ventral groove

region (Holberton and Ward, 1981). Despite the loss of crossbridges, disc microtubules

remained curved, implying that disc curvature is governed by other structures such as the

microribbons (Holberton, 1981; Holberton and Ward, 1981). We observed the loss of

crossbridges in the S3 fraction, yet curvature of the microtubule/microribbon structure

remained intact (Figure 2E, F).   Holberton argued that mild dissolution of crossbridges results in

a flat disc that is more energetically stable with fully extended CBs, and that outer ventral disc

structures may help to hold the lattice in the spiral conformation (Crossley and Holberton,

1983b). Our lab has observed that the ability of the ventral disc to form a dome is crucial for

functional attachment of *Giardia* to surfaces (Woessner and Dawson, 2012). Therefore, the

crossbridges may be critical for both the structure and function of the ventral disc.

Lastly, the disc is a "hyperstable" structure in that drugs that normally affect MT dynamic

instability have no effect on ventral disc microtubules (Dawson et al., 2007). No turnover of any

disc-associated protein has been reported (Hagen et al., 2011). The large number of unique

microtubule-associated proteins and protein complexes decorating the disc spiral, the disc

margin, and the dense bands may also contribute to the observed hyperstability of the plus and

minus ends of the disc MTs.

Evolution of a complex microtubule organelle for attachment

The ubiquity and diversity of unique cytoskeletal organelles in microbial eukaryotes

underscores the fact that cytoskeletal variation is the norm rather than the exception. The

complex ventral disc spiral MT array and associated structures (e.g., lateral crest) have evolved

only in *Giardia* species, by the cooption of existing proteins or structures or through the

invention of novel MT-binding proteins. Because several components of the disc are MT-

associated proteins (e.g., those with CAP-Gly or kinesin domains) or are related to flagellar root

structures (SF-assemblins (Weber et al., 1993)), portions of the ventral disc may be derived from

ancestral flagellar structures. However, the sheer number of non-homologous and ankyrin

repeat domain proteins in the disc suggests that much of the complexity of the ventral disc may

have evolved through the invention of novel cytoskeletal proteins rather than the elaboration

and modification of an existing cytoskeletal structure like an axoneme.

Given a high-resolution structure and a growing inventory of DAPs, the ongoing

development of molecular genetic and biochemical tools in *Giardia* (Carpenter and Cande,

2009; Dawson et al., 2007; Touz et al., 2005) will be central toward understanding the roles of

specific DAPs in the assembly and functions of the complex disc architecture as well as their

roles in modulating disc conformational dynamics that promote attachment to the host.

Acknowledgements

This work was supported by NIH R01AI077571to SCD and Center for Comparative Medicine at

UC Davis School of Medicine T32 AI060555 Animal Models of Infectious Diseases Training

Program to CN. We graciously thank Dr. Michael Paddy of the UC Davis MCB Microscopy Core

for helpful advice and use of the SDC and SIM microscopes. We also thank Patricia Kysar of the

UC Davis MCB Electron Microscopy Lab for advice and training on TEM and Michelle Salemi and

Brett Phinney at the UCD Proteomics Core for help with mass spectroscopy.

Figure legends

Figure 1. Ventral disc substructures support the microtubule array. A schematic of the ventral

disc indicating the primary structure elements is shown in panel A: disc body, oz: overlap zone,

lc: lateral crest, vg: ventral groove region, db: dense bands, dm: disc margin (A), MT:

microtubule, MR: microribbon, CB: crossbridge (B,C). In B, a negative-stained cytoskeletal

preparation of the ventral disc highlights the primary structural elements as well as the eight

cytoplasmic axonemes (CF = caudal flagella; VF = ventral flagella; PF = posteriolateral flagella;

AF = anterior flagella) and basal bodies (BB). In C, transmission electron microscopy of thin

sections of whole embedded trophozoites show both microribbons (MR) associated with each

MT of the spiral array and the associated lateral crest (LC). Negative staining of the disc (D)

highlights the microribbons (MR) linked with regularly spaced crossbridges (CB). In E, the

overlap zone (OZ) of the MT spiral array, along with the MR-CB complexes are shown in cross

section (N = nucleus).

Figure 2: Biochemical fractionation readily extracts disc and axoneme proteins for shotgun

proteomic analysis and mass spectrometry. Extraction of *Giardia* trophozoites with detergent

and high salt (A) removed membrane and cytosol from the microtubule cytoskeleton (P1).

Subsequent extractions with TRIS-EDTA and HEPES to further disrupted the disc resulting in the

S2, P2, S3 and P3 fractions. Immunostaining of P1 fraction (B, C) retained the disc (green, anti-

delta-giardin) and flagella (purple, anti-alpha-tubulin). Negative staining of the P1 fraction (D)

confirms the loss of the nuclei (N) but retention of the disc (body, vg, oz, lc), flagella (AF, PF)

and the funis (fn); however, greater dissociation of the disc from the axonemes is evident in the

S3 fraction as shown by delta-giardin immunostaining (E, F). SDS-PAGE resolves proteins that

are enriched in each pellet (P1, P3) and supernatant (S2, S3) fraction. Following mass

spectrometry of each fraction (P1, P3 and S2, S3), the Venn Diagram comparison indicates some

overlap between protein the various fractions (H).

Figure 3. Many disc-associated proteins (DAPs) also localize to other cytoskeletal structures.

Subcellular localizations of 87 DAP-GFP fusions identified by fractionation and mass

spectrometry (Figure 2) were categorized into 42 that associated exclusively to the disc (disc

only, B) or localizations of both the disc and other cytoskeletal structures including the flagella

(N=31), the median body (N=12), and lateral crest (N=4). The Venn diagram indicates DAP-GFP

fusions with overlapping localization to the disc, flagella, median body, and lateral crest (A).

Representative localizations are shown for the 12 DAP-fusions localizing to the disc and median

body (mb, C); the 11 localizing to the disc, median body and flagella (mb and fl, D); and the 18

localizing to the disc and flagella only (N=18, E). Two DAPs localized to the lateral crest and disc

(F) and two DAPs localized to the lateral crest and flagella (2). The flagella category includes

localizations to either the basal bodies and /or to the cytoplasmic or membrane bound regions

of the flagella.

Figure 4. Disc-associated proteins localize to specific structurally distinct regions of the

ventral disc. A Venn Diagram (A) shows categorization of the 54 new DAPs by their localization

to distinct regions of the disc including the MT spiral array (body), disc margin (DM), ventral

groove (VG), dense bands (DB), and overlap zone (OZ). In B, representative DAPs defining each

localization category are shown with their functional annotations.

Figure 5. The majority of disc associated proteins (DAPs) lack known microtubule binding domains

PFAM domains are summarized and ranked by abundance for the 87 confirmed *Giardia* DAPs.

CAP_GLY and kinesin motor domains are the only known MT-binding motifs. The five DAPs with

Nek kinase domains that also possess ankyrin repeat domains are not included in the final

summary.

Figure 6. The overall composition and structural integrity of the ventral disc is insensitive to microtubule drugs and high salt extraction. The area and overall structure of the disc body,

overlap zone (oz) and central bare area (ba) are not affected by treatment with either MT

stabilizing or depolymerizing drugs such as Taxol or nocodazole (A). In contrast, other

cytoskeletal elements, including the flagellar, are dynamic and sensitive to MT drugs (B). SDS-

PAGE of fractions (pellet, P, and supernatant S) from 0 to 2M KCl extractions shows that salt

also has little effect on extracted proteins (C). Furthermore, the GFP-tagged disc protein MBP

remains associated with the disc even after extraction with 1M KCl (images in D green), as do α-

tubulin and the microribbon protein δ-giardin as seen via Western blot (D).

Bibliography

1. Russell, J.J., et al., *Non-model model organisms.* BMC Biol, 2017. **15**(1): p. 55.

2. Hu, K., et al., *Cytoskeletal components of an invasion machine--the apical complex of Toxoplasma gondii.* PLoS Pathog, 2006. **2**(2): p. e13.

3. Preisner, H., et al., *The Cytoskeleton of Parabasalian Parasites Comprises Proteins that Share Properties Common to Intermediate Filament Proteins.* Protist, 2016. **167**(6): p. 526-543.

4. Hagen, K.D., et al., *Novel structural components of the ventral disc and lateral crest in Giardia intestinalis.* PLoS Negl Trop Dis, 2011. **5**(12): p. e1442.

5. Hampl, V., et al., *Phylogenomic analyses support the monophyly of Excavata and resolve relationships among eukaryotic "supergroups".* Proc Natl Acad Sci U S A, 2009. **106**(10): p. 3859-64.

6. Einarsson, E., S. Ma'ayeh, and S.G. Svard, *An up-date on Giardia and giardiasis.* Curr Opin Microbiol, 2016. **34**: p. 47-52.

7. Nosala, C. and S.C. Dawson, *The Critical Role of the Cytoskeleton in the Pathogenesis of Giardia.* Curr Clin Microbiol Rep, 2015. **2**(4): p. 155-162.

8. Elmendorf, H.G., S.C. Dawson, and J.M. McCaffery, *The cytoskeleton of Giardia lamblia.* Int J Parasitol, 2003. **33**(1): p. 3-28.

9. Feely, D.E., J.V. Schollmeyer, and S.L. Erlandsen, *Giardia spp.: distribution of contractile proteins in the attachment organelle.* Exp Parasitol, 1982. **53**(1): p. 145-54.

10. Crossley, R. and D.V. Holberton, *Assembly of 2.5 nm filaments from giardin, a protein associated with cytoskeletal microtubules in Giardia.* J Cell Sci, 1985. **78**: p. 205-31.

11. Crossley, R. and D.V. Holberton, *Characterization of proteins from the cytoskeleton of Giardia lamblia.* J Cell Sci, 1983. **59**: p. 81-103.

12. Holberton, D.V., *Fine structure of the ventral disk apparatus and the mechanism of attachment in the flagellate Giardia muris.* J Cell Sci, 1973. **13**(1): p. 11-41.

13. Holberton, D.V., *Arrangement of subunits in microribbons from Giardia.* J Cell Sci, 1981. **47**: p. 167-85.

14. Friend, D.S., *The fine structure of Giardia muris.* J Cell Biol, 1966. **29**(2): p. 317-32.

15. Cheissin, E.M., *Ultrastructure of Lamblia Duodenalis. I. Body Surface, Sucking Disc and Median Bodies.* J Protozool, 1964. **11**: p. 91-8.

16. Schwartz, C.L., et al., *A detailed, hierarchical study of Giardia lamblia's ventral disc reveals novel microtubule-associated protein complexes.* PLoS One, 2012. **7**(9): p. e43783.

17. Brown, J.R., et al., *A detailed look at the cytoskeletal architecture of the Giardia lamblia ventral disc.* J Struct Biol, 2016. **194**(1): p. 38-48.

18. Feely, D.E., Hoberton, D.V., Erlandsen, S.L., *The Biology of Giardia*, in *Giardiasis*, E.A. Meyer, Editor. 1990, Elsevier: Amsterdam. p. 11-50.

19. Dawson, S.C., et al., *Kinesin-13 regulates flagellar, interphase, and mitotic microtubule dynamics in Giardia intestinalis.* Eukaryot Cell, 2007. **6**(12): p. 2354-64.

20. Holberton, D.V. and A.P. Ward, *Isolation of the cytoskeleton from Giardia. Tubulin and a low-molecular-weight protein associated with microribbon structures.* J Cell Sci, 1981. **47**: p. 139-66.

21. Crossley, R. and D.V. Holberton, *Selective extraction with Sarkosyl and repolymerization in vitro of cytoskeleton proteins from Giardia.* J Cell Sci, 1983. **62**: p. 419-38.

22.   Weisbrich, A., et al., *Structure-function relationship of CAP-Gly domains.* Nat Struct Mol

Biol, 2007. **14**(10): p. 959-67.

23.   Palm, J.E., et al., *Identification of immunoreactive proteins during acute human giardiasis.*

J Infect Dis, 2003. **187**(12): p. 1849-59.

24.   Weiland, M.E., et al., *Annexin-like alpha giardins: a new cytoskeletal gene family in Giardia*

lamblia. Int J Parasitol, 2005. **35**(6): p. 617-26.

25.   Weiland, M.E., et al., *Characterisation of alpha-1 giardin: an immunodominant Giardia*

lamblia annexin with glycosaminoglycan-binding activity. Int J Parasitol, 2003. **33**(12): p.

1341-51.

26.   Bauer, B., et al., *Functional identification of alpha 1-giardin as an annexin of Giardia*

lamblia. FEMS Microbiol Lett, 1999. **173**(1): p. 147-53.

27.   Peattie, D.A., *The giardins of Giardia lamblia: genes and proteins with promise.* Parasitol

Today, 1990. **6**(2): p. 52-56.

28.   Dawson, S.C. and A.R. Paredez, *Alternative cytoskeletal landscapes: cytoskeletal novelty*

and evolution in basal excavate protists. Curr Opin Cell Biol, 2013. **25**(1): p. 134-41.

29.   Woessner, D.J. and S.C. Dawson, *The Giardia median body protein is a ventral disc protein*

that is critical for maintaining a domed disc conformation during attachment. Eukaryot

Cell, 2012. **11**(3): p. 292-301.

30.   Andersson, J.O., et al., *A genomic survey of the fish parasite Spironucleus salmonicida*

indicates genomic plasticity among diplomonads and significant lateral gene transfer in

eukaryote genome evolution. BMC Genomics, 2007. **8**: p. 51.

31. Li, J., A. Mahajan, and M.D. Tsai, *Ankyrin repeat: a unique motif mediating protein-protein interactions.* Biochemistry, 2006. **45**(51): p. 15168-78.

32. Davis, J.Q. and V. Bennett, *Brain ankyrin. A membrane-associated protein with binding sites for spectrin, tubulin, and the cytoplasmic domain of the erythrocyte anion channel.* J Biol Chem, 1984. **259**(21): p. 13550-9.

33. Bennett, V. and J. Davis, *Erythrocyte ankyrin: immunoreactive analogues are associated with mitotic structures in cultured cells and with microtubules in brain.* Proc Natl Acad Sci U S A, 1981. **78**(12): p. 7550-4.

34. Morrissette, N., *Targeting Toxoplasma tubules: tubulin, microtubules, and associated proteins in a human pathogen.* Eukaryot Cell, 2015. **14**(1): p. 2-12.

35. Long, S., et al., *A conserved ankyrin repeat-containing protein regulates conoid stability, motility and cell invasion in Toxoplasma gondii.* Nat Commun, 2017. **8**(1): p. 2236.

36. Manning, G., et al., *The minimal kinome of Giardia lamblia illuminates early kinase evolution and unique parasite biology.* Genome biology, 2011. **12**(7): p. R66.

37. Palm, D., et al., *Developmental changes in the adhesive disk during Giardia differentiation.* Mol Biochem Parasitol, 2005. **141**(2): p. 199-207.

38. Tumova, P., J. Kulda, and E. Nohynkova, *Cell division of Giardia intestinalis: assembly and disassembly of the adhesive disc, and the cytokinesis.* Cell Motil Cytoskeleton, 2007. **64**(4): p. 288-98.

39. Davids, B.J., et al., *Giardia lamblia aurora kinase: a regulator of mitosis in a binucleate parasite.* Int J Parasitol, 2008. **38**(3-4): p. 353-69.

40. Davids, B.J., et al., *An atypical proprotein convertase in Giardia lamblia differentiation.* Mol Biochem Parasitol, 2011. **175**(2): p. 169-80.

41. Ellis, J.G., M. Davila, and R. Chakrabarti, *Potential involvement of extracellular signal-regulated kinase 1 and 2 in encystation of a primitive eukaryote, Giardia lamblia. Stage-specific activation and intracellular localization.* J Biol Chem, 2003. **278**(3): p. 1936-45.

42. Touz, M.C., J.T. Conrad, and T.E. Nash, *A novel palmitoyl acyl transferase controls surface protein palmitoylation and cytotoxicity in Giardia lamblia.* Mol Microbiol, 2005. **58**(4): p. 999-1011.

43. Carpenter, M.L. and W.Z. Cande, *Using morpholinos for gene knockdown in Giardia intestinalis.* Eukaryot Cell, 2009. **8**(6): p. 916-9.

44. Weber, K., et al., *SF-assemblin, the structural protein of the 2-nm filaments from striated microtubule associated fibers of algal flagellar roots, forms a segmented coiled coil.* J Cell Biol, 1993. **121**(4): p. 837-45.

1. Russell, J.J., et al., *Non-model model organisms.* BMC Biol, 2017. **15**(1): p. 55.

2. Hu, K., et al., *Cytoskeletal components of an invasion machine--the apical complex of Toxoplasma gondii.* PLoS Pathog, 2006. **2**(2): p. e13.

3. Preisner, H., et al., *The Cytoskeleton of Parabasalian Parasites Comprises Proteins that Share Properties Common to Intermediate Filament Proteins.* Protist, 2016. **167**(6): p. 526-543.

4. Hagen, K.D., et al., *Novel structural components of the ventral disc and lateral crest in Giardia intestinalis.* PLoS Negl Trop Dis, 2011. **5**(12): p. e1442.

5. Hampl, V., et al., *Phylogenomic analyses support the monophyly of Excavata and resolve relationships among eukaryotic "supergroups".* Proc Natl Acad Sci U S A, 2009. **106**(10): p. 3859-64.

6. Morrissette, N., *Targeting Toxoplasma tubules: tubulin, microtubules, and associated proteins in a human pathogen.* Eukaryot Cell, 2015. **14**(1): p. 2-12.

7. Einarsson, E., S. Ma'ayeh, and S.G. Svard, *An up-date on Giardia and giardiasis.* Curr Opin Microbiol, 2016. **34**: p. 47-52.

8. Nosala, C. and S.C. Dawson, *The Critical Role of the Cytoskeleton in the Pathogenesis of Giardia.* Curr Clin Microbiol Rep, 2015. **2**(4): p. 155-162.

9. Feely, D.E., J.V. Schollmeyer, and S.L. Erlandsen, *Giardia spp.: distribution of contractile proteins in the attachment organelle.* Exp Parasitol, 1982. **53**(1): p. 145-54.

10. Crossley, R. and D.V. Holberton, *Assembly of 2.5 nm filaments from giardin, a protein associated with cytoskeletal microtubules in Giardia.* J Cell Sci, 1985. **78**: p. 205-31.

11. Crossley, R. and D.V. Holberton, *Characterization of proteins from the cytoskeleton of Giardia lamblia.* J Cell Sci, 1983. **59**: p. 81-103.

12. Holberton, D.V., *Fine structure of the ventral disk apparatus and the mechanism of attachment in the flagellate Giardia muris.* J Cell Sci, 1973. **13**(1): p. 11-41.

13. Holberton, D.V., *Arrangement of subunits in microribbons from Giardia.* J Cell Sci, 1981. **47**: p. 167-85.

14. Friend, D.S., *The fine structure of Giardia muris.* J Cell Biol, 1966. **29**(2): p. 317-32.

15. Cheissin, E.M., *Ultrastructure of Lamblia Duodenalis. I. Body Surface, Sucking Disc and Median Bodies.* J Protozool, 1964. **11**: p. 91-8.

16. Schwartz, C.L., et al., *A detailed, hierarchical study of Giardia lamblia's ventral disc reveals novel microtubule-associated protein complexes.* PLoS One, 2012. **7**(9): p. e43783.

17. Brown, J.R., et al., *A detailed look at the cytoskeletal architecture of the Giardia lamblia ventral disc.* J Struct Biol, 2016. **194**(1): p. 38-48.

18. Feely, D.E., Hoberton, D.V., Erlandsen, S.L., *The Biology of Giardia*, in *Giardiasis*, E.A. Meyer, Editor. 1990, Elsevier: Amsterdam. p. 11-50.

19. Dawson, S.C., et al., *Kinesin-13 regulates flagellar, interphase, and mitotic microtubule dynamics in Giardia intestinalis.* Eukaryot Cell, 2007. **6**(12): p. 2354-64.

20. Li, J., A. Mahajan, and M.D. Tsai, *Ankyrin repeat: a unique motif mediating protein-protein interactions.* Biochemistry, 2006. **45**(51): p. 15168-78.

21. Dawson, S.C. and S.A. House, *Imaging and analysis of the microtubule cytoskeleton in Giardia.* Methods Cell Biol, 2010. **97**: p. 307-39.

22. Holberton, D.V. and A.P. Ward, *Isolation of the cytoskeleton from Giardia. Tubulin and a low-molecular-weight protein associated with microribbon structures.* J Cell Sci, 1981. **47**: p. 139-66.

23. Crossley, R. and D.V. Holberton, *Selective extraction with Sarkosyl and repolymerization in vitro of cytoskeleton proteins from Giardia.* J Cell Sci, 1983. **62**: p. 419-38.

24. Nosala, C., K.D. Hagen, and S.C. Dawson, *'Disc-o-Fever': Getting Down with Giardia's Groovy Microtubule Organelle.* Trends Cell Biol, 2018. **28**(2): p. 99-112.

25. Aurrecoechea, C., et al., *GiardiaDB and TrichDB: integrated genomic resources for the eukaryotic protist pathogens Giardia lamblia and Trichomonas vaginalis.* Nucleic Acids Res, 2009. **37**(Database issue): p. D526-30.

26. Nohria, A., R.A. Alonso, and D.A. Peattie, *Identification and characterization of gamma giardin and the gamma giardin gene from Giardia lamblia.* Mol Biochem Parasitol, 1992. **56**(1): p. 27-37.

27. Woessner, D.J. and S.C. Dawson, *The Giardia median body protein is a ventral disc protein that is critical for maintaining a domed disc conformation during attachment.* Eukaryot Cell, 2012. **11**(3): p. 292-301.

28. Palm, J.E., et al., *Identification of immunoreactive proteins during acute human giardiasis.* J Infect Dis, 2003. **187**(12): p. 1849-59.

29. Weiland, M.E., et al., *Annexin-like alpha giardins: a new cytoskeletal gene family in Giardia lamblia.* Int J Parasitol, 2005. **35**(6): p. 617-26.

30. Weiland, M.E., et al., *Characterisation of alpha-1 giardin: an immunodominant Giardia lamblia annexin with glycosaminoglycan-binding activity.* Int J Parasitol, 2003. **33**(12): p. 1341-51.

31. Bauer, B., et al., *Functional identification of alpha 1-giardin as an annexin of Giardia lamblia.* FEMS Microbiol Lett, 1999. **173**(1): p. 147-53.

32. Peattie, D.A., *The giardins of Giardia lamblia: genes and proteins with promise.* Parasitol Today, 1990. **6**(2): p. 52-56.

33. Manning, G., et al., *The minimal kinome of Giardia lamblia illuminates early kinase evolution and unique parasite biology.* Genome biology, 2011. **12**(7): p. R66.

34. O'Regan, L., J. Blot, and A.M. Fry, *Mitotic regulation by NIMA-related kinases.* Cell Div, 2007. **2**: p. 25.

35. Morrison, H.G., et al., *Genomic minimalism in the early diverging intestinal parasite Giardia lamblia.* Science, 2007. **317**(5846): p. 1921-6.

36. Weisbrich, A., et al., *Structure-function relationship of CAP-Gly domains.* Nat Struct Mol Biol, 2007. **14**(10): p. 959-67.

37. Pfannenschmid, F., et al., *Chlamydomonas DIP13 and human NA14: a new class of proteins associated with microtubule structures is involved in cell division.* J Cell Sci, 2003. **116**(Pt 8): p. 1449-62.

38. Fritz-Laylin, L.K., et al., *The genome of Naegleria gruberi illuminates early eukaryotic versatility.* Cell, 2010. **140**(5): p. 631-42.

39. Touz, M.C., J.T. Conrad, and T.E. Nash, *A novel palmitoyl acyl transferase controls surface protein palmitoylation and cytotoxicity in Giardia lamblia.* Mol Microbiol, 2005. **58**(4): p. 999-1011.

40. Carpenter, M.L. and W.Z. Cande, *Using morpholinos for gene knockdown in Giardia intestinalis.* Eukaryot Cell, 2009. **8**(6): p. 916-9.

41. Long, S., et al., *A conserved ankyrin repeat-containing protein regulates conoid stability, motility and cell invasion in Toxoplasma gondii.* Nat Commun, 2017. **8**(1): p. 2236.

42. Goellner, B. and H. Aberle, *The synaptic cytoskeleton in development and disease.* Dev Neurobiol, 2012. **72**(1): p. 111-25.

43. Davis, J.Q. and V. Bennett, *Brain ankyrin. A membrane-associated protein with binding sites for spectrin, tubulin, and the cytoplasmic domain of the erythrocyte anion channel.* J Biol Chem, 1984. **259**(21): p. 13550-9.

44. Bennett, V. and J. Davis, *Erythrocyte ankyrin: immunoreactive analogues are associated with mitotic structures in cultured cells and with microtubules in brain.* Proc Natl Acad Sci U S A, 1981. **78**(12): p. 7550-4.

45. Weber, K., et al., *SF-assemblin, the structural protein of the 2-nm filaments from striated microtubule associated fibers of algal flagellar roots, forms a segmented coiled coil.* J Cell Biol, 1993. **121**(4): p. 837-45.

46. Francia, M.E., et al., *Cell division in Apicomplexan parasites is organized by a homolog of the striated rootlet fiber of algal flagella.* PLoS Biol, 2012. **10**(12): p. e1001444.

47. Hansen, W.R., et al., *Giardia lamblia attachment force is insensitive to surface treatments.* Eukaryot Cell, 2006. **5**(4): p. 781-3.

48. House, S.A., et al., *Giardia flagellar motility is not directly required to maintain attachment to surfaces.* PLoS Pathog, 2011. **7**(8): p. e1002167.

49. Dawson, S.C. and A.R. Paredez, *Alternative cytoskeletal landscapes: cytoskeletal novelty and evolution in basal excavate protists.* Curr Opin Cell Biol, 2013. **25**(1): p. 134-41.

1.

Andersson, J. O., Sjogren, A. M., Horner, D. S., Murphy, C. A., Dyal, P. L., Svard, S. G.,

Logsdon, J. M., Jr., Ragan, M. A., Hirt, R. P. and Roger, A. J. (2007). A genomic survey of the fish

parasite Spironucleus salmonicida indicates genomic plasticity among diplomonads and

significant lateral gene transfer in eukaryote genome evolution. *BMC Genomics* **8**, 51.

Bauer, B., Engelbrecht, S., Bakker-Grunwald, T. and Scholze, H. (1999). Functional

identification of alpha 1-giardin as an annexin of *Giardia lamblia*. *FEMS Microbiol Lett* **173**, 147-

53.

Bennett, V. and Davis, J. (1981). Erythrocyte ankyrin: immunoreactive analogues are

associated with mitotic structures in cultured cells and with microtubules in brain. *Proc Natl Acad*

Sci U S A **78**, 7550-4.

Brown, J. R., Schwartz, C. L., Heumann, J. M., Dawson, S. C. and Hoenger, A. (2016). A

detailed look at the cytoskeletal architecture of the Giardia lamblia ventral disc. *J Struct Biol* **194**,

38-48.

Carpenter, M. L. and Cande, W. Z. (2009). Using morpholinos for gene knockdown in

Giardia intestinalis. *Eukaryot Cell* **8**, 916-9.

Chaaban, S. and Brouhard, G. J. (2017). A microtubule bestiary: structural diversity in

tubulin polymers. *Mol Biol Cell* **28**, 2924-2931.

Cheissin, E. M. (1964). Ultrastructure of Lamblia Duodenalis. I. Body Surface, Sucking Disc

and Median Bodies. *J Protozool* **11**, 91-8.

Crossley, R. and Holberton, D. V. (1983a). Characterization of proteins from the

cytoskeleton of *Giardia lamblia*. *J Cell Sci* **59**, 81-103.

Crossley, R. and Holberton, D. V. (1983b). Selective extraction with Sarkosyl and repolymerization in vitro of cytoskeleton proteins from *Giardia*. *J Cell Sci* **62**, 419-38.

Crossley, R. and Holberton, D. V. (1985). Assembly of 2.5 nm filaments from giardin, a protein associated with cytoskeletal microtubules in *Giardia*. *J Cell Sci* **78**, 205-31.

Davis, J. Q. and Bennett, V. (1984). Brain ankyrin. A membrane-associated protein with binding sites for spectrin, tubulin, and the cytoplasmic domain of the erythrocyte anion channel. *J Biol Chem* **259**, 13550-9.

Dawson, S. C. and House, S. A. (2010). Imaging and analysis of the microtubule cytoskeleton in *Giardia*. *Methods Cell Biol* **97**, 307-39.

Dawson, S. C. and Paredez, A. R. (2013). Alternative cytoskeletal landscapes: cytoskeletal novelty and evolution in basal excavate protists. *Curr Opin Cell Biol* **25**, 134-41.

Dawson, S. C., Sagolla, M. S., Mancuso, J. J., Woessner, D. J., House, S. A., Fritz-Laylin, L. and Cande, W. Z. (2007). Kinesin-13 regulates flagellar, interphase, and mitotic microtubule dynamics in Giardia intestinalis. *Eukaryot Cell* **6**, 2354-64.

Einarsson, E., Ma'ayeh, S. and Svard, S. G. (2016). An up-date on Giardia and giardiasis. *Curr Opin Microbiol* **34**, 47-52.

Elmendorf, H. G., Dawson, S. C. and McCaffery, J. M. (2003). The cytoskeleton of *Giardia lamblia*. *Int J Parasitol* **33**, 3-28.

Feely, D. E., Hoberton, D.V., Erlandsen, S.L. (1990). The Biology of *Giardia*. In *Giardiasis*, vol. 3 (ed. E. A. Meyer), pp. 11-50. Amsterdam: Elsevier.

Feely, D. E., Schollmeyer, J. V. and Erlandsen, S. L. (1982). *Giardia* spp.: distribution of contractile proteins in the attachment organelle. *Exp Parasitol* **53**, 145-54.

Friend, D. S. (1966). The fine structure of *Giardia muris*. *J Cell Biol* **29**, 317-32.

Hagen, K. D., Hirakawa, M. P., House, S. A., Schwartz, C. L., Pham, J. K., Cipriano, M. J., De La Torre, M. J., Sek, A. C., Du, G., Forsythe, B. M. et al. (2011). Novel structural components of the ventral disc and lateral crest in Giardia intestinalis. *PLoS Negl Trop Dis* **5**, e1442.

Hampl, V., Hug, L., Leigh, J. W., Dacks, J. B., Lang, B. F., Simpson, A. G. and Roger, A. J. (2009). Phylogenomic analyses support the monophyly of Excavata and resolve relationships among eukaryotic "supergroups". *Proc Natl Acad Sci U S A* **106**, 3859-64.

Holberton, D. V. (1973). Fine structure of the ventral disk apparatus and the mechanism of attachment in the flagellate *Giardia muris*. *J Cell Sci* **13**, 11-41.

Holberton, D. V. (1981). Arrangement of subunits in microribbons from *Giardia*. *J Cell Sci* **47**, 167-85.

Holberton, D. V. and Ward, A. P. (1981). Isolation of the cytoskeleton from *Giardia*. Tubulin and a low-molecular-weight protein associated with microribbon structures. *J Cell Sci* **47**, 139-66.

Hu, K., Johnson, J., Florens, L., Fraunholz, M., Suravajjala, S., DiLullo, C., Yates, J., Roos, D. S. and Murray, J. M. (2006). Cytoskeletal components of an invasion machine--the apical complex of Toxoplasma gondii. *PLoS Pathog* **2**, e13.

Li, J., Mahajan, A. and Tsai, M. D. (2006). Ankyrin repeat: a unique motif mediating protein-protein interactions. *Biochemistry* **45**, 15168-78.

Long, S., Anthony, B., Drewry, L. L. and Sibley, L. D. (2017). A conserved ankyrin repeat-containing protein regulates conoid stability, motility and cell invasion in Toxoplasma gondii. *Nat Commun* **8**, 2236.

Manning, G., Reiner, D. S., Lauwaet, T., Dacre, M., Smith, A., Zhai, Y., Svard, S. and Gillin,

F. D. (2011). The minimal kinome of Giardia lamblia illuminates early kinase evolution and unique

parasite biology. *Genome Biol* **12**, R66.

Morrissette, N. (2015). Targeting Toxoplasma tubules: tubulin, microtubules, and

associated proteins in a human pathogen. *Eukaryot Cell* **14**, 2-12.

Nosala, C. and Dawson, S. C. (2015). The Critical Role of the Cytoskeleton in the

Pathogenesis of Giardia. *Curr Clin Microbiol Rep* **2**, 155-162.

Nosala, C., Hagen, K. D. and Dawson, S. C. (2018). 'Disc-o-Fever': Getting Down with

Giardia's Groovy Microtubule Organelle. *Trends Cell Biol* **28**, 99-112.

Palm, J. E., Weiland, M. E., Griffiths, W. J., Ljungstrom, I. and Svard, S. G. (2003).

Identification of immunoreactive proteins during acute human giardiasis. *J Infect Dis* **187**, 1849-

59.

Peattie, D. A. (1990). The giardins of *Giardia lamblia*: genes and proteins with promise.

Parasitol Today **6**, 52-56.

Preisner, H., Karin, E. L., Poschmann, G., Stuhler, K., Pupko, T. and Gould, S. B. (2016).

The Cytoskeleton of Parabasalian Parasites Comprises Proteins that Share Properties Common to

Intermediate Filament Proteins. *Protist* **167**, 526-543.

Russell, J. J., Theriot, J. A., Sood, P., Marshall, W. F., Landweber, L. F., Fritz-Laylin, L.,

Polka, J. K., Oliferenko, S., Gerbich, T., Gladfelter, A. et al. (2017). Non-model model organisms.

BMC Biol **15**, 55.

Schwartz, C. L., Heumann, J. M., Dawson, S. C. and Hoenger, A. (2012). A detailed,

hierarchical study of Giardia lamblia's ventral disc reveals novel microtubule-associated protein

complexes. *PLoS One* **7**, e43783.

Touz, M. C., Conrad, J. T. and Nash, T. E. (2005). A novel palmitoyl acyl transferase

controls surface protein palmitoylation and cytotoxicity in *Giardia lamblia*. *Mol Microbiol* **58**, 999-

1011.

Weber, K., Geisler, N., Plessmann, U., Bremerich, A., Lechtreck, K. F. and Melkonian, M.

(1993). SF-assemblin, the structural protein of the 2-nm filaments from striated microtubule

associated fibers of algal flagellar roots, forms a segmented coiled coil. *J Cell Biol* **121**, 837-45.

Weiland, M. E., McArthur, A. G., Morrison, H. G., Sogin, M. L. and Svard, S. G. (2005).

Annexin-like alpha giardins: a new cytoskeletal gene family in *Giardia lamblia*. *Int J Parasitol* **35**,

617-26.

Weiland, M. E., Palm, J. E., Griffiths, W. J., McCaffery, J. M. and Svard, S. G. (2003).

Characterisation of alpha-1 giardin: an immunodominant *Giardia lamblia* annexin with

glycosaminoglycan-binding activity. *Int J Parasitol* **33**, 1341-51.

Weisbrich, A., Honnappa, S., Jaussi, R., Okhrimenko, O., Frey, D., Jelesarov, I.,

Akhmanova, A. and Steinmetz, M. O. (2007). Structure-function relationship of CAP-Gly

domains. *Nat Struct Mol Biol* **14**, 959-67.

Woessner, D. J. and Dawson, S. C. (2012). The Giardia median body protein is a ventral

disc protein that is critical for maintaining a domed disc conformation during attachment.

Eukaryot Cell **11**, 292-301.

A

CB
MR
CB
MT

dm
oz
db
disc
body
vg
lc

B

AF
oz
db
BB
disc
body
vg
CF VF PF
lc

C

MR
MT
LC

D

CB
MR

E

N
OZ
lc

A

P1 1M KCl
1X PHEM pH6.9
2% TX100

S2 TRIS-EDTA, pH7.7
CB
LC
BB
DM

S3 HEPES-EDTA, pH 8.7
MR
P3
disc MTs
MAPs

B P1

C P1

D
body bb oz
vg AF
fn lc
PF
N P1

E S3

F S3

G
kDa P1 S2 S3 P3

H
S3 S2
0 24
P1 1 3 1 P3
10 17 1 9
48 35 1
3 1
3

A

flagella   median body

0   0

17   0   0

disc   42   11   0   0   lateral crest

12   0   0

0   3

total DAPs =87

B   disc only (42)

DAP4410

C   disc/median body (12)

DAP5188   ◄mb

D   disc/mb/flagella (11)

DAP41512   fl ◄   ◄ mb

E   disc/flagella (18)

DAP10893

F   disc/lateral crest (2)

DAP23492

G   lateral crest/flagella (2)

DAP15218