Using Weibull accelerated failure time regression model to predict survival time and life expectancy

Enwu Liu[1,2*]

1 Musculoskeletal Health and Ageing Research Program, Mary MacKillop Institute for Health Research, Australian Catholic University, Melbourne, Victoria, Australia
2 College of Medicine and Public Health, Flinders University, Adelaide, South Australia, Australia

* enwu.liu@acu.edu.au

Abstract

Predict mean time to failure (MTTF) or mean time between failures(MTBF) and median survival time are quite common in Engineering reliability researches. In medical literature most prediction models are used to predict probabilities during a certain period. In this paper we introduced detailed calculations to predict different survival times using Weibull accelerated failure time regression model and assessed the accuracy of the point predictions. The method to construct confidence interval for the predicted survival time was also discussed.

Weibull distribution

₁

The Weibull distribution is also called type III extreme value distribution. [1] The distribution has three parameters, the location parameter μ, the scale parameter ρ and the shape parameter γ. The location parameter μ was predetermined as the minimum value in the distribution, we usually choose 0 as the minimum value for survival analysis, therefore, the distribution is reduced to a two-parameter distribution.

The cumulative distribution function(CDF) for a two parameter Weibull distributed random variable $T \sim W(\rho, \gamma)$ is given as

$$F_T(t; \rho, \gamma) = 1 - \exp\left[-(\frac{t}{\rho})^\gamma\right] \tag{1}$$

for $t \geq 0, \rho > 0, \gamma > 0$ and $F(t; \rho, \gamma) = 0$ for $t < 0$

The probability density function(PDF) of the Weibull distribution is given as

$$f_T(t; \rho, \gamma) = F^{'}(t; \rho, \gamma) = \frac{\gamma}{\rho}(\frac{t}{\rho})^{\gamma-1} \exp\left[-(\frac{t}{\rho})^\gamma\right] \tag{2}$$

The survival function of the Weibull distribution is given as

$$S_T(t) = 1 - F_T(t) = \exp\left[-(\frac{t}{\rho})^\gamma\right] \tag{3}$$

The mean survival time or mean time to failure(MTTF) is given as

$$E(T) = \int_0^\infty S(t)dt \qquad \text{(Darth Vader rule)}$$

$$= \int_0^\infty \exp\left[-(\frac{t}{\rho})^\gamma\right] dt \qquad \text{let } (\frac{t}{\rho})^\gamma = u \Rightarrow t = \rho u^{\frac{1}{\gamma}}$$

$$= \int_0^\infty e^{-u} \rho \frac{1}{\gamma} u^{\frac{1}{\gamma}-1} du$$

$$= \rho\frac{1}{\gamma}\Gamma(\frac{1}{\gamma}) \qquad \text{note } \frac{1}{\gamma}\Gamma(\frac{1}{\gamma}) = \Gamma(\frac{1}{\gamma}+1)$$

$$= \rho\Gamma(\frac{1}{\gamma}+1) \tag{4}$$

Log-Weibull distribution

The log-Weibull distribution is also called Gumbel distribution, or type I extreme value distribution. [2] Let $T \sim W(\rho, \gamma)$ and $Y = g(T) = logT$ is a one-one transformation from support $\mathcal{T} = \{t|t > 0\}$ to $\mathcal{Y} = \{y| -\infty < y < \infty\}$. The inverse of Y is

$$T = g^{-1}(Y) = e^Y$$

and the Jacobian is given as

$$|J| = |\frac{dg^{-1}(Y)}{dY}| = e^Y$$

The probability density function of Y is then derived from (2)

$$f_Y(y) = f_T(g^{-1}(y))|J| = \frac{\gamma}{\rho}(\frac{e^y}{\rho})^{\gamma-1}\exp\left[-(\frac{e^y}{\rho})^\gamma\right]e^y = \frac{\gamma}{\rho}\frac{e^{y(\gamma-1)}}{\rho^{(\gamma-1)}}e^y\exp\left[-\frac{e^{\gamma y}}{\rho^\gamma}\right]$$

$$= \gamma\frac{e^{\gamma y}}{\rho^\gamma}\exp\left[-\frac{e^{\gamma y}}{\rho^\gamma}\right] \qquad \text{note } (\rho^\gamma = e^{\gamma\log\rho})$$

$$= \gamma\frac{e^{\gamma y}}{e^{\gamma log\rho}}\exp\left[-\frac{e^{\gamma y}}{e^{\gamma log\rho}}\right] = \gamma e^{\gamma y - \gamma\log\rho}\exp\left[-e^{\gamma y - \gamma log\rho}\right]$$

$$= \gamma e^{\gamma(y-\log\rho)}\exp\left[-e^{\gamma(y-\log\rho)}\right] \qquad (\text{let } \gamma = \frac{1}{b} \text{ and } \log\rho = a)$$

$$= \frac{1}{b}\exp(\frac{y-a}{b})\exp\left[-\exp(\frac{y-a}{b})\right] \text{ where } -\infty < y < \infty \tag{5}$$

It shows log-Weibull distribution has a Gumbel distribution $G(a,b)$, where $a = \log\rho$ and $b = \frac{1}{\gamma}$. The cumulative distribution function of the log-Weibull distribution can be derived from the above PDF or by the definition directly.

$$F_Y(y) = P(Y \le y) = P(log(T) \le y) = P(T \le e^y) = F_T(e^y)$$

By Eq(1) we get

$$F_Y(y) = F_T(e^y) = 1 - \exp\left[-(\frac{e^y}{a})^\gamma\right] = 1 - \exp\left[-\frac{e^{\gamma y}}{a^\gamma}\right]$$

$$= 1 - \exp\left[-\frac{e^{\gamma y}}{e^{\gamma\log\rho}}\right] = 1 - \exp\left[-e^{\gamma(y-\log\rho)}\right]$$

$$= 1 - \exp\left[-\exp(\frac{y-a}{b})\right] \text{ where } \gamma = \frac{1}{b}, \log\rho = a \tag{6}$$

The survival function of $Y = \log(T)$ is given by

$$S_Y(y) = 1 - F_Y(y) = \exp\left[-\exp(\frac{y-a}{b})\right] \qquad (7)$$

The hazard function is given by

$$h_Y(y) = \frac{f_Y(y)}{S_Y(y)} = \frac{1}{b}\exp(\frac{y-a}{b})$$

Weibull AFT regression model

Let T be the survival time. Suppose we have a random sample of size n from a target population. For a subject $i(i = 1, 2, ..., n)$, we have observed values of covariates $x_{i1}, x_{i2}, ..., x_{ip}$ and possibly censored survival time t_i. We can write Weibull accelerated failure time (AFT) model as

$$\log(t_i) = \beta_0 + \beta_1 x_{i1} + ... + \beta_p x_{ip} + \sigma\epsilon_i = \boldsymbol{x'_i}\boldsymbol{\beta} + \sigma\epsilon_i, \qquad i = 1, 2, ..., n \qquad (8)$$

where $\boldsymbol{\beta} = (\beta_0, ..., \beta_p)$ are the regression coefficients of interest, σ is a scale parameter and $\epsilon_1, ...\epsilon_n$ are i.i.d distributed according to Gumbel distribution with the PDF

$$f_\epsilon(x) = \exp(x)\exp\left[-\exp(x)\right] \qquad (9)$$

and the CDF

$$F_\epsilon(x) = 1 - \exp\left[-\exp(x)\right] \qquad (10)$$

Note this is equal to $a = 0, b = 1$ in Eq(5) and Eq(6). We denote it as $G(0, 1)$ distribution or a standard Gumbel distribution.

Now, let us find the PDF of T from Eq(8)

$$\log(T) = \boldsymbol{x'}\boldsymbol{\beta} + \sigma\epsilon$$
$$\Rightarrow T = e^{\boldsymbol{x'}\boldsymbol{\beta} + \sigma\epsilon}$$
$$\Rightarrow \epsilon = g^{-1}(T) = \frac{\log(T) - \boldsymbol{x'}\boldsymbol{\beta}}{\sigma} \qquad (11)$$
$$\Rightarrow |J| = |\frac{d(g^{-1}(T))}{dT}| = \frac{1}{\sigma T} \qquad (12)$$

Put Eq(11) and Eq(12) to Eq(9) we get

$$f_T(t) = f_\epsilon(g^{-1}(t))|J| = \exp(\frac{\log(t) - \boldsymbol{x'}\boldsymbol{\beta}}{\sigma})\exp\left[-\exp(\frac{\log(t) - \boldsymbol{x'}\boldsymbol{\beta}}{\sigma})\right]\frac{1}{\sigma t} \qquad (13)$$

$$= (\frac{t}{\exp(\boldsymbol{x'}\boldsymbol{\beta})})^{\frac{1}{\sigma}}\exp\left[-(\frac{t}{\exp(\boldsymbol{x'}\boldsymbol{\beta})})^{\frac{1}{\sigma}}\right]\frac{1}{\sigma t}$$

$$= \frac{1/\sigma}{\exp(\boldsymbol{x'}\boldsymbol{\beta})}(\frac{t}{\exp(\boldsymbol{x'}\boldsymbol{\beta})})^{\frac{1}{\sigma}-1}\exp\left[-(\frac{t}{\exp(\boldsymbol{x'}\boldsymbol{\beta})})^{\frac{1}{\sigma}}\right]$$

Compare Eq(13) with Eq (2) and let $\gamma = \frac{1}{\sigma}$ and $\rho = \exp(\boldsymbol{x'}\boldsymbol{\beta})$, we can see T has a Weibull distribution $T \sim W(\exp(\boldsymbol{x'}\boldsymbol{\beta}), \frac{1}{\sigma})$.

Refer to Eq(3), now the survival function of $T \sim W(\exp(\boldsymbol{x'}\boldsymbol{\beta}), \frac{1}{\sigma})$ can be written as

$$S_T(t) = \exp\left[-(\frac{t}{\exp(\boldsymbol{x'}\boldsymbol{\beta})})^{\frac{1}{\sigma}}\right] \qquad (14)$$

Refer to Eq(4) the expected survival time of $W(\exp(\boldsymbol{x}'\boldsymbol{\beta}), \frac{1}{\sigma})$ is given as

$$E(T) = \exp(\boldsymbol{x}'\boldsymbol{\beta})\Gamma(\sigma + 1) \tag{15}$$

Since most statistical software use $\log(T)$ to calculate the parameters, let us show distribution and characteristics of the $\log(T)$. Let

$$Y = \log(T) = \boldsymbol{x}'\boldsymbol{\beta} + \sigma\epsilon$$

$$\Rightarrow \epsilon = g^{-1}(Y) = \frac{Y - \boldsymbol{x}'\boldsymbol{\beta}}{\sigma} \tag{16}$$

$$\Rightarrow |J| = |\frac{d(g^{-1}(Y))}{dY}| = \frac{1}{\sigma} \tag{17}$$

Put Eq(16) and Eq(17) to Eq(9) we get

$$f_Y(y) = f_\epsilon(g^{-1}(Y))|J| = \frac{1}{\sigma}\exp(\frac{y - \boldsymbol{x}'\boldsymbol{\beta}}{\sigma})\exp\left[-\exp(\frac{y - \boldsymbol{x}'\boldsymbol{\beta}}{\sigma})\right] \tag{18}$$

Compare Eq(18) to Eq(5) we can see Y i.e. $\log(T)$ has a $G(\boldsymbol{x}'\boldsymbol{\beta}, \sigma)$ distribution. We also can see the use of error term ϵ which has a $G(0, 1)$ distribution is the similar as the error term in simple linear regression that has a $N(0, \sigma^2)$ distribution.

Refer to Eq(13) and Eq(18), we can see in Weibull AFT model, T has a Weibull $W(\exp(\boldsymbol{x}'\boldsymbol{\beta}, \frac{1}{\sigma}))$ distribution, and $\log(T)$ has a Gumbel $G(\boldsymbol{x}'\boldsymbol{\beta}, \sigma)$ distribution.

From Eq (7) the survival function of Y i.e. $\log(T)$ is given as

$$S_Y(y) = \exp\left[-\exp(\frac{y - \mathbf{x}'\beta}{\sigma})\right] \tag{19}$$

And the expectation of Y i.e $\log(T)$ is calculated as

$$
\begin{aligned}
E(Y) = E(\log(T) &= \int_{-\infty}^{\infty} y_i f_Y(y_i)dy_i \\
&= \int_{-\infty}^{\infty} y\frac{1}{\sigma}\exp(\frac{y - \boldsymbol{x}'\boldsymbol{\beta}}{\sigma})\exp\left[-\exp(\frac{y - \boldsymbol{x}'\boldsymbol{\beta}}{\sigma})\right]dy \quad (\text{let } z = \exp(\frac{y - \boldsymbol{x}'\boldsymbol{\beta}}{\sigma})) \\
&= \frac{1}{\sigma}\int_0^{\infty}(\sigma\log z + \boldsymbol{x}'\boldsymbol{\beta})z\exp(-z)d(\sigma\log z + \boldsymbol{x}'\boldsymbol{\beta}) \quad (y = \sigma\log z + \boldsymbol{x}'\boldsymbol{\beta}) \\
&= \int_0^{\infty}(\sigma\log z + \boldsymbol{x}'\boldsymbol{\beta})\exp(-z)dz = \sigma\int_0^{\infty}\log z\exp(-z)dz + \boldsymbol{x}'\boldsymbol{\beta}\int_0^{\infty}\exp(-z)dz \\
&= \sigma\int_0^{\infty}\frac{\partial}{\partial\alpha}\left[z^\alpha e^{-z}\right]_{\alpha=0}dz + \boldsymbol{x}'\boldsymbol{\beta} \\
&= \sigma\frac{\partial}{\partial\alpha}\left[\int_0^{\infty}z^\alpha e^{-z}dz\right]_{\alpha=0} \quad (\text{differentiating under the integral, note } \frac{dz^\alpha}{d\alpha} = \log(z)z^\alpha) \\
&\quad + \boldsymbol{x}'\boldsymbol{\beta} \\
&= \sigma\Gamma(1)' + \boldsymbol{x}'\boldsymbol{\beta} \\
&= \boldsymbol{x}'\boldsymbol{\beta} - \sigma\xi \quad (\text{where } \Gamma(1)' = -\xi \approx -0\cdot 57721, \ \xi \text{ is the Euler-Mascheroni Constant })
\end{aligned}
$$

Note, by Jensen's inequality, $E(\log(T)) \leq \log(E(T))$ since $\log(x)$ is a concave down function, we should not use $\exp(\boldsymbol{x}'\boldsymbol{\beta} - \sigma\xi)$ to calculate the expected survival time. Eq (15) is the correct formula to be used to calculate the expected survival time.

Parameter estimation for the Weibull AFT model

The parameters of Weibull AFT model can be estimated by the maximum likelihood method. The likelihood function of the n observed $\log(t)$ time, $y_1, y_2, ..., y_n$ is given by

$$L(\beta, \sigma; y_i) = \prod_{i=1}^{n} [f_Y(y_i)]^{\delta_i} [S_Y(y_i)]^{1-\delta_i}$$
$$= \prod_{i=1}^{n} \left\{ \frac{1}{\sigma} \exp(\frac{y_i - \boldsymbol{x'\beta}}{\sigma}) \exp\left[-\exp(\frac{y_i - \boldsymbol{x'\beta}}{\sigma})\right] \right\} \left\{ \exp\left[-\exp(\frac{y_i - \boldsymbol{x'\beta}}{\sigma})\right] \right\} \quad (20)$$

where δ_i is the event indicator for ith subject with $\delta_i = 1$ if an event has occured and $\delta_i = 0$ if the event has not occurred. The maximum likelihood estimates $p + 1$ parameters $\sigma, \beta_1...., \beta_p$. We can take log of the likelihood functin and use Newton-Raphson method to calculate these parameters. Most statistical software can do the calculations

Predict mean survival time by the Weibull AFT model

In reliability researches mean survival time is called mean time to failure(MTTF) or mean time between failures(MTBF). [3]

Suppose we want to predict a person i's mean survival time t_i by the Weibull AFT model. First we use MLE method of Eq(20) to calculate the $\hat{\boldsymbol{\beta}}$ and $\hat{\sigma}$ then by the invariance property of MLE, we use Eq(15) to calculate the predicted MTTF directly.

$$t_i = E\hat{(t_i)} = \exp(\boldsymbol{x'_i\hat{\beta}})\Gamma(\hat{\sigma} + 1) \quad (21)$$

After we calculate the mean time to failure(MTTF), we can use the Delta method to calculate the confidence interval for the MTTF. We will treat the predicted MTTF as a function of $\hat{\boldsymbol{\beta}}$ and $\hat{\sigma}$. The standard error of the MTTF can be calculated as

$$SE = \left\{ \begin{pmatrix} \frac{\partial E\hat{(t_i)}}{\partial \hat{\boldsymbol{\beta}}} \\ \frac{\partial E\hat{(t_i)}}{\partial \hat{\sigma}} \end{pmatrix}^t \Sigma_{\hat{\sigma}\hat{\boldsymbol{\beta}}} \begin{pmatrix} \frac{\partial E\hat{(t_i)}}{\partial \hat{\boldsymbol{\beta}}} \\ \frac{\partial E\hat{(t_i)}}{\partial \hat{\sigma}} \end{pmatrix} \right\}^{\frac{1}{2}} \quad (22)$$

where $\Sigma_{\hat{\sigma}\hat{\boldsymbol{\beta}}}$ is the variance-covariance matrix of $\hat{\boldsymbol{\beta}}$ and $\hat{\sigma}$. It can be estimated by the observed Fisher information of the Weibull AFT model. The $(1-\alpha)\%$ confidence interval is given as

$$\hat{t_i} - z_{1-\frac{\alpha}{2}}SE < t_i < \hat{t_i} + z_{1-\frac{\alpha}{2}}SE \quad (23)$$

where α is the type I error, z is the quantile of the standard normal distribution.

Predict median survival time by the Weibull AFT model

Another important statistic in survival analysis is the median survival time or percentile survival time. The pth percentile of survival time is calculated from the survival function. For the individual i the pth percentile of survival time is calculated as

$$S_T(t_i(p)) = \frac{100 - p}{100}$$

For Weibull AFT model we use Eq(14) to calculate pth survival time of an individual i.

$$S_T(t_i) = \exp\left[-(\frac{t_i}{\exp(\boldsymbol{x'\beta})})^{\frac{1}{\sigma}}\right] = \frac{100 - p}{100}$$

$$\Rightarrow (\frac{t_i}{\exp(\boldsymbol{x'\beta})})^{\frac{1}{\sigma}} = -\log(\frac{100 - p}{100})$$

$$\Rightarrow t_i = \left[-\log(\frac{100 - p}{100})\right]^{\sigma} \exp(\boldsymbol{x'_i\beta})$$

After we get $\hat{\boldsymbol{\beta}}$ and $\hat{\sigma}$ from Eq(20) and use the the invariance property of MLE, the median survival time is estimated by

$$t_i(50) = (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x'_i\hat{\beta}}) \tag{24}$$

Still, we can treat the estimated survival time percentile as a function of $\hat{\sigma}$ and $\hat{\boldsymbol{\beta}}$ when p is fixed, we use Delta method to calculate the standard error of predicted pth survival time. The method is the same as in Eq(22) and Eq(23)

Minimum prediction error survival time(MPET)

Both mean and median survival time estimates are biased when small sample is used and the model incorporate censor. [3] Henderson et al purpose a method to find the optimum prediction time with the minimum prediction error. [4] They define that if a observed survival time t falls in the interval $\frac{p}{k} < t < kp$ where p is the predicted survival time and $k > 1$ then the prediction is accurate. The probability of prediction error E_k condition on the predicted time p is given as

$$P(E_k|p) = P(T < p/k) + P(T > kp) \qquad \text{(observed time falls outside the bounds)}$$
$$= 1 - S_T(p/k) + S_T(kp)$$

It can be shown when

$$f_T(p/k) = k^2 f(kp) \tag{25}$$

the probability of prediction error $P(E_k|p)$ will achieve the minimum value.

Now let us calculate the minimum prediction error for Weibull AFT model. From Eq(13) we get

$$f_T(p/k) = \frac{1/\sigma}{\exp(\boldsymbol{x'\beta})}(\frac{p/k}{\exp(\boldsymbol{x'\beta})})^{\frac{1}{\sigma} - 1} \exp\left[-(\frac{p/k}{\exp(\boldsymbol{x'\beta})})^{\frac{1}{\sigma}}\right]$$

$$k^2 f_T(kp) = k^2 \frac{1/\sigma}{\exp(\boldsymbol{x'\beta})}(\frac{kp}{\exp(\boldsymbol{x'\beta})})^{\frac{1}{\sigma} - 1} \exp\left[-(\frac{kp}{\exp(\boldsymbol{x'\beta})})^{\frac{1}{\sigma}}\right]$$

From Eq(25) and cancel the common parts, we get

$$k^{1 - \frac{1}{\sigma}} \exp\left[-(\frac{p/k}{\exp(\boldsymbol{x'\beta})})^{\frac{1}{\sigma}}\right] = k^{1 + \frac{1}{\sigma}} \exp\left[-(\frac{kp}{\exp(\boldsymbol{x'\beta})})^{\frac{1}{\sigma}}\right]$$

take log from both sides we get

$$(1 - \frac{1}{\sigma})\log(k) - (\frac{p/k}{\exp(\boldsymbol{x'\beta})})^{\frac{1}{\sigma}} = (1 + \frac{1}{\sigma})\log(k) - (\frac{kp}{\exp(\boldsymbol{x'\beta})})^{\frac{1}{\sigma}}$$

Rearrange these terms we get

$$p = \left[\frac{\frac{2}{\sigma} \log(k)}{k^{\frac{1}{\sigma}} - k^{-\frac{1}{\sigma}}} \right]^{\sigma} \exp(\boldsymbol{x'\beta}) \qquad (26)$$

Here p is the minimum prediction error survival time. We may use Delta method to get the stand error of the minimum prediction error survival time. Bootstrap methods also could be used to get a confidence interval.

An example to predict the survival time

We use a published larynx cancer dataset [5] to demonstrate the calculations of the predictions. The data set includes 90 male larynx cancer patients with five variables: stage of disease (1=stage 1, 2=stage2, 3=stage 3, 4=stage 4), time to death or on-study time in months, age at diagnosis of larynx cancer, year of diagnosis of larynx cancer and death indicator (0=alive, 1=dead).The author added a variable ID into the dataset and changed the variable name delta to death. The data set can be downloaded from *here*. The larynx cancer data have the following structure.

```
ID stage time age diagyr death
  1     1  0.6  77     76     1
  2     1  1.3  53     71     1
  3     1  2.4  45     71     1
  4     1  2.5  57     78     0
  5     1  3.2  58     74     1
 ...  ...  ...  ...
 46     2  6.2  74     72     1
 47     2  7.0  62     73     1
 48     2  7.5  50     73     0
 49     2  7.6  53     73     0
 50     2  9.3  61     71     0
 51     3  0.3  49     72     1
 52     3  0.3  71     76     1
 ...  ...  ...  ...
 89     4  3.8  84     74     1
 90     4  4.3  48     76     0
```

We will use two predictors to predict the survival time, one predictor is the stage of disease, another predictor is the age at diagnosis of larynx cancer. Since stage is a categorical variable, we will create three dummy variables for the stage and make stage 1 as the default reference group. The Weibull AFT model can be written as

$$\log(T) = \beta_0 + \beta_1 * stage2 + \beta_2 * stage3 + \beta_3 * stage4 + \beta_4 * age + \sigma\epsilon \quad \epsilon \sim G(0,1)$$

Most statistical software can run the Weibull regression model, we use R here.

```
library(survival)
larynx<-read.csv("D:/larynx.csv")
wr <- survreg(Surv(time, death) ~ factor(stage) + age,
data = larynx,dist="w")
summary(wr)
```

We got the following results

```
Call:
survreg(formula = Surv(time, death) ~ factor(stage) + age, data = larynx,
```

```
     dist = "w")
                  Value Std. Error      z        p
(Intercept)      3.5288    0.9041  3.903 9.50e-05
factor(stage)2 -0.1477    0.4076 -0.362 7.17e-01
factor(stage)3 -0.5866    0.3199 -1.833 6.68e-02
factor(stage)4 -1.5441    0.3633 -4.251 2.13e-05
age             -0.0175    0.0128 -1.367 1.72e-01
Log(scale)      -0.1223    0.1225 -0.999 3.18e-01
Scale= 0.885
Weibull distribution
Loglik(model)= -141.4   Loglik(intercept only)= -151.1
Chisq= 19.37 on 4 degrees of freedom, p= 0.00066
Number of Newton-Raphson Iterations: 5
n= 90
```
₁₁₈ ₁₁₉ ₁₂₀ ₁₂₁ ₁₂₂ ₁₂₃ ₁₂₄ ₁₂₅ ₁₂₆ ₁₂₇ ₁₂₈ ₁₂₉ ₁₃₀ ₁₃₁

Suppose we try to predict the survival time of a patient ID=46 who was at larynx cancer stage 2 and aged 74. We use Eq(21) to calculate his expected survival time (mean, or MTTF)

$$
\begin{aligned}
MTTF_{46} = E(\hat{t}_{46}) &= \exp(\boldsymbol{x}_i'\hat{\boldsymbol{\beta}})\Gamma(\hat{\sigma}+1) \\
&= exp(3.5288 - 0.1477 * 1 - 0.5866 * 0 - 1.5441 * 0 - 0.0175 * 74) * \Gamma(1.885) \\
&= 7.7(months)
\end{aligned}
$$

We use Eq(24) to predict his median survival time

$$
\begin{aligned}
Median_{46} &= (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}_i'\hat{\boldsymbol{\beta}}) \\
&= \log(2)^{0.885} exp(3.5288 - 0.1477 * 1 - 0.5866 * 0 - 1.5441 * 0 - 0.0175 * 74) \\
&= 5.8(months)
\end{aligned}
$$

Now let us calculate the standard error of the predicted median time. By Eq(22)

$$
\begin{aligned}
SE &= \left\{ \begin{pmatrix} \frac{\partial E(\hat{t}_i)}{\partial \hat{\boldsymbol{\beta}}} \\ \frac{\partial E(\hat{t}_i)}{\partial \hat{\sigma}} \end{pmatrix}^t \Sigma_{\hat{\sigma}\hat{\boldsymbol{\beta}}} \begin{pmatrix} \frac{\partial E(\hat{t}_i)}{\partial \hat{\boldsymbol{\beta}}} \\ \frac{\partial E(\hat{t}_i)}{\partial \hat{\sigma}} \end{pmatrix} \right\}^{\frac{1}{2}} \\
&= \left\{ \begin{pmatrix} \frac{\partial (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}_i'\hat{\boldsymbol{\beta}})}{\partial \hat{\boldsymbol{\beta}}} \\ \frac{\partial (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}_i'\hat{\boldsymbol{\beta}})}{\partial \hat{\sigma}} \end{pmatrix}^t \Sigma_{\hat{\sigma}\hat{\boldsymbol{\beta}}} \begin{pmatrix} \frac{\partial (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}_i'\hat{\boldsymbol{\beta}})}{\partial \hat{\boldsymbol{\beta}}} \\ \frac{\partial (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}_i'\hat{\boldsymbol{\beta}})}{\partial \hat{\sigma}} \end{pmatrix} \right\}^{\frac{1}{2}} \\
&= \left\{ \begin{pmatrix} (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) \\ (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) * stage2 \\ (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) * stage3 \\ (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) * stage4 \\ (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) * age \\ (\log 2)^{\hat{\sigma}} \log(\log 2) \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) \end{pmatrix}^t \Sigma_{\hat{\sigma}\hat{\boldsymbol{\beta}}} \begin{pmatrix} (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) \\ (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) * stage2 \\ (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) * stage3 \\ (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) * stage4 \\ (\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) * age \\ (\log 2)^{\hat{\sigma}} \log(\log 2) \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) \end{pmatrix} \right\}^{\frac{1}{2}} \quad (27)
\end{aligned}
$$

First, we need to find the variance-covariance matrix $\Sigma_{\hat{\sigma}\hat{\boldsymbol{\beta}}}$, this can be calculated by the observed Fisher information of the Weibull AFT model, most software can calculate this variance-covariance matrix, we use R to do this job.

`wr$var`

We get

₁₃₂ ₁₃₃ ₁₃₄ ₁₃₅ ₁₃₆

```
             (Intercept)    stage2    stage3  stage4       age Log(scale)
(Intercept)        0.817 -0.09049 -0.08479 -0.0444 -0.01114    0.02591
stage2            -0.090  0.16611  0.05319  0.0507  0.00057    0.00016
stage3            -0.085  0.05319  0.10237  0.0567  0.00042   -0.00731
stage4            -0.044  0.05068  0.05668  0.1320 -0.00020   -0.01070
age               -0.011  0.00057  0.00042 -0.0002  0.00016   -0.00026
Log(scale)         0.026  0.00016 -0.00731 -0.0107 -0.00026    0.01501
```

Note, from R what we got is the variance-covariance of coefficients and log(scale), we need to change the log(scale) to scale. We need do some extra calculations. Use formulas in page 401 of the book by John Klein [5] and set $g_1(\sigma, \boldsymbol{\beta}) = \boldsymbol{\beta}$ and $g_2(\sigma, \boldsymbol{\beta}) = \sigma = e^{\log(\sigma)}$, i.e $\sigma = e^{\theta}$ is a function of θ and $\theta = \log(\sigma)$, we get

$$Cov(\beta_0, \sigma,) = Cov(\beta_0, e^{\log(\sigma)}) = Cov(\beta_0, \log(\sigma)) * \sigma = 0.02292735$$

$$Cov(\beta_1, \sigma,) = Cov(\beta_1, e^{\log(\sigma)}) = Cov(\beta_1, \log(\sigma)) * \sigma = 0.0001403178$$

$$Cov(\beta_2, \sigma,) = Cov(\beta_2, e^{\log(\sigma)}) = Cov(\beta_2, \log(\sigma)) * \sigma - 0.006469443$$

$$Cov(\beta_3, \sigma,) = Cov(\beta_3, e^{\log(\sigma)}) = Cov(\beta_3, \log(\sigma)) * \sigma = -0.009470604$$

$$Cov(\beta_4, \sigma,) = Cov(\beta_4, e^{\log(\sigma)}) = Cov(\beta_4, \log(\sigma)) * \sigma = -0.0002297781$$

$$Cov(\sigma) = Cov(e^{\log(\sigma)}) = (e^{\log(\sigma)})^2 Cov(\log(\sigma)) = \sigma^2 Var(log(\sigma)) = 0.0117501$$

Now we will use these six values to replace the last row and column of the variance-covariance matrix of the coefficients and log(scale). We get

```
       [,1]     [,2]     [,3]    [,4]     [,5]     [,6]
[1,]  0.817 -0.09049 -0.08479 -0.0444 -0.01114  0.02293
[2,] -0.090  0.16611  0.05319  0.0507  0.00057  0.00014
[3,] -0.085  0.05319  0.10237  0.0567  0.00042 -0.00647
[4,] -0.044  0.05068  0.05668  0.1320 -0.00020 -0.00947
[5,] -0.011  0.00057  0.00042 -0.0002  0.00016 -0.00023
[6,]  0.023  0.00014 -0.00647 -0.0095 -0.00023  0.01175
```

This is the $\Sigma_{\hat{\sigma}\hat{\beta}}$ matrix we needed.

If we use SAS software we can directly get the variance-covarince matrix of $\hat{\boldsymbol{\beta}}$ and $\hat{\sigma}$ by using the following statements

```
proc lifereg data=larynx order=data COVOUT outest=est;
class stage;
model time*death(0)=stage age/dist=weibull;
run;
proc print data=est;
run;
```

The right side vector of the Eq(27) is calculated as

$$
\begin{pmatrix}
(\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) \\
(\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) * stage2 \\
(\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) * stage3 \\
(\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) * stage4 \\
(\log 2)^{\hat{\sigma}} \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}}) * age \\
(\log 2)^{\hat{\sigma}} \log(\log 2) \exp(\boldsymbol{x}'\hat{\boldsymbol{\beta}})
\end{pmatrix}
$$

$$
=
\begin{pmatrix}
(\log 2)^{0.885} \exp(3.5288 - 0.1477 * 1 - 0.5866 * 0 - 1.5441 * 0 - 0.0175 * 74) \\
(\log 2)^{0.885} \exp(3.5288 - 0.1477 * 1 - 0.5866 * 0 - 1.5441 * 0 - 0.0175 * 74) * 1 \\
(\log 2)^{0.885} \exp(3.5288 - 0.1477 * 1 - 0.5866 * 0 - 1.5441 * 0 - 0.0175 * 74) * 0 \\
(\log 2)^{0.885} a \exp(3.5288 - 0.1477 * 1 - 0.5866 * 0 - 1.5441 * 0 - 0.0175 * 74) * 0 \\
(\log 2)^{0.885} \exp(3.5288 - 0.1477 * 1 - 0.5866 * 0 - 1.5441 * 0 - 0.0175 * 74) * 74 \\
(\log 2)^{0.885} \log\log(\log 2) \exp(3.5288 - 0.1477 * 1 - 0.5866 * 0 - 1.5441 * 0 - 0.0175 * 74)
\end{pmatrix}
$$

$$
=
\begin{pmatrix}
5.8383 \\
5.8383 \\
0 \\
0 \\
432.03 \\
-7.7915
\end{pmatrix}
$$

Now we have everything to calculate the standard error of the median survival time.

$$
SE = \left\{
\begin{pmatrix}
5.8383 \\
5.8383 \\
0 \\
0 \\
432.03 \\
-7.7915
\end{pmatrix}^{t}
\begin{pmatrix}
0.817 & -0.0905 & -0.0848 & -0.0444 & -0.0111 & 0.0229 \\
-0.090 & 0.1661 & 0.0532 & 0.0507 & 0.00057 & 0.00014 \\
-0.0859 & 0.0532 & 0.1024 & 0.0567 & 0.00042 & -0.0065 \\
-0.044 & 0.0507 & 0.0567 & 0.1320 & -0.00020 & -0.0095 \\
-0.011 & 0.00057 & 0.0004 & -0.00020 & 0.00016 & -0.00023 \\
0.023 & 0.00014 & -0.0065 & -0.0095 & -0.00023 & 0.01175
\end{pmatrix}
\begin{pmatrix}
5.8383 \\
5.8383 \\
0 \\
0 \\
432.03 \\
-7.7915
\end{pmatrix}
\right\}^{\frac{1}{2}}
$$

$$
= 2.156133
$$

Therefore, the 95% confidence interval is given as

$$
95\% CI : (5.83 - 1.96 * 2.16 < Median_{46} < 5.83 + 1.96 * 2.16) = (1.60 \text{ to } 10.01) \text{ months}.
$$

Which means we are 95% confident that the survival time will be within 1.60 to 10.01 months. We can also use the R build in function *predict* to predict the median survival time.

```
Median46<-predict(wr, newdata=data.frame(stage=2,age=74),type="quantile",
p=0.5,se.fit=TRUE)
Median46
```

We get

```
$fit
 5.838288
$se.fit
2.095133
```

The standard error is a little bit different from our calculations which might be due to the round off error.

We use Eq(26) to calculate minimum prediction error survival time (MPET) and we fix $k=2$.

$$
MPET_{46} = \left[\frac{\frac{2}{\sigma} \log(k)}{k^{\frac{1}{\sigma}} - k^{-\frac{1}{\sigma}}} \right]^{\sigma} \exp(\boldsymbol{x}'\boldsymbol{\beta})
$$

$$
= \left[\frac{\frac{2}{\sigma} \log(k)}{2^{\frac{1}{0.885}} - k^{-\frac{1}{0.885}}} \right]^{0.885} * \exp(3.5288 - 0.1477 * 1 - 0.5866 * 0 - 1.5441 * 0 - 0.0175 * 74)
$$

$$
= 7.4 (months)
$$

It seems the three prediction methods are all quite close to the real survival time of the patient ID=46 which was 6.2 months.

Note in R build in *predict* function for Weibull AFT model type= "response" calculates $\exp(\boldsymbol{x'\hat{\beta}})$ without the $\Gamma(1+\hat{\sigma})$ and type="lp" calculates $\boldsymbol{x'\hat{\beta}}$ only, we should not use them to predict MTTF. There is no software to calculate the minimum prediction error survival time.

Assess the point prediction accuracy

Parkes [6] suggested a simple method to measure the accuracy of the predicted survival time. Let t be the observed survival time and p is the predicted time, if $p/k \leq t \leq kp$ then the point prediction p is defined as "accurate", outside the interval is defined as" inaccurate". Christakis and Lamont purposed a 33 percent rule to measure the accuracy, where they divided the observed time by the predicted survival time, and defined the prediction is "accurate" if this quotient was between 0.67 and 1.33. Values less than 0.67 or greater than 1.33 were defined as "error". [7] This method in fact is just fix $k = 3$ in Parkes' method. We choose $k = 2$ for our accuracy assessment. The accurate rate is defined as the proportion of "accurate" prediction over the total sample size. The results were presented in table 1.

Discussion

In this paper, we introduced how to use Weibull AFT model to predict survival times. Mean survival time (mean time to failure time, mean time between failures), median survival time and minimum prediction error survival time were used to predict the survival time and the prediction accuracy was assessed by Parke's method. When we fixed $k = 2$ the accuracy was 55.6% for median, 50% for MTTF and 51.1% for MPET. If we fixed $k = 3$ as suggested by Christakis and Lamont the accuracy rate was increased to 77.8%,66.7 % and 67.8%, respectively. The sample we used is quite small and we only used two predictors, with bigger sample size and more predictors the accuracy rate might be even higher. In this sample we did not observe that minimum prediction error time had better accuracy rate than median survival time.

The parametric survival models have advantages in predicting survival time than the semi-parametric Cox regression model. The Cox regression model which can be specified as $S_i(t|x_i) = S_0(t)^{\exp(\boldsymbol{x'_i\beta})}$ cannot predict time directly. What it can do is to specify a certain time first then to calculate the probability for that period of time. The disadvantage of the parametric survival models is that we have to make stronger assumptions than semi-parametric models. [8]

Currently, most clinical prediction models calculate a patient's probability of having or developing a certain disease or risk scores that based on the calculated probabilities. [9]However, provide a probability seems difficult to be understand by general population and probability itself can be defined by quite different ways. [10] In practice, the time axis remains the most natural measure for both clinicians and patients. It is much easier to understand a survival time rather than a subjective assessment of probability of survival to a certain time point. [4] Predicting survival time can provide a practical and concrete guide to clinicians and health care providers to manage their patients and help families and patients make suitable plans for the remaining lifespan.

References

1. Gorgoso-Varela JJ, Rojo-Alboreca A. Use of Gumbel and Weibull functions to model extreme values of diameter distributions in forest stands. Annals of forest

science. 2014;71(7):741-50.

2. Lai C-D. Generalized Weibull Distributions. Generalized Weibull Distributions: Springer; 2014. p. 23-75.

3. Ho L, Silva A. Unbiased estimators for mean time to failure and percentiles in a Weibull regression model. International Journal of Quality & Reliability Management.2006;23(3):323-39.

4. Henderson R, Jones M, Stare J. Accuracy of point predictions in survival analysis. Statistics in medicine. 2001;20(20):3083-96.

5. Klein JP, Moeschberger ML. Survival analysis: techniques for censored and truncated data. Springer Science & Business Media; 2005.

6. Parkes CM. Accuracy of predictions of survival in later stages of cancer. British medical journal. 1972;2(5804):29.

7. Christakis NA, Smith JL, Parkes CM, Lamont EB. Extent and determinants of error in doctors' prognoses in terminally ill patients: prospective cohort studyCommentary: Why do doctors overestimate? Commentary: Prognoses should be based on proved indices not intuition. Bmj. 2000;320(7233):469-73.

8. Nardi A, Schemper M. Comparing Cox and parametric models in clinical studies. Statistics in medicine. 2003;22(23):3597-610.

9. Lee Y-h, Bang H, Kim DJ. How to establish clinical prediction models. Endocrinology and Metabolism. 2016;31(1):38-44.

10. Saunders S. What is probability? Quo vadis quantum mechanics? Springer; 2005. p. 209-38.

Table 1. Prediction results and accuracy (last digit in predicted time: 0 inaccurate, 1, accurate)

ID	stage	age	death	time	Median(95% CI)	MTTF	MPET
1	1	77	1	0.6	6.42(3.16,9.68),0	8.47,0	8.11,0
2	1	53	1	1.3	9.77(4.07,15.46),0	12.9,0	12.34,0
3	1	45	1	2.4	11.23(3.07,19.39),0	14.84,0	14.19,0
4	1	57	0	2.5	9.11(4.32,13.89),0	12.03,0	11.5,0
5	1	58	1	3.2	8.95(4.36,13.54),0	11.82,0	11.3,0
6	1	51	0	3.2	10.11(3.88,16.34),0	13.36,0	12.78,0
7	1	76	1	3.3	6.54(3.29,9.78),1	8.62,0	8.25,0
8	1	63	0	3.3	8.2(4.37,12.03),0	10.83,0	10.36,0
9	1	43	1	3.5	11.63(2.72,20.54),0	15.36,0	14.7,0
10	1	60	1	3.5	8.64(4.4,12.89),0	11.41,0	10.91,0
11	1	52	1	4	9.94(3.98,15.89),0	13.13,0	12.55,0
12	1	63	1	4	8.2(4.37,12.03),0	10.83,0	10.36,0
13	1	86	1	4.3	5.49(1.96,9.01),1	7.24,1	6.92,1
14	1	48	0	4.5	10.66(3.52,17.79),0	14.08,0	13.47,0
15	1	68	0	4.5	7.52(4.12,10.91),1	9.92,0	9.49,0
16	1	81	1	5.3	5.99(2.63,9.34),1	7.9,1	7.56,1
17	1	70	0	5.5	7.26(3.96,10.56),1	9.58,1	9.16,1
18	1	58	0	5.9	8.95(4.36,13.54),1	11.82,0	11.3,1
19	1	47	0	5.9	10.84(3.38,18.31),1	14.33,0	13.7,0
20	1	75	1	6	6.65(3.41,9.89),1	8.78,1	8.39,1
21	1	77	0	6.1	6.42(3.16,9.68),1	8.47,1	8.11,1
22	1	64	0	6.2	8.06(4.34,11.77),1	10.64,1	10.18,1
23	1	77	1	6.4	6.42(3.16,9.68),1	8.47,1	8.11,1
24	1	67	1	6.5	7.65(4.19,11.1),1	10.1,1	9.66,1
25	1	79	0	6.5	6.2(2.9,9.5),1	8.18,1	7.83,1
26	1	61	0	6.7	8.49(4.4,12.58),1	11.21,1	10.73,1
27	1	66	0	7	7.78(4.25,11.31),1	10.27,1	9.83,1
28	1	68	1	7.4	7.52(4.12,10.91),1	9.92,1	9.49,1
29	1	73	0	7.4	6.89(3.65,10.12),1	9.09,1	8.69,1
30	1	56	0	8.1	9.27(4.28,14.26),1	12.24,1	11.71,1
31	1	73	0	8.1	6.89(3.65,10.12),1	9.09,1	8.69,1
32	1	58	0	9.6	8.95(4.36,13.54),1	11.82,1	11.3,1
33	1	68	0	10.7	7.52(4.12,10.91),1	9.92,1	9.49,1
34	2	86	1	0.2	4.73(0.68,8.78),0	6.25,0	5.97,0
35	2	64	1	1.8	6.95(2.37,11.54),0	9.18,0	8.78,0
36	2	63	1	2	7.07(2.4,11.75),0	9.34,0	8.93,0
37	2	71	0	2.2	6.15(1.96,10.34),0	8.12,0	7.77,0
38	2	67	0	2.6	6.6(2.22,10.97),0	8.71,0	8.33,0
39	2	51	0	3.3	8.72(2.28,15.17),0	11.52,0	11.02,0
40	2	70	1	3.6	6.26(2.03,10.49),1	8.26,0	7.9,0
41	2	72	0	3.6	6.05(1.89,10.2),1	7.98,0	7.63,0
42	2	81	1	4	5.17(1.13,9.21),1	6.82,1	6.52,1
43	2	47	0	4.3	9.36(1.98,16.73),0	12.36,0	11.82,0
44	2	64	0	4.3	6.95(2.37,11.54),1	9.18,0	8.78,0
45	2	66	0	5	6.71(2.28,11.15),1	8.86,1	8.48,1

Table 1 continued Prediction results and accuracy (last digit in predicted time: 0 inaccurate, 1, accurate)

ID	stage	age	death	time	Median(95% CI)	MTTF	MPET
46	2	74	1	6.2	5.84(1.73,9.94),1	7.7,1	7.37,1
47	2	62	1	7	7.2(2.43,11.97),1	9.51,1	9.09,1
48	2	50	0	7.5	8.88(2.22,15.54),1	11.73,1	11.22,1
49	2	53	0	7.6	8.42(2.38,14.47),1	11.13,1	10.64,1
50	2	61	0	9.3	7.33(2.46,12.2),1	9.67,1	9.25,1
51	3	49	1	0.3	5.83(2.41,9.24),0	7.69,0	7.36,0
52	3	71	1	0.3	3.97(2.19,5.75),0	5.24,0	5.01,0
53	3	57	1	0.5	5.07(2.68,7.45),0	6.69,0	6.4,0
54	3	79	1	0.7	3.45(1.56,5.34),0	4.55,0	4.35,0
55	3	82	1	0.8	3.27(1.32,5.23),0	4.32,0	4.13,0
56	3	49	1	1	5.83(2.41,9.24),0	7.69,0	7.36,0
57	3	60	1	1.3	4.81(2.68,6.94),0	6.35,0	6.07,0
58	3	64	1	1.6	4.48(2.58,6.39),0	5.92,0	5.66,0
59	3	74	1	1.8	3.76(1.96,5.56),0	4.97,0	4.75,0
60	3	72	1	1.9	3.9(2.12,5.68),0	5.14,0	4.92,0
61	3	53	1	1.9	5.43(2.6,8.27),0	7.17,0	6.86,0
62	3	54	1	3.2	5.34(2.63,8.05),1	7.05,0	6.74,0
63	3	81	1	3.5	3.33(1.4,5.27),1	4.39,1	4.2,1
64	3	52	0	3.7	5.53(2.56,8.49),1	7.3,1	6.98,1
65	3	66	0	4.5	4.33(2.49,6.17),1	5.71,1	5.47,1
66	3	54	0	4.8	5.34(2.63,8.05),1	7.05,1	6.74,1
67	3	63	0	4.8	4.56(2.61,6.51),1	6.02,1	5.76,1
68	3	59	1	5	4.89(2.69,7.1),1	6.46,1	6.18,1
69	3	49	0	5	5.83(2.41,9.24),1	7.69,1	7.36,1
70	3	69	0	5.1	4.11(2.32,5.89),1	5.42,1	5.19,1
71	3	70	1	6.3	4.04(2.26,5.82),1	5.33,1	5.1,1
72	3	65	1	6.4	4.41(2.54,6.27),1	5.82,1	5.56,1
73	3	65	0	6.5	4.41(2.54,6.27),1	5.82,1	5.56,1
74	3	68	1	7.8	4.18(2.38,5.98),1	5.52,1	5.28,1
75	3	78	0	8	3.51(1.64,5.38),0	4.63,1	4.43,1
76	3	69	0	9.3	4.11(2.32,5.89),0	5.42,1	5.19,1
77	3	51	0	10.1	5.63(2.52,8.73),1	7.43,1	7.11,1
78	4	65	1	0.1	1.69(0.77,2.61),0	2.23,0	2.14,0
79	4	71	1	0.3	1.52(0.7,2.34),0	2.01,0	1.92,0
80	4	76	1	0.4	1.4(0.61,2.18),0	1.84,0	1.76,0
81	4	65	1	0.8	1.69(0.77,2.61),0	2.23,0	2.14,0
82	4	78	1	0.8	1.35(0.56,2.13),1	1.78,0	1.7,0
83	4	41	1	1	2.57(0.3,4.84),0	3.4,0	3.25,0
84	4	68	1	1.5	1.6(0.74,2.47),1	2.12,1	2.03,1
85	4	69	1	2	1.58(0.73,2.42),1	2.08,1	1.99,1
86	4	62	1	2.3	1.78(0.78,2.79),1	2.35,1	2.25,1
87	4	74	0	2.9	1.44(0.65,2.24),0	1.91,1	1.82,1
88	4	71	1	3.6	1.52(0.7,2.34),0	2.01,1	1.92,1
89	4	84	1	3.8	1.21(0.42,2.01),0	1.6,0	1.53,0
90	4	48	0	4.3	2.28(0.57,3.99),1	3.01,1	2.87,1
Accuracy rate(%)					55.6%(50/90)	50%(45/90)	51.1%(46/90)