Life history evolution and phenotypic plasticity in parasitic eyebrights (*Euphrasia*,

Orobanchaceae)

Alex D. Twyford[1], Natacha Frachon[2], Edgar L. Y. Wong[3], Chris Metherell[4], Max R. Brown[1]

[1]University of Edinburgh, Institute of Evolutionary Biology, West Mains Road, Edinburgh, EH3 9JT,

UK.

[2]Royal Botanic Garden Edinburgh, 20A Inverleith Row, Edinburgh, EH3 5LR, UK.

[3]Department of Plant Sciences, University of Oxford, South Parks Road, Oxford, OX1 3RB, UK.

[4]Botanical Society of Britain and Ireland, 57 Walton Rd, Bristol BS11 9TA.

Email for correspondence: Alex.Twyford@ed.ac.uk

Running head: Life history of parasitic eyebrights

Abstract

Parasite lifetime reproductive success is determined by both genetic variation and phenotypically

plastic life history traits that respond to host quality and external environment. In this study we use the

generalist parasitic plant genus *Euphrasia* to test aspects of life history theory. We evaluate species

life history strategies when grown on a common host, document phenotypic plasticity when grown on

many different hosts, and relate our observations to trait differences recorded in the wild. We find

Euphrasia in a common garden and in the field exhibit a range of life history strategies that differ

between species that transition rapidly to flower at the expense of early season growth, and those that

invest in vegetative growth and delay flowering. However, aspects of life history show extensive

phenotypic plasticity in response to host quality, and demonstrate the costs of attaching to a low-

quality host. Our experiments suggest life history strategies in a generalist parasitic plant are the

product of natural selection on traits related to growth and flowering, while host quality may be a

primary determinant of lifetime reproductive success.

Keywords: flowering time; host range; life history evolution; parasitic plants; phenotypic plasticity

Background

Parasitism is a ubiquitous feature of the natural world, with parasitic organisms present in every ecosystem and found to exploit all free-living organisms [1, 2]. The life history of parasitic organisms must be synchronised with the host to maximise a parasite's individual growth, survival and fecundity. However, the life history of external parasites (exoparasites) must not only be timed to their host, but to variable and often stochastic external environmental conditions [3]. Such responses will primarily be plastic when responding to local conditions or fluctuating environments, or genetic in response to more predictable or spatially structured variation [4]. Understanding how genetic and plastic differences contribute to life history trait variation in parasite species and populations is essential for understanding parasite speciation, host-parasite interactions, and transmission dynamics [5].

Parasitic plants are a group of c. 4500 species of at least 12 separate evolutionary origins that have evolved a modified feeding organ, the haustorium, which allows them to attach to a host plant and extract nutrients and water [6, 7]. Parasitic plants are incredibly morphologically diverse and present a broad range of life history strategies [8], with the host species range a critical trait that determines parasite interactions and performance in the wild. Generalist parasitic plants can attach to a broad range of hosts with the grassland parasite *Rhinanthus* found to attach to over 50 co-occurring grass and herbaceous species [9]. All generalist parasitic plants are exoparasites where the leaves, stems, roots and flowers grow outside the host and only the haustorium invades and grows within the host [10]. The range of hosts and their visible external growth make them ideal for investigating host-parasite interactions. Generalist parasitic plants have been used in experimental studies for over 125 years [11], with these studies revealing that parasite growth, reproduction and metabolism are determined by host functional group (herb, grass, legume etc.) and host species [12]. Despite this large body of experimental work manipulating host type, most studies have not simultaneously used the replication of parasite species, host species and individuals, to identify the different life history

strategies of parasite species and how phenotypic plasticity in life history traits determines parasite

performance.

In this study, we explore life history trait evolution in parasitic eyebrights (*Euphrasia*,

Orobanchaceae). *Euphrasia* represents an ideal study system for investigating the recent evolution of

parasite life history strategies. These taxa are hemiparasites that are photosynthetically competent and

can complete their lifecycle without a host, but grow much better after attachment to one of many

suitable hosts. In the United Kingdom, the 21 *Euphrasia* species demonstrate recent postglacial

divergence with many species indistinguishable at DNA barcoding loci [13], presenting similar and

overlapping morphological variation [14, 15], and found to readily hybridise [16]. Despite this

shallow species divergence, *Euphrasia* demonstrates substantial ecological divergence, with many

taxa restricted to specific habitats such as coastal turf, mountain scree or open grassland. This

ecological diversity would be expected to exert strong selection on aspects of the life history of a

generalist parasitic annual plant species such as *Euphrasia*. This may include selection on growth to

match seasonal water availability and to exploit local hosts, or selection on flowering time in response

to local competition.

In our experiments, we first assess the extent of life history trait variation among *Euphrasia* species

and their hybrids when grown on a single host in standardised growing conditions. This experiment

addresses whether there is life history trait divergence among recently diverged parasite species. We

then inspect plasticity of a single focal *Euphrasia* population grown on many different hosts. This

experiment tests whether they are truly generalists by growing them on a wide range of hosts, and also

includes a treatment without a host to observe the phenotype when free-living. Finally, we relate our

trait observations made in a common garden to recordings made on herbarium specimens collected in

the wild. This comparison will help us understand whether life history trait differences hold in both

the common garden and in nature. Overall, our joint observations of phenotypic variation between

closely related species, and the extent of host-induced plasticity within species, both in an experiment

and in the wild, provides new insights into life history variation in parasitic taxa.

Materials and Methods

    (a) Experimental design and plant cultivation

We performed two common garden experiments to investigate phenotypic variation in parasitic

Euphrasia. Our 'species differences' (SpD) experiment observed life history trait differences of five

species and six natural hybrids when grown on clover (*Trifolium repens*), which is often used as a

host in parasitic plant experiments because the parasite grows vigorously and has high survival [17].

This experiment includes multiple populations of three widespread and closely related species, *E.*

arctica, *E. confusa* and *E. nemorosa*, and a single population of two more distinct taxa, *E. micrantha*

and *E. pseudokerneri*. Our 'phenotypic plasticity' (PP) experiment assessed the impact of eight hosts

(*Arabidopsis thaliana*, *Equisetum arvense, Festuca rubra*, *Holcus lanatus*, *Marchantia polymorpha*,

Pinus sylvaticus, Plantago lanceolata, and *Trifolium repens*), and growing without a host, on parasite

phenotype. These hosts were chosen to represent both typical hosts encountered in the wild, as well as

novel hosts, and a broad representation of phylogenetic diversity and functional groups (full details in

Table S1). Both common garden experiments took place in parallel in 2016. Wild-collected open-

pollinated *Euphrasia* seeds were contributed by plant recorders as part of the 'Eye for Eyebrights'

(E4E) public engagement project and as such included a scattered geographic sample across Great

Britain (Table S2). Host seeds were sourced from commercial suppliers and from wild-collected

material (Table S1).

Reliable cultivation of *Euphrasia* can be challenging due to low seed germination, variation in time to

establishment, the requirement of seed stratification, and high seedling mortality when transplanted

[17, 18]. Here, we develop cultivation protocols that combine winter germination cues that improve

germination and mimic nature, but also use highly standardised and replicated pot conditions that

avoid transplanting *Euphrasia* and thus maximise survivorship. We planted a single *Euphrasia* seed

into 9cm plastic pots filled with Melcourt Sylvamix® Special growing media in December, with

plants left outside overwinter to experience natural seed stratification. Hosts were planted in seed

trays in April. *Euphrasia* plants were moved to an unheated and well-ventilated greenhouse at the

Royal Botanic Garden Edinburgh (RBGE) in the spring once the cotyledons were fully expanded, and

a single seedling from each host introduced. Hosts that died within ten days of planting were replaced.

Twenty or more replicates were grown for each host-parasite combination. Plants were subsequently

grown to flowering with regular watering, randomised at weekly intervals, and foreign weed seedlings

removed.

(b) Trait measurements and statistical analyses

We measured a suite of seven morphological traits at first flowering. These traits are characters

related to life history variation, indicators of plant vigour, or characters used in taxonomy. In addition

to date of first flowering, we measured: corolla length, cauline leaf length:internode length ratio

('internode ratio'), leaf teeth on the lower floral leaf, nodes to flower, number of branches and plant

height. All length measurements were made to the nearest millimetre, and followed protocols from

reference [14]. For the many hosts experiment, we also recorded early season growth (height six

weeks after transplantation) and height at the end of season after senescence. We did not make direct

observations of below-ground attachment due to the fine root structure of *Euphrasia*, and instead used

height at first flowering to infer likely attachment [19].

We analysed data for the two experiments separately but with similar statistical approaches. For the

SpD experiment we used generalised linear mixed effects models to fit species as a fixed effect and

population as a random effect. For the PP experiment we used generalised linear mixed effects models

with presence or absence of host as a fixed effect and host species identity as a random effect. We

used a Gaussian distribution for all models except where the response variable were positive integer

counts, where we used a Poisson distribution. To calculate the significance of a fixed or random effect

we used Likelihood Ratio Tests where the full model was compared with a nested model where one

effect had been dropped. When a random effect was dropped, we implemented generalised linear

models using R base functions. For the trait number of branches the generalised linear mixed effects

models failed to converge so we used MCMCglmm [20], with the best fitting of either a full or nested

model chosen based on a lower Deviance Information Criterion (DIC) value. For the many hosts,

models for number of branches failed to converge in every model run, so this model was omitted.

Using the flexible variance structures accommodated by MCMCglmm we were able to calculate

"variance explained" by random effects in each model. In the case of a Gaussian model, variance

explained by a variable is the fraction of the total variance accounted for by the variable. In the case

of a Poisson model the process is more complex due to the mean-variance relationship [21]. The

models were run for 91,000 iterations with a burn-in of 21,000 and a thinning interval of 70. The

priors were standard inverse-Wishart for the (co)variances and parameter expanded where there was

poor mixing. We calculated r^2 values from Pearson Correlation Coefficients between all traits and

used Principal Component Analyses to quantify phenotypic clustering of individuals. All analyses

were done in R version 3.4.3, with the packages lme4 [22] and MCMCglmm for generalised linear

mixed effects models and ggplot2 for data visualisation [23].

(c) Trait variation in the wild

We tested how phenotypes in the experiments relate to phenotypes in nature, by comparing results

from experiment SpD to phenotypic measurements made on herbarium specimens of the same

populations sampled in the wild. Pressed plants submitted by field collectors varied in quality, and

these sometimes missed the base of the plant, therefore we were unable to measure height. We

measured traits on three individuals from each collection sheet for a given population. The results

were then averaged for graphical representation, but pooled for statistical analysis. We used

generalised linear mixed effects models to fit variables as a function of where individuals were

measured (i.e. common garden or measured on herbarium specimens). This two levelled categorical

factor was implemented as a fixed effect. Adding species as a random effect shows how much of the

variability of the trait was explained by species given where individuals were measured. Count data

was analysed with a Poisson distribution, in all other cases a Gaussian distribution was used. Analyses

were performed in MCMCglmm.

Results

Species differences

Our first common garden experiment reveals extensive morphological trait variation across *Euphrasia*

species. From the 222 individuals surviving to flowering on their clover host, we see a 3-fold

difference in mean height, while other traits also proved variable (Figure 1a; Table S3). A large

degree of this variation is separated by species and by population (Table 1). The species with the most

distinct lift-history strategy is *E. micrantha*, which flowers early in the season from a low node on the

plant, and flowers while it is short. It also forms a distinct cluster in the PCA analysis (Figure S1). *E.*

pseudokerneri is relatively distinct, flowering once it has grown tall and from a high node on the

plant, but shows little separation in the PCA analysis. The morphologically similar *E. arctica*, *E.*

confusa and *E. nemorosa* are partly distinct, with *E. nemorosa* initiating flowering 14 days later and

from 3.3 nodes higher than *E. arctica*, but overlaps in other traits and in overall multi-trait phenotype

(Figure S1). Hybrids have intermediate morphology that is consistent with the phenotype of their

parental progenitors, for example hybrids involving *E. nemorosa* flower later in the season and from a

later node than *E. arctica* hybrids (Figure 1a).

Correlation analyses reveal clear suites of traits that are related, with strong correlations present in 6

of 21 pairwise comparisons ($R^2 > 0.4$, Table 2a). Plants flowering at a late node are more likely to be

tall, more highly branched, as well as having many teeth on the lower floral leaf. The relationship of

traits is also supported in the PCA analysis, with many traits contributing to multiple principal

components (Table S4). These height and flowering node related traits are largely uncorrelated with

cauline internode-leaf ratio and corolla length.

Phenotypic plasticity

Our phenotypic plasticity experiment shows substantial morphological variation across 194 *E. arctica*

plants grown with 8 different host species, and the 22 plants grown without a host. Plants growing on

clover transitioned to flower quickly, grew tall by the time of first flowering, and produced large

flowers (Figure 1b, Table S5). This contrasts with *Euphrasia* with no host, which flowered on average

53 days later, were extremely short at first flowering, and produced small flowers. Despite this, the

presence or absence of host was not significant (Table 1), due to *Euphrasia* grown with *Marchantia*

and *Pinus* growing similarly to the no host treatment (Figure 1b). The other hosts, *Arabidopsis*,

Equisetum, *Festuca*, *Holcus* and *Plantago* conferred some benefit for *Euphrasia* and fell between

clover and no host. Our comparison of growth across host treatments measured through the year

showed that height at the end of the season is poorly predicted from early season growth ($r^2 = 0.22$),

but strongly correlated with height at first flowering ($r^2 = 0.69$). Plants that flowered early were more

likely to grow larger by the end of season ($r^2 = -0.30$) and become more highly branched ($r^2 = 0.62$).

Across host treatments, there was a significant negative correlation between Julian days to flower and

most other traits (Table 2b). We find that late flowering individuals are likely to be smaller at first

flowering, are less highly branched, have less highly toothed leaves, and have smaller flowers (Table

S4). While strongly correlated, the magnitude of change for different traits varied substantially. Days

to flower differed considerable depending on host, with a 3.8x greater difference than seen between

means of different *Euphrasia* species (Fig. 1aiii and 1biii). In contrast corolla length and node to

flower proved less variable depending on host, with a 1.4x and 1.2x change between means,

respectively.

Variation in the wild

The comparison between common garden and wild-collected herbarium specimens revealed a single

trait, nodes to flower, is strongly correlated ($r^2=0.78$) and not significantly different ($p_{MCMC}=0.71$)

between environments (Figure 2). All other traits did differ significantly between environments

($p_{MCMC} < 0.05$), with *Euphrasia* plants in the common garden having corollas on average 1.4mm

longer, with 0.2 more teeth on the lower floral leaves, an increase in cauline:internode ratio of 1.0mm,

and 4 more pairs of branches. Despite differing between environments, trait values were highly

correlated for corolla length ($r^2 = 0.97$, $p_{MCMC} < 0.001$) and cauline internode:leaf length ratio ($r^2 =$

0.63, $p_{MCMC} < 0.001$), but not for number of leaf teeth ($r^2 = 0.13$, $p_{MCMC} = 0.034$) and number of

branches ($r^2 = 0.047$, $p_{MCMC} < 0.001$).

Discussion

Our study sheds new light on life history evolution and phenotypic plasticity in generalist parasitic

plants. We find different life history strategies between recently divergent parasitic *Euphrasia* species,

with some species rapidly transitioning to flower at the expense of growth-related traits, while others

delay flowering and invest in early-season vegetative growth. However, many traits related to life

history differences are phenotypically plastic and show major changes in response to host quality. In

addition to our experimental observations, the consistency between life history traits in a common

garden and in the wild suggest our experimental observations generalise to patterns observed in

nature. Overall our study highlights the value in integrating multiple common garden experiments and

field collections to study life history strategies in parasitic plants, and demonstrates the rapid

evolution of life history differences in a postglacial radiation of parasites.

Life history variation in a parasitic plant

Our study finds evidence for different life history strategies between *Euphrasia* species in Britain. *E.*

arctica, *E. micrantha* and hybrids such as *E. arctica* x *E. confusa*, transition rapidly to flower, flower

while they are short, and produce their first flower from a low node on the plant. This contrasts with

E. pseduokerneri, *E. nemorosa* and hybrids involving *E. nemorosa* that delay flowering until later in

the season, grow tall before flowering, and produce their first flower from a late node on the main

axis. These different life history strategies correspond to the known of ecology of these species, with

E. nemorosa flowering late in tall mixed grassland, while *E. micrantha* flowers early in patchy

heathland where competition is less intense [14]. Our observations are consistent with the classic life

history trade-off between growth and reproduction [24, 25]. For *Euphrasia* growing in the wild, early

reproduction allows the plants to reliably complete their lifecycle before summer competition,

herbivory, mowing, summer drought and other seasonal abiotic and biotic stresses. However early

flowering involves reproducing at the expense of early season growth and at a time when the resource

budget is constrained by relatively few haustorial connections. These trait trade-offs pose an

interesting comparison to the well-studied *Mimulus guttatus* (*syn. Erythranthe guttata*), a non-

parasitic relative in the Lamiales that shares the same basic plant architecture. In *M. guttatus* multiple

traits related to growth and reproduction are correlated both within and between populations, due to

genetic trade-offs between time to flower and fecundity [26, 27]. In *Euphrasia* the genetics

underpinning this life history trade-off have yet to be characterised, and may be a consequence of

multiple independent loci or trade-offs at individual loci [28].

While much life history variation is captured by differences in time to flower and growth-related

traits, we also see evidence for flower size representing a separate axis of variation across *Euphrasia*

species. In our common garden *E. micrantha* has small corollas, while *E. arctica* and *E. nemorosa*

have larger corollas, and corolla size is not correlated with other traits. *Euphrasia* species are well-

known to have flower size variation, with a continuum between small flowered species that are highly

selfing (e.g. *E. micrantha*, corolla size = 4.5 - 6.5mm, inbreeding coefficient F_{IS} > 0.88, [29]), and

large flowered species that are highly outcrossing (e.g. *E. rostkoviana* flower size 8 - 12mm, F_{IS} =

0.17 - 0.25 [30]). Such continuous variation in outcrossing rate has been documented in species of

Datura [31], *Mimulus* [32] and *Nicotiana* [33]. Small flowers have shorter anther-stigma separation

and thus increased potential for autogamous selfing [32], while also having reduced attractiveness to

pollinators and thus receiving less outcross pollen [34]. In addition to differences in corolla size

between *Euphrasia* species, corolla size also shows a change of up to two millimetres in response to

host quality, with this change potentially affect the mating system [35]. This suggests host quality

represents a previously unaccounted factor affecting the mating system of parasitic plants.

Our comparisons of *Euphrasia* species in a common garden also sheds light on the distinctiveness of

these recently divergent species. *Euphrasia* is one of the most taxonomically complex plant genera,

with the 21 currently described British species presenting complex and often overlapping

morphological variation [13, 14, 36]. Our study suggests varying degrees of morphological

distinctiveness of *Euphrasia* species. We see *E. micrantha* is clearly morphologically distinct and *E.*

pseudokerneri mostly distinct, while the closely related species *E. arctica*, *E. confusa* and *E.*

nemorosa differ in life history traits such as nodes to flower, but overlap in many other traits and are

not clearly separated in the PCA. Moreover, our study is likely to overestimate the distinctiveness of

taxa by only including a subset of UK species. We suspect adaptive divergence between closely

related *E. arctica*, *E. confusa* and *E. nemorosa* is a consequence of differential natural selection for

local ecological conditions such as soil water availability or mowing. Selection appears to be

operating at a fine spatial scale, with significant life history trait differences evident between

populations within species. These taxa may be genetically cohesive, either showing genome-wide

divergence, or divergence in genomic regions underlying life history differences [37]. Alternatively,

these taxa may be polytopic and not genetically cohesive [38]. Genomic sequencing of natural

populations is hoped to resolve this issue.

Phenotypic plasticity in response to host quality

Our many hosts experiment shows *Euphrasia* benefit from growing with a range of hosts. *Euphrasia*

with a high quality host such as clover rapidly transitions to flowering. Differences in growth are

established early in the season, and early flowering plants go on to grow the tallest, are more highly

branched and have the potential to produce many extra flowers. At the other extreme, growing with a

poor host or without a host results in late-flowering minimally vegetative plants. Most hosts fall

between these extremes and offer moderate return to the parasite. *Euphrasia* parasitizing *Arabidopsis*

grew relatively tall despite the host senescing early in the growth season, similar to an experiment

where hemiparasitic *Melampyrum arvense* grew vigorously on short-lived *Capsella bursa pastoris*

[12]. *Euphrasia* on *Equisetum* performed surprisingly well, similar to the commonly encountered

grass *Holcus lanatus*. This could be through direct benefits via haustorial attachment to *Equisetum*

roots, or indirect benefits through association with *Equisetum* fungal symbionts [39]. This could be

tested by observing below ground attachment, which would also be interesting for *Pinus*, where poor

Euphrasia performance may be caused by lack of attachment or defence to parasite attack. Our results

point to *Euphrasia* being a true generalist parasite in what it can attach to, but much more specialised

in what it can benefit from. The effects of hosts on the growth of *Euphrasia* are consistent with

Melampyrum, where there is a general trend that legumes are the best host, followed by non-

leguminous herbaceous species, and grasses, with considerable variation among species within

functional groups [12]. Overall host quality may be attributed to root architecture, germination time

and resource availability, as well as the presence of mechanisms to defend against parasite attack,

such as cell wall thickening, localised host dieback, and chemical defence [7, 9]. Many of these

attributes of host quality appear to be shared between related hemiparasites in the Orobanchaceae,

though contrasting results from *Rhinanthus* suggest there are not universal 'good' hosts for all

parasitic plants [12].

The availability of suitable *Euphrasia* hosts in the wild will vary in space and time, and theory

predicts such fluctuating environments should select for trait plasticity [40]. As expected, we see

considerable plasticity in response to host quality. Only three pairs of trait correlations are consistent

across both *Euphrasia* common garden experiments (between height, number of branches and leaf

teeth), with the positive correlation between node to flower, height, and number of branches seen

between species breaking down when *Euphrasia* are grown on different hosts. The most notable

plasticity is seen in flowering time, with plants on the best hosts rapidly transitioning to flower within

~100 days of germination, while plants with a more typical host flower a month later. Phenotypic

plasticity in flowering time in response to resource availability is well documented in many plant

groups, particularly *Arabidopsis* (e.g. [41]), but has largely been overlooked in studies of parasitic

plants, which are more likely to look at fitness related traits such as number of flowers or biomass [12,

42]. We expect date of first flowering to be critical for the life time reproductive success of parasitic

plants in the wild.

In contrast to traits showing extensive plasticity, we also see evidence of developmental constraint in

number of nodes to flower. This trait showed the least plasticity with different hosts, is consistent

between populations within species, and between the common garden and the field. This suggests that

the developmental event of transitioning to flower is genetically determined, with changes in

flowering time altered by plasticity in internode length and not nodes to flower. This may explain why

nodes to flower is such an important diagnostic trait for species identification in *Euphrasia* and related

species in the Rhinantheae [43]. Despite nodes to flower changing little in response to host quality,

our overall impression is that *Euphrasia* show considerable plasticity and little developmental

constraint in many aspects of growth, or that trait values are affected by complex interactions related

to host-parasite attachment. In particular, the variation between individuals on a given host, suggests

other variation, such as genetic background in host and parasite, as well as the timing of attachment,

may be crucial in determining performance.

Conclusion

Despite over a century of experimental studies in parasitic plants, our understanding of the evolution

of life history strategies in these diverse organisms is extremely limited. Our results in *Euphrasia*

provide strong support for the rapid evolution of distinct life history strategies in response to local

ecological conditions, with phenotypic plasticity further altering plant growth in response to host

availability. We anticipate that future studies that test life time reproductive success of many parasitic

plant species grown on many different host species will give further insight into the complex nature of

host-parasite interactions.

Data accessibility

Phenotypic data from both common garden experiments and from herbarium collections, as well as

the R scripts used for data analysis, are deposited in Dryad XXXXXX (accession to be given on

acceptance).

Authors' contributions

A.D.T conceived and designed the research. A.D.T., N.F. and E. L. Y.W. carried out the experiments.

C.M. identified the plants. A.D.T. and M.B. analysed the data. A.D.T. and M.B. wrote the manuscript.

All authors read and approved the manuscript.

Competing interests

We have no competing interests.

Funding statement

A.D.T. is supported by a Natural Environment Research Council (NERC) Fellowship NE/L011336/1.

N.F. acknowledges funding from the Scottish Government's Rural and Environment Science and

Analytical Services Division (RESAS). M. B. is funded by BBSRC EASTBIO studentship.

Acknowledgements

We are indebted to the *Euphrasia* research community and to botanical recorders for providing seed

and herbarium specimens from natural populations. We also thank horticulture staff at the RBGE for

help with plant care. Jarrod Hadfield provided statistical advice and Susanne Wicke provided useful

information about parasitic plant biology.

Legends to figures and tables.

Figure 1. Trait variation in a common garden experiment of (a) diverse *Euphrasia* species and hybrids grown on clover, (b) *Euphrasia arctica* grown on many different hosts. The edges of the boxplots show the first and third quartiles, the solid lines the median, the whiskers the highest and lowest values within 1.5-fold of the inter-quartile range and the dots each individual measurement.

Figure 2. Relationship between morphological trait measurements made in the common garden and on wild-collected herbarium specimens. Bars represent the standard error of measurements.

Table 1. Summary of generalised linear mixed effects models of the influences on life history, plant vigour, and taxonomic traits for *Euphrasia* in a common garden. Table summarises model outputs for the many species and the many hosts experiment. The percentage variance explained by random effects are reported in brackets along with the 95% credibility interval. †Models for many branches were implemented with a different statistical approach in MCMCglmm (see methods). *** $P < 0.001$, ** $P < 0.01$, * $P < 0.05$

Table 2. Correlation coefficients for seven phenotypic traits measured in a common garden experiment for (a) Five *Euphrasia* species and 6 hybrids, (b) *Euphrasia arctica* grown with 8 hosts and without a host. *** $P < 0.001$, ** $P < 0.01$, * $P < 0.05$

Supporting information

Figure S1. Principal components analysis of morphological variation of *Euphrasia* in a common

garden for (a) five species and six hybrids, (b) five species, (c) *Euphrasia arctica* with nine host

treatments.

Figure S2. Relationship between early and midseason growth and flowering related traits and end of

season height. (a) Height at first flowering, (b) height 6-weeks after germination, (c) Julian days to

flower, (d) number of branches. Length measurements are reported in mm.

Table S1. Species and collection details for hosts used in the common garden experiment.

Table S2. Collection details for *Euphrasia* species used in the common garden experiment.

Table S3. Summary of trait values for many *Euphrasia* species and hybrids grown on a clover host.

Values are mean +/- one standard error. Length measurements are in mm.

Table S4. Summary statistics for the principal components analyses of (a) five species and six

hybrids, (b) five species, (c) *Euphrasia arctica* with nine host treatments.

Table S5. Summary of trait values for *Euphrasia arctica* grown on many different hosts. Values are

mean +/- one standard error. Length measurements are in mm.

Table 1

	Many species		Many hosts	
	Species	Population	Host/no host	Host
Branches†	$DIC_{full} = 867.36$; $DIC_{-spp} = 869.11$	$DIC_{full} = 867.36$; $DIC_{-pop} = 905.68$	n/a	n/a
Corolla length	$\chi^2(1) = 22.38^{*}$	$\chi^2(1) = 42.89^{***}$ (27.7%, 16.4 - 61.2%)	$\chi^2(1) = 3.19$	$\chi^2(1) = 28.79^{***}$ (13.1%, 5.6 - 52.3%)
Height	$\chi^2(1) = 20.10^{*}$	$\chi^2(1) = 67.68^{***}$ (49.1%, 28.8 - 75.5%)	$\chi^2(1) = 0$	$\chi^2(1) = 91.15^{***}$ (54.1%, 35.9 - 85.5%)
Internode ratio	$\chi^2(1) = 24.5^{**}$	$\chi^2(1) = 27.39^{***}$ (57.2%, 37.3 - 80.2%)	$\chi^2(1) = 1.38$	$\chi^2(1) = 6.94^{**}$ (4.6%, 0.8 - 23.0%)
Julian days to flower	$\chi^2(1) = 8.34$	$\chi^2(1) = 1.34$ (21.2%, 7.7 - 64.6%)	$\chi^2(1) = 2.52$	$\chi^2(1) = 32.50^{***}$ (37.4%, 13.4 - 66.3%)
Node to flower	$\chi^2(1) = 26.66^{**}$	$\chi^2(1) = 0.71$ (10.1%, 0.4 - 31.6%)	$\chi^2(1) = 0.38$	$\chi^2(1) = 0$ (1%, 0.18 - 9.8%)
Number of leaf teeth	$\chi^2(1) = 23.43^{*}$	$\chi^2(1) = 0$	$\chi^2(1) = 1.49$	$\chi^2(1) = 5.46^{*}$

Table 2

(a) Many species	Corolla length	Height	Internode ratio	Julian days to flower	Leaf teeth	Nodes to flower
Branches	0.068***	0.370***	0.014	0.003	0.433***	0.601***
Corolla length		0.102***	-0.026*	-0.016	-0.039***	-0.002
Height			0.060***	0.085***	0.317***	0.395***
Internode ratio				0.042*	0.014	0.006
Julian days to flower					0.003	0.062***
Leaf teeth						0.424***
(b) Many hosts	Corolla length	Height	Internode ratio	Julian days to flower	Leaf teeth	Nodes to flower
Branches	0.275***	0.696***	-0.089***	-0.327***	-0.482***	-0.062***
Corolla length		0.253***	0.010	-0.165***	0.287***	-0.028*
Height			-0.228***	-0.231***	0.480***	-0.035**
Internode ratio				0.001	0.028*	0.000
Julian days to flower					-0.478***	0.281***
Leaf teeth						-0.057***

Literature cited

1.   Price P.W. 1980 *Evolutionary biology of parasites*, Princeton University Press.

2.   Windsor D.A. 1998 Controversies in parasitology, Most of the species on Earth are parasites. *International Journal for Parasitology* **28**, 1939-1941. (doi: 10.1016/S0020-7519(98)00153-2).

3.   Fain A. 1994 Adaptation, specificity and host-parasite coevolution in mites (ACARI). *International Journal for Parasitology* **24**, 1273-1283. (doi: 10.1016/0020-7519(94)90194-5).

4.   Reed T.E., Waples R.S., Schindler D.E., Hard J.J., Kinnison M.T. 2010 Phenotypic plasticity and population viability: the importance of environmental predictability. *Proceedings of the Royal Society of London B: Biological Sciences* **277**, 3391-3400.

5.   Birget P.L.G., Repton C., O'Donnell A.J., Schneider P., Reece S.E. 2017 Phenotypic plasticity in reproductive effort: malaria parasites respond to resource availability. *Proceedings of the Royal Society B: Biological Sciences* **284** (doi: 10.1098/rspb.2017.1229).

6.   Westwood J.H., Yoder J.I., Timko M.P. 2010 The evolution of parasitism in plants. *Trends in plant science* **15**, 227-235.

7.   Twyford A. D. 2018 Parasitic plants. *Current Biology* **In Press.**

8.   Schneeweiss G M. 2006 Correlated evolution of life history and host range in the nonphotosynthetic parasitic flowering plants *Orobanche* and *Phelipanche* (Orobanchaceae). *Journal of Evolutionary Biology* **20**, 471-478. (doi:10.1111/j.1420-9101.2006.01273.x).

9.   Cameron D.D., Coats A.M., Seel W.E. 2006 Differential resistance among host and non-host species underlies the variable success of the hemi-parasitic plant *Rhinanthus minor*. *Annals of Botany* **98**(6), 1289-1299. (doi:10.1093/aob/mcl218).

10.   Twyford Alex D. 2017 New insights into the population biology of endoparasitic Rafflesiaceae. *American Journal of Botany* **104**, 1433-1436. (doi:10.3732/ajb.1700317).

11.   Koch L. 1891 Zur Entwicklungsgeschichte der Rhinanthaceen (II *Euphrasia officinalis* L.). *Jahrb Wiss Bot* **22**, 1-34.

12.   Matthies D. 2017 Interactions between a root hemiparasite and 27 different hosts: Growth,

biomass allocation and plant architecture. *Perspectives in Plant Ecology, Evolution and Systematics*

24, 118-137. (doi:https://doi.org/10.1016/j.ppees.2016.12.006).

13.   Wang X., Gussarova G., Ruhsam M., de Vere N., Metherell C., Hollingsworth P.M., Twyford

A.D. 2018 DNA barcoding a taxonomically complex hemiparasitic genus reveals deep divergence

between ploidy levels but lack of species-level resolution. *AoB PLANTS* **10**, ply026-ply026.

(doi:10.1093/aobpla/ply026).

14.   Metherell C., Rumsey F.J. 2018 *Eyebrights (Euphrasia) of the UK and Ireland*. Bristol,

Botanical Society of Britain and Ireland.

15.   Yeo P. 1968 The evolutionary significance of the speciation of *Euphrasia* in Europe.

Evolution **22**, 736-747.

16.   Stace C.A., Preston C.D., Pearman D.A. 2015 *Hybrid flora of the British Isles*, Botanical

Society of Britain and Ireland.

17.   Zopfi H.J. 1998 The genetic basis of ecotypic variants of *Euphrasia rostkoviana* Hayne

(Scrophulariaceae) in relation to grassland management. *Flora* **193**, 41-58.

(doi:https://doi.org/10.1016/S0367-2530(17)30814-9).

18.   Yeo P. 1961 Germination, seedlings, and the formation of haustoria in *Euphrasia*. *Watsonia*

5, 1-22.

19.   Yeo P. 1964 The growth of *Euphrasia* in cultivation. *Watsonia* **6**(1), 1-24.

20.   Hadfield J.D. 2010 MCMC methods for multi-response generalized linear mixed models: the

MCMCglmm R package. *Journal of Statistical Software* **33**, 1-22.

21.   Nakagawa S., Schielzeth H. 2010 Repeatability for Gaussian and non‐Gaussian data: a

practical guide for biologists. *Biological Reviews* **85**(4), 935-956.

22.   Bates D., Mächler M., Bolker B., Walker S. 2014 Fitting linear mixed-effects models using

lme4. *arXiv preprint arXiv:14065823*.

23.   Wickham H. 2016 *ggplot2: elegant graphics for data analysis*, Springer.

24.   Roff D.A. 1992 The evolution of life histories: theory and analysis. *New York: Ghapman &*

Hall.

25.   Stearns S.C. 1992 *The evolution of life histories*.

26.   Friedman J., Twyford A.D., Willis J.H., Blackman B.K. 2015 The extent and genetic basis of

phenotypic divergence in life history traits in *Mimulus guttatus*. *Molecular Ecology* **24**, 111-122.

(doi:10.1111/mec.13004).

27.   Mojica J.P., Lee Y.W., Willis J.H., Kelly J.K. 2012 Spatially and temporally varying

selection on intrapopulation quantitative trait loci for a life history trade-off in *Mimulus guttatus*.

Molecular Ecology **21**, 3718-3728.

28.   Hall M.C., Lowry D.B., Willis J.H. 2010 Is local adaptation in *Mimulus guttatus* caused by

trade-offs at individual loci? *Molecular Ecology* **19**, 2739-2753.

29.   Stone H. 2013 Evolution and conservation of tetraploid *Euphrasia* L. in Britain. PhD Thesis,

University of Edinburgh.

30.   French G., Ennos R., Silverside A., Hollingsworth P. 2005 The relationship between flower

size, inbreeding coefficient and inferred selfing rate in British *Euphrasia* species. *Heredity* **94**, 44-51.

31.   Motten A.F., Stone J.L. 2000 Heritability of stigma position and the effect of stigma-anther

separation on outcrossing in a predominantly self-fertilizing weed, *Datura stramonium* (Solanaceae).

American Journal of Botany **87**, 339-347.

32.   Karron J.D., Jackson R.T., Thumser N.N., Schlicht S.L. 1997 Outcrossing rates of individual

Mimulus ringens genets are correlated with anther–stigma separation. *Heredity* **79**, 365.

(doi:10.1038/hdy.1997.169).

33.   Breese E. 1959 Selection for differing degrees of out-breeding in *Nicotiana rustica*. *Annals of*

Botany **23**(2), 331-344.

34.   Mitchell R., Karron J., Holmquist K., Bell J. 2004 The influence of *Mimulus ringens* floral

display size on pollinator visitation patterns. *Functional Ecology* **18**, 116-124.

35.   Luo Y., Widmer A. 2013 Herkogamy and its effects on mating patterns in *Arabidopsis*

thaliana. *PLOS ONE* **8**, e57902. (doi:10.1371/journal.pone.0057902).

36.   French G., Hollingsworth P., Silverside A., Ennos R. 2008 Genetics, taxonomy and the

conservation of British *Euphrasia*. *Conservation genetics* **9**, 1547-1562.

37.   Twyford A.D., Friedman J. 2015 Adaptive divergence in the monkey flower *Mimulus*

guttatus is maintained by a chromosomal inversion. *Evolution* **69**, 1476-1486.

(doi:10.1111/evo.12663).

38.   Hollingsworth P.M., E. Neaves L., Twyford A.D. 2017 Using DNA Sequence Data to

Enhance Understanding and Conservation of Plant Diversity at the Species Level. In *Plant*

Conservation Science and Practice: The Role of Botanic Gardens (eds. Oldfield S., Blackmore S.),

pp. 23-48. Cambridge, Cambridge University Press.

39.   Bouwmeester H.J., Roux C., Lopez-Raez J.A., Becard G. 2007 Rhizosphere communication

of plants, parasitic plants and AM fungi. *Trends in plant science* **12**.

(doi:10.1016/j.tplants.2007.03.009).

40.   Caswell H. 1983 Phenotypic plasticity in life-history traits: demographic effects and

evolutionary consequences. *American Zoologist* **23**, 35-46.

41.   Zhang J., Lechowicz M.J. 1994 Correlation between time of flowering and phenotypic

plasticity in *Arabidopsis thaliana* (Brassicaceae). *American Journal of Botany* **81**, 1336-1342.

(doi:10.2307/2445409).

42.   Ahonen R., Puustinen S., Mutikainen P. 2005 Host use of a hemiparasitic plant: no trade-offs

in performance on different hosts. *Journal of Evolutionary Biology* **19**, 513-521. (doi:10.1111/j.1420-

9101.2005.01024.x).

43.   Jonstrup A., Hedrén M., Andersson S. 2016 Host environment and local genetic adaptation

determine phenotype in parasitic *Rhinanthus angustifolius*. *Botanical Journal of the Linnean Society*

180, 89-103. (doi:10.1111/boj.12352).

a Nodes to flower

b Standard corolla length

c Mean number of leaf teeth

d Mean cauline leaf: internode ratio

Taxa

- E. arctica
- E. confusa
- E. nemorosa
- E. pseudokerneri
- E. anglica x nemorosa
- E. anglica x rostkoviana
- E. arctica x confusa
- E. arctica x nemorosa