Full title: Multivariate methods for testing hypotheses of temporal community dynamics

Running header: Testing hypotheses of temporal community dynamics

Authors and affiliations:

Hannah L. Buckley[1]*, Nicola J. Day[2], Bradley S. Case[1], Gavin Lear[3], Aaron M. Ellison[4]

[1] School of Science, Auckland University of Technology, Auckland, New Zealand

[2] Biology Department, Wilfrid Laurier University, Waterloo, Ontario, Canada

[3] School of Biological Sciences, University of Auckland, Auckland, New Zealand

[4] Harvard Forest, Harvard University, Petersham, Massachusetts, USA

* Corresponding author: Hannah.Buckley@aut.ac.nz

ABSTRACT

For ecological research to make important contributions towards understanding and managing temporally-variable global change processes, such as responses to land-use and climatic change, we must have effective and comparable ways to quantify and analyse compositional change over time in biological communities. These changes are the sum of local colonisation and extinction events, or changes in the biomass and relative abundance of taxa within and among samples. We conducted a quantitative review of currently available methods for the analysis of multivariate datasets collected at temporal intervals. This review identified the need for the application of quantitative, hypothesis-based approaches to understand temporal change in community composition, particularly for small datasets with less than 15 temporal replicates. To address this gap, we: (1) conceptually present how temporal patterns in community dynamics can be framed as specific, testable hypotheses; (2) provide three fully-worked case-studies, complete with R code, demonstrating multivariate analysis methods for temporal hypothesis testing and pattern visualisation; and (3) present a road map for testing specific, quantitative hypotheses relating to the underlying mechanisms of temporal community dynamics.

KEYWORDS

Beta diversity, biodiversity, community variation, compositional change, multivariate analysis, species turnover, temporal change, temporal variability, time series, zeta diversity

INTRODUCTION

A key task for ecologists is to monitor, quantify, analyse, and predict temporal changes in the structure of ecological communities as a function of ongoing changes in climate, land use, and other environmental drivers. Ecological communities comprise populations of multiple taxa. The structure of communities—the number of taxa ('richness') and their relative abundance ('composition')—changes through time because of fluctuations of individual

populations and local colonisations and extinctions of taxa (Micheli et al. 1999, Legendre and

Gauthier 2014). These temporal dynamics are governed by both intrinsic factors, including

intraspecific and interspecific interaction, colonisation, and extinction, and extrinsic factors,

which are processes that relate to periodic disturbances and directionally changing

environmental conditions (Connell and Slatyer 1977, Brown and Lawson 2010).

Understanding the temporal dynamics of community structure can illuminate fundamental

ecological processes, including effects of individual life histories on rates of ecosystem

change, the relative importance of biotic and abiotic factors in determining community

structure, or how taxa and the networks in which they are embedded respond to

environmental change (Tilman 1999, Hobbs et al. 2007, Kampichler et al. 2012). Temporal

changes in taxon richness and composition also can provide both simple summaries of

ecosystem change and information that can be translated rapidly into action, either to protect

and restore declining taxa, or to staunch unsustainable growth of unwanted organisms.

Representations of community structure as multivariate datasets that include

information on multiple taxa can be more sensitive to changes in explanatory or predictor

variables (e.g., pH, temperature, soil type), than univariate summary statistics, such as taxon

richness, diversity, or biomass (Micheli et al. 1999, Chao et al. 2014), and hence more

informative about causal processes. However, quantitative methods for investigating and

testing hypotheses about temporal dynamics of multivariate measures of community structure

(henceforth 'community dynamics') also are more complex than methods used for univariate

data (Schaefer et al. 2005, Legendre and Legendre 2012). To date, methods for the analysis

of spatial multivariate community structure data (e.g., Whittaker 1960, Legendre and

Legendre 2012) have received far more attention than methods for analysis of temporal

multivariate data. Well-known concepts used to describe spatial changes in community

structure—grain, extent, nestedness, beta diversity, taxonomic turnover—have been

transferred to the temporal domain (Azeria and Kolasa 2008, Anderson et al. 2011, Legendre

and Gauthier 2014), and some well-developed multivariate methods for large spatial datasets

have been used to detect temporal changes in community structure (Legendre and Gauthier

2014). However, although temporal studies of compositional change are commonly

undertaken, it is not always clear which analytical methods are appropriate or available for

any given research question, data type, or community structure; indeed, there may not be one

ideal method for all situations and a range of methods may be required to answer a given

question (Whittaker 1960). Comparisons of multivariate ecological data through time pose

unique, non-trivial challenges because these analyses must also account for spatial variation

in these data, (Philippi et al. 1998, Micheli et al. 1999, Collins et al. 2000, Schaefer et al.

2005).

In this paper, we: (1) review empirical studies that have undertaken analyses of

multivariate temporal community datasets, to gain insight into how researchers have been

applying analysis methods over the past several decades, (2) provide recommendations for

the greater use of quantitative methods for testing specific temporal hypotheses, and (3)

provide guidance for researchers, including those new to community dynamics research, in

analysis decisions, particularly for small temporal datasets (fewer than 15 temporal

replicates).

How to use this paper

This paper does not necessarily need to be read from start to finish; we suggest readers

identify the section(s) most appropriate for their interests. The remainder of this paper is

structured into five main sections: (A) a review of the recent literature; (B) a conceptual

description of the types of hypotheses that can be addressed in analyses of temporal

community dynamics data; (C) three worked case study examples illustrating hypothesis-

driven analyses of temporal community data, including code for the software programme 'R'

(R Core Development Team 2017) to allow readers to easily implement methods described in

this paper; (D) A "road map" to analysing temporal community data, including descriptions,

and how to identify and present, analyses appropriate for testing particular hypotheses; (E) A

brief discussion of recent advances in multivariate analysis of community dynamics.

A. LITERATURE REVIEW

DATA COLLECTION AND ANALYSIS

We reviewed and described the multivariate analyses of community dynamics in studies

published up until September 2017 to provide an overview of available methods, their

benefits and drawbacks, and what they can reveal about community dynamics. Specifically,

we asked: Are data collected from different taxonomic groups, habitats, spatial extents, and

time-scales analysed in particular (or different) ways, i.e., are there consistencies in analysis

approaches due to discipline-specific preferences?

We searched the literature for studies investigating changes in ecological

communities over time. We searched the ISI Web of Science Core Collection on 5 September

2017 for articles using the following search statement: ["temporal" AND "composition*"

AND "communit*" AND ("community dynamic*" OR "temporal dynamic*" OR "temporal

community varia*" OR "community change" OR "compositional change")]. We refined this

search to include only document types listed as 'articles' and published in 'English'. This

search yielded 692 articles for review. Of these, we excluded 241 papers that lacked

community data (i.e. those that included fewer than three taxa), had no temporal replication

or pooled all temporal data, used only artificial communities or simulated datasets, or used

space-for-time substitution (e.g., using chronosequences) to infer community ages. Although

we did not constrain the temporal search window, our search terms only returned papers after

1990, indicating that the specific terms we used were not part of the compositional change

literature prior to the 1990s.

We extracted and collated key attributes relating to each study in the remaining 451

papers from the ISI database (e.g., publication year, number of citations). Specific study

attributes were determined from the body text of each manuscript, including information on

(i) habitat type (e.g., estuarine); (ii) habitat location (e.g., U.S.A.); (iii) taxonomic identity

(e.g., vertebrate; fish); (iv) manipulation (e.g., natural variation, experimental resource

manipulation); (v) spatial extent (e.g., local, global); (vi) temporal grain and extent (e.g.,

minutes, months); (vii) how composition was measured (e.g., presence-absence, relative

abundances of taxa); (viii) how time was included in each analysis (i.e., explicitly as a

predictor variable, or where other variables represent time, e.g., climatic factors recorded at

different times); and (ix) the key perceived research question (e.g., exploration of yearly

dynamics) (Data S1). We categorised organisms being studied into the following taxonomic

groups: plants (Kingdom Plantae); vertebrates (Phylum Chordata); invertebrates (Phyla

Annelida, Arthropoda, Cnidaria, Ctenophora, Echinodermata, Mollusca, Platyhelminthes, or

Porifera); microeukaryotes (algae and Kingdoms Fungi and Protista); prokaryotes (Domains

Bacteria and Archaea); and viruses (non-cellular pathogenic organisms). The data types

collated and included papers are provided in the supplementary material (Data S1).

LITERATURE REVIEW RESULTS AND DISCUSSION

The 451 papers identified by our search, when compared by year to the number of papers in

community ecology, showed that every year since 1990, publications on temporal community

dynamics consistently have comprised approximately 24% of studies within community

ecology. Thus, studies on temporal dynamics have increased from less than five publications

per year in the early 1990s to more than 30 per year in the last five years (Appendix S1: Fig.

S1). Few of the 451 reviewed publications (20 papers; 4% of those reviewed) contained more

than one independent dataset; those instances were treated as multiple studies that were

analysed as separate data points (increasing our total sample size n to 475). The majority of

studies (70%) were conducted within North America and Europe (Fig. 1), likely reflecting the

relative availability of research funding and resources.

Reviewed studies generally focused on organisms, scales, and habitats that were

conducive to studying temporal community dynamics. For instance, prokaryotes and

invertebrates were studied most frequently (Fig. 1), being relatively dynamic and easy to

study over smaller temporal and spatial scales; by contrast, research on plants and vertebrate

fauna together accounted for less than one-third of all studies of community dynamics.

Although studies spanned a broad range of habitat types, taxa, spatial, and temporal scales,

there were two-to-three times more studies at small spatial extents (micro, point sample, and

local scales) across the 14 habitat categories identified from the articles (Appendix S1: Fig.

S2A). Similarly, the most common temporal extent (i.e., the time between the collection of

the first and last samples) of individual studies was years, and only a minority of studies on

plant and vertebrate communities extended for a decade or more (Appendix S1: Fig. S2B).

Smaller organisms were often measured at finer temporal resolutions (grain) than larger

organisms, and many studies on short-lived prokaryote and viral communities lasted more

than one year. The temporal grain of sampling generally was smaller for communities of

microorganisms than for communities of vertebrate animals and plants, for which the most

common grain for repeated sample collection was years rather than months or seasons

(Appendix S1: Fig. S2B).

Researchers frequently used a range of different analytical methods, even within one

study, to enable them to understand the complexities of observed temporal changes. Up to

eight different analytical methods were used (Fig. 2A), with about two-thirds of the studies

applying ≤ 3 methods, and most using two. Although the number of studies addressing

community dynamics have increased dramatically since the 1990s and have begun to

incorporate more computationally-complex methods over time (Appendix S1: Fig. S3A),

descriptive methods (i.e., visual, non-statistical comparisons) and ordination have remained

the most commonly-used analyses, comprising 33% and 22% of all applied methods,

respectively (Fig. 2B). In fact, 134 out of 451 studies (30%) used only descriptive approaches

to "analyse" their datasets, while another 35% of studies used descriptive methods in addition

to at least one other method. We suspect that the overall ease of use of descriptive and

ordination methods, their accessibility via readily-available software programs, and their

focus on visually-presenting complex multivariate data, together have driven this trend. The

use of the raw dissimilarity values (18%), time-lag analysis (9%), and cluster analysis (7%)

comprised the next most common types of methods; the remaining 12 methods were used

infrequently (Fig. 2B). Several methods captured by this review were not multivariate (see

Appendix 2), including univariate linear methods, temporal stability (*a.k.a.* coefficient of

variation), frequency change (testing for significant changes in the frequency of individual

taxa over time), and multiplicative change (percent cover change statistic for a single taxon or

taxon group used in subsequent regression analyses). Since 2005, the use of raw dissimilarity

methods has increased considerably. In the same decade, a steady stream of alternative

methods that incorporate time explicitly have been applied, albeit with poor uptake across the

wider literature in this area. These methods often are 'one-off' analyses that have been

applied only a handful of times; we refer to these as 'minor' methods (Appendix S1: Fig. S3).

Preferred ordination methods changed from detrended correspondence analysis (DCA) and

principal components analysis (PCA) in the early 1990s, to canonical correspondence

analysis (CCA) in the late 1990s, and, since the early 2000s, to non-metric multidimensional

scaling (NMDS) (Appendix S1: Fig. S3B).

Three-quarters of all analyses compiled were from observational studies aimed at

describing patterns of community change or related to monitoring community composition,

rather than exploring environmental effects or studying the effects of biotic processes or

interactions (Appendix S1: Fig. S4). In fewer cases, researchers were interested in the effect

of environmental change, such as seasonality or climatic cycles over longer time periods.

Our review also shows how common it is for researchers to analyse datasets with low

temporal replication. Across research aims and for most methods, most time series had fewer

than 15 observations (mean = 14.9, median = 7, mode = 4; Fig. 3); the few studies that used

network analyses and machine-learning methods used 50 or more observations taken through

time. Overall, neither the number of temporal replicates, nor any other study attributes we

extracted during our review (e.g., habitat, taxonomic group), appeared to explain well the

differences in how researchers analysed their data or their motivations for doing so.

Typically, researchers will have some notion of how their communities of interest

might be changing, which motivated them to undertake temporal sampling in the first place

(Bestelmeyer et al. 2011). In some cases, long-term datasets exist because previous

researchers began data collection with a rationale and method that is different from the

current research questions, and so we can sometimes be limited by past sampling designs.

Nonetheless, our review highlights that most studies describe temporal community dynamics

within their studied communities without explicit, quantitative hypotheses and specific,

testable predictions about the expected drivers of observed community changes (Appendix

S1: Fig. S4). Clearly, there are many benefits to entering the analysis pathway with a strong *a*

priori prediction of the compositional pattern(s) expected and the biological hypotheses that

underpin those predictions.

B. HYPOTHESES OF TEMPORAL COMMUNITY DYNAMICS

Quantitative and specific predictions of the nature and variability in temporal community

dynamics allows deeper understanding of the patterns and processes involved. However, to

achieve this, *a priori* expectations of change need to be developed and paired with the

appropriate analytical method(s) to achieve rigorous tests. Here, we describe in detail the

possible observed changes in community composition (taxon presence or abundance) in one

sampling unit (site, host, etc.) over time (Fig. 4) or in a set of sampling units over time (Figs.

5-7) and what some of the ecological research questions are that may underpin such

hypotheses. In each case, we give predictions for how pairwise dissimilarity values, the basis

for most of the commonly-used quantitative measures of compositional change identified in

the literature review, would change. Many of these hypotheses are not mutually exclusive

although some could be used as multiple alternatives to be tested simultaneously, depending

on the research question(s) of interest. Note that 'dissimilarity measures', also called

'distance measures' or 'beta-diversity', are the inverse of 'similarity measures'. These

metrics quantify the compositional distance between two samples (pairwise dissimilarity).

Simple predictions for a single sample unit: Persistence and directional change

Aside from the 'null' prediction of no change in composition ('persistence'; Fig. 4b), one of

the simplest predictions of compositional dynamics is that of 'directional' change, whereby

certain taxa either increase or decrease so that the dissimilarity of a sampling unit to itself at

previous sampling occasions increases (or decreases) over time (Fig. 4c); when composition

turns over as taxa are replaced, the magnitude of change in dissimilarity can be high. Such a

hypothesis could arise from many ecological processes, e.g., has the community undergone

'succession', whereby there is a predictable replacement of individual or groups of taxa over

time; has it undergone 'degradation', whereby rare taxa are lost from the system over time;

has it had an 'invasion' event, where the relative dominance of one taxon or functional group

increases; is it changing in composition in response to a changing biotic or abiotic variable,

such as predation or temperature?

Simple patterns for multiple sample units: Directional change, divergence and convergence

Where a set of sample units are all undergoing the same directional changes in composition

over time, pairwise 'spatial' dissimilarity among that set of sampling units, would not change

significantly over time, because all sampling units are moving in the same direction at the

same rate in compositional space because the same taxa are changing in the same ways (Fig.

5b). We also might ask whether sample units are becoming more dissimilar to each other, due

to interactions between the changing factors described above and characteristics of the

community, such as initial composition or differences in habitat conditions. In this case

'divergence' represents the hypothesis where a set of sampling units are all becoming more

dissimilar in their composition, so that their pairwise dissimilarities increase over time; this

term therefore does not apply to change within one sampling unit (Fig. 5c).

Two very similar patterns predict decreasing pairwise dissimilarity values over time.

In contrast to 'divergence', under 'convergence' taxonomic composition of two or more

sampling units becomes more similar over time (Fig. 5e) in a predictable way because of

increases or decreases in taxon presence or abundance; this is also called 'homogenisation'

(Fig. 5e), where there is a concurrent decrease in diversity and dissimilarity becomes

relatively low. In contrast, convergence simply implies that between sample-unit dissimilarity

declines over time, but not necessarily to a very low level. In the cases where taxa are lost,

the predicted overall decline in dissimilarly is large. These hypotheses of change would be

relevant if we are interested in the effects of an experimental treatment designed to increase

the growth of a particular set of taxa, or under invasion, such as the biotic homogenisation of

cities due to urbanisation, or loss of native taxa due to competition with the invading taxa.

Note that the predicted effect of invasion on dissimilarity values depends on whether we are

considering the trajectory of one sample unit (divergence) or many sample units

(convergence) and the among-sample unit variation in other interacting factors.

More complex temporal patterns: Perturbations, resilience, and cyclical change

More complex patterns in temporal dynamics involve compositional shifts that change in

their magnitude or direction over time. Natural or manipulative experiments asking questions

about the effects of disturbance on community composition can lead to complex predictions

of change. 'Perturbation' is a pattern for which the change in composition is initially high but

lessens over time and can be detected by comparison to undisturbed treatments or areas (Fig.

6a). The perturbation can occur as either a 'press perturbation' (Fig. 6b), such as the removal

of a predator or a continuous nutrient addition treatment, or a 'pulse perturbation' (Fig. 6c),

such as a disturbance event or a population density reduction; these two perturbations result

in different rates of change in pairwise temporal dissimilarities. Post-perturbation, a

community either could remain in a state that is dissimilar to un-perturbed sample units

('tipping point'), or could return to its previous state over time ('resilience'). The prediction

of a 'pivot point' or 'threshold' can be used to test for a significant compositional shift for

one or more sampling units over a relatively short period of time, meaning that temporal

pairwise dissimilarity among times for each sampling unit is low before, and also after, the

pivot point, but is high for temporal pairs across the pivot point (Fig. 6c).

Questions about the scale of temporal dynamics in a community, such as the presence

of seasonal cycles, lead to predictions of 'cyclical' change or 'periodicity', which represent

shifts in composition that repeatedly diverge and re-converge back to a previous state;

dissimilarity increases at short temporal scales but decreases at larger temporal scales, and

will alternate between high and low at increasing temporal scales that correspond to some

cyclical process, e.g., warming events (Fig. 4d).

Predictability of change: Instability and stabilisation

Questions about the predictability of compositional changes over time resulting from

interactions between causal-change processes, such as temperature shifts throughout a year,

and biotic or abiotic differences among sampling units, could also lead to hypotheses

regarding the variability in relative compositional change. For instance, 'instability' predicts

significantly increasing temporal variation in dissimilarity for a set of sampling units because

relative increases and decreases among taxa are not correlated or consistent; change is not

directional although mean dissimilarity would also likely increase with the variance (Fig. 5d).

In contrast, 'stabilisation' predicts the opposite: significantly decreasing temporal variation in

dissimilarity as initially significant compositional change lessens over time; pairwise

dissimilarity is initially high at short temporal scales (the opposite of Fig. 5d).

Taxon dynamics: Synchronicity

In many cases, we are interested in asking if different components of the community are

responding in similar or different ways over time. Where individual taxa, or particular groups

of taxa (e.g., functional groups), change in similar ways at the same time ('synchronicity'),

we can detect this because among-taxon-group dissimilarity increases, but within-taxon-

group dissimilarity does not change (this involves comparing the pairwise sampling unit

dissimilarities separately for different taxonomic subsets) (Fig. 7a). Synchronicity may result

in directional or other types of compositional dynamics patterns within one sampling unit or a

set of sampling units (Fig. 7b).

Hypotheses of biological processes causing dynamics may be very specific, such as a

decrease in certain prey taxa in response to a predator, or more general, such as a shift

towards a grassland community in response to fire disturbance of a woodland or an increase

in annual plant taxa in spring compared to winter. More specific and directional (increases or

decreases) hypotheses lead to more powerful tests of compositional change; we strongly

encourage researchers to generate such hypotheses *a priori* and to pair these with predictions

that are as quantitative as possible, e.g., define the predicted timescales of variation.

C. WORKED CASE STUDY EXAMPLES

Here, we provide three worked examples to demonstrate the hypothesis-testing approach we

advocate for different types of community dynamics. We used open-access datasets to test

for: (1) significant divergence in temporal community dynamics among disturbance

treatments imposed upon plant communities in experimental grasslands, (2) neutral

community dynamics for a fully-censused forest plot, and (3) seasonal (cyclical) temporal

dynamics in marine aquatic bacterial communities. We present a brief description of each

dataset and describe the results and interpretation for the relevant exploratory data analysis

(EDA; see section 'D. Road map for analysing temporal community dynamics') and

hypothesis-testing analyses. More detailed descriptions, explanations and R code for each

case study are provided as supporting information (Data S2-4).

TESTING FOR DIVERGENT COMMUNITY DYNAMICS WITHIN A CONTROLLED EXPERIMENT

Porensky et al. (2016a, 2016b) measured plant species composition annually from 2007 to

2014 in quadrats on 24 transects to which six grazing treatments were applied since 1982

(four transects in each treatment): none (exclosure), light, moderate, heavy, heavy-to-light

(lightly grazed since 2007; 'new light') and heavy-to-none (new exclosure since 2007). They

calculated the mean cover in 25 quadrats for each species on transects. We tested the

hypothesis that increasing grazing intensity causes increasing divergence (directional change;

as detailed in section B) in species composition relative to the control (no-grazing) treatment.

EDA (as described in section D)) showed that the 24 transects were dominated across

all measurements by ten common species out of a total of 92 species (Data S2: row and

column summaries, relative abundance plots and rank abundance curves). Species turnover

(section D 4.0) through time was high; mostly 30% or greater for time period-treatment

combinations (Fig. 8). Principal coordinates analysis (PCoA; section D 1.4 Ordination)

suggested that the main compositional differences corresponded to the grazing treatments,

and that transect composition changed over time, but not markedly (Data S2: PCoA analysis

and figure). PERMANOVA (section D 1.4a) showed that transects were significantly

compositionally distinct and changed significantly over time in different ways (Data S2:

PERMANOVA outputs – all three coefficients for a two-way PERMANOVA with transect,

year and their interaction were significant $P < 0.05$). Partitioning spatial and temporal beta

diversity (section D 1.2) showed that two of the control (exclosure) treatments differed

significantly (Fig. 9). Principal Response curves (section D 1.4b) showed that the grazed

treatments differed from the control and that the relative abundances of four species in

particular were influential in driving these differences among treatments, but did not show

increasing divergence over time (Fig. 10). This result is consistent with the analysis of

temporal coherence (icc = -0.04; see section D 1.4a and Data S2 for detailed description of

this statistic), which showed that transects were not synchronous in their temporal changes

(Data S2: temporal coherence analysis). Thus, we conclude that treatment differences in

turnover and shifts in composition were not consistent or directional and so we reject our

hypothesis that increasingly grazed transects diverged more over time relative to the control.

TESTING FOR NEUTRAL COMMUNITY DYNAMICS ACROSS SPATIAL SCALES

The 50-ha forest census plot on Barro Colorado Island, Panama, has been censused eight

times between 1982 and 2015 (Condit 1998, Hubbell et al. 1999,

http://ctfs.si.edu/webatlas/datasets/bci/, 2005). This plot was established in part to test for

neutral community dynamics (Hubbell 1979) by fully censusing all woody stems greater than

1 cm diameter at breast height every five years. Under this hypothesis, we expected to see no

directional or other patterns in temporal community dynamics ('persistence'; as described in

section B.). EDA showed that the dataset consisted of 324 species generating a highly right-

skewed abundance distribution (most species were rare) (Data S3: EDA analyses and section

D). A line graph plot of species' abundance values over time showed very little change in

total abundance over time (Data S3: line graph of abundance; Appendix S2: Univariate linear

methods). Two analyses that allowed us to test for stability in community composition in

time, i.e., change in species composition is lower than expected if individual trees were

randomly distributed in time (keeping their spatial positions fixed), showed that change in

species composition was lower than expected: (1) Zeta diversity analysis (Data S3: zeta

diversity and section D 2.0) showed that the presence of species across census times was

more consistent than expected (Fig. 11) and (2) total turnover in species abundances (Data

S3: total turnover rates analyses and section D 4.0) at each of the seven time periods was

lower than expected (Fig. 12), particularly when the plot data were at larger spatial scales.

Thus, we conclude that the observed lack of pattern is consistent with our hypothesis of

neutral community dynamics.

TESTING FOR SEASONAL COMMUNITY DYNAMICS

Gilbert et al. (2012) measured bacterial taxonomic composition monthly over six years using

pyrosequencing of the 16S rRNA gene for water samples from a single site off the coast of

Plymouth, United Kingdom (source data can be obtained from the journal's online

supplementary file). We tested the hypothesis that subsets of the community would show

cyclical patterns of temporal change due to seasonality. EDA (section D) showed that the

dataset consisted of 74 operational taxonomic units (OTUs) across 62 temporal samples.

Most taxa were common through time. Proportional turnover between temporal samples

varied from 4% to 48% (Data S4: turnover calculation and section D 4.0), which is relatively

low, and was consistent with the detrended correspondence analysis (DCA) result showing a

short gradient length of 1.8 for axis 1 (Data S4: DCA and section D. EDA). Similarly, PCoA

graphs (section D. EDA) showed little consistent change within years (Fig. 13). However, 13

of the OTUs were deemed to be highly variable in that they changed by more than 500 reads

across the sampling period (mean number of reads = 296 ± 829 S.D..; Data S4: row and

column summaries, relative abundance distributions and abundance plots and section D.

EDA).

Time-lag analysis (section D 1.3) showed that although there was a significant

increase in dissimilarity within increasing time lag, variability was high, showing that

temporal changes did not occur in a simple divergence over time pattern (Fig. 14). Temporal

synchrony analysis (sensu Loreau and de Mazancourt 2008) showed that some subsets of

OTUs in the community were significantly changing in similar ways (Synchrony = 0.27; $P <$

0.05; Data S4: temporal synchrony analysis and section D 3.0). When patterns in these

variable taxa were examined through time using zeta diversity analysis (Hui and McGeoch

2014) (Data S4: zeta diversity analysis; section D 2.0), which is based on the presence of

species in samples, winter samples shared significantly more species across sets of temporal

samples than autumn, spring and summer (Fig. 15; ANOVA, $P < 0.01$; Appendix S2:

Univariate linear methods). Further analysis of the most variable taxa using the raw

dissimilarity values based on abundance data calculated for temporal samples within seasons

showed clear cyclical patterns (Fig. 16; Data S4; section D 1.1). Thus, we conclude that the

patterns are consistent with our hypothesis of seasonal dynamics in community composition,

largely driven by a small subset of OTUs that varied in their abundance, rather than presence,

over time.

D. ROAD MAP FOR ANALYSING TEMPORAL COMMUNITY DYNAMICS

The two most common sets of methods seen in our review of the temporal community

dynamics literature were descriptive (mostly tables and figures that show changes in taxon

presence or abundance) and ordination-based methods. Although these techniques are useful

for illustrating temporal trends and qualitatively exploring patterns in the data, they are used

less often for testing specific quantitative hypotheses of patterns and processes in community

dynamics. We recommend that researchers augment such descriptions with methods that

quantify temporal dynamics in taxonomic composition in more powerful ways, thus allowing

more quantitative and potentially comparable descriptions of spatiotemporal patterns and

explicit hypothesis-testing of mechanisms underlying observed patterns in community

dynamics.

Once the hypotheses to be tested have been generated, we suggest that researchers

undertake five steps: (Step 1) perform exploratory data analysis (EDA), (Step 2) choose

multivariate methods that can test the *a priori* temporal hypotheses, (Step 3) test for statistical

significance, (Step 4) report on the analysis workflow, and (Step 5) consider how to present

the results of the selected methods in the text and in figures.

STEP 1: EXPLORATORY DATA ANALYSIS (EDA)

As with any data analysis, the first step in the analysis of temporal community dynamics is an

exploratory data analysis (EDA) phase (Zuur et al. 2010). This generates basic information

that can be used to both error-check the data and to gain an understanding of key aspects of

the dataset (Table 1). R code for the relevant EDA steps is provided in three worked

examples, described previously (Data S2-4).

a. Count the number of taxa, the number of samples and the number of empty

samples and missing data points

The ratio of the number of taxa and samples is an important consideration because some

multivariate methods are more susceptible than others to the influence of rare taxa (Lepš and

Šmilauer 2003) and these may be of more or less interest, depending on the research

hypotheses. Whilst long-term data and monitoring are highly valued (e.g., Hobbs et al. 2007),

it is unfortunately common for long-term datasets to contain missing values; these values

should be denoted as missing (i.e., 'NA' in R) and not given zero or other numerical values.

The importance of missing values and how to deal with them during data analysis depend on

the number of missing data points and how they are distributed across the dataset.

Multivariate analyses based on compositional dissimilarities, e.g., some ordination methods,

cannot be applied to datasets containing samples where no taxa are present (McCune and

Grace 2002). However, empty samples may be important for analyses where we are

interested in understanding responses to disturbance and local extinctions. The ratio of the

number of explanatory variables to the number of independent replicates is also important for

temporal analyses because it affects the degrees of freedom available for statistical testing

(Johnstone and Titterington 2009).

b. Examine taxon frequency distributions and response curves

Some multivariate analyses assume specific frequency distributions of taxa (a histogram of

the number of individuals or their relative abundance across samples for each taxon), e.g.,

CCA assumes that taxa have optima along environmental gradients ('unimodal' responses).

Examining the shapes of taxon frequency distributions is necessary for determining whether

taxa are responding strongly to environmental gradients in the dataset and whether their

distributions conform to analysis assumptions. Where many taxa are involved, the common

ones should be examined.

c. Determine the ratio of temporal to spatial variation in composition

The degree of spatial and temporal replication is important in determining the method of

analysis because few sites with many measurements over time would be analysed differently

from many sites measured only a few times. If the data have a nested structure, e.g., samples

are nested within sites, this needs to be accounted for in the analysis, particularly if one wants

to report significance values. For example, if plant species composition was recorded in

quadrats that were nested within transects that were nested within sites, then quadrats are not

fully independent replicates. Similarly, information from technical replicates (such as PCR

products) should be kept separate at this stage because they provide useful information about

variability at different scales in the dataset. It is inadvisable to average or pool data until the

significant sources of variation are understood. For instance, if we want to test hypotheses of

temporal dynamics, but there is a high degree of spatial variation among samples in the

dataset, this can alter analysis choices or how results are presented and interpreted (Thrush et

al. 2008). The relative amount of change in taxonomic composition can be estimated from the

differences in sample scores on DCA ordination axes, e.g., the 'gradient length' statistic

quantifies the turnover in taxonomic composition (Lepš and Šmilauer 2003); this relatively

descriptive analysis would ideally be conducted as part of the EDA process prior to selecting

an analysis pathway appropriate for testing a quantitative hypothesis of temporal dynamics,

such as divergence or convergence.

d.　Determine the optimal dissimilarity measure

A community dataset may be an abundance matrix (number of individuals), relative

abundance matrix (e.g., percent cover, sequence abundances), rank-abundance matrix

(ordinal abundance ranks), or a taxon-occurrence (presence-absence) matrix. These different

response variables can influence the choice of analysis methods, including which distance

measure to use to calculate pairwise community dissimilarity values for samples (see

Magurran 2004, or Legendre and Legendre 2012 for a comprehensive description). To ensure

that abundance data from different taxa are comparable using dissimilarity, data should be

standardised prior to analysis if they are measured on different scales (Legendre and

Legendre 2012). For example, taxon abundance values within each sample can be divided by

the total abundance within the sample, or can be standardised across all samples so that each

taxon has a mean of zero and standard deviation of one (Borcard et al. 2011). Alternatively, a

distance measure that incorporates a standardisation can be used, e.g., Hellinger distance

(Legendre and Gallagher 2001, Anderson et al. 2011, Legendre and Legendre 2012). The

commonly-used Euclidean distance measure is not usually appropriate for taxon abundance

data, which are often characterised by non-linear relationships among taxa, abundance values

measured on different scales, or by the 'double-zero problem', which is the idea that samples

that are missing the same taxa and therefore, have lower dissimilarity when using this

distance measure (Borcard et al. 2011). Comparing results based on presence-absence data

with those generated using relative abundance data allows us to evaluate the role of taxon

abundances in generating temporal patterns (e.g., Angeler and Johnson 2012). Estimating

abundances of each population within the community has its own important considerations,

i.e., variation in taxon detection probabilities, which have been addressed elsewhere (e.g.,

animals: Gaston and McArdle 1994, metagenomics data: Amend et al. 2010). Many distance

measures cannot be computed if there are empty samples or missing values. Deleting entire

rows or estimating missing data so that the dataset can be analysed may be an option in some

cases (McCune and Grace 2002). However, other measures, such as the Gower coefficient,

can be calculated while ignoring comparisons among samples with missing data (Legendre

and Legendre 2012; coefficient S15: 278).

STEP 2: CHOOSE ANALYSES FOR TESTING HYPOTHESES OF TEMPORAL DYNAMICS

In this section, we describe a range of methods for testing hypotheses of community

dynamics, including their outputs, highlighting in each case, the relevant ecological questions

and the kinds of hypotheses that those analyses can be used to address: pairwise dissimilarity-

based methods (section 1, below), zeta diversity (section 2), synchrony (section 3), turnover

rates (section 4) and multivariate regression modelling (section 5). Further, each of the

described methods has been cross-referenced with (i) which hypotheses it is best suited to

test; (ii) whether or not it allows the identification of individual taxon effects on, or responses

to, temporal change; (iii) whether or not explanatory and spatial variables can be explicitly

incorporated; (iv) the type of statistical testing normally used; and (v) any requirements for

the length of the time series (number of temporal samples) (Fig. 17). The selection of an

appropriate analytical method also can be influenced by the kind of output desired (Table 2).

For instance, some analyses result in outputs that require more effort to interpret than others,

whilst others show differences in taxon responses or generate outputs that can be added to a

geographic map. Further, as our literature review highlighted, in many cases several different

analytical methods often are appropriate for asking questions about temporal dynamics and

presenting the results at the desired level of complexity; it is also the case that many analyses

can be used to test a wide range of hypotheses, so our focus is on describing the strengths of

each analysis method and the minimum requirements for use, rather than on making explicit

recommendations. Thus, we recommend that, at this point, readers use Tables 2 and 3 to help

identify the methods of interest and then refer to the relevant sub-sections of section C that

describe those particular methods.

Our literature review revealed that in numerous cases, researchers performed and

compared separate analyses on their community matrices from each sample time, e.g.,

separate nestedness analyses at each sample time (Azeria et al. 2006); we do not specifically

address these applications here in any depth because these methods do not explicitly evaluate

community dynamics through time. We also focus on methods that are truly multivariate

rather than methods that are applied separately to abundances of individual taxa; however, if

less than ten taxa are of interest, it may be that hypothesis testing via univariate approaches

(e.g., separate linear regressions of taxon abundances) will be adequate for drawing synthetic

conclusions about community change. In the supplementary material (Appendix S2), we

summarise univariate methods that were commonly used in the literature review articles:

univariate linear methods, temporal stability (*a.k.a.* coefficient of variation), frequency

change, multiplicative change, and Markov chain modelling. A few studies used alternative

methods such as cluster analysis and network analysis; however, we excluded those because

we think the methods presented here provide equally good, or better, alternatives, but we

nonetheless provide an explanation of them (Appendix S2) because they appeared in our

literature review. Some of the analytical methods described below can be implemented in

several different software packages (e.g., PC-ORD, PRIMER, CANOCO), but, for simplicity

and widest accessibility, we illustrate their implementation in R (Data S2-4), except for

Moran's Eigenvector Maps (MEMs), which is given in Legendre & Gauthier (2014), and

Multivariate Autoregressive State Space modelling (MARSS), which is presented in the

documentation associated with the R package 'MARSS' (Holmes et al. 2012).

1. Methods based on pairwise dissimilarity values

Pairwise dissimilarity methods can be used to test nearly the full range of questions and

hypotheses in community dynamics (Anderson et al. 2011; Figs. 4-7). Pairwise dissimilarity

methods are the most frequent way ecologists quantify the difference in the identities of taxa

occurring in two samples. The result is a single number that measures the distance in

compositional space between the two samples. Dissimilarity can also be represented as

similarity, i.e., 1 − dissimilarity (Whittaker 1960, Anderson et al. 2011, Magurran 2013).

When applied to multiple samples, it results in a matrix of all possible comparisons (a

triangular dissimilarity matrix). For example, Collins (1992) measured the between-year

percent change in plant species composition for grassland plots that had been repeatedly

measured over a nine-year period. Dissimilarity methods are more desirable than univariate

methods, which analyse the abundance of each taxon separately, because they consider the

entire community at once and directly quantify change in community structure.

The development of dissimilarity indices is an active area of research. Magurran

(2013) is a useful general discussion of dissimilarity measures, Anderson et al. (2011)

provides a practical guide to selecting an appropriate index and issues of importance, such as

whether to include joint absences in the calculation. Different authors often use different

names for the same dissimilarity measure, so we recommend using a comprehensive and

consistent guide, such as Legendre and Legendre (2012; Table 7.4: 324). Legendre and De

Cáceres (2013) also provide comprehensive descriptions of most dissimilarity measures and

rationales for choosing one. Several recent advances are specifically relevant to temporal data

(e.g., Podani and Schmera 2011, Baselga 2012, Shimadzu et al. 2015) and some contributions

aid in improving dissimilarity measures for use in certain circumstances, such as imperfect

detection of taxa (e.g., Chao et al. 2006) or which indices are most appropriate for calculating

dissimilarity (Legendre and De Cáceres 2013). For example, the unique fraction (UniFrac)

dissimilarity metric was designed for microbial communities where phylogenetic information

is available from sequencing that has been used to determine taxa in samples (Lozupone and

Knight 2005, Lozupone et al. 2007). This metric measures phylogenetic distance between

samples as the fraction of branch lengths leading to descendants from each sample. UniFrac

can be used with abundance data, which weights the branch lengths prior to calculating

distances (weighted UniFrac), or with presence-absence data (unweighted UniFrac Lozupone

et al. 2007).

In the context of temporal community dynamics, dissimilarity is calculated in two

different ways: (1) separately among spatial samples at each of a series of time points or (2)

among samples taken at different times at the same location(s). Dissimilarity matrices are the

basis for several methods recommended in our guide, including ordination analyses and

Moran's Eigenvector Maps (MEMs). It is important to realise that it is not easy to use such

methods to simultaneously assess both temporal and spatial variation in the same analysis;

however, temporal and spatial dissimilarity can be computed and assessed in parallel to test

hypotheses and make inferences about spatiotemporal community dynamics (e.g., Legendre

and Gauthier 2014, Lamy et al. 2015).

1.1.   *Using the raw dissimilarity values*

There are two ways in which raw dissimilarity values for samples are typically used.

Either of these approaches could be used for different taxonomic subsets of the community to

test hypotheses of taxonomic variation in dynamics. First, compositional dissimilarity can be

calculated among samples taken at different locations to obtain a measure of spatial

heterogeneity in community composition. To use these values to analyse temporal

community dynamics, pairwise dissimilarity (spatial dissimilarity) must be calculated

separately for each sampling time, and the means of these spatial dissimilarity values can

then be compared across time visually in a table (e.g., Collins 1992), graph (e.g., Barros et al.

2014), or using a correlation or regression analysis where time is the predictor variable and

amount of spatial variability at each site is the response (e.g., Barros et al. 2014). The mean

dissimilarity can be divided by the number of taxa to generate an independent measure of

compositional change (Cleland et al. 2013). Such comparisons over time could be done

separately for different groups of samples, such as different experimental treatments, to test

for predictors of differences in temporal change in composition. Bagchi et al. (2017) illustrate

predictions of how temporal differences in dissimilarity can include non-linear patterns and

thus how applying linear and non-linear regression modelling to pairwise dissimilarity values

can be used to test more complex hypotheses of temporal dynamics. Regression approaches

can also include mixed-effects models that account for nesting and other types of non-

independence in the data, or non-linear models (e.g., generalised additive models) that may

be useful to predict cyclical or threshold patterns in community dynamics (e.g., Shenhav et

al. 2017). However, these modelling methods will apply well only in situations with longer

time series (i.e., greater than 25 temporal replicates) and the more complex the prediction of

temporal change, the greater replication is likely required. Alternatively, a Mantel test can be

used to compare pairs of dissimilarity matrices computed at different times and to test for

significant difference over time (see section in the main text 'Test for statistical

significance'). This approach is only useful for a small number of spatiotemporal

observations (i.e., ten or fewer) because the comparisons are pairwise.

Second, if a set of samples were all taken at the same site at different times, 'temporal

heterogeneity' can be calculated using the mean dissimilarity of the entire sequence of

temporal samples to obtain a single measure of within-site temporal variability (Collins and

Smith 2006). If multiple sites are sampled through time, this method can be applied at

different spatial scales (Collins 1992, Collins and Xia 2015) to examine spatial scale-

dependent temporal dynamics (Collins and Smith 2006). Regression approaches that relate

mean temporal dissimilarity values to other explanatory variables for sites or samples could

be used to explore spatial variability in temporal change. Alternatively, the 'dissimilarity-

overlap' method, first described by Bashan et al. (2016) and further explained by Kalyuzhny

and Shnerb (2017), can be applied to raw dissimilarity values to test for similarity among

samples in community dynamics. This method uses the relationship between the pairwise

dissimilarity values and the proportion of individuals in pairs of samples that overlap

taxonomically to detect if changes in taxonomic composition (degree of overlap) alters

relative abundances in a consistent way across samples (dissimilarities). It can be used for

very large numbers of samples and taxa and could be applied to different groups of taxa to

test for synchronicity.

1.2.   Beta diversity decomposition

'Beta diversity' is term used for several different types of measures in ecology, e.g.,

(Whittaker 1960, Baselga 2010, Legendre and De Cáceres 2013), including as a synonym for

compositional dissimilarity. It always describes some measure of how taxonomic

composition changes across a set of samples. A number of recent advances in the calculation

of beta diversity have illustrated that, if calculated in particular ways, it can be 'partitioned'

into different components that relate to different predictions of how taxa or their abundances

change in space or time (Baselga 2010, Ellison 2010, Legendre and De Cáceres 2013). Beta

diversity decomposition methods can be applied to testing almost any hypothesis of

compositional change (Fig. 17). The outputs are focussed on species replacement and

whether samples are subsets of each other (Table 2), so detecting patterns such as

convergence or divergence is very easy. However, depending on what subsets of the data are

used in separate calculations, beta diversity decomposition is equally able to detect rapid

shifts in composition, e.g., perturbation, tipping point, or increases or decreases in temporal

variability, e.g., 'instability' or 'stability'. Two approaches so far have been usefully applied

to quantifying community dynamics measured in the field (Baselga 2010, Baselga and Orme

2012, Legendre and De Cáceres 2013, Lamy et al. 2015).

First, when among-sample dissimilarity is measured as Sørensen's dissimilarity [$(b +$

$c)/2a + b + c)$, where a is the number of shared presences and b and c are the number of

unshared presences in each of the two samples], the Simpson's dissimilarity index (also

known as 'species turnover': $b / (b + a)$) can be subtracted from this value, resulting in a

measure of 'nestedness' (Baselga 2010, 2012, Baselga and Orme 2012). This measure of

nestedness reflects the degree to which the species composition of smaller samples (lower

species richness) is encompassed by the species composition of larger samples (higher

richness); samples may differ in space or time. In contrast, 'species turnover' measures the

degree to which samples do not share species: how the identities of species differ among

samples. It is useful to separate these two components (nestedness and turnover) to obtain a

more informative quantification of beta diversity that reflects differences in the underlying

processes. Such separation can be done either pairwise or for multiple sets of samples

(Baselga 2010, 2017), and has been used to decompose temporal beta diversity in tests of

hypotheses unique to temporal data (e.g., Baselga and Orme 2012).

Second, and in contrast, Legendre & De Cáceres (2013) defined beta diversity as the

sums of squares in the variation of taxon abundance (or presence) corrected for the number of

samples. They partitioned this measure of beta diversity into two different components:

'Species contribution to beta diversity' (SCBD) and 'local contribution to beta diversity'

(LCBD), which describe, respectively, the relative contributions of different taxa and sites to

overall beta diversity (Table 2; Legendre and Gauthier 2014). This method allows for tests of

predictions of patterns in temporal community dynamics related to different taxa in the

dataset. For example, Lamy et al. (2015) decomposed beta diversity (based on Bray-Curtis

dissimilarity) of fish communities at each of 13 reef sites into contributions from biomass

replacement and biomass differences over time. They demonstrated that total biomass in the

community changed little, but that the observed significant temporal changes in fish

composition resulted from differential changes in the relative biomass of species.

Spatiotemporal patterns in the two components of fish community beta diversity were

visualised using principal coordinates analysis (PCoA).

1.3.   Time-lag analysis

A time-lag analysis involves relating the amount change in composition to the amount of

change in time across increasing temporal distances, called 'lags' (Table 2). For example,

Collins & Xia (Collins and Xia 2015) compared plant species composition of repeatedly-

measured grassland transects across all time lags between 1989-2008 (from a 1- to 19-year

difference). There are two ways of assessing the time-lag effect on community dissimilarity:

graphically ('time-decay curve') and statistically (time-lag regression analysis). For

meaningful inference, time-lag graphs require at least five samples and statistical time-lag

analysis requires at least ten samples for each temporal sequence. For both approaches, the

pairwise compositional dissimilarity (temporal beta diversity) among temporal samples is

calculated first.

To generate a time-decay curve, pairwise temporal dissimilarity values are plotted

against temporal distance between samples (time lag; pairwise Euclidean distances in sample

times) to show how community dissimilarity changes as samples become further apart in time

(Collins 2000, Collins et al. 2000, Hallett et al. 2016). This graph can be then used to visually

assess whether community composition is converging (becoming more similar) or diverging

(becoming less similar) over time (e.g., Collins 2000). For example, Bar-Massada & Hadar

(2017) plotted temporal beta diversity (as Bray-Curtis dissimilarity) of plants in grazed and

ungrazed treatments in Mediterranean herb communities, revealing that composition diverged

over time, regardless of grazing treatment. Alternatively, temporal community dissimilarity

can be graphed against dissimilarity in environmental variables (rather than time), if the

primary research focus is on response of the community to environmental change rather than

time *per se*.

Time-lag regression analysis was developed specifically for data with greater than ten

time points to assess directional, cyclic or stochastic changes within communities (Collins

2000, Collins et al. 2000) and has been evaluated against other temporal analysis methods

(Angeler et al. 2009). For time-lag regression analysis, the pairwise compositional distances

are plotted against the square-root of the time lag, which reduces the bias that results from

having fewer data points at larger time lags (Collins 2000, Collins et al. 2000). A regression

line is then fitted to determine the pattern of compositional change. A significantly positive

relationship indicates that compositional changes in communities is directional, because

sampling points that are further apart in time are more dissimilar than those that are closer

together in time (i.e., Fig. 2 in Collins 2000). A negative slope shows that some convergence

towards the taxonomic composition at one of the earlier time points has occurred, because

measurements close together in time are more dissimilar in composition compared to those

further apart in time. No slope, or a non-significant regression, implies stochastic changes in

the community over time. Non-linear regressions also can be used to describe more complex

temporal patterns (Magalhães et al. 2007).

A robust measure of significance in time-lag regression analysis can be determined by

running Mantel tests on the dissimilarity matrix and the matrix of pairwise distances among

samples in time, with Monte Carlo permutations (Bêche and Resh 2007, Day et al. 2017).

Time-lag regression analysis is appealing because it is intuitive, produces one statistic (the

regression coefficient), and can be easily visualised (e.g., freshwater macroinvertebrates:

Bêche and Resh 2007, gut bacteria: Dimitriu et al. 2013, and plants: Collins and Xia 2015).

However, time-lag analysis only provides one value for each sampling unit, so further

analyses may be required to investigate changes within treatments or groups.

1.4.   Ordination

Ordination summarises multivariate community data by optimising relationships between

high-dimensional samples and taxa in low-dimensional space to detect important ecological

gradients in communities (Table 2). This dimensionality reduction means that they are not

especially useful for directly testing, or visualising, complex changes in community

composition over time, such as tipping points, cyclical patterns or synchronicity. However,

more simple patterns, such as divergence or convergence, can be more easily detected.

Ordinations are either unconstrained, meaning that they do not incorporate explanatory

variables, or constrained, which assess the effect of explanatory variables, including time

(e.g., Kent 2012, Legendre and Legendre 2012, Palmer 2019). Ordination methods, including

constrained ordination and using the ordination scores from an unconstrained ordination in a

regression model (e.g., general linear model, mixed-effects model) are only appropriate for

testing hypotheses of community dynamics in cases where EDA shows that the ordination

axes used strongly represent temporal compositional change and therefore explain a

substantial proportion of the variation. This is because the amount of temporal compositional

variation represented by the ordination axis scores will be low if there is not a strong

temporal gradient along the axes. Ordination methods allow only weak inference of the

relationship between taxa and community dynamics if the temporal pattern is complex

relative to the ordination axes because some ordinations, e.g., DCA, only provide an

ordination axis score for taxa, and others do not provide 'species scores' at all, e.g., PCoA

and NMDS. PCoA can be used to calculate the amount of variance explained by the

ordination axes, which is important if a time-series analysis is to be performed using the axis

scores. Where species scores are not provided, weighted averaging can be used to plot points

for each taxon on the ordination diagrams to relate differences in site composition to

differences in taxon presence or relative abundance if you wish to examine the associations

between samples and taxa (McCune and Grace 2002). One advantage of PCoA is that the

first PCoA axis represents the gradient explaining the greatest amount of variation in

composition, followed by the second axis that explains the next greatest amount, and so on;

this is not the case for NMDS (McCune and Grace 2002). Therefore, it is recommended that

NMDS scores are not used in community dynamics analyses unless the amount of variation

explained by each axis is clearly stated in order to validate their use in further analyses. 'Co-

inertia analysis' can be coupled with ordination in a 'partial triadic analysis' to investigate

community dynamics in relation to environmental variables. This method may be particularly

useful when testing for temporal stability (Kidé et al. 2015).

a.   Using ordination site scores to assess compositional change

Ordination axis scores for samples taken at different times can be used to quantify and test

hypotheses of compositional change. Ordination diagrams are visually appealing, but for

illustrating compositional change through time, it is difficult to interpret the magnitude and

significance of temporal dynamics without statistical testing, such as permutational analysis

of variance (PERMANOVA: Anderson 2001), unless temporal trends in the data are very

strong along the ordination gradients in composition. This difficulty is amplified if there are

many samples or if the sampling design includes a large number of sites, environmental

factors, or temporal replicates. Note that PERMANOVA requires the assumption of

homogeneity of dispersion to be met (this can be tested using the "betadisper" function in the

R package vegan; Oksanen et al. 2017). Further, ordination diagrams and their statistics

computed for different matrices (different sets of taxa or sites) are not comparable; results of

individual ordinations cannot be compared among them. The partial exception to this is

Procrustes analysis, which can measure the distortion, or 'lack of fit' between two

ordinations, or dissimilarity matrices (Jackson 1995, Peres-Neto and Jackson 2001). It can be

used to identify which taxa are most important in differentiating between the two ordination

patterns.

A more quantitative, and often more flexible, approach is to use the ordination axis

'site' scores in linear models with time and other explanatory variables of interest entering as

predictor variables (e.g., Berthon et al. 2014, Broadway et al. 2015). For example, Cotton et

al. (2015) used DCA scores as a response variable in a linear mixed model to investigate the

importance of carbon dioxide, ozone, and time on arbuscular mycorrhizal fungal community

composition. This is like doing a constrained ordination, such as RDA, dbRDA, or CCA,

where time is used as a constraining explanatory variable in the ordination; these methods

will test the significance of time and any other explanatory variables. However, it is more

difficult within a constrained ordination to test for non-linear, cyclical or other complex

temporal patterns, and in this case, using the ordination axis scores in a more complex model

may be more useful.

Ordination axis scores for samples can be used to quantify temporal changes in other

ways. For example, Quinn et al. (1991) developed a 'malleability index' by regressing the

pairwise Euclidean distances between the first three dimensions of ordination space for

temporal samples, against environmental variables. One potential problem with this method

is caused by the tendency of ordinations to distort the relative distances (dissimilarity)

between samples, depending on, for example, differences in the number of rare taxa, and so

values from distance-based indices may not be comparable, even in different parts of the

same ordination.

Temporal coherence measures the degree to which a response variable is similarly

variable in time across a set of sites. In the case of measuring temporal coherence in

community composition, the multivariate response variable is taxonomic composition in the

form of ordination axis site scores (Angeler and Johnson 2012). Temporal coherence

compares the amount of temporal variation in the ordination axis site scores across a set of

sites by using statistics generated by an ANOVA of the site scores (using year and site as

factors) to calculate the 'intra-class correlation coefficient', which ranges from $-1/(n-1)$ to

+1, where n is the number of sites sampled, given that $n > 1$ (for calculation details see Zar

1996 pp 398, Rusak et al. 1999). High positive values of this metric indicate increased

coherence of the different sites in composition over time. The measure can be recalculated

while dropping each time in sequence (an ordered jackknife) to determine the relative effect

of each time on the overall value (Rusak et al. 1999). Different sets of sites or treatments can

be compared using a Fisher's Z test (Angeler and Johnson 2012) or randomisation test in the

case of low sample sizes (Rusak et al. 1999).

b. Principal Response Curves (PRC)

The method of principal response curves (PRC) was first developed by van den Brink & ter

Braak (1999) for comparing compositional change in treatment sites relative to a control site.

It is similar to redundancy analysis (RDA) in that a multivariate regression is performed on

the sample × taxon matrix constrained by variables of interest, but PRC is explicitly

constrained by time (Lepš and Šmilauer 2003, ter Braak and Šmilauer 2015). It was

developed to eliminate the need for cluttered biplots that make it difficult to see temporal

trends within and between treatments. To overcome this problem, PRC uses a single axis,

plotted against time, to express trajectories of compositional change (Lepš and Šmilauer

2003: 145); further axes can be calculated if they are of interest. Taxon weights also can be

calculated and plotted for each PRC to show the change in the relative abundance (e.g.,

percent cover) of each taxon in the treatments relative to the control. The significance of the

PRC is tested using Monte Carlo simulations (Van den Brink and Ter Braak 1999). As with

RDA, PRC yields a measure of the percent variance explained for the overall taxonomic

composition and how much is attributable to each variable (treatment and time). For example,

Sugiura et al. (2013) used PRC to show that plant invasion in forests on a small island in the

north-western Pacific had little effect on the temporal dynamics of invertebrate communities.

The potential drawbacks of this method include its requirement for a 'baseline' or 'control'

treatment level to be set for comparison, an attribute that many observational datasets will not

have. Furthermore, PRC allows for only one 'treatment' variable and each analysis focusses

on the change at one site, relative to the control site. However, Auber et al. (2017) have

recently generalised this method so that it can be used to analyse compositional change at

multiple sites, relative to a baseline or control, but this application removes the ability to

consider an environmental or treatment effect. Thus, a principal response curve analysis

yields a line graph showing the relative change in taxonomic composition for each

'treatment' along one compositional gradient relative to a control (Table 2).

1.5.   Moran's eigenvector maps (MEMs)

Moran's eigenvector maps (MEMs; also known as 'principal coordinates of neighbour

matrices', PCNM) is a method designed to characterise the spatial scales of variation in

univariate or multivariate data (Borcard and Legendre 2002, Dray et al. 2006, Legendre and

Gauthier 2014). However, MEMs have recently been applied to detect temporal scales of

variation in communities (Angeler et al. 2009, Caruso et al. 2013, Legendre and Gauthier

2014). For instance, if a cyclical pattern in compositional change, such as seasonality, was

detected as significant for a dataset, this can be illustrated using line graphs showing the

temporal pattern (Table 3; Legendre and Gauthier 2014). This method is particularly useful

for detecting very complex temporal patterns, such as seasonality (periodicity), cyclical

changes or tipping points, but requires relatively large temporal sample sizes (Fig. 17). For

example, Kampichler et al. (2014) used MEMs to detect temporal patterns in multiple

datasets that had at least 20 annual measurements of bird communities.

First, a distance matrix is generated. Then, to focus on neighbour relationships, a

threshold value is chosen (see Borcard et al. 2011 for advice). For values below this

threshold, the distance value is retained; values above this threshold are assigned an

arbitrarily large value. Legendre & Gauthier (2014) recommend using one time interval as

the threshold for regular time series or the length of the longest lag for irregular series. A

PCoA is then applied to this modified distance matrix to generate a large number of

eigenvectors that describe a wide range of temporal scales of variation.

The eigenvectors from the PCoA can be used as explanatory variables, most often in a

redundancy analysis (RDA) or multiple regression. Combined with RDA, MEMs can be used

to determine whether there is cyclic or synchronised temporal change in composition and

which taxa, or taxon groups, are the drivers of it (Fig. 17). MEMs-RDA scores also can be

plotted as a function of time to assess visually the intervals of fluctuating community

dynamics (Angeler et al. 2009, Kampichler et al. 2014). In a comparison of different methods

to assess community dynamics, Angeler et al. (2009) observed that MEMs-RDA consistently

explained a greater proportion of variation in data with low taxonomic richness and linear

changes in time relative to time-lag regression analysis, Mantel tests, or RDA. However,

power was higher for taxon-rich communities because there was a greater probability of more

taxa fitting the temporal models. As a scale-detection method, MEMs could be applied to

determining the relevant time intervals for community monitoring. This method can be

combined with variance partitioning to show the relative variance in composition explained

by temporal, spatial and other explanatory variables (Legendre and Gauthier 2014).

1.6.   Compositional pivot days

Compositional pivot days is a dissimilarity-based method developed by, and unique to,

Lellouch et al. (2014). This method is appropriate for situations where we want to test for the

occurrence of fast changes in taxonomic composition across one point in time, such as the

application of an experimental treatment or the effects of a rapid change in environmental

conditions (Table 2). It works by comparing pairwise distance values for pairs of temporal

observations taken at the same sample location to identify the time points where rapid shifts

in composition occurred. A 'pivot day' is a sampled time point that is identified by pairwise

similarities being low before and after that time point, but relatively higher across it.

Compositional pivot days are identified either visually from the pairwise distance matrix plot,

or using clustering (see Lellouch et al. 2014 for details). This method may be a good option

for a preliminary identification of thresholds or rapid shifts in community structure, but does

not provide a statistical test for the significance of the threshold relative to background. The

power of this analysis will increase as the number of sampled time points increases relative to

the number of pivot days.

1.7.   Community trajectory analysis

Community trajectory analysis quantitatively compares the temporal change in multivariate

compositional dissimilarity space among samples using geometric calculations (de Caceres et

al. (2019). These comparisons are based on the relative amounts, rates, and directions of

sample movement in compositional space, thus, the method allows flexible testing of a wide

range of very specific, quantitative hypotheses of change, such as the divergence or

convergence of samples over time. This method is especially useful for a small number of

spatial replicates because the output can be graphical; plots use arrows on an ordination

diagram to show how samples have moved in compositional space over time; however, larger

numbers of samples could be graphed separately. We do not illustrate this method here

because comprehensive R code and explanations are presented in the R package 'vegeclust'

(De Cáceres et al. 2010) and demonstrative examples are given in de Caceres et al. (2019).

2. Zeta diversity

Zeta diversity, a measure of similarity, analyses change in community composition among

samples in the dataset by calculating the average number of taxa that are shared between

samples that are grouped in pairs, triples, quadruples, etc. (Table 3). The method was

developed and introduced by Hui & McGeoch (2014), extended to temporal data by

McGeoch et al. (2017), and comes with its own R package, 'zetadiv' (Latombe et al. 2017b).

Zeta diversity estimates compositional turnover from multiple ($n \geq 2$) samples, rather than the

pairwise comparisons used to estimate beta diversity. What is obtained by the method is a

series of zeta values (similarity) for different orders of zeta: $zeta_1$ to $zeta_n$, where n is the total

number of time points. Each of these values represents the average of the number of taxa

shared by each set of times in the dataset. For example, $zeta_2$ (zeta order 2) represents the

average number of taxa shared by all possible pairs of samples in the dataset (equivalent to 1

− beta diversity or mean similarity), $zeta_3$ represents the average number of taxa shared by all

possible subsets of three samples, and so on. $Zeta_1$ (zeta order 1) represents the average alpha

diversity (taxon richness) of all the samples in the dataset.

If there is complete turnover in community composition in time, at some point, $zeta_i$

(where $i \neq 1$) will equal zero when there are no taxa shared by a given sized subset of

samples. In contrast, if community composition never changes through time, $zeta_n = zeta_1$.

Therefore, the rate of decline in zeta with increasing 'zeta order' tells us about the rate of

change in community composition in time ('zeta decline'). 'Zeta decay' can be considered

analogous to time-lag graphs and time-lag regression analyses, which we described earlier for

pairwise dissimilarity measures (Table 3). This means it experiences the same issue of few

time points at higher lags, and thus higher uncertainty. The temporal scale(s) of community

dynamics can be illustrated by plotting the change in sample zeta diversity of a particular zeta

order against the temporal lag of samples, meaning that this method is useful for detecting

both directional changes and cyclical patterns. For instance, if a multi-year time series in

composition is cyclical (e.g., shows seasonality), zeta would seasonally decrease and then

increase. These analyses can be done for different sets of samples to make spatial or temporal

comparisons and, if necessary, zeta values can be normalised using the gamma diversity

(total number of taxa in the dataset) to account for differences in taxon richness across

samples (McGeoch et al. 2017). Further, different methods of creating the subsets for

calculating zeta can be used, such as nearest neighbours or relative to a time-zero sample,

further facilitating hypothesis testing (McGeoch et al. 2017). Latombe et al. (2017a) used

zeta diversity to extend generalised dissimilarity modelling (Ferrier et al. 2007). This enables

the user to assess taxonomic turnover at different orders of zeta and understand drivers of

turnover in rare and common taxa (Latombe et al. 2017a). This method could be extended to

assess temporal turnover at different orders of zeta by adding time as an environmental

variable.

3. Synchrony

Although other methods can be performed on taxonomic subsets to look for similarities and

differences in the temporal dynamics among groups of taxa ("synchronicity"), a useful

measure of the 'temporal synchrony' in multivariate taxon abundances was developed by

Loreau & Mazancourt (2008) and comes with its own R package, 'synchrony' (Gouhier and

Guichard 2014). This standardised value measures how closely changes in taxon abundances

within a community track one another in time. It varies between zero (complete asynchrony

of taxon abundances in the community) and one (perfect synchrony; all taxon abundances

changing in the same way) and, by exploring different sized 'time windows', can be used to

detect scales of temporal variation. Overall significance tests of synchronicity in community

change can be obtained by performing null model analysis (e.g., Pedersen et al. 2017).

A similar method (Houlahan et al. 2007) uses the sum of the temporal variances and

pairwise covariances of all taxa in the community to distinguish if the community shows

'compensatory dynamics' over time. This method results in either a positive, negative or zero

community-level covariance value, indicating that taxa are increasing or decreasing in

synchrony (positive community covariance), some taxa are increasing, but this is

compensated for by decreases in other taxa (negative community covariance), or that no

change is evidence (zero community covariance) (Houlahan et al. 2007, Hallett et al. 2016).

4. Turnover Rates

The degree of temporal variation in community composition can be assessed by calculating

the turnover rate of the community (a.k.a. 'temporal turnover' or 'species turnover'), which is

a measure of the rate of change in taxonomic composition for a 'site' over time (e.g., Aguirre

et al. 2003). Turnover rates can be computed to measure the magnitude of compositional

change, but in a way that does not result in an understanding of the direction of change or the

taxa involved in compositional shifts. This method can detect when, and how rapidly,

temporal shifts occur in a temporal sequence to detect complex temporal patterns such as

perturbations, tipping points and pivot points (Fig. 17). Turnover rates are calculated using

combinations of colonisation, immigration, extinction, mortality, recruitment, and survival.

One of the commonly used methods is that of Diamond (1969) where the percentage turnover

rate, T, is calculated as $T = 100 \times (E + C) / (S1 + S2)$, where E is the number of taxa that went

extinct between two time points, C is the number of taxa that colonised between the same two

time points, $S1$ is the number of taxa present at time 1 and $S2$ is the number of taxa present at

time 2. T gives the percent change in taxon identities over the time period and varies from 0,

representing no turnover, to 1, indicating complete turnover (Diamond 1969). This measure

can be calculated only for a single location at two time points; pairwise turnovers must be

calculated separately for datasets with multiple time periods or multiple sites. T values can be

compared through time, at different time intervals, or at different spatial scales (e.g.,

Boulinier et al. 2001, Tworek 2004). They also can be used in a regression-style analysis to

assess the effects of a treatment variable or other explanatory factor on compositional

turnover (e.g., Aguirre et al. 2003, Tworek 2004). These methods are only useful for

communities that have relatively high rates of temporal turnover in taxa. The degree of

compositional turnover also is tied to the temporal extent of the study for the given system

and the average lifespan of taxa. For example, complete turnover in prokaryotes can be

observed over a period of days (e.g., Kara and Shade 2009), whilst many decades may be

required to detect any species turnover in some plant communities (e.g., Kardol et al. 2010).

However, because it can be standardised by the number of taxa in the community, T is a

potentially comparable measure of compositional change among different datasets, if they are

on similar timescales and encompass communities with a similar distribution of life-histories.

5. Multivariate regression modelling

Multivariate autoregressive state-space models (MARSS) are multivariate linear regression

methods that model the stochastic temporal dynamics in matrices of species' abundances in

relation to abiotic or biotic covariates, while accounting for potential observation

(measurement) error (Hampton et al. 2013). The models are autoregressive in that previous

measurements in time are assumed to be correlated to future measurement, and these

autocorrelations are formalised by modelling the current values of the response matrix as a

weighted linear sum of previous values at a specified temporal lag (Penny and Harrison

2006). MARSS models can be used to disentangle temporal variation in species abundances

into the relative contributions of inter- and intraspecific interactions, abiotic variations, and of

demographic stochasticity (Mutshinda et al. 2009). Effect sizes of these different components

are estimated from the multivariate model structure through the use of conditional least-

squares, maximum-likelihood, or Bayesian estimation methods (Hampton et al. 2013).

Observation errors, and other complexities such as spatially-replicated time series and non-

uniform sampling, can be formally modelled within the state-space framework (Holmes et al.

2012). Further, MARSS models can be used to compute several metrics that indicate the

degree to which communities are possibly becoming less stable over time (Ives et al. 2003).

This level of flexibility means that they are able to test for even the most complex temporal

patterns in communities, and can simultaneously reveal similarities and differences among

taxa, and account for environmental factors or other covariates. However, they do require

relatively long time series (ideally more than 25 temporal replicates), and, like any statistical

model, more independent replication is required for more complex models. There are several

examples of the use of MARSS models in freshwater and marine ecology, particularly in

terms of estimating aquatic food web dynamics (Beisner et al. 2003, Hampton et al. 2008,

Thibaut et al. 2012); there have been relatively few applications for terrestrial communities

(e.g., Yamamura et al. 2006, Mutshinda et al. 2009). The 'MARSS' R package (Holmes et al.

2012) enables the fitting of MARSS models and includes a range of ecological examples, so

we have not included example code in this paper.

Multispecies N mixture models and joint species distribution models are other

regression methods that can be used to model explicit, quantitative hypotheses of community

change, with a focus on interactions among taxa (e.g., Dorazio et al. 2015, Ovaskainen et al.

2017). These methods can allow for useful additions, such as accounting for uncertainty in

the detection of taxa; however, due to their complexity and high computational requirements,

they have yet to be implemented for datasets with large numbers of species.

STEP 3: TEST FOR STATISTICAL SIGNIFICANCE

Whichever multivariate analysis has been selected, most ecologists interested in community

dynamics seek a way to determine if the changes that they see over time in the results of

those analyses are greater than, or less than, they would expect by chance alone. Some of the

methods presented above have their own built-in significance tests, often based on

randomisations of the community matrix. Commonly-used randomisation tests include the

Mantel test (Legendre and Legendre 2012) and permutational multivariate analysis of

variance (PERMANOVA; Anderson 2001, Legendre and Legendre 2012). Null models,

whereby community matrices are randomised under sensible criteria (Gotelli and Graves

1996, Gotelli 2000, 2001, Gotelli and Entsminger 2001, Ulrich and Gotelli 2007, 2010), can

be used to test the significance of any observed change in compositional dissimilarity, and

can easily be incorporated into analyses described above including zeta diversity. The

primary consideration in applying any randomisation test is what assumptions are being made

about taxon niches and whether they are valid. If the interest is in testing the effects of

interactions among taxa on their co-occurrence, but the randomisation procedure (null model)

allows taxa to occur in samples where they are unable to occur for other reasons, such as

habitat differences, a difference among samples may be falsely detected (an inflated Type I

error rate). Thus, if strong inferences are to be made about compositional differences, we

must be sure that the differences among the groups of sites that we are testing are the only

difference. Similarly, the sample size needs to be high enough to allow the detection of

differences among sets of sites for the given level of compositional variability (Type II error

rate).

Schaefer, et al. (2005) introduced a novel method for detecting significant shifts in

community composition using randomisations under null models of taxon mean abundances

through time. They demonstrate that this approach, which uses both compositional distance

(pairwise dissimilarities) and the coefficient of variation (CV), could detect significant

dynamics in community composition, even where only small amounts of change had

occurred. In this method, temporal changes in communities are assessed by using a

compositional distance measure that is selected by the user. This quantifies the multivariate

distance between the randomly generated communities (null model randomisations) and

observed 'reference' samples (e.g., 'Time 0' samples, compared to randomisations of 'Time

1' samples). If the observed distance is greater than under randomisation, then we can infer

statistically significant temporal change. Because this can be done in an ordination context,

increases and decreases of particular taxa driving the change can be determined from the

ordination plots. Limitations of this method are that it requires a relatively large degree of

temporal change in the data compared to spatial variation (i.e., more temporal, than spatial,

variation) and it is relatively computationally intensive; however, we provide the R code for a

worked example in the supplementary material (Data S2).

STEP 4: REPORT ON THE ANALYSIS WORKFLOW

Our literature review suggested a lack of consistent reporting of critical aspects of the data

analysis workflow. Such reporting of ensures that the results are robust, represent the

variation deemed important for the research aim and are reproducible. The number of taxa,

samples, and independent replicates in space and time all should be clearly reported in the

Methods or Results section of the paper (see Hurlbert 1984 for guidance on statistical

independence). The dissimilarity measure selected, and the rationale and procedures used to

arrive at that decision, should be reported in the Methods section of the paper. We encourage

researchers to refer to a specific source, such as Legendre & Legendre (2012), or to give the

formula applied to maximise the repeatability and transparency of their research; these

indices can be inherent in particular methods or built-in as defaults in software, but

nevertheless should still be reported. Different methods of dissimilarity or ordination (or

other analysis types) that were compared, e.g., in the EDA phase, should be listed in the

Methods section of the paper stating that their results were consistent with each other. The

choice of dissimilarity or ordination method is subjective, but rules of thumb can be used for

assessing different methods, so that the method selected optimises the reduction of

dimensionality in the data and thus maximises the variation accounted for by the analysis (see

McCune and Grace 2002 for comprehensive advice).

STEP 5: PRESENT TEMPORAL PATTERNS AND RESULTS

Some of the analysis methods that we describe in this paper do not produce simple visual

outputs that clearly show temporal trends in community datasets and visualising temporal

dynamics can be difficult for species-rich datasets or complex sampling designs. However,

there are a variety of useful methods that can be used to complement a hypothesis-driven

analysis (Table 3). Many descriptive methods work best where a small group of taxa is

measured only a few times. However, there are examples of descriptive methods being used

effectively for larger, more complex examples. Pereira et al. (2017) used a heat map to show

compositional changes of 808 bacterial genera, at a single location, sampled 24 times over

two years. Certain descriptive methods are useful for only specific types of temporal patterns,

e.g., 'time cores' is a method that identifies 'core' groups of taxa that remain present over

time (Vallès et al. 2014). Others, such as Venn diagrams and measures of overlap in

composition, are useful only for small numbers of temporal comparisons. We recommend

and provide R code (Data S2) for time cores and Venn diagrams because these methods are

suitable for assessing hypotheses of taxon replacement while specifically identifying the taxa

driving compositional shifts (Tables 2 and 3). Relative abundance distributions (RADs, *a.k.a.*

'rank abundance curves'), can provide relatively sophisticated visualisations of community

dynamics, and rank abundances can be used to quantitatively measure the shift in taxon ranks

within samples taken at different time points (Hallett et al. 2016). RADs are plots of the

relative abundances (on the *y*-axis) of all species in a community ranked (on the *x*-axis) from

highest to lowest. They are closely related to *k*-dominance curves, in which species ranks are

shown on the *x*-axis, but cumulative proportional abundance (dominance) is plotted on the *y*-

axis (Lambshead et al. 1983). If samples taken at different times are plotted separately, the

set of RADs can illustrate changes in community structure over time. RADs also can be used

to explore community dynamics if a dissimilarity measure, such as Bray-Curtis, is calculated

for the species' ranks at each time point. The resulting temporal dissimilarity of the RADs

can be explored by plotting rank dissimilarity against time lag or by ordinating rank

dissimilarities in the same way that spatial compositional dissimilarities are examined. For

example, temporal changes in RAD dissimilarity can illustrate changes in species evenness

over time as might be expected if compositional homogenisation and a decline in evenness

occurs because of invasion by an increasingly dominant species (Ruhl 2008).

E. DISCUSSION

There is an ever-expanding set of analyses that use a wide variety of multivariate and

univariate methods to quantitatively address temporal questions about communities that

continue to be developed (e.g., De Cáceres et al. 2019). One of the key reasons for the

proliferation of methods appears to be variability in the length of the time series. For

example, if the time series is long enough, methods such as temporal MEMs could be used;

however, consistent with other reviews (e.g., Vaughn and Young 2010) and databases

(Dornelas et al. 2018), most datasets captured by our literature had relatively few temporal

measurements (mean = 14.9, median = 7, mode = 4), requiring very different methods. In

other cases, the ability to implement some methods in temporal community ecology still

requires development, such as the application of time-ordered and time-aggregated network

analysis for spatiotemporal community datasets (Blonder et al. 2012).

Some analysis types are well established for spatial data, but can be applied easily to

temporal data, potentially yielding useful and comparable results for explicit temporal

analyses. Further research is required to generate standard extensions and approaches using

such methods for temporal data. For example, nestedness (*sensu* Patterson and Atmar 1986)

is usually applied to spatial data and the analysis is repeated for different time points;

however, there is no reason that we cannot apply this method across times to look at whether

sites are becoming more or less nested over time. For instance, if younger sites were nested

within older sites, this may indicate that sites were becoming more homogeneous over time.

Our understanding of temporal dynamics in biological communities, and the methods

we use to investigate them, have improved greatly over the past 25 years, probably because

of increases in computational power and in dataset size. We can continue to improve our

understanding by moving away from merely describing community changes, to linking these

changes to understanding and testing the processes driving community change and

biodiversity; that is, moving from hypothesis-generating methods towards hypothesis-testing

methods enabled by well-defined research questions with specific, testable hypotheses that

result in clear predictions for observed temporal patterns (e.g., Figs 4-7). Even for

exploratory studies of compositional variation through time, we emphasise the benefits of

making *a priori* predictions for how we think composition should change through time.

Interactions between community composition and environmental or other factors

across space and time are complicated. For example, Brown (2003) showed that temporal

variability in species composition of stream invertebrates was reduced by high spatial

heterogeneity, which may be because there are increased niches or refugia in changing

environmental conditions. Manipulative experiments are useful, but not always possible,

especially for long-lived taxa. Therefore, combining current knowledge of intrinsic and

extrinsic factors controlling community change into null models and simulations may be the

way forward (Brown and Lawson 2010). Carefully considering sample sizes and appropriate

replication of particular factors are also important if we are to properly test hypotheses of

species coexistence over time (Philippi et al. 1998). There is a need for us to make useful

predictions about how communities may continue to respond to ecosystem disturbances, such

as climate change, invasive species, and habitat fragmentation (Hobbs et al. 2007,

Kampichler et al. 2012). Many methods commonly used to analyse temporal community

dynamics are not quantitative, meaning conclusions can only be narrative descriptions, rather

than clear tests of hypotheses. Many of the methods presented here push us to quantitatively

assess dynamics, which naturally leads to clearer hypothesis testing.

There is a wide variety of methods available for analysing temporal dynamics of

biological communities; however, many studies in our literature review using these methods

are descriptive, showing that a large body of literature in community ecology is not focussed

on testing quantitative hypotheses of change. Surprisingly few studies in our review

presented hypotheses that were more complex than simply predicting that composition would

have changed in some way over time, instead of specifying predictions of quantitative and

directional patterns of change in community composition. Even additional searching beyond

our original search terms that included more specific terminology for compositional change

(succession, turnover, synchron*, resilience, homogenization, divergence, convergence)

revealed few empirical papers that tested specific, quantitative hypotheses. This search

resulted in an additional 1,514 papers of which we read 10% (151 papers). Of these, 60

papers met the inclusion criteria of our original review and only 13 papers tested hypotheses

that were more sophisticated than the simple prediction that composition would have changed

in some way over time, e.g., directional change or that the abundances of some taxa would

change more than others.

This lack of specific, quantitative hypothesis testing is not surprising because to

predict change in composition in a quantitative way, we require an in-depth understanding of

a system and how different components of the diversity differ in their ecology or other factors

that could cause directional, cyclical, or other differential change in time. In many cases,

particularly in the observational, rather than experimental, community ecology literature, this

is not the situation. This means that, although such studies may state hypotheses, these

hypotheses are not very specific or quantitative. In most cases, they predict some change in

response to time or another factor that is correlated with time (such as disturbance or

pollution), but the types of temporal change (divergence, convergence, synchrony, etc.) are

not specified. We strongly recommend that researchers generate *a priori* hypotheses of

compositional change (e.g., Figs. 4-7; Bagchi et al. 2017). These predictions should be used

to select appropriate quantitative data analysis methods (Fig. 17). The comprehensive

description and hypothesis testing of temporal community patterns requires the use of

multiple data analysis methods, but by using standard exploratory data analysis methods

followed by testing *a priori* hypotheses, the methods selected can be explicitly focussed on

the most appropriate tests for the situation. We hope that this synthesis will allow researchers

to make informed decisions about their choice of data analysis for their research question and

pave the way for more uniformity in reporting of these analyses and/or innovation in

comparability. Our understanding of patterns and drivers of changes in species communities

would benefit by having more longer-term and fine-scale studies.

ACKNOWLEDGEMENTS

We thank Audrey Lustig for initial database and R code development. This paper is a

contribution from the Harvard Forest LTER program. We thank members of the Forest

Ecology Research Group at Wilfrid Laurier University, Glenda Mendieta-Leiva, Brian

Inouye, Miguel De Cáceres, and an anonymous reviewer for comments on the manuscript.

We thank the authors of Porensky et al. (2016a, 2016b) and Gilbert et al. (2012) for making

their data publicly available. The BCI forest dynamics research project was founded by S.P.

Hubbell and R.B. Foster and is now managed by R. Condit, S. Lao, and R. Perez under the

Center for Tropical Forest Science and the Smithsonian Tropical Research in Panama.

Numerous organizations have provided funding, principally the U.S. National Science

Foundation, and hundreds of field workers have contributed.

FUNDING DISCLOSURE

H. L. Buckley and B. S. Case were supported by a Charles Bullard Fellowship at Harvard

Forest. A. M. Ellison's work on this project was supported by the Harvard Forest.

SUPPORTING INFORMATION

Data S1. Description of attributes, and associated data, extracted from papers in the literature

review.

Data S2. Example R code for plant community dataset

Data S3. Example R code for BCI dataset

Data S4. Example R code for marine bacterial community dataset (from high throughput

sequencing)

Appendix S1. Supplementary figures from the literature review.

Appendix S2. Descriptions of methods encountered in the literature review that are not

included in the main text.

1199

REFERENCES CITED

Aguirre, L. F., L. Lens, R. van Damme, and E. Matthysen. 2003. Consistency and variation in

the bat assemblages inhabiting two forest islands within a neotropical savanna in

Bolivia. Journal of Tropical Ecology 19:367–374.

Amend, A. S., K. A. Seifert, and T. D. Bruns. 2010. Quantifying microbial communities with

454 pyrosequencing: does read abundance count? Molecular Ecology 19:5555–5565.

Anderson, M. J. 2001. A new method for non-parametric multivariate analysis of variance.

Austral Ecology 26:32–46.

Anderson, M. J., T. O. Crist, J. M. Chase, M. Vellend, B. D. Inouye, A. L. Freestone, N. J.

Sanders, H. V. Cornell, L. S. Comita, K. F. Davies, S. P. Harrison, N. J. B. Kraft, J. C.

Stegen, and N. G. Swenson. 2011. Navigating the multiple meanings of β diversity: A

roadmap for the practicing ecologist. Ecology Letters 14:19–28.

Anderson, M. J., and T. J. Willis. 2003. Canonical analysis of principal coordinates: A useful

method of constrained ordination for ecology. Ecology 84:511–525.

Angeler, D. G., and R. K. Johnson. 2012. Patterns of temporal community turnover are

spatially synchronous across boreal lakes: Temporal coherence in invertebrate

community turnover. Freshwater Biology 57:1782–1793.

Angeler, D. G., O. Viedma, and J. M. Moreno. 2009. Statistical performance and information

content of time lag analysis and redundancy analysis in time series modeling. Ecology

90:3245–3257.

Auber, A., M. Travers-Trolet, M. C. Villanueva, and B. Ernande. 2017. A new application of

principal response curves for summarizing abrupt and cyclic shifts of communities

over space. Ecosphere 8:e02023.

Azeria, E. T., A. Carlson, T. Pärt, and C. G. Wiklund. 2006. Temporal dynamics and

nestedness of an oceanic island bird fauna. Global Ecology and Biogeography

15:328–338.

Azeria, E. T., and J. Kolasa. 2008. Nestedness, niche metrics and temporal dynamics of a

metacommunity in a dynamic natural model system. Oikos 117:1006–1019.

Bagchi, S., N. J. Singh, D. D. Briske, B. T. Bestelmeyer, M. P. McClaran, and K. Murthy.

2017. Quantifying long-term plant community dynamics with movement models:

implications for ecological resilience. Ecological Applications 27:1514–1528.

Bar-Massada, A., and L. Hadar. 2017. Grazing and temporal turnover in herbaceous

communities in a Mediterranean landscape. Journal of Vegetation Science 28:270–

280.

Barros, F., H. Blanchet, K. Hammerstrom, P.-G. Sauriau, and J. Oliver. 2014. A framework

for investigating general patterns of benthic β-diversity along estuaries. Estuarine,

Coastal and Shelf Science 149:223–231.

Baselga, A. 2010. Partitioning the turnover and nestedness components of beta diversity:

Partitioning beta diversity. Global Ecology and Biogeography 19:134–143.

Baselga, A. 2012. The relationship between species replacement, dissimilarity derived from

nestedness, and nestedness: Species replacement and nestedness. Global Ecology and

Biogeography 21:1223–1232.

Baselga, A. 2017. Partitioning abundance-based multiple-site dissimilarity into components:

Balanced variation in abundance and abundance gradients. Methods in Ecology and

Evolution 8:799–808.

Baselga, A., and C. D. L. Orme. 2012. betapart : an R package for the study of beta diversity:

Betapart package. Methods in Ecology and Evolution 3:808–812.

Bashan, A., T. E. Gibson, J. Friedman, V. J. Carey, S. T. Weiss, E. L. Hohmann, and Y.-Y.

Liu. 2016. Universality of human microbial dynamics. Nature 534:259–262.

Bêche, L. A., and V. H. Resh. 2007. Short-term climatic trends affect the temporal variability

of macroinvertebrates in California 'Mediterranean' streams. Freshwater Biology

52:2317–2339.

Beisner, B. E., A. R. Ives, and S. R. Carpenter. 2003. The effects of an exotic fish invasion on

the prey communities of two lakes. Journal of Animal Ecology 72:331–342.

Berthon, V., B. Alric, F. Rimet, and M.-E. Perga. 2014. Sensitivity and responses of diatoms

to climate warming in lakes heavily influenced by humans. Freshwater Biology

59:1755–1767.

Bestelmeyer, B. T., A. M. Ellison, W. R. Fraser, K. B. Gorman, S. J. Holbrook, C. M. Laney,

M. D. Ohman, D. P. C. Peters, F. C. Pillsbury, A. Rassweiler, R. J. Schmitt, and S.

Sharma. 2011. Analysis of abrupt transitions in ecological systems. Ecosphere

2:art129.

Blonder, B., T. W. Wey, A. Dornhaus, R. James, and A. Sih. 2012. Temporal dynamics and

network analysis. Methods in Ecology and Evolution 3:958–972.

Borcard, D., F. Gillet, and P. Legendre. 2011. Numerical Ecology with R. Springer, New

York.

Borcard, D., and P. Legendre. 2002. All-scale spatial analysis of ecological data by means of

principal coordinates of neighbour matrices. Ecological Modelling 153:51–68.

Boulinier, T., J. D. Nichols, J. E. Hines, J. R. Sauer, C. H. Flather, and K. H. Pollock. 2001.

Forest fragmentation and bird community dynamics: Inference at regional scales.

Ecology 82:1159–1169.

ter Braak, C. J. F., and P. Šmilauer. 2015. Topics in constrained and unconstrained

ordination. Plant Ecology 216:683–696.

Broadway, K. J., M. Pyron, J. R. Gammon, and B. A. Murry. 2015. Shift in a large river fish

assemblage: Body-size and trophic structure dynamics. PLOS ONE 10:e0124954.

Brown, B. L. 2003. Spatial heterogeneity reduces temporal variability in stream insect

communities. Ecology Letters 6:316–325.

Brown, B. L., and R. L. Lawson. 2010. Habitat heterogeneity and activity of an omnivorous

ecosystem engineer control stream community dynamics. Ecology 91:1799–1810.

Caruso, T., V. Trokhymets, R. Bargagli, and P. Convey. 2013. Biotic interactions as a

structuring force in soil communities: Evidence from the micro-arthropods of an

Antarctic moss model system. Oecologia 172:495–503.

Chao, A., R. L. Chazdon, R. K. Colwell, and T.-J. Shen. 2006. Abundance-based similarity

indices and their estimation when there are unseen species in samples. Biometrics

62:361–371.

Chao, A., N. J. Gotelli, T. C. Hsieh, E. L. Sander, K. H. Ma, R. K. Colwell, and A. M.

Ellison. 2014. Rarefaction and extrapolation with Hill numbers: a framework for

sampling and estimation in species diversity studies. Ecological Monographs 84:45–

67.

Chapman, J. I., and R. W. McEwan. 2016. Thirty years of compositional change in an old-

growth temperate forest: The role of topographic gradients in oak-maple dynamics.

PLOS ONE 11:e0160238.

Clavero, M., and E. Garcia-Berthou. 2006. Homogenization dynamics and introduction

routes of invasive freshwater fish in the Iberian Peninsula. Ecological Applications

16:2313–2324.

Cleland, E. E., S. L. Collins, T. L. Dickson, E. C. Farrer, K. L. Gross, L. A. Gherardi, L. M.

Hallett, R. J. Hobbs, J. S. Hsu, L. Turnbull, and K. N. Suding. 2013. Sensitivity of

grassland plant community composition to spatial vs. temporal variation in

precipitation. Ecology 94:1687–1696.

Collins, S. L. 1992. Fire frequency and community heterogeneity in tallgrass prairie

vegetation. Ecology 73:2001–2006.

Collins, S. L. 2000. Disturbance frequency and community stability in native tallgrass prairie.

The American Naturalist 155:311–325.

Collins, S. L., F. Micheli, and L. Hartt. 2000. A method to determine rates and patterns of

variability in ecological communities. Oikos 91:285–293.

Collins, S. L., and M. D. Smith. 2006. Scale-dependent interaction of fire and grazing on

community heterogeneity in tallgrass prairie. Ecology 87:2058–2067.

Collins, S. L., and Y. Xia. 2015. Long-term dynamics and hotspots of change in a desert

grassland plant community. The American Naturalist 185:E30–E43.

Condit, R. 1998. Tropical Forest Census Plots. Springer-Verlag and R. G. Landes Company,

Berlin, Germany, and Georgetown, Texas.

Connell, J. H., and R. O. Slatyer. 1977. Mechanisms of succession in natural communities

and their role in community stability and organization. The American Naturalist

111:1119–1144.

Cotton, T. E. A., A. H. Fitter, R. M. Miller, A. J. Dumbrell, and T. Helgason. 2015. Fungi in

the future: interannual variation and effects of atmospheric change on arbuscular

mycorrhizal fungal communities. New Phytologist 205:1598–1607.

Day, N. J., and H. L. Buckley. 2013. Twenty-five years of plant community dynamics and

invasion in New Zealand tussock grasslands. Austral Ecology 38:688–699.

Day, N. J., S. Carrière, and J. L. Baltzer. 2017. Annual dynamics and resilience in post-fire

boreal understory vascular plant communities. Forest Ecology and Management

401:264–272.

De Cáceres, M., L. Coll, P. Legendre, R. B. Allen, S. K. Wiser, M.-J. Fortin, R. Condit, and

S. Hubbell. 2019. Trajectory analysis in community ecology. Ecological

Monographs:e01350.

De Cáceres, M., X. Font, and F. Oliva. 2010. The management of vegetation classifications

with fuzzy clustering: Fuzzy clustering in vegetation classifications. Journal of

Vegetation Science 21:1138–1151.

Diamond, J. M. 1969. Avifaunal equilibria and species turnover rates on the Channel Islands

of California. Proceedings of the National Academy of Sciences 64:57–63.

Dimitriu, P. A., G. Boyce, A. Samarakoon, M. Hartmann, P. Johnson, and W. W. Mohn.

2013. Temporal stability of the mouse gut microbiota in relation to innate and

adaptive immunity: Mouse gut microbiota dynamics. Environmental Microbiology

Reports 5:200–210.

Dorazio, R. M., E. F. Connor, and R. A. Askins. 2015. Estimating the Effects of Habitat and

Biological Interactions in an Avian Community. PLOS ONE 10:e0135987.

Dornelas, M., L. H. Antão, F. Moyes, A. E. Bates, A. E. Magurran, D. Adam, A. A.

Akhmetzhanova, W. Appeltans, J. M. Arcos, H. Arnold, N. Ayyappan, G. Badihi, A.

H. Baird, M. Barbosa, T. E. Barreto, C. Bässler, A. Bellgrove, J. Belmaker, L.

Benedetti-Cecchi, B. J. Bett, A. D. Bjorkman, M. Błażewicz, S. A. Blowes, C. P.

Bloch, T. C. Bonebrake, S. Boyd, M. Bradford, A. J. Brooks, J. H. Brown, H.

Bruelheide, P. Budy, F. Carvalho, E. Castañeda-Moya, C. A. Chen, J. F. Chamblee, T.

J. Chase, L. Siegwart Collier, S. K. Collinge, R. Condit, E. J. Cooper, J. H. C.

Cornelissen, U. Cotano, S. Kyle Crow, G. Damasceno, C. H. Davies, R. A. Davis, F.

P. Day, S. Degraer, T. S. Doherty, T. E. Dunn, G. Durigan, J. E. Duffy, D. Edelist, G.

J. Edgar, R. Elahi, S. C. Elmendorf, A. Enemar, S. K. M. Ernest, R. Escribano, M.

Estiarte, B. S. Evans, T.-Y. Fan, F. Turini Farah, L. Loureiro Fernandes, F. Z.

Farneda, A. Fidelis, R. Fitt, A. M. Fosaa, G. A. Daher Correa Franco, G. E. Frank, W.

R. Fraser, H. García, R. Cazzolla Gatti, O. Givan, E. Gorgone-Barbosa, W. A. Gould,

C. Gries, G. D. Grossman, J. R. Gutierréz, S. Hale, M. E. Harmon, J. Harte, G.

Haskins, D. L. Henshaw, L. Hermanutz, P. Hidalgo, P. Higuchi, A. Hoey, G. Van

Hoey, A. Hofgaard, K. Holeck, R. D. Hollister, R. Holmes, M. Hoogenboom, C.

Hsieh, S. P. Hubbell, F. Huettmann, C. L. Huffard, A. H. Hurlbert, N. Macedo

Ivanauskas, D. Janík, U. Jandt, A. Jażdżewska, T. Johannessen, J. Johnstone, J. Jones,

F. A. M. Jones, J. Kang, T. Kartawijaya, E. C. Keeley, D. A. Kelt, R. Kinnear, K.

Klanderud, H. Knutsen, C. C. Koenig, A. R. Kortz, K. Král, L. A. Kuhnz, C.-Y. Kuo,

D. J. Kushner, C. Laguionie-Marchais, L. T. Lancaster, C. Min Lee, J. S. Lefcheck, E.

Lévesque, D. Lightfoot, F. Lloret, J. D. Lloyd, A. López-Baucells, M. Louzao, J. S.

Madin, B. Magnússon, S. Malamud, I. Matthews, K. P. McFarland, B. McGill, D.

McKnight, W. O. McLarney, J. Meador, P. L. Meserve, D. J. Metcalfe, C. F. J.

Meyer, A. Michelsen, N. Milchakova, T. Moens, E. Moland, J. Moore, C. Mathias

Moreira, J. Müller, G. Murphy, I. H. Myers-Smith, R. W. Myster, A. Naumov, F.

Neat, J. A. Nelson, M. Paul Nelson, S. F. Newton, N. Norden, J. C. Oliver, E. M.

Olsen, V. G. Onipchenko, K. Pabis, R. J. Pabst, A. Paquette, S. Pardede, D. M.

Paterson, R. Pélissier, J. Peñuelas, A. Pérez-Matus, O. Pizarro, F. Pomati, E. Post, H.

H. T. Prins, J. C. Priscu, P. Provoost, K. L. Prudic, E. Pulliainen, B. R. Ramesh, O.

Mendivil Ramos, A. Rassweiler, J. E. Rebelo, D. C. Reed, P. B. Reich, S. M.

Remillard, A. J. Richardson, J. P. Richardson, I. van Rijn, R. Rocha, V. H. Rivera-

Monroy, C. Rixen, K. P. Robinson, R. Ribeiro Rodrigues, D. de Cerqueira Rossa-

Feres, L. Rudstam, H. Ruhl, C. S. Ruz, E. M. Sampaio, N. Rybicki, A. Rypel, S. Sal,

B. Salgado, F. A. M. Santos, A. P. Savassi-Coutinho, S. Scanga, J. Schmidt, R.

Schooley, F. Setiawan, K.-T. Shao, G. R. Shaver, S. Sherman, T. W. Sherry, J.

Siciński, C. Sievers, A. C. da Silva, F. Rodrigues da Silva, F. L. Silveira, J. Slingsby,

T. Smart, S. J. Snell, N. A. Soudzilovskaia, G. B. G. Souza, F. Maluf Souza, V.

Castro Souza, C. D. Stallings, R. Stanforth, E. H. Stanley, J. Mauro Sterza, M.

Stevens, R. Stuart-Smith, Y. Rondon Suarez, S. Supp, J. Yoshio Tamashiro, S.

Tarigan, G. P. Thiede, S. Thorn, A. Tolvanen, M. Teresa Zugliani Toniato, Ø.

Totland, R. R. Twilley, G. Vaitkus, N. Valdivia, M. I. Vallejo, T. J. Valone, C. Van

Colen, J. Vanaverbeke, F. Venturoli, H. M. Verheye, M. Vianna, R. P. Vieira, T.

Vrška, C. Quang Vu, L. Van Vu, R. B. Waide, C. Waldock, D. Watts, S. Webb, T.

Wesołowski, E. P. White, C. E. Widdicombe, D. Wilgers, R. Williams, S. B.

Williams, M. Williamson, M. R. Willig, T. J. Willis, S. Wipf, K. D. Woods, E. J.

Woehler, K. Zawada, and M. L. Zettler. 2018. BioTIME: A database of biodiversity

time series for the Anthropocene. Global Ecology and Biogeography 27:760–786.

Dray, S., P. Legendre, and P. R. Peres-Neto. 2006. Spatial modelling: a comprehensive

framework for principal coordinate analysis of neighbour matrices (PCNM).

Ecological Modelling 196:483–493.

Ellison, A. 2010. Partitioning diversity. Ecology 91:1962–1963.

Ferrier, S., G. Manion, J. Elith, and K. Richardson. 2007. Using generalized dissimilarity

modelling to analyse and predict patterns of beta diversity in regional biodiversity

assessment. Diversity and Distributions 13:252–264.

Franca, A., S. Caredda, F. Sanna, F. Fava, and G. Seddaiu. 2016. Early plant community

dynamics following overseeding for the rehabilitation of a Mediterranean

silvopastoral system. Grassland Science 62:81–91.

Gaston, K. J., and B. H. McArdle. 1994. The temporal variability of animal abundances:

measures, methods and patterns. Philosophical Transactions of the Royal Society B:

Biological Sciences 345:335–358.

Gilbert, J. A., J. A. Steele, J. G. Caporaso, L. Steinbrück, J. Reeder, B. Temperton, S. Huse,

A. C. McHardy, R. Knight, and I. Joint. 2012. Defining seasonal marine microbial

community dynamics. The ISME journal 6:298.

Gotelli, N. J. 2000. Null model analysis of species co-occurrence patterns. Ecology 81:2606–

2621.

Gotelli, N. J. 2001. Research frontiers in null model analysis. Global Ecology and

Biogeography 10:337–343.

Gotelli, N. J., and G. L. Entsminger. 2001. Swap and fill algorithms in null model analysis:

rethinking the knight's tour. Oecologia 129:281–291.

Gotelli, N. J., and G. R. Graves. 1996. Null Models in Ecology. Smithsonian Institution

Press, Washington and London.

Gouhier, T. C., and F. Guichard. 2014. Synchrony: Quantifying variability in space and time.

Methods in Ecology and Evolution 5:524–533.

Hallett, L. M., S. K. Jones, A. A. M. MacDonald, M. B. Jones, D. F. B. Flynn, J. Ripplinger,

P. Slaughter, C. Gries, and S. L. Collins. 2016. codyn: An R package of community

dynamics metrics. Methods in Ecology and Evolution 7:1146–1151.

Hampton, S. E., E. E. Holmes, L. P. Scheef, M. D. Scheuerell, S. L. Katz, D. E. Pendleton,

and E. J. Ward. 2013. Quantifying effects of abiotic and biotic drivers on community

dynamics with multivariate autoregressive (MAR) models. Ecology 94:2663–2669.

Hampton, S. E., L. R. Izmest'Eva, M. V. Moore, S. L. Katz, B. Dennis, and E. A. Silow.

2008. Sixty years of environmental change in the world's largest freshwater lake -

Lake Baikal, Siberia. Global Change Biology 14:1947–1958.

Hobbs, R. J., S. Yates, and H. A. Mooney. 2007. Long-term data reveal complex dynamics in

grassland in relation to climate and disturbance. Ecological Monographs 77:545–568.

Holmes, E. E., E. J. Ward, and K. Wills. 2012. MARSS: Multivariate Autoregressive State-

space models for analyzing time-series data. R Journal 4:11–19.

Houlahan, J. E., D. J. Currie, K. Cottenie, G. S. Cumming, S. K. M. Ernest, C. S. Findlay, S.

D. Fuhlendorf, U. Gaedke, P. Legendre, J. J. Magnuson, B. H. McArdle, E. H.

Muldavin, D. Noble, R. Russell, R. D. Stevens, T. J. Willis, I. P. Woiwod, and S. M.

Wondzell. 2007. Compensatory dynamics are rare in natural ecological communities.

Proceedings of the National Academy of Sciences 104:3273–3277.

Hubbell, S. P. 1979. Tree dispersion, abundance, and diversity in a tropical dry forest.

Science 203:1299–1309.

Hubbell, S. P., R. Condit, and R. B. Foster. 2005. Barro Colorado Forest Census Plot Data.

Hubbell, S. P., R. B. Foster, S. T. O'Brien, K. E. Harms, R. Condit, B. Wechsler, S. J.

Wright, and S. Loo de Lao. 1999. Light-gap disturbances, recruitment limitation, and

tree diversity in a neotropical forest. Science 283:554–557.

Hui, C., and M. A. McGeoch. 2014. Zeta diversity as a concept and metric that unifies

incidence-based biodiversity patterns. The American Naturalist 184:684–694.

Hurlbert, S. H. 1984. Pseudoreplication and the design of ecological field experiments.

Ecological Monographs 54:187–211.

Ives, A. R., B. Dennis, K. L. Cottingham, and S. R. Carpenter. 2003. Estimating community

stability and ecological interactions from time-series data. Ecological monographs

73:301–330.

Jackson, D. A. 1995. PROTEST: A PROcrustean Randomization TEST of community

environment concordance. Écoscience 2:297–303.

Johnstone, I. M., and D. M. Titterington. 2009. Statistical challenges of high-dimensional

data. Philosophical Transactions of the Royal Society A: Mathematical, Physical and

Engineering Sciences 367:4237–4253.

Kalyuzhny, M., and N. M. Shnerb. 2017. Dissimilarity-overlap analysis of community

dynamics: Opportunities and pitfalls. Methods in Ecology and Evolution 8:1764–

1773.

Kampichler, C., D. G. Angeler, R. T. Holmes, A. Leito, S. Svensson, H. P. van der Jeugd,

and T. Wesołowski. 2014. Temporal dynamics of bird community composition: An

analysis of baseline conditions from long-term data. Oecologia 175:1301–1313.

Kampichler, C., C. A. M. van Turnhout, V. Devictor, and H. P. van der Jeugd. 2012. Large-

scale changes in community composition: Determining land use and climate change

signals. PLoS ONE 7:e35272.

Kara, E., and A. Shade. 2009. Temporal dynamics of South End Tidal Creek (Sapelo Island,

Georgia) bacterial communities. Applied and Environmental Microbiology 75:1058–

1064.

Kardol, P., D. E. Todd, P. J. Hanson, and P. J. Mulholland. 2010. Long-term successional

forest dynamics: species and community responses to climatic variability. Journal of

Vegetation Science.

Kent, M. 2012. Vegetation Description and Data Analysis: A practical approach. Second

Edition. Wiley-Blackwell, Chichester, UK.

Kidé, S. O., C. Manté, L. Dubroca, H. Demarcq, and B. Mérigot. 2015. Spatio-temporal

dynamics of exploited groundfish species assemblages faced to environmental and

fishing forcings: insights from the Mauritanian exclusive economic zone. PloS one

10:e0141566.

Lambshead, P. J. D., H. M. Platt, and K. M. Shaw. 1983. The detection of differences among

assemblages of marine benthic species based on an assessment of dominance and

diversity. Journal of Natural History 17:859–874.

Lamy, T., P. Legendre, Y. Chancerelle, G. Siu, and J. Claudet. 2015. Understanding the

spatio-temporal response of coral reef fish communities to natural disturbances:

Insights from beta-diversity decomposition. PLOS ONE 10:e0138696.

Latombe, G., C. Hui, and M. A. McGeoch. 2017a. Multi-site generalised dissimilarity

modelling: using zeta diversity to differentiate drivers of turnover in rare and

widespread species. Methods in Ecology and Evolution 8:431–442.

Latombe, G., M. A. McGeoch, D. A. Nipperess, and C. Hui. 2017b. zetadiv: Functions to

Compute Compositional Turnover Using Zeta Diversity. R package version 1.0.

https://CRAN.R-project.org/package=zetadiv.

Legendre, P., and M. De Cáceres. 2013. Beta diversity as the variance of community data:

dissimilarity coefficients and partitioning. Ecology Letters 16:951–963.

Legendre, P., and E. Gallagher. 2001. Ecologically meaningful transformations for ordination

of species data. Oecologia 129:271–280.

Legendre, P., and O. Gauthier. 2014. Statistical methods for temporal and space-time analysis

of community composition data. Proceedings of the Royal Society B: Biological

Sciences 281:20132728–20132728.

Legendre, P., and L. F. J. Legendre. 2012. Numerical Ecology. Third Edition. Elsevier,

Radarweg, The Netherlands.

Lellouch, L., S. Pavoine, F. Jiguet, H. Glotin, and J. Sueur. 2014. Monitoring temporal

change of bird communities with dissimilarity acoustic indices. Methods in Ecology

and Evolution 5:495–505.

Lepš, J., and P. Šmilauer. 2003. Multivariate Analysis of Ecological Data Using CANOCO.

Cambridge University Press, Cambridge.

Lewis, S. L., and M. A. Maslin. 2015. Defining the Anthropocene. Nature 519:171–180.

Loreau, M., and C. de Mazancourt. 2008. Species synchrony and its drivers: neutral and

nonneutral community dynamics in fluctuating environments. The American

Naturalist 172:E48–E66.

Lozupone, C. A., M. Hamady, S. T. Kelley, and R. Knight. 2007. Quantitative and qualitative

diversity measures lead to different insights into factors that structure microbial

communities. Applied and Environmental Microbiology 73:1576–1585.

Lozupone, C., and R. Knight. 2005. UniFrac: a new phylogenetic method for comparing

microbial communities. Applied and Environmental Microbiology 71:8228–8235.

Magalhães, M. F., P. Beja, I. J. Schlosser, and M. J. Collares-Pereira. 2007. Effects of multi-

year droughts on fish assemblages of seasonally drying Mediterranean streams.

Freshwater Biology 52:1494–1510.

Magurran, A. E. 2004. Measuring Biological Diversity. Blackwell, Malden.

Magurran, A. E. 2013. Measuring Biological Diversity. John Wiley and Sons, Oxford.

McCune, B., and J. B. Grace. 2002. Analysis of Ecological Communities. MjM Software

Design, Gleneden Beach.

McGeoch, M. A., G. Latombe, N. R. Andrew, S. Nakagawa, D. A. Nipperess, M. Roige, E. .

Marzinelli, A. H. Campbell, A. Vergés, T. Thomas, P. D. Steinberg, K. E. Selwood,

and C. Hui. 2017. The application of zeta diversity as a continuous measure of

compositional change in ecology. bioRxiv 216580.

Micheli, F., K. L. Cottingham, J. Bascompte, O. N. Bjornstad, G. L. Eckert, J. M. Fischer, T.

H. Keitt, B. E. Kendall, J. L. Klug, and J. A. Rusak. 1999. The dual nature of

community variability. Oikos 85:161–169.

Mutshinda, C. M., R. B. O'Hara, and I. P. Woiwod. 2009. What drives community

dynamics? Proceedings of the Royal Society B: Biological Sciences 276:2923–2929.

Noll, M., D. Matthies, P. Frenzel, M. Derakshani, and W. Liesack. 2005. Succession of

bacterial community structure and diversity in a paddy soil oxygen gradient.

Environmental Microbiology 7:382–395.

Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin,

R. B. O'Hara, G. L. Simpson, P. Solymos, M. H. H. Stevens, E. Szoecs, and H.

Wagner. 2017. vegan: Community Ecology Package. R package version 2.4-2.

http://CRAN.R-project.org/package=vegan.

O'Sullivan, D. J., P. D. Cotter, O. O'Sullivan, L. Giblin, P. L. H. McSweeney, and J. J.

Sheehan. 2015. Temporal and spatial differences in microbial composition during the

manufacture of a Continental-type cheese. Applied and Environmental Microbiology

81:2525–2533.

Ovaskainen, O., G. Tikhonov, D. Dunson, V. Grøtan, S. Engen, B.-E. Sæther, and N. Abrego.

2017. How are species interactions structured in species-rich communities? A new

method for analysing time-series data. Proceedings of the Royal Society B: Biological

Sciences 284:20170768.

Palmer, M. 2019. Ordination Methods for Ecologists. http://ordination.okstate.edu/.

Patterson, B. D., and W. Atmar. 1986. Nested subsets and the structure of insular mammalian

faunas and archipelagos. Biological journal of the Linnean society 28:65–82.

Pedersen, E. J., P. L. Thompson, R. A. Ball, M.-J. Fortin, T. C. Gouhier, H. Link, C. Moritz,

H. Nenzen, R. R. E. Stanley, Z. E. Taranu, A. Gonzalez, F. Guichard, and P. Pepin.

2017. Signatures of the collapse and incipient recovery of an overexploited marine

ecosystem. Royal Society Open Science 4:170215.

Penny, W., and L. Harrison. 2006. Multivariate autoregressive models. Statistical Parametric

Mapping: The Analysis of Functional Brain Images, eds K. J. Friston, JT Ashburner,

SJ Kiebel, TE Nichols, and WD Penny (London: Elsevier):534–540.

Pereira, R. P. A., J. Peplies, M. G. Höfle, and I. Brettar. 2017. Bacterial community dynamics

in a cooling tower with emphasis on pathogenic bacteria and Legionella species using

universal and genus-specific deep sequencing. Water Research 122:363–376.

Peres-Neto, P. R., and D. A. Jackson. 2001. How well do multivariate data sets match? The

advantages of a Procrustean superimposition approach over the Mantel test.

Oecologia 129:169–178.

Philippi, T. E., P. M. Dixon, and B. E. Taylor. 1998. Detecting trends in species composition.

Ecological Applications 8:300–308.

Podani, J., and D. Schmera. 2011. A new conceptual and methodological framework for

exploring and explaining pattern in presence - absence data. Oikos 120:1625–1638.

Porensky, L. M., K. E. Mueller, D. J. Augustine, and J. D. Derner. 2016a. Thresholds and

gradients in a semi-arid grassland: long-term grazing treatments induce slow,

continuous and reversible vegetation change. Journal of Applied Ecology 53:1013–

1022.

Porensky, L. M., K. E. Mueller, D. J. Augustine, and J. D. Derner. 2016b. Data from:

Thresholds and gradients in a semi-arid grassland: long-term grazing treatments

induce slow, continuous and reversible vegetation change.

https://doi.org/10.5061/dryad.3m8p1.

Quinn, M. A., S. E. Halbert, and L. Williams. 1991. Spatial and temporal changes in aphid

(Homoptera: Aphididae) species assemblages collected with suction traps in Idaho.

Journal of Economic Entomology 84:1710–1716.

R Core Development Team. 2017. R: A language and environment for statistical computing

v. 3.3.3. R Foundation for Statistical Computing, Vienna.

Rayner, T. S., B. J. Pusey, and R. G. Pearson. 2009. Spatio-temporal dynamics of fish

feeding in the lower Mulgrave River, north-eastern Queensland: The influence of

1567   seasonal flooding, instream productivity and invertebrate abundance. Marine and

1568   Freshwater Research 60:97.

1569 Rojas-Sandoval, J., E. Meléndez-Ackerman, and D. S. Fernández. 2012. Plant community

1570   dynamics of a tropical semi-arid system following experimental removals of an exotic

1571   grass. Applied Vegetation Science 15:513–524.

1572 Ruhl, H. A. 2008. Community change in the variable resource habitat of the abyssal

1573   Northeast Pacific. Ecology 89:991–1000.

1574 Rusak, J. A., N. D. Yan, K. M. Somers, and D. J. McQueen. 1999. The temporal coherence of

1575   zooplankton population abundances in neighboring north-temperate lakes. The

1576   American Naturalist 153:46–58.

1577 Schaefer, J., K. Gido, and M. Smith. 2005. A test for community change using a null model

1578   approach. Ecological Applications 15:1761–1771.

1579 Shenhav, L., O. Furman, I. Mizrahi, and E. Halperin. 2017. Modeling the temporal dynamics

1580   of the gut microbial community in adults and infants. bioRxiv:212993.

1581 Shertzer, K. W., E. H. Williams, and J. C. Taylor. 2009. Spatial structure and temporal

1582   patterns in a large marine ecosystem: Exploited reef fishes of the southeast United

1583   States. Fisheries Research 100:126–133.

1584 Shi, S., D. Zou, Q. Wang, X. Xia, T. Zheng, C. Wu, and M. Gao. 2016. Responses of

1585   ammonia-oxidizing bacteria community composition to temporal changes in

1586   physicochemical parameters during food waste composting. RSC Adv. 6:9541–9548.

1587 Shimadzu, H., M. Dornelas, and A. E. Magurran. 2015. Measuring temporal turnover in

1588   ecological communities. Methods in Ecology and Evolution 6:1384–1394.

1589 Stead, T. K., J. M. Schmid-Araya, and A. G. Hildrew. 2003. All creatures great and small:

1590   patterns in the stream benthos across a wide range of metazoan body size. Freshwater

1591   Biology 48:532–547.

Sugiura, S., T. Tsuru, and Y. Yamaura. 2013. Effects of an invasive alien tree on the diversity

and temporal dynamics of an insect assemblage on an oceanic island. Biological

Invasions 15:157–169.

Thibaut, L. M., S. R. Connolly, and H. P. A. Sweatman. 2012. Diversity and stability of

herbivorous fishes on coral reefs. Ecology 93:891–901.

Thrush, S. F., J. Halliday, J. E. Hewitt, and A. M. Lohrer. 2008. The effects of habitat loss,

fragmentation, and community homogenization on resilience in estuaries. Ecological

Applications 18:12–21.

Tilman, D. 1999. The ecological consequences of changes in biodiversity: A search for

general principles. Ecology 80:1455–1474.

Tworek, S. 2004. Factors Affecting Temporal Dynamics of Avian Assemblages in a

Heterogeneous Landscape. Acta Ornithologica 39:155–163.

Ulrich, W., and N. J. Gotelli. 2007. Null model analysis of species nestedness patterns.

Ecology 88:1824–1831.

Ulrich, W., and N. J. Gotelli. 2010. Null model analysis of species associations using

abundance data. Ecology 91:3384–3397.

Vallès, Y., A. Artacho, A. Pascual-García, M. L. Ferrús, M. J. Gosalbes, J. J. Abellán, and M.

P. Francino. 2014. Microbial succession in the gut: Directional trends of taxonomic

and functional change in a birth cohort of Spanish infants. PLoS Genetics

10:e1004406.

Van den Brink, P. J., and C. J. Ter Braak. 1999. Principal response curves: Analysis of time-

dependent multivariate responses of biological community to stress. Environmental

Toxicology and Chemistry 18:138–148.

Vaughn, K. J., and T. P. Young. 2010. Contingent conclusions: Year of initiation influences

ecological field experiments, but temporal replication is rare. Restoration Ecology

18:59–64.

Whittaker, R. H. 1960. Vegetation of the Siskiyou Mountains, Oregon and California.

Ecological Monographs 30:279.

Xiong, J., J. Zhu, K. Wang, X. Wang, X. Ye, L. Liu, Q. Zhao, M. Hou, L. Qiuqian, and D.

Zhang. 2014. The temporal scaling of bacterioplankton composition: high turnover

and predictability during shrimp cultivation. Microbial Ecology 67:256–264.

Yamamura, K., M. Yokozawa, M. Nishimori, Y. Ueda, and T. Yokosuka. 2006. How to

analyze long-term insect population dynamics under climate change: 50-year data of

three insect pests in paddy fields. Population Ecology 48:31–48.

Zar, J. H. 1996. Biostatistical Analysis. Third. Prentice Hall, New Jersey.

Zuur, A. F., E. N. Ieno, and C. S. Elphick. 2010. A protocol for data exploration to avoid

common statistical problems: Data exploration. Methods in Ecology and Evolution

1:3–14.

1630

1631

Appendix S1

Figure S1. Proportion of studies identified using our search terms that were published each year until September 2017. No studies prior to 1991 were captured by our search terms. The total number of studies in each year is the value above the bars.

(A)

(B)

Figure S2. (A) The numbers of small and large spatial scale studies by habitat category. Small spatial scale (blue bars) studies were considered to be those carried out at micro, point sample, and local scales, while large spatial scale studies (purple bars) were those at regional, continental, or global scales. (B) The proportion of studies of different taxa within categories defined by the temporal grain (black circles) and extent (purple dots). Temporal grain is the minimum time between sampling events within the study and the temporal extent is the maximum time between sampling events within the study.

(A)

(B)

Figure S3. (A) The proportion of uses of each type of analysis across all sampled publications between 1990 and 2017. 'Minor methods' include compositional pivot days, frequency change, Markov chain modelling, multiplicative change, Moran's eigenvector maps (MEMs), Nestedness analysis, and synchrony. (B) The popularity of the different ordination methods used across the 27 years. 'Minor methods' are: dbRDA, DCCA, pCCA, pRDA, Procrustes, RA and multiple co-inertia analysis. Descriptions of analysis types are detailed the S1 Text supplementary material.

Figure S4. The distribution of studies across the different research aims. Research aims vary from understanding compositional dynamics at fine, medium and coarse temporal scales, to investigating the effect on species composition over time of species interactions, environmental conditions, and ecological disturbance, to understanding successional change in communities. Scales of temporal dynamics were classified as fine (days), medium (weeks, months, or seasons), and coarse (annual, years, or decades).

Appendix S2. Other methods encountered in the literature review, not explained in the text

UNIVARIATE LINEAR METHODS

Univariate linear methods, including analysis of variance (ANOVA), analysis of covariance (ANCOVA), repeated measures ANOVA (rmANOVA), simple correlation (e.g., Pearson's and Spearman's correlation coefficients), and regression (linear regression, generalised linear models, generalised linear mixed effects models, logistic regression, *etc.*), are used to examine temporal changes in a univariate response (dependent) variable. The independent variable is either time, or an environmental variable that changes over time, and the response variable usually is the (relative) abundance of a focal taxon or functional group.

Univariate linear methods are applied most frequently to community dynamics by constructing separate models for each species. These methods of assessing compositional dynamics are limited because multiple analyses, conducted in parallel on selected taxa, are required to make inferences regarding composition. For even modest numbers of taxa, these multiple analyses can become unwieldy and require *P*-value corrections for multiple tests (Benjamini and Hochberg 1995, Zar 1996). For example, Bugalho et al. (2013) used repeated-measures mixed-effects modelling to assess the effects of deer exclusion over time on the abundance of seedlings and saplings of forest tree species. In this case, the univariate modelling approach was usefully applied because fewer than ten species were analysed. Alternatively, abundance can be analysed for all species in the same model as long as there are enough degrees of freedom in the model to include species as a fixed or random effect term (e.g., Buckley and Freckleton 2010); see Harrell (2001) for a rule of thumb stating that we should have at least ten independent samples for each predictor variable in our model. Generalised linear models are more flexible in that they can handle designs with unbalanced data and nested sampling (e.g., plots within sites) (Bolker et al. 2009, Zuur et al. 2009). R code is provided in Appendix S4.

CLUSTER ANALYSIS (A DISSIMILARITY BASED METHOD)

Cluster analysis is a catch-all term applied to dissimilarity-based methods that either group samples together (agglomerative methods) or split all samples into sub-groups (divisive methods); the clustering is based on the distances between clusters (Legendre and Legendre

2012). There are two approaches for using cluster analysis to assess community dynamics:

(1) identifications of transitions between clusters over time, and (2) examination of

differences of clusters of repeated observations. In the first approach, a separate cluster

analysis is computed for each temporal sample of community data (e.g., Day and Buckley

2013). In the second approach, all samples taken at different times are included in a single

cluster analysis (e.g., Araujo et al. 2008). If compositional shifts are small relative to the total

compositional variation in the dataset, the first approach will not be able to detect temporal

differences. However, if compositional shifts are relatively large, it yields a clear way of

showing shifts in 'community types': groups of samples of similar composition that can be

characterised using a method such as indicator species analysis (e.g., Day and Buckley 2013).

The first method can potentially be applied to a larger number of time points if all that is

desired is to report the percent of sites that changed cluster groups each time. For example,

cluster analysis of 34 phytoplankton species in Laoshan Bay, China revealed distinct groups

of species occurred at different times over a six-month period (Xu et al. 2010).

TEMPORAL STABILITY (A.K.A. COEFFICIENT OF VARIATION)

The coefficient of variation (CV) is used to measure temporal stability of the abundance or

biomass of an individual taxon, or group of taxa, across all times (not space); this is not a

multivariate method (Grossman et al. 1990, Brown and Lawson 2010, Hector et al. 2010).

The CV is the ratio of the standard deviation to the mean, and often is multiplied by 100 to

obtain a percentage. The mean of the CV values for all taxa (e.g., abundance or biomass) is

used as an aggregate measure of temporal stability for a whole community; smaller values

imply greater stability. Because the CV is a ratio, changes in community composition can be

compared even if there are differences in total abundances among taxa or in taxon richness

among taxon groups. Such values are often presented in tables or the text as a measure of

variation or are sometimes used as a response variable against other (spatial) variables of

interest (e.g., Doxa et al. 2012). For example, CV can be calculated for disturbed and

undisturbed communities to assess if the disturbance has resulted in a more temporally

variable community. Grossman et al. (1990) categorised particular values of CV to assess

stability of freshwater fish communities: a community was 'highly stable' if it had a CV <

25%, 'moderately stable' if 25% < CV < 50%, 'moderately fluctuating' if 50% < CV < 75%

or 'highly fluctuating' if CV > 75%. The statistical significance of CV values could be

obtained by comparing the observed values to those from communities generated using

randomisations under sensible null models. Note that temporal CV is necessarily be measured
on a per site basis to exclude spatial variation.

FREQUENCY CHANGE

Changes in the frequency of individual taxa within a community over time can be assessed by
subtracting their frequency at each sampling occasion from a prior sampling occasion; this is
not a multivariate method. Permutation tests can be used to test the significance of these
changes in frequencies (Kapfer et al. 2011). Community samples (with their given taxon
frequencies) are randomised among sampling times and the change in taxon frequency
recalculated with each permutation. A significant change in the observed taxon frequency is
inferred if the observed change is greater than the 95[th] percentile change from the
permutations. Permutations should be spatially restricted so that only samples from
homogeneous habitat types are compared; otherwise the chance of spuriously detecting
change is greatly increased. The outcome of these tests is a list of taxa that had significant
increases or decreases in frequencies, and the amount, between sampling times. For example,
Kapfer et al. (2011) used permutation tests to assess changes in frequencies of 85 plant
species from two time points 50 years apart. This method could be used on data from
multiple sampling occasions, but would quickly become cumbersome with many temporal
observations. It can only be used to compare individual taxa over time, does not apply to rare
taxa (taxa need to occur in enough sites to make the calculations tractable), and results in a
(potentially) large table of taxa with significant (or not) changes, therefore, it has limited
applicability to taxon-rich datasets where changes in rare taxa are of interest.

MULTIPLICATIVE CHANGE

The change in percent cover of a taxon, or group of taxa, at a single site can be used to obtain
a 'growth rate' using a linear regression equation; this is not a multivariate method (Debinski
et al. 2010, Collins and Xia 2015). This rate of change can then be used in a regression or
ANOVA to assess predictors of change for each species ('multiplicative change'). For
example, different sites or sets of sites, can be compared in their relative change in the
percent cover of different functional groups (Debinski et al. 2010), as long as communities
consist of taxa with similar ecology and life-histories, e.g., grassland plant communities in
different moisture regimes. This method appears to be most useful for studies where (1) there
is little temporal replication, or at least where differences among temporal replicates are not

of interest, because the growth rate is calculated using only two temporal samples, and (2)

few species, species groups, are of interest because a different analysis must be performed for

each. However, it would be possible to calculate the change in percent cover for different

time periods within the overall study, if one was interested in investigating finer-scale trends.

MARKOV CHAIN MODELLING

Markov chain modelling uses a series of transition probabilities (a 'Markov chain'), which

describe how the occupancy of a taxon or functional group in a site changes from time t to

time $t + 1$ (Hill et al. 2004). The user varies these probabilities in space and/or time to create

a set of different hypothesis-driven models that predict the observed occupancy. The model

set can then be evaluated against the observed data using log-linear modelling and AIC

model selection to determine the influential and significant processes on taxon site occupancy

through time (Hill et al. 2004). Community dynamics usually are examined by comparing

predicted changes in individual taxa, making the method easier to apply to communities of

few taxa (or groups of taxa). Because models often are parameterised using the modelled

data, dependence of samples needs to be dealt with through sampling design or post-hoc

statistical analyses (Hill et al. 2004). This method is most appropriate for sessile communities

where patch occupancy is easy to replicate and define (e.g., Hill et al. 2004, Pueyo and

Beguería 2007). It best describes directional dynamics, such as secondary succession (e.g.,

Baasch et al. 2010, Jiménez-Franco et al. 2011) and it is relatively computationally difficult.

However, it makes detailed, taxon-level predictions that can be useful for the analysis of

experimental studies where researchers want to explore specific predictions from multiple

scenarios, such as taxon extinctions (e.g., Wootton 2004) or restoration manipulations

(Baasch et al. 2010).

MACHINE LEARNING AND DATA MINING METHODS

Machine learning is branch of computer science that deals with the development of learning

algorithms that are used to explore large, multivariate datasets, and has the primary aim of

generating accurate, predictive models (Carbonell et al. 1983). Common machine-learning

methods include artificial neural networks and decision-tree methods which can be used with

either univariate or multivariate response variables, and Bayesian versions can incorporate

prior knowledge (Cutler et al. 2007, Olden et al. 2008). Whilst the field of machine learning

focusses on making predictions from data, many of its methods also are used in a data-mining

capacity to discern spatiotemporal patterns from complex and noisy datasets and can be used

for classification, regression, clustering, and data reduction (Olden et al. 2008).

Machine-learning methods are attractive because they do not have many of the

restrictive assumptions, such as linearity or data normality, that underpin many traditional

statistical methods (Olden et al. 2008). Thus, machine-learning methods are becoming

increasingly popular for exploratory and predictive analyses of community dynamics (e.g.,

Kampichler et al. 2010). For example, Vallès et al. (2014) used self-organising maps, an

implementation of artificial neural networks (Kohonen 1982), to show that complex

microbial communities sampled from infants' guts formed distinct clusters at each of five

sample times (based on relative abundance data), suggesting that successional processes were

occurring within the first year of these infants' lives. Sheaves et al. (2010) used a multivariate

classification and regression tree analysis (De'ath 2002) to investigate spatial and temporal

differences in fish assemblages; results showed that despite considerable spatial variation in

community structure among different estuaries, there nonetheless were clear patterns of

temporal change in species' abundances and assemblage composition. These methods are

very flexible and can handle both univariate and multivariate input datasets with large

numbers of predictor variables, including space and time, allowing simultaneous

investigation of spatial and temporal community dynamics. Standardised approaches for

characterising different types of temporal patterns and causal processes in community

datasets will hopefully emerge from further research. R code is not provided, but R packages

are available (the MultivariateRandomForest package: Rahman 2017, and rpart package:

Therneau and Atkinson 2018).

NETWORK ANALYSIS

Ecological network analysis enables the investigation of the complex associations among

species, and among species and their environment, through space and time (Newman 2010,

Blonder et al. 2012). This method recognises that an ecological community is a complex

biological system comprised of interconnected units whose associations can be modelled

mathematically using constructs such as vertices (representing taxa) and edges, which are the

connections between the vertices (representing ecological interactions) (Blonder et al. 2012).

In the context of investigating community dynamics, patterns of species' associations and

community memberships are most typically visualised as a topological network diagram

created from taxon abundance data collected at multiple time points (*sensu* "time-aggregated

networks" (Blonder et al. 2012). From the network topology, a variety of statistics can be

calculated that measure variability in community composition and interactions at particular

times, which can then be compared. This method is most applicable for multi-trophic

communities or where interaction networks can be clearly defined for the community; it is

less useful where a single trophic level is of interest and interactions are not quantified, e.g., a

plant community, but see Cram et al. (2015) for an example of spatiotemporal microbial

networks. In R, the package 'timeordered' is the easiest way to implement a network analysis

of community dynamics data. R code is not provided.

REFERENCES CITED

Araujo, H. M. P., D. A. Nascimento-Vieira, S. Neumann-Leitão, R. Schwamborn, A. P. O.

Lucas, and J. P. H. Alves. 2008. Zooplankton community dynamics in relation to the

seasonal cycle and nutrient inputs in an urban tropical estuary in Brazil. Brazilian

Journal of Biology 68:751–762.

Baasch, A., S. Tischew, and H. Bruelheide. 2010. Twelve years of succession on sandy

substrates in a post-mining landscape: a Markov chain analysis. Ecological

Applications 20:1136–1147.

Benjamini, Y., and Y. Hochberg. 1995. Controlling the false discovery rate: a practical and

powerful approach to multiple testing. Journal of the Royal Statistical Society. Series

B (Methodological) 57:289–300.

Blonder, B., T. W. Wey, A. Dornhaus, R. James, and A. Sih. 2012. Temporal dynamics and

network analysis. Methods in Ecology and Evolution 3:958–972.

Bolker, B. M., M. E. Brooks, C. J. Clark, S. W. Geange, J. R. Poulsen, M. H. H. Stevens, and

J.-S. S. White. 2009. Generalized linear mixed models: a practical guide for ecology

and evolution. Trends in Ecology & Evolution 24:127–135.

Brown, B. L., and R. L. Lawson. 2010. Habitat heterogeneity and activity of an omnivorous

ecosystem engineer control stream community dynamics. Ecology 91:1799–1810.

Buckley, H. L., and R. P. Freckleton. 2010. Understanding the role of species dynamics in

abundance-occupancy relationships. Journal of Ecology 98:645–658.

Bugalho, M. N., I. Ibáñez, and J. S. Clark. 2013. The effects of deer herbivory and forest type

on tree recruitment vary with plant growth stage. Forest Ecology and Management

308:90–100.

Carbonell, J. G., R. S. Michalski, and T. M. Mitchell. 1983. An overview of machine

learning. Pages 3–23 Machine Learning: An artificial intelligence approach. Tioga

Publishing Company, Palo Alto.

Collins, S. L., and Y. Xia. 2015. Long-term dynamics and hotspots of change in a desert

grassland plant community. The American Naturalist 185:E30–E43.

Cram, J. A., L. C. Xia, D. M. Needham, R. Sachdeva, F. Sun, and J. A. Fuhrman. 2015.

Cross-depth analysis of marine bacterial networks suggests downward propagation of

temporal changes. The ISME journal 9:2573.

Cutler, D. R., T. C. Edwards, K. H. Beard, A. Cutler, K. T. Hess, J. Gibson, and J. J. Lawler.

2007. Random forests for classification in ecology. Ecology 88:2783–2792.

Day, N. J., and H. L. Buckley. 2013. Twenty-five years of plant community dynamics and

invasion in New Zealand tussock grasslands. Austral Ecology 38:688–699.

De'ath, G. 2002. Multivariate regression trees: A new technique for modeling species-

environment relationships. Ecology 83:1105–1117.

Debinski, D. M., H. Wickham, K. Kindscher, J. C. Caruthers, and M. Germino. 2010.

Montane meadow change during drought varies with background hydrologic regime

and plant functional group. Ecology 91:1672–1681.

Doxa, A., M. L. Paracchini, P. Pointereau, V. Devictor, and F. Jiguet. 2012. Preventing biotic

homogenization of farmland bird communities: The role of high nature value

farmland. Agriculture, Ecosystems & Environment 148:83–88.

Grossman, G. D., J. F. Dowd, and M. Crawford. 1990. Assemblage stability in stream fishes:

A review. Environmental Management 14:661–671.

Harrell, F. E. 2001. Regression Modeling Strategies: With Applications to Linear Models,

Logistic Regression, and Survival Analysis. Springer, New York.

Hector, A., Y. Hautier, P. Saner, L. Wacker, R. Bagchi, J. Joshi, M. Scherer-Lorenzen, E. M.

Spehn, E. Bazeley-White, M. Weilenmann, M. C. Caldeira, P. G. Dimitrakopoulos, J.

A. Finn, K. Huss-Danell, A. Jumpponen, C. P. H. Mulder, C. Palmborg, J. S. Pereira,

A. S. . Siamantziouras, A. C. Terry, A. Y. Troumbis, B. Schmid, and M. Loreau.

2010. General stabilizing effects of plant diversity on grassland productivity through

population asynchrony and overyielding. Ecology 91:2213–2220.

Hill, M. F., J. D. Witman, and H. Caswell. 2004. Markov chain analysis of succession in a

rocky subtidal community. The American Naturalist 164:E46–E61.

Jiménez-Franco, M. V., J. E. Martínez, and J. F. Calvo. 2011. Territorial occupancy

dynamics in a forest raptor community. Oecologia 166:507–516.

Kampichler, C., R. Wieland, S. Calmé, H. Weissenberger, and S. Arriaga-Weiss. 2010.

Classification in conservation biology: A comparison of five machine-learning

methods. Ecological Informatics 5:441–450.

Kapfer, J., J.-A. Grytnes, U. Gunnarsson, and H. J. B. Birks. 2011. Fine-scale changes in

vegetation composition in a boreal mire over 50 years: Fine-scale changes in boreal

mire. Journal of Ecology 99:1179–1189.

Kohonen, T. 1982. Self-organized formation of topologically correct feature maps. Biological

Cybernetics 43:59–69.

Legendre, P., and L. F. J. Legendre. 2012. Numerical Ecology. Third Edition. Elsevier,

Radarweg, The Netherlands.

Newman, M. E. J. 2010. Networks: An introduction. Oxford University Press, New York.

Olden, J. D., J. J. Lawler, and N. L. Poff. 2008. Machine learning methods without tears: A

primer for ecologists. The Quarterly Review of Biology 83:171–193.

Pueyo, Y., and S. Beguería. 2007. Modelling the rate of secondary succession after farmland

abandonment in a Mediterranean mountain area. Landscape and Urban Planning

83:245–254.

Rahman, R. 2017. MultivariateRandomForest: Models Multivariate Cases Using Random

Forests. R package version 1.1.5, https://CRAN.R-

project.org/package=MultivariateRandomForest.

Sheaves, M., R. Johnston, and R. Connolly. 2010. Temporal dynamics of fish assemblages of

natural and artificial tropical estuaries. Marine Ecology Progress Series 410:143–157.

Therneau, T., and B. Atkinson. 2018. rpart: Recursive Partitioning and Regression Trees. R

package version 4.1-12. https://CRAN.R-project.org/package=rpart.

Vallès, Y., A. Artacho, A. Pascual-García, M. L. Ferrús, M. J. Gosalbes, J. J. Abellán, and M.

P. Francino. 2014. Microbial succession in the gut: Directional trends of taxonomic

and functional change in a birth cohort of Spanish infants. PLoS Genetics

10:e1004406.

Wootton, T. J. 2004. Markov chain models predict the consequences of experimental

extinctions. Ecology Letters 7:653–660.

Xu, H., G.-S. Min, J.-K. Choi, K. A. S. Al-Rasheid, X. Lin, and M. Zhu. 2010. Temporal

dynamics of phytoplankton communities in a semi-enclosed mariculture pond and

their responses to environmental factors. Chinese Journal of Oceanology and

Limnology 28:295–303.

Zar, J. H. 1996. Biostatistical Analysis. Third. Prentice Hall, New Jersey.

Zuur, A. F., E. N. Ieno, N. J. Walker, A. A. Saveliev, and G. M. Smith. 2009. Mixed Effects

Models and Extensions in Ecology with R. Springer, New York.

Hannah L. Buckley, Nicola J. Day, Bradley S. Case, Gavin Lear, Aaron M. Ellison. 2019. Multivariate methods for testing hypotheses of temporal community dynamics. *Ecological Monographs.*

Data S1

Data extracted as part of the literature review (as .csv file)

Author(s) [of the material provided in DataS1.zip]

Hannah L. Buckley
School of Science
Auckland University of Technology
Auckland
New Zealand
Hannah.Buckley@aut.ac.nz

Nicola J. Day
Biology Department
Wilfrid Laurier University
Waterloo, Ontario
Canada

Bradley S. Case
School of Science
Auckland University of Technology
Auckland
New Zealand
Bradley.Case@aut.ac.nz

Gavin Lear
School of Biological Sciences
University of Auckland
Auckland
New Zealand

Aaron M. Ellison
Harvard Forest
Harvard University ,
Petersham, Massachusetts

USA

File list (files found within DataS1.zip)

```
Literature review data attributes.txt
Literature review data.csv
```

Description

Please write a prose description of the files, including variable descriptions, use, *etc.*

Literature review data attributes.txt – This text file contains attributes of the data extracted during the quantitative literature review, providing a description of the data found in the associated spreadsheet (Data S1.csv).

Literature review data.csv – This is a .csv spreadsheet of the data extracted as part of the quantitative literature review carried out in this study.

Hannah L. Buckley, Nicola J. Day, Bradley S. Case, Gavin Lear, Aaron M. Ellison. 2019. Multivariate methods for testing hypotheses of temporal community dynamics. *Ecological Monographs.*

Data S2

Worked case study example 1 R code files

Author(s) [of the material provided in DataS2.zip]

Hannah L. Buckley
School of Science
Auckland University of Technology
Auckland
New Zealand
Hannah.Buckley@aut.ac.nz

Nicola J. Day
Biology Department
Wilfrid Laurier University
Waterloo, Ontario
Canada
njday.ac@gmail.com

Bradley S. Case
School of Science
Auckland University of Technology
Auckland
New Zealand
Bradley.Case@aut.ac.nz

Gavin Lear
School of Biological Sciences
University of Auckland
Auckland
New Zealand
g.lear@auckland.ac.nz

Aaron M. Ellison
Harvard Forest
Harvard University

Petersham, Massachusetts
USA
aellison@fas.harvard.edu

File list (files found within DataS2.zip)

`Worked example 1.txt`

Description

`Worked example 1.txt` – R code for the worked case study example 1: Testing for significant temporal community dynamics within a controlled experiment. Further description and instruction can be found within the R script as annotation.

Hannah L. Buckley, Nicola J. Day, Bradley S. Case, Gavin Lear, Aaron M. Ellison. 2019. Multivariate methods for testing hypotheses of temporal community dynamics. *Ecological Monographs.*

Data S3

Worked case study example 2 R code file

Author(s) [of the material provided in DataS3.zip]

Hannah L. Buckley
School of Science
Auckland University of Technology
Auckland
New Zealand
Hannah.Buckley@aut.ac.nz

Nicola J. Day
Biology Department
Wilfrid Laurier University
Waterloo, Ontario
Canada
njday.ac@gmail.com

Bradley S. Case
School of Science
Auckland University of Technology
Auckland
New Zealand
Bradley.Case@aut.ac.nz

Gavin Lear
School of Biological Sciences
University of Auckland
Auckland
New Zealand
g.lear@auckland.ac.nz

Aaron M. Ellison
Harvard Forest
Harvard University

Petersham, Massachusetts
USA
aellison@fas.harvard.edu

File list (files found within DataS3.zip)

```
Worked example 2.txt
```

Description

`Worked example 2.txt` – R code for the worked case study example 2: Community dynamics analysis of the BCI dataset. Further description and instruction can be found within the R script as annotation.

Hannah L. Buckley, Nicola J. Day, Bradley S. Case, Gavin Lear, Aaron M. Ellison. 2019. Multivariate methods for testing hypotheses of temporal community dynamics. *Ecological Monographs.*

Data S4

Worked case study example 3 R code files

Author(s) [of the material provided in DataS4.zip]

Hannah L. Buckley
School of Science
Auckland University of Technology
Auckland
New Zealand
Hannah.Buckley@aut.ac.nz

Nicola J. Day
Biology Department
Wilfrid Laurier University
Waterloo, Ontario
Canada
njday.ac@gmail.com

Bradley S. Case
School of Science
Auckland University of Technology
Auckland
New Zealand
Bradley.Case@aut.ac.nz

Gavin Lear
School of Biological Sciences
University of Auckland
Auckland
New Zealand
g.lear@auckland.ac.nz

Aaron M. Ellison
Harvard Forest
Harvard University

Petersham, Massachusetts
USA
aellison@fas.harvard.edu

File list (files found within DataS4.zip)

```
Worked example 3.txt
```

Description

`Worked example 3.txt` – R code for the worked case study example 3: Cyclical dynamics example from Gilbert et al. (2012). Further description and instruction can be found within the R script as annotation.