Systems Biology Understanding of the Effects of Lithium on Cancer.

Weihao Ge[1,2] and Eric Jakobsson[1,2,3,*]

[1]National Center for Supercomputing Applications, [2]Center for Biophysics and Computational Biology, [3]Department of Molecular and Integrative Physiology University of Illinois at Urbana-Champaign, Urbana, IL, USA

Correspondence

*Eric Jakobsson jake@illinois.edu

Abstract

Lithium has many widely varying biochemical and phenomenological effects, suggesting that a systems biology approach is required to understand its action. Multiple lines of evidence point to lithium as a significant factor in development of cancer, showing that understanding lithium action is of high importance. In this paper we undertake first steps towards a systems approach by analyzing mutual enrichment between the interactomes of lithium-sensitive enzymes and the pathways associated with cancer. This work integrates information from two important databases, STRING and KEGG pathways. We find that for the majority of cancer pathways the mutual enrichment is many times greater than chance, reinforcing previous lines of evidence that lithium is an important influence on cancer.

Keywords

Lithium, systems biology, biochemical pathways, biochemical networks,

Introduction

Clinical and Epidemiological Context for Lithium and Cancer: By far the most common medical use of lithium is as a first line therapy for bipolar disorder, including associated depression as well as mania.[1] A comprehensive review of the literature confirms that lithium is also effective against unipolar depression with unique anti-suicidal effectiveness, and may also be useful against cancer and neurodegenerative disease.[2]

One line of evidence for the possible use of lithium as an anticancer agent is epidemiological. A retrospective study showed that psychiatric patients undergoing lithium therapy for bipolar disorder had a much lower incidence of cancer than a matched group not receiving lithium therapy.[3] More recent studies of similar design, one conducted nationwide across Sweden, and another across Taiwan, achieved the same result.[4][5] On the other hand another nationwide study, this time from Denmark, showed no correlation of lithium with colorectal adenocarcinoma.[6] On closer look, the Denmark study does not contradict the Swedish study. The Swedish study also found that for the entire population lithium was not correlated with cancer incidence, but in addition found that bipolar individuals not treated with lithium had a higher incidence of cancer than the general population. Lithium-treated bipolar patients, on the other hand, had essentially the same cancer incidence as the general population.

One piece of experimental evidence for lithium's potential as a cancer therapy is that inhibition of GSK3 was observed to inhibit prostate cancer cell proliferation[7]. A detailed study of molecular mechanisms by which lithium inhibition of GSK3B inhibits proliferation of prostate tumor cells in culture was presented by Sun et al.[8] The work was subsequently extended to an animal model.[9] A clinical trial for the

effect of lithium coupled with prostatectomy on men has been conducted but as of this
writing results have not yet been published.[10]
With respect to other cancers, lithium has been found to be lethal to
neuroblastoma cells but not to normal nerve cells.[11] The experimentally determined
effective dose was 12 mM, a level which would be lethal if achieved systemically in a
human or model organism but perhaps could be induced locally. A similar effect was
found in ovarian cancer cells,[12] although a subsequent similar study on ovarian cancer
cells suggests only a more modest benefit.[13] It is not clear from our reading of the two
ovarian cancer papers why the results are significantly different from each other.
With respect to colorectal cancer, it has been found that lithium inhibits
proliferation of a colorectal cancer cell line.[14] PTEN overexpression and lithium
administration were shown to cooperatively inhibit proliferation of colorectal cancer
cells.[15] Another study on colon cancer cells showed that lithium specifically induced a
reversal of the epithelial-to-mesenchymal transition characteristic of the cancer cells.[16]
Two studies with relatively small sample size suggested a possible link between
lithium and tumors of the upper urinary tract.[17] [18] However a large-scale study involving
all urinary tract cancers in Denmark over a multi-year period found no correlation with
lithium use.[19]
Because lithium therapy is systemic rather than topical or local, it follows that
lithium might inhibit metastasis. Evidence that this is the case for colon cancer comes
from observation of inhibition of metastasis-inducing factors by lithium and by
observation on reduced metastasis in model animals given lithium therapy.[20]
Autophagy is a key cellular process in the inhibition of cancer.[21] Lithium has
been shown to induce autophagy, due to its inhibition of inositol monophosphatase.[22]
The full range of lithium effects on autophagy is complicated,[23] as might be expected
because of lithium's lack of specificity.[2]
Because of the promising indications as cited above, lithium has been suggested
as one of a number of drugs commonly used for other reasons, to be repurposed for
cancer.[24]
76
Biochemical Context for Lithium and Cancer: Much of lithium's biochemical action
may be summarized by noting that it inhibits, or competes with, action of
magnesium.[2] Examples of such competition, that appear specially relevant for cancer,
are given below.
Lithium appears to inhibit β-adrenergic and muscarinic receptor coupling to G
proteins by competing with magnesium, which facilitates such coupling. [25] [26] [27] [28] [29]
Overexpression of G protein coupled receptors is common theme of many cancers,[30]
suggesting that inhibition of receptor-G protein coupling may inhibit cancer progression.
A particularly important example is substitution of a lithium ion for a magnesium
ion acting as a cofactor in inositol monophosphatase, mentioned earlier in this paper in
connection with autophagy. In this protein the binding site for lithium is not revealed in
crystallography nor in solution NMR but can be identified in magic angle spinning solid
state NMR, which is more suitable for systems with large internal motion.[31]
One mode of action with many consequences is lithium inhibition of glycogen
synthase kinase 3 beta (GSK3B), mentioned earlier in this paper in connection with
prostate cancer. Lithium inhibition of GSK3B was initially shown in vitro and in intact

93	cells,[32] and in the context of embryonic development.[33] It was later shown that lithium
94	exerted its inhibitory effect on GSK3B by competing with magnesium for an essential
95	binding site.[34] There are two closely related forms of GSK3, termed alpha (GSK3A) and
96	beta (GSK3B), which are equivalently inhibited by lithium.[35] The two forms of GSK3
97	have substantial functional redundancy.[36] However some of their physiological
98	properties are different, as demonstrated by the fact that GSK3B knockout mice are not
99	viable,[37] but GSK3A knockout mice survive.[38] The very widespread nature of GSK3B
100	effects is related to the large number of transcription factors that it regulates.[39] It
101	functionally modulates cellular threshold for apoptosis,[40] it is central to mediating
102	mitochondrial response to stress;[41] it facilitates immune responses by enabling the
103	nuclear export of NF-ATc;[42] it regulates inflammation;[43] it regulates cardiac hypertrophy
104	and development,[44] to name just a few. Based on microarray studies of brain cells in
105	animals, lithium alters gene expression patterns significantly,[45] to be expected due to the
106	large number of transcription factors regulated by GSK3B. Mice heterozygous for
107	GSK3B exhibit similar behavioral traits to wild type littermates treated with 1mM lithium
108	(a concentration that inhibits about 25% of GSK3 activity, in line with 1 of the 4 alleles
109	of GSK3 inactivated in the GSK3B heterozygous mice)[46]
110	In addition to inhibiting the activity of GSK3B, lithium also inhibits its
111	transcription.[47] Of all kinases, GSK3 appears to have the largest number of known
112	substrates, over 100 known[48] and about 500 predicted by theory based on scanning and
113	interpreting relevant motif sequences in the human genome.[49] Lithium will thus to some
114	extent modulate activity along all pathways containing the hundreds of GSK3 substrates.
115	So far, to our knowledge there are no published counterexamples to the hypothesis that
116	lithium will exert an inhibitory effect on all proteins with essential magnesium binding
117	sites, of which there are estimated to be over three thousand.[50]
118	Multiple other studies, in addition to those mentioned above, have addressed the
119	potential efficacy of GSK3 inhibition as a therapeutic strategy against cancer.[51 52 53 54 55]
120	[56 57 58 59 60 61 62 63]
121	A widespread mechanism of lithium action is as a modulator of magnesium action
122	in interacting with phosphate groups. The primary energy source for cells and the
123	substrate for phosphorylating enzymes is not bare ATP, but rather magnesium-associated
124	ATP (MgATP).[64] NMR studies show that lithium associates with MgATP.[65] Based on
125	this admittedly small amount of data, we hypothesize that lithium associates with all
126	magnesium-phosphate complexes and will thus modulate to some extent all
127	phosphorylation reactions and all ATP-splitting processes. This is a reasonable
128	interpretation of early work by Willis and Fang, in which lithium was found to increase
129	the activity of the sodium-potassium pump without itself being transported
130	significantly.[66] We have noted earlier in this paper the inhibitory effect of lithium on
131	GSK3 by the mechanism of competing with Mg. Here we note that lithium also inhibits
132	the activity of GSK3 by a second method, that is, by increasing phosphorylation.[67]
133	Depending on context of relevant protein-protein interactions, lithium's effect on
134	phosphorylation of a particular protein may be to either increase it or decrease it. For
135	example, lithium decreases phosphorylation of tau-protein, presumably because it inhibits
136	GSK3B, which is implicated in the phosphorylation of the tau-protein.[68]
137	Because lithium affects many different biological molecules and processes[2], it is
138	essential to utilize the tools of systems biology[69] if a comprehensive understanding of

lithium action and its prospects for therapy are to be obtained. Important concepts for organizing biological information in a systems context are pathways and networks. A very useful tool for obtaining data about known pathways is the KEGG database.[70] An equally useful and complementary tool is the STRING database of interacting proteins.[71] In the present paper we investigate further the possible linkages among 1) lithium, 2) affective disorders, and 3) neurodegenerative disorders by analyzing the mutual enrichment between STRING-derived interactomes of lithium-sensitive enzymes, and the KEGG pathways associated with cancer.

Methods

Analysis was performed on the interactomes of lithium-sensitive genes, as identified by prior literature search[2]. This search suggested BDNF, BPNT1, DISC1, DIXDC1, FBP1, FBP2, GSK3A, GSK3B, inositol monophosphatases (IMPA1, IMPA2, and IMPAD1), INPP1, and PGM1 as key to understanding the broad biological actions of lithium. The interactomes of these genes were extracted from the STRING database (https://string-db.org). For each key gene, we adjust confidence level and order of neighbors (nearest only or next nearest included), so that each set contains a few hundred genes. This size is large enough for statistically reliable enrichment analysis. Very similar sets were merged; in particular FBP1 and FBP2 were merged into one set, and the inositol monophosphatases were merged into one set. On the other hand, GSK3A and GSK3B showed sufficient differences to be considered separately. Overall, we consider 10 distinct lithium-sensitive entities.

Disease Association

We used the R-package KEGGgraph[72] [73] to identify the genes associated with the pathways of interest.

 P-value calculation

The fundamental question we address is whether there is significant overlap or mutual enrichment between the interactomes of lithium-sensitive genes and the pathways or gene sets implicated in various cancers.

For each of the 10 lithium sets, an ensemble of 1000 null sets are generated by random selection from the human genome. Each null set is the same size as the corresponding lithium set. Then we used the R-package STRINGdb[74] to perform KEGG pathway enrichment analysis. This operation is a particular example of the powerful technique of gene-annotation enrichment analysis.[75] In gene-annotation enrichment analysis a test list of genes (often derived from gene expression experiments) is compared to an organized database of gene annotations, often referred to as a gene ontology[76], an array of gene lists corresponding to different biological functions, molecular functions, or locations in the cell. The output of the gene-annotation enrichment analysis is expressed as the likelihood that the list overlaps could have occurred by chance (p-value). A very low p-value implies that the degree of overlap is highly significant statistically and very likely is significant biologically. In our study the gene lists we are comparing are the interactomes of

lithium sensitive enzymes on the one hand, and KEGG pathways and Kegg pathways
associated with cancer on the other hand. For each KEGG term retrieved, a null
distribution of uncorrected p-value is generated by the 1000 null sets. This gives us
a measure of the false discovery rate, since any overlap between the null sets and
the KEGG pathways is purely accidentally. Then the fraction of null set uncorrected
p-values smaller than or equal to the lithium-sensitive set uncorrected p-value
would be the empirical p-value. For a detailed discussion of empirical p-value
determination see Ge et al[77].

Key Gene Prediction

We predict key genes by counting how many times a gene appears in the
cross section of interactomes and pathways associated with a particular disease. In
this way, we predict which genes might be most important in disease-related
pathways. Then, the genes are scored by the sum of mean counts over all diseases. A
higher ranking indicates a gene would be associated with an important factor in
many diseases.

<div align="center">

Results

</div>

Fig.1 shows mutual lithium interactome enrichment with specific cancer pathways,
represented by heatmaps. Each area on the heatmap is a color-coded representation of the
degree of mutual enrichment between the genes in the interactome of the indicated
lithium sensitive enzyme and the genes in the indicated pathway. The darker the shade,
the more significant the mutual enrichment of the interactome-pathway combination is.
The light areas on the heatmap represent situations where a lithium-sensitive interactome
has little or no mutual enrichment with a cancer pathway. The dark areas, deep orange
and red, represent situations where enrichment is very strong—far greater than could be
expected by chance.   Three genes stand out as being not strongly connected to cancer
pathways: BPNT1, DISC1, and PGM1. Of the cancer pathways, breast cancer stands out
as being not likely to be strongly influenced by lithium levels. For the remainder of the
genes and the remainder of the cancers, the relationship between the lithium-sensitive
interactome and the cancer phenome is strong.   For example, for the DIXDC1
interactome, the strong correlation we see with cancer pathways is reinforced by data
suggesting DIXDC1 implication in colon cancer,[78] lung cancer,[79] gastric cancer,[80]
prostate cancer,[81] glioma,[82] and pancreatic cancer.[83] It should be noted that DIXDC1 has
not been noted in the literature to be directly inhibited by lithium. Rather its modulation
by lithium is a consequence of its proximity to GSK3B in functional protein-protein
networks. Consequently, there is a significant overlap between its interactome and that
of GSK3B. However, there is sufficient difference to warrant the two interactomes being
considered as separate entities for purposes of the analysis presented here.

 Figure

Figure 1. *Visual representation of mutual enrichment patterns between cancer-*
associated pathways and the interactomes of lithium-sensitive gene products.
Calibration of p-value vs. color is indicated by a vertical scale to the right of the heat
map. Red or dark orange indicates very strong enrichment while lighter color indicates
weak or, if white, no enrichment. Three genes stand out as being not strongly connected
to cancer pathways: BPNT1, DISC1, and PGM1. Of the cancer pathways, breast cancer
stands out as being not likely to be strongly influenced by lithium levels. For the
remainder of the genes and the remainder of the cancers, the relationship between the
lithium-sensitive interactome and the cancer phenome is strong.

For each of the cancer-associated pathways we wished to compute a single number
representing the relative likely sensitivity of the disease to lithium, in order to contribute
to prioritizing which diseases are most likely to benefit from clinical trials with lithium.
There is a significant literature on combining p-values,[84] with choices among methods
depending on the detailed structure of the data. We adopt a relatively simple approach,
which is to compute the geometric mean of the individual p-values for each pathway-
interactome mutual enrichment value.

$$p_{mean} = (p_1 \text{ x } p_2 \text{ x } p_3 \text{ x} \ldots \ldots p_n)^{1/n} \qquad \text{Equation (1)}$$

The method of averaging in Equation (1) ensures that both strong and weak enrichments
contribute significant weight to the mean. Note that all of the p-values that go into
Equation (1) are corrected for false discovery rate by random resampling. Thus, no
further false discovery rate correction is necessary for computing p_{mean}. Note also that our
method is bounded at the low end of p-values by the number of null samples it is
reasonable to compute, given compute time constraints. For one thousand null sets as
used in this paper, the computed p-value will be zero when none of the thousand null sets

246   shows the degree of enrichment of the test sets. For purposes of computing the p_{mean} in
247   equation (1) we substitute 10^{-4} for zero for each of these cases.
248

Cancer-related KEGG pathway	p_{mean}	$1/p_{mean}$
Colorectal_cancer	0.00049903	2003.89209
Pancreatic_cancer	0.00053973	1852.76828
Proteoglycans_in_cancer	0.00054323	1840.83131
Renal_cell_carcinoma	0.00056283	1776.72794
Pathways_in_cancer	0.00056569	1767.76501
Chronic_myeloid_leukemia	0.00085134	1174.61894
Non-small_cell_lung_cancer	0.00090896	1100.15513
Endometrial_cancer	0.00091244	1095.95823
Prostate_cancer	0.00091481	1093.12212
MicroRNAs_in_cancer	0.00093106	1074.04618
Melanoma	0.00093303	1071.77346
Viral_carcinogenesis	0.00099762	1002.38273
Glioma	0.00122264	817.904951
Acute_myeloid_leukemia	0.00125638	795.934615
Bladder_cancer	0.00150566	664.158631
Small_cell_lung_cancer	0.00352993	283.291551
Thyroid_cancer	0.00567253	176.288152
Basal_cell_carcinoma	0.03711938	26.9401078
Chemical_carcinogenesis	0.05277248	18.9492716
Transcriptional_misregulation_in_cancer	0.14953148	6.68755483
Central_carbon_metabolism_in_cancer	1	1
Choline_metabolism_in_cancer	1	1
Breast_cancer	1	1

Table 1. *Rank order of significance of enrichment between lithium-sensitive*
interactomes and KEGG cancer-associated pathways. It is seen that for the great
majority of pathways the mutual enrichment is very significant, with p-values
significantly below .01. Breast cancer is unusual; it appears there is no enrichment
beyond chance. The table also displays a "lithium sensitivity index", which is $1/p_{mean}$

Table 1 shows in rank order the significance of enrichment between lithium-
sensitive interactomes and KEGG cancer-associated pathways. It is seen that for the
great majority of pathways the mutual enrichment is very significant, with p-values
significantly below .01 Breast cancer is unusual; it appears there is no enrichment
beyond chance. The table also displays a "lithium sensitivity index", which is $1/p_{mean}$
We should note that sensitivity to lithium does not necessarily imply a *beneficial*
sensitivity. There are some indications for some cancers that lithium might be beneficial,
as described in the Introduction section of this paper, but because of the complexity of the
feedback relationships in these pathways, a complicated relationship between lithium
ingestion and cancer incidence is very possible.

267
Figure 2. *This figure visualizes the strength of the projected lithium influence on cancer*
pathways. The logarithms of the lithium sensitivity indices ($1/p_{mean}$) are shown in boxplot
format together with the corresponding results when the lithium interactomes are
replaced with random gene sets. Essentially this figure shows the signal-to-noise ratio of
our results and demonstrates conclusively that lithium ingestion is overwhelmingly likely
to modulate the incidence of a wide range of cancers.
274
Figure 2 visualizes the strength of the projected lithium influence on cancer
pathways. In this figure the logarithms of the lithium sensitivity indices ($1/p_{mean}$) are
shown in boxplot format together with the corresponding results when the lithium
interactomes are replaced with random gene sets. Essentially this figure shows the
signal-to-noise ratio of our results and suggests that lithium ingestion is overwhelmingly
likely to influence the incidence of a wide range of cancers.
281
282

Summary and Discussion

We have conducted a pathway and network analysis exploring the role of lithium in multiple cancers. The results show that for the large majority of such cancers, there is high mutual enrichment between the interactomes of lithium-sensitive enzymes and the pathways associated with those diseases, indicating that lithium is very likely to affect the incidence and course of the disease. Our results are consistent with a variety of lines of evidence from both epidemiology and from experiment, cited in the Introduction section of this paper, suggesting possible influence of lithium on the incidence and progression of cancer.

We hope that the results described in this paper will contribute to prioritizing and designing clinical trials of lithium for cancer. To provide context for such prioritization and design, it is essential to take into account the ways in which lithium is unique, both as a pharmaceutical and as an ion that is ubiquitous in the environment, and therefore ubiquitous in the water and food we ingest[2]:

1. Unlike other ions, lithium is not regulated by selective membrane transport processes. Therefore, lithium concentration in both extracellular and intracellular compartments, rather than being roughly constant, is roughly proportional to lithium ingestion.

2. Unlike other pharmaceuticals, lithium is wildly nonselective in its biochemical effects. The major underlying mechanism for the lack of selectivity is lithium's general propensity to modulate the many biochemical processes and structures that involve magnesium.

3. Unlike other pharmaceuticals, lithium is an essential nutrient. The question with lithium is not whether it should be ingested or not, but rather how much. Extreme lithium deprivation results in failure to thrive, while too much lithium is toxic.

In the light of all these factors, we suggest that the correct question to ask with respect to lithium and a particular disease is not, "Should lithium be administered for this particular disease?" but rather, "What is the optimum blood level of lithium for this individual, given his or her disease history, status, genetic propensities, and other medications?" Unlike other pharmaceuticals that are far more specific and inhibit or activate one gene or a small number of genes, the model for lithium action is that it alters the balance between a large number of interacting processes and pathways. Thus, a dose-response curve for lithium is likely to be highly nonlinear and not always monotonic.

There are just a few well-established markers for optimum concentrations. For a patient with a reliable diagnosis of bipolar disorder a common target for optimality would be blood concentration of 0.8-1 mM. Significantly higher concentrations will result in acute toxicity, while significantly lower will result in loss of effectiveness. However, this level has some side effects when sustained for years or decades, namely an increased risk of kidney damage and lowered thyroid activity.

At the other end of the dosage scale, epidemiological evidence is compelling that geographical variations in concentration of lithium in the drinking water are correlated with a variety of health and wellness markers.

Another important marker is provided by a study showing that over a four-year period a lithium level of .25-.4 mM of lithium (1/4 to 1/2 of the bipolar therapeutic dose)

did not incur any renal damage[85]. This study suggests that clinical studies exploring low
to medium-dose lithium could be undertaken with relatively minimal concerns for side
effects.
Li et al have suggested GSK3B inhibitors for prostate cancer management.[86] One
possible piece of low-hanging fruit for a clinical trial would be low- to medium-dose
lithium, or other GSK3B inhibitors, for men undergoing active surveillance (AS) for
advance of prostate cancer. From studies of AS outcomes, a large fraction of patients on
AS ultimately require invasive treatment, as reviewed by Dall'Era et al[87]. When this
need arises it typically comes after only a few years. Thus, a trial of lithium in this
context would produce statistically significant results in a short time and, especially for
lithium, would be relatively inexpensive. One of us (EJ) conducted an informal one-
person trial on himself after being diagnosed with prostate cancer in 2014, ingesting
lithium supplements sufficient to bring his blood lithium to .3-.4mM while undergoing
AS by Memorial Sloan Kettering Cancer Center. (MSK did not prescribe the lithium but
agreed to include lithium level measurement in periodic blood tests.) In October 2017 EJ
was told that there was no longer a need for AS, as his markers were not progressing.
One case, important as it is to EJ, does not have statistical significance. We need clinical
trials with significant numbers of people.
We will be happy to collaborate on further specific pathway or network analysis
relevant to any of the cancers for which lithium, or other more specific drugs targeting
lithium-sensitive genes, may be a promising component of therapy.

350   **Author Contributions**
351   The work was planned jointly in conversations between EJ and WG. WG did the
352 computations and prepared the figures and tables. WG wrote the first draft of the
353 Methods and Results sections. EJ wrote the first draft of the Introduction and
354 Conclusions sections. Both authors shared in the final refinement of the manuscript.

355   **Conflict of Interest**
356   The authors have no personal, professional, or financial relationships that could
357 be construed as a conflict of interest with the work described in this paper.

[1] Post, Robert M. "Treatment of bipolar depression: evolving recommendations." *Psychiatric Clinics* 39, no. 1 (2016): 11-33.

[2] Jakobsson, Eric, Orlando Argüello-Miranda, See-Wing Chiu, Zeeshan Fazal, James Kruczek, Santiago Nunez-Corrales, Sagar Pandit, and Laura Pritchet. "Towards a Unified Understanding of Lithium Action in Basic Biology and its Significance for Applied Biology." *The Journal of membrane biology* 250, no. 6 (2017): 587-604.

[3] Cohen, Y., A. Chetrit, P. Sirota, and B. Modan. "Cancer morbidity in psychiatric patients: influence of lithium carbonate treatment." *Medical Oncology* 15, no. 1 (1998): 32-36.

[4] Martinsson L, Westman J, Hällgren J, Ösby U, Backlund L. Lithium treatment and cancer incidence in bipolar disorder. Bipolar disorders. 2016 Feb 1;18(1):33-40.

[5] Huang, Ru-Yu, Kun-Pin Hsieh, Wan-Wen Huang, and Yi-Hsin Yang. "Use of lithium and cancer risk in patients with bipolar disorder: population-based cohort study." *The British Journal of Psychiatry* (2016): bjp-bp.

[6] Pottegård A, Ennis ZN, Hallas J, Jensen BL, Madsen K, Friis S. Long-term use of lithium and risk of colorectal adenocarcinoma: a nationwide case–control study. British journal of cancer. 2016 Mar 1;114(5):571-5.

[7] Mazor, Michal, Yoshiaki Kawano, Hanneng Zhu, Jonathan Waxman, and Robert M. Kypta. "Inhibition of glycogen synthase kinase-3 represses androgen receptor activity and prostate cancer cell growth." *Oncogene* 23, no. 47 (2004): 7882-7892.

[8] Sun, Aijing, Ilanchezian Shanmugam, Jiawu Song, Paul F. Terranova, J. Brantley Thrasher, and Benyi Li. "Lithium suppresses cell proliferation by interrupting E2F–DNA interaction and subsequently reducing S–phase gene expression in prostate cancer." *The Prostate* 67, no. 9 (2007): 976-988.

[9] Zhu, Qing, Jun Yang, Suxia Han, Jihong Liu, Jeffery Holzbeierlein, J. Brantley Thrasher, and Benyi Li. "Suppression of glycogen synthase kinase 3 activity reduces tumor growth of prostate cancer in vivo." *The Prostate* 71, no. 8 (2011): 835-845.

[10] https://clinicaltrials.gov/ct2/show/NCT02198859

[11] Duffy, David J., Aleksandar Krstic, Thomas Schwarzl, Desmond G. Higgins, and Walter Kolch. "GSK3 inhibitors regulate MYCN mRNA levels and reduce neuroblastoma cell viability through multiple mechanisms, including p53 and Wnt signaling." *Molecular cancer therapeutics* 13, no. 2 (2014): 454-467.

[12] Cao, Qi, Xin Lu, and You-Ji Feng. "Glycogen synthase kinase-3β positively regulates the proliferation of human ovarian cancer cells." *Cell research* 16, no. 7 (2006): 671-677.

[13] Novetsky, Akiva P., Dominic M. Thompson, Israel Zighelboim, Premal H. Thaker, Matthew A. Powell, David G. Mutch, and Paul J. Goodfellow. "Lithium and inhibition of GSK3β as a potential therapy for serous ovarian cancer."*International journal of gynecological cancer: official journal of the International Gynecological Cancer Society* 23, no. 2 (2013): 361.

[14] Li, H., Huang, K., Liu, X., Liu, J., Lu, X., Tao, K., Wang, G. and Wang, J., 2014. Lithium chloride suppresses colorectal cancer cell survival and proliferation through ROS/GSK-3β/NF-κB signaling pathway. *Oxidative medicine and cellular longevity*, *2014*.

[15] de Araujo, Wallace Martins, Bruno Kaufmann Robbs, Lilian G. Bastos, Waldemir F. de Souza, Flávia CB Vidal, João PB Viola, and Jose A. Morgado-Diaz. "PTEN overexpression cooperates with lithium to reduce the malignancy and to increase cell death by apoptosis via PI3K/Akt suppression in colorectal cancer cells." *Journal of cellular biochemistry* 117, no. 2 (2016): 458-469.

[16] Costabile, V., Duraturo, F., Delrio, P., Rega, D., Pace, U., Liccardo, R., Rossi, G.B., Genesio, R., Nitsch, L., Izzo, P. and De Rosa, M., 2015. Lithium chloride induces mesenchymal-to-epithelial reverting transition in primary colon cancer cell cultures. *international journal of oncology*, *46*(5), pp.1913-1923.

[17] Rookmaaker, Maarten B., Heleen AJM van Gerven, Roel Goldschmeding, and Walther H. Boer. "Solid renal tumours of collecting duct origin in patients on chronic lithium therapy." *Clinical kidney journal* 5, no. 5 (2012): 412-415.

[18] Zaidan, Mohamad, Fabien Stucker, Bénédicte Stengel, Viorel Vasiliu, Aurélie Hummel, Paul Landais, Jean-Jacques Boffa, Pierre Ronco, Jean-Pierre Grünfeld, and Aude Servais. "Increased risk of solid renal tumors in lithium-treated patients." *Kidney international* 86, no. 1 (2014): 184-190.

[19] Pottegård, Anton, Jesper Hallas, Boye L. Jensen, Kirsten Madsen, and Søren Friis. "Long-term lithium use and risk of renal and upper urinary tract cancers." *Journal of the American Society of Nephrology* (2015): ASN-2015010061.

[20] Maeng, Yong-Sun, Rina Lee, Boram Lee, Seung-il Choi, and Eung Kweon Kim. "Lithium inhibits tumor lymphangiogenesis and metastasis through the inhibition of TGFBIp expression in cancer cells." *Scientific reports* 6 (2016).

[21] Zhong, Zhenyu, Elsa Sanchez-Lopez, and Michael Karin. "Autophagy, inflammation, and immunity: a troika governing cancer and its treatment." *Cell* 166, no. 2 (2016): 288-298.

[22] Sarkar, Sovan, R. Andres Floto, Zdenek Berger, Sara Imarisio, Axelle Cordenier, Matthieu Pasco, Lynnette J. Cook, and David C. Rubinsztein. "Lithium induces autophagy by inhibiting inositol monophosphatase." *J Cell Biol* 170, no. 7 (2005): 1101-1111.

[23] Motoi, Yumiko, Kohei Shimada, Koichi Ishiguro, and Nobutaka Hattori. "Lithium and autophagy." *ACS chemical neuroscience* 5, no. 6 (2014): 434-442.

[24] Sleire, Linda, Hilde Elisabeth Førde-Tislevoll, Inger Anne Netland, Lina Leiss, Bente Sandvei Skeie, and Per Øyvind Enger. "Drug repurposing in cancer." *Pharmacological research* 124 (2017): 74-91.

[25] Avissar, Sofia, Dennis L. Murphy, and Gabriel Schreiber. "Magnesium reversal of lithium inhibition of β-adrenergic and muscarinic receptor coupling to G proteins." *Biochemical pharmacology* 41, no. 2 (1991): 171-175.

[26] Mota de Freitas, Duarte, M. Margarida CA Castro, and Carlos FGC Geraldes. "Is competition between Li+ and Mg2+ the underlying theme in the proposed mechanisms for the pharmacological action of lithium salts in bipolar disorder?." *Accounts of chemical research* 39, no. 4 (2006): 283-291.

[27] Yoshikawa, Tomoko, and Sato Honma. "Lithium lengthens circadian period of cultured brain slices in area specific manner." *Behavioural Brain Research* 314 (2016): 30-37.

[28] Bauer, Michael, and Michael Gitlin. "What Is Lithium and How Does It Work?." In *The Essential Guide to Lithium Treatment*, pp. 33-43. Springer International Publishing, 2016.

[29] Wang, Hoau-Yan, and Eitan Friedman. "Effects of lithium on receptor-mediated activation of G proteins in rat brain cortical membranes." *Neuropharmacology* 38, no. 3 (1999): 403-414.

[30] Dorsam, Robert T., and J. Silvio Gutkind. "G-protein-coupled receptors and cancer." *Nature reviews cancer* 7, no. 2 (2007): 79.

[31] Haimovich, A., Eliav, U. and Goldbourt, A., 2012. Determination of the lithium binding site in inositol monophosphatase, the putative target for lithium therapy, by magic-angle-spinning solid-state NMR. *Journal of the American Chemical Society*, *134*(12), pp.5647-5651.

[32] Stambolic, Vuk, Laurent Ruel, and James R. Woodgett. "Lithium inhibits glycogen synthase kinase-3 activity and mimics wingless signalling in intact cells." *Current Biology* 6, no. 12 (1996): 1664-1669.

[33] Klein, Peter S., and Douglas A. Melton. "A molecular mechanism for the effect of lithium on development." *Proceedings of the National Academy of Sciences* 93, no. 16 (1996): 8455-8459.

[34] Ryves, W. Jonathan, and Adrian J. Harwood. "Lithium inhibits glycogen synthase kinase-3 by competition for magnesium." *Biochemical and biophysical research communications* 280, no. 3 (2001): 720-725.

[35] Freland, Laure, and Jean-Martin Beaulieu. "Inhibition of GSK3 by lithium, from single molecules to signaling networks." *Front Mol Neurosci* 5 (2012).

[36] Doble, Bradley W., Satish Patel, Geoffrey A. Wood, Lisa K. Kockeritz, and James R. Woodgett. "Functional redundancy of GSK-3α and GSK-3β in Wnt/β-catenin signaling shown by using an allelic series of embryonic stem cell lines." *Developmental cell* 12, no. 6 (2007): 957-971.

[37] Hoeflich, K. P., Luo, J., Rubie, E. A., Tsao, M. S., Jin, O., and Woodgett, J. R. (2000). Requirement for glycogen synthase kinase-3beta in cell survival and NF-kappaB activation. *Nature* 406, 86–90

[38] MacAulay, K., Doble, B. W., Patel, S., Hansotia, T., Sinclair, E. M., Drucker, D. J., Nagy, A., and Woodgett, J. R. (2007). Glycogen synthase kinase 3alpha-specific regulation of murine hepatic glycogen metabolism. *Cell Metab.* 6, 329–337

[39] Grimes, Carol A., and Richard S. Jope. "The multifaceted roles of glycogen synthase kinase 3β in cellular signaling." *Progress in neurobiology* 65, no. 4 (2001): 391-426.

[40] Beurel, Eléonore, and Richard S. Jope. "The paradoxical pro-and anti-apoptotic actions of GSK3 in the intrinsic and extrinsic apoptosis signaling pathways." *Progress in neurobiology* 79, no. 4 (2006): 173-189.

[41] Juhaszova, Magdalena, Dmitry B. Zorov, Suhn-Hee Kim, Salvatore Pepe, Qin Fu, Kenneth W. Fishbein, Bruce D. Ziman et al. "Glycogen synthase kinase-3β mediates convergence of protection signaling to inhibit the mitochondrial permeability transition pore." *The Journal of clinical investigation* 113, no. 11 (2004): 1535-1549.

[42] Beals, Chan R., Colleen M. Sheridan, Christoph W. Turck, Phyllis Gardner, and Gerald R. Crabtree. "Nuclear export of NF-ATc enhanced by glycogen synthase kinase-3." *Science* 275, no. 5308 (1997): 1930-1933.

[43] Beurel, Eléonore, Suzanne M. Michalek, and Richard S. Jope. "Innate and adaptive immune responses regulated by glycogen synthase kinase-3 (GSK3)." *Trends in immunology* 31, no. 1 (2010): 24-31.

[44] Hardt, Stefan E., and Junichi Sadoshima. "Glycogen synthase kinase-3β a novel regulator of cardiac hypertrophy and development." *Circulation research* 90, no. 10 (2002): 1055-1063.

[45] McQuillin, Andrew, Mie Rizig, and Hugh MD Gurling. "A microarray gene expression study of the molecular pharmacology of lithium carbonate on mouse brain mRNA to understand the neurobiology of mood stabilization and treatment of bipolar affective disorder." *Pharmacogenetics and genomics* 17, no. 8 (2007): 605-617.

[46] O'Brien, W. Timothy, Amber DeAra Harper, Fernando Jové, James R. Woodgett, Silvia Maretto, Stefano Piccolo, and Peter S. Klein. "Glycogen synthase kinase-3β haploinsufficiency mimics the behavioral and molecular effects of lithium." *The Journal of neuroscience* 24, no. 30 (2004): 6791-6798.

[47] Mendes, Camila Teixeira, Fábio Borges Mury, Eloísa de Sá Moreira, Fernando Lopes Alberto, Orestes Vicente Forlenza, Emmanuel Dias-Neto, and Wagner Farid Gattaz. "Lithium reduces Gsk3b mRNA levels: implications for Alzheimer disease." *European archives of psychiatry and clinical neuroscience* 259, no. 1 (2009): 16-22.

[48] Beurel, Eleonore, Steven F. Grieco, and Richard S. Jope. "Glycogen synthase kinase-3 (GSK3): regulation, actions, and diseases." *Pharmacology & therapeutics* 148 (2015): 114-131.

[49] Linding, Rune, Lars Juhl Jensen, Gerard J. Ostheimer, Marcel ATM van Vugt, Claus Jørgensen, Ioana M. Miron, Francesca Diella et al. "Systematic discovery of in vivo phosphorylation networks." *Cell* 129, no. 7 (2007): 1415-1426.

[50] Piovesan, Damiano, Giuseppe Profiti, Pier Luigi Martelli, and Rita Casadio. "The human" magnesome": detecting magnesium binding sites on human proteins." *BMC bioinformatics* 13, no. 14 (2012): 1.

[51] Martinez, Ana, Ana Castro, Isabel Dorronsoro, and Mercedes Alonso. "Glycogen synthase kinase 3 (GSK-3) inhibitors as new promising drugs for diabetes, neurodegeneration, cancer, and inflammation." *Medicinal research reviews* 22, no. 4 (2002): 373-384.

[52] Luo, Jia. "Glycogen synthase kinase 3β (GSK3β) in tumorigenesis and cancer chemotherapy." *Cancer letters* 273, no. 2 (2009): 194-200.

[53] Ougolkov, Andrei V., Martin E. Fernandez-Zapico, Doris N. Savoy, Raul A. Urrutia, and Daniel D. Billadeau. "Glycogen synthase kinase-3β participates in nuclear factor κB–mediated gene transcription and cell survival in pancreatic cancer cells." *Cancer research* 65, no. 6 (2005): 2076-2081.

[54] Ding, Qingqing, Xianghuo He, Jung-Mao Hsu, Weiya Xia, Chun-Te Chen, Long-Yuan Li, Dung-Fang Lee et al. "Degradation of Mcl-1 by β-TrCP mediates glycogen synthase kinase 3-induced tumor suppression and chemosensitization." *Molecular and cellular biology* 27, no. 11 (2007): 4006-4017.

[55] Skov, Søren, Marianne Terndrup Pedersen, Lars Andresen, Per Thor Straten, Anders Woetmann, and Niels Ødum. "Cancer cells become susceptible to natural killer cell killing after exposure to histone deacetylase inhibitors due to glycogen synthase kinase-3–dependent expression of MHC class I–related chain A and B." *Cancer research* 65, no. 23 (2005): 11136-11145.

[56] Luo, Jia. "Glycogen synthase kinase 3β (GSK3β) in tumorigenesis and cancer chemotherapy." *Cancer letters* 273, no. 2 (2009): 194-200.

[57] Liao, Xinbo, J. Brantley Thrasher, Jeffery Holzbeierlein, Scott Stanley, and Benyi Li. "Glycogen synthase kinase-3β activity is required for androgen-stimulated gene expression in prostate cancer." *Endocrinology* 145, no. 6 (2004): 2941-2949.

[58] Liao, Xinbo, Liping Zhang, J. Brantley Thrasher, Jie Du, and Benyi Li. "Glycogen synthase kinase-3β suppression eliminates tumor necrosis factor-related apoptosis-inducing ligand resistance in prostate cancer." *Molecular cancer therapeutics* 2, no. 11 (2003): 1215-1222.

[59] Sun, Aijing, Changlin Li, Ruibao Chen, Yiling Huang, Qi Chen, Xiangjun Cui, Huafeng Liu, J. Brantley Thrasher, and Benyi Li. "GSK-3β controls autophagy by modulating LKB1-AMPK pathway in prostate cancer cells." *The Prostate* 76, no. 2 (2016): 172-183.

[60] Darrington, R. Siobhan, Victor M. Campa, Marjorie M. Walker, Nora Bengoa-Vergniory, Irantzu Gorrono-Etxebarria, Pinar Uysal-Onganer, Yoshiaki Kawano, Jonathan Waxman, and Robert M. Kypta. "Distinct expression and activity of GSK-3α and GSK-3β in prostate cancer." *International journal of cancer* 131, no. 6 (2012): E872-E883.

[61] Salim, Tavga, Anita Sjölander, and Janna Sand-Dejmek. "Nuclear expression of Glycogen synthase kinase-3β and lack of membranous β-catenin is correlated with poor survival in colon cancer." *International journal of cancer* 133, no. 4 (2013): 807-815.

[62] Miyashita, Katsuyoshi, Mitsutoshi Nakada, Abbas Shakoori, Yasuhito Ishigaki, Takeo Shimasaki, Yoshiharu Motoo, Kazuyuki Kawakami, and Toshinari Minamoto. "An emerging strategy for cancer treatment targeting aberrant glycogen synthase kinase 3β." *Anti-Cancer Agents in Medicinal Chemistry (Formerly Current Medicinal Chemistry-Anti-Cancer Agents)* 9, no. 10 (2009): 1114-1122.

[63] Walz, Amy, Andrey Ugolkov, Sunandana Chandra, Alan Kozikowski, Benedito A. Carneiro, Thomas V. O'Halloran, Francis J. Giles, Daniel D. Billadeau, and Andrew P. Mazar. "Molecular pathways: Revisiting glycogen synthase kinase-3β as a target for the treatment of cancer." *Clinical Cancer Research* (2017).

[64] Gout, Elisabeth, Fabrice Rébeillé, Roland Douce, and Richard Bligny. "Interplay of Mg2+, ADP, and ATP in the cytosol and mitochondria: unravelling the role of Mg2+ in cell respiration." *Proceedings of the National Academy of Sciences* 111, no. 43 (2014): E4560-E4567.

[65] Briggs, Katharine T., Gary G. Giulian, Gong Li, Joseph PY Kao, and John P. Marino. "A Molecular Model for Lithium's Bioactive Form." *Biophysical Journal* 111, no. 2 (2016): 294-300.

[66] Willis, J. S., and L. S. T. Fang. "Li+ stimulation of ouabain-sensitive respiration and (Na+-K+)-ATPase of kidney cortex of ground squirrels." *Biochimica et Biophysica Acta (BBA)-Biomembranes* 219, no. 2 (1970): 486-489.

[67] Jope, Richard S. "Lithium and GSK-3: one inhibitor, two inhibitory actions, multiple outcomes." *Trends in pharmacological sciences* 24, no. 9 (2003): 441-443.

[68] Hong, Ming, Daniel CR Chen, Peter S. Klein, and Virginia M-Y. Lee. "Lithium reduces tau phosphorylation by inhibition of glycogen synthase kinase-3." *Journal of Biological Chemistry* 272, no. 40 (1997): 25326-25332.

[69] Kitano, Hiroaki. "Systems biology: a brief overview." *Science* 295, no. 5560 (2002): 1662-1664.

[70] Kanehisa, Minoru, Miho Furumichi, Mao Tanabe, Yoko Sato, and Kanae Morishima. "KEGG: new perspectives on genomes, pathways, diseases and drugs." *Nucleic acids research* 45, no. D1 (2016): D353-D361.

[71] Szklarczyk, Damian, John H. Morris, Helen Cook, Michael Kuhn, Stefan Wyder, Milan Simonovic, Alberto Santos et al. "The STRING database in 2017: quality-controlled protein–protein association networks, made broadly accessible." *Nucleic acids research* (2016): gkw937.

[72] Zhang JD and Wiemann S (2009). "KEGGgraph: a graph approach to KEGG PATHWAY in R and Bioconductor." *Bioinformatics*, pp. 1470–1471.

[73] Zhang JD (2017). *KEGGgraph: Application Examples*. R package version 1.38.0.

[74] Franceschini, A (2013). "STRING v9.1: protein-protein interaction networks, with increased coverage and integration." *Nucleic Acids Research (Database issue)*, **41**

[75] Huang, Da Wei, Brad T. Sherman, and Richard A. Lempicki. "Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists." *Nucleic acids research* 37, no. 1 (2008): 1-13.

[76] Thomas, P. D. "Expansion of the gene ontology knowledgebase and resources: the gene ontology consortium." *Nucleic Acids Res* 45 (2017): D331-D338.

[77] Ge, Weihao, Zeeshan Fazal, and Eric Jakobsson. "Using Optimal F-Measure and Random Resampling in Gene Ontology Enrichment Calculations." *bioRxiv* (2017): 218248.

[78] Wang, Lei, Xi-Xi Cao, Qi Chen, Teng-Fang Zhu, Hong-Guang Zhu, and Li Zheng. "DIXDC1 targets p21 and cyclin D1 via PI3K pathway activation to promote colon cancer cell proliferation." *Cancer science* 100, no. 10 (2009): 1801-1808.

[79] Xu, Zhonghai, Di Liu, Chuifeng Fan, Lan Luan, Xiupeng Zhang, and Enhua Wang. "DIXDC1 increases the invasion and migration ability of non-small-cell lung cancer cells via the PI3K-AKT/AP-1 pathway." *Molecular carcinogenesis* 53, no. 11 (2014): 917-925.

[80] Tan, Cong, Fan Qiao, Ping Wei, Yayun Chi, Weige Wang, Shujuan Ni, Qifeng Wang et al. "DIXDC1 activates the Wnt signaling pathway and promotes gastric cancer cell invasion and metastasis." *Molecular carcinogenesis* 55, no. 4 (2016): 397-408.

[81] Zhong, Jiateng, Yufei Liu, Qingli Xu, Jian Yu, and Muchun Zhang. "Inhibition of DIXDC1 by microRNA-1271 suppresses the proliferation and invasion of prostate cancer cells." *Biochemical and biophysical research communications* 484, no. 4 (2017): 794-800.

[82] Quan, Junjie, Jianqiang Qu, and Le Zhou. "MicroRNA-539 inhibits glioma cell proliferation and invasion by targeting DIXDC1." *Biomedicine & Pharmacotherapy* 93 (2017): 746-753.

[83] Li, Xiaohong, Ying Xiao, Shaoqing Fan, Mingbing Xiao, Xiaotong Wang, Xiaolin Zhu, Xudong Chen et al. "Overexpression of DIXDC1 correlates with enhanced cell growth and poor prognosis in human pancreatic ductal adenocarcinoma." *Human pathology* 57 (2016): 182-192.

[84] Loughin, Thomas M. "A systematic comparison of methods for combining p-values from independent tests." *Computational statistics & data analysis* 47, no. 3 (2004): 467-485.

[85] Aprahamian, Ivan, Franklin S. Santos, Bernardo dos Santos, Leda Talib, Breno S. Diniz, Márcia Radanovic, Wagner F. Gattaz, and Orestes V. Forlenza. "Long-term, low-dose lithium treatment does not impair renal function in the elderly: a 2-year randomized, placebo-controlled trial followed by single-blind extension." *The Journal of clinical psychiatry* 75, no. 7 (2014): 672-678.

[86] Li, Benyi, James Brantley Thrasher, and Paul Terranova. "Glycogen synthase kinase-3: A potential preventive target for prostate cancer management." In *Urologic Oncology: Seminars and Original Investigations*, vol. 33, no. 11, pp. 456-463. Elsevier, 2015.

[87] Dall'Era, Marc A., Peter C. Albertsen, Christopher Bangma, Peter R. Carroll, H. Ballentine Carter, Matthew R. Cooperberg, Stephen J. Freedland, Laurence H. Klotz, Christopher Parker, and Mark S. Soloway. "Active surveillance for prostate cancer: a systematic review of the literature." *European urology* 62, no. 6 (2012): 976-983.