Hydrophobic organic contaminants are not linked to microplastic uptake in Baltic Sea herring

*Ogonowski M.[1,2], Wenman, V.[1], Danielsson S.[3] and Gorokhova E.[1]

[1] Department of Environmental Science and Analytical Chemistry, Stockholm University, SE-10691,

Stockholm, Sweden

[2] AquaBiota Water Research, Löjtnantsgatan 25, SE-115 50, Stockholm, Sweden

[3] Swedish Museum of Natural History, Department of Environmental Science and Monitoring, P. O. Box

50 007, SE-104 05, Stockholm Sweden

*corresponding author:

martin.ogonowski@aces.su.se

Declarations of interest: none

Abstract

It is commonly accepted that microplastic (MP) ingestion can lead to lower food intake and bioaccumulation of hydrophobic organic contaminants (HOCs) in aquatic organisms. However, causal links between MP and contaminant levels in biota are poorly understood and *in situ* data are virtually non-existent. Here, we investigated whether HOC concentrations in herring muscle tissue (*Clupea harengus membras*) are related to MP ingestion using fish caught along the West coast of the Baltic Sea. The MP occurrence exhibited a large geographic variability, with MP found in 28% of the fish examined. The population average was 2.4 MP ind^{-1}; however, when only individuals containing MP were considered, the average MP burden was 8.4 MP ind^{-1}. We also found that MP burden decreased with reproductive stage of the fish but increased with its body size. To predict MP abundance in fish guts, we constructed a mass-balance model using literature data on MP in the water column and physiological rates on ingestion and gut evacuation for clupeids of a similar size. The model output was in agreement with the observed values, thus supporting the validity of the results. Contaminant concentrations in the muscle tissue were unrelated to the MP levels in fish, suggesting a lack of direct links between the levels of HOCs and MP ingestion. Thus, despite their ubiquity, MP are unlikely to have a measurable impact on food intake or the total body burden of hydrophobic contaminants in Baltic herring.

Keywords: *Microplastic, Baltic Sea, herring, hydrophobic organic contaminants, marine monitoring*

Introduction

Plastic debris, including microplastics (MP < 5 mm), can be ingested by various aquatic organisms across several trophic levels (Lusher et al. 2013, 2015, Cole et al. 2013). Due to the importance of commercial fish and shellfish species for human consumption, the ingestion and presence of MP in these animals has become a matter of concern (EFSA Panel on Contaminants in the Food Chain (CONTAM) 2016). To address this concern and to provide a quantitative assessment of MP ingestion in various fish species, an active research is ongoing (Lusher et al. 2013, Foekema et al. 2013, Rummel et al. 2016, Budimir et al. 2018, Beer et al. 2018).

A commonly held paradigm states that MP ingestion can lead to decreased nutritional status (Cole et al. 2015, Ogonowski et al. 2016) and bioaccumulation of hydrophobic organic chemicals (HOCs) (Oliveira et al. 2012, Besseling et al. 2013, Rochman et al. 2013, Wardrop et al. 2016) that sorb to the MP particles in the water and desorb in the gut lumen (Mato et al. 2001, Rusina et al. 2010, Rochman et al. 2014). However, some experimental and modeling studies indicate that plastic polymers could also have a net cleaning effect acting as passive samplers while in the digestive system and thereby relieve the animals of HOCs (Gouin et al. 2011, Herzke et al. 2016, Koelmans et al. 2016). The relative importance of microplastics as vectors for contaminant transport remains unresolved, possibly also due to the lack of field data linking HOC concentrations in biota to ingested MP.

Here, we studied MP ingestion by Baltic Sea herring (*Clupea harengus membras* L.), a commercially exploited fish and a keystone species in the Baltic food web. Being facultative pelagic filter-feeders (Huse and Toresen 1996), herring stand a high risk of ingesting MP along with zooplankton prey and hence accumulating MP-associated contaminants. It is also a sentinel species in the Swedish National Monitoring Program for Contaminants in Marine Biota and, thus, a potential indicator species for MP monitoring in the Baltic Sea (Beer et al. 2018).

If MP ingestion indeed contributes significantly to HOC bioaccumulation in contaminated environments, then one would see a positive correlation between the amount of MP ingested over time and HOC concentrations in the herring tissues. However, there are no reliable methods to estimate accumulated MP exposure using field samples, because MP do not accumulate to any significant extent in the fish digestive system (Lusher et al. 2013, Jovanović 2017). Although gut contents reflect only a recent ingestion history (Ahlbeck et al. 2012), the MP burden determined by gut content analysis is commonly used as a reflection of the feeding habits and habitats of the

fish. Another area of concern with respect to the interpretation of MP counts in environmental

samples, including fish guts, is analytical accuracy and reliability of MP extraction and

determination (Dehaut et al. 2016). Therefore, to increase the reliability of the MP gut content

data, it is important to verify whether the recorded MP body burden is within ecologically plausible

rates of ingestion and gut evacuation. To compare the observed MP abundance in the fish gut with

the intake that can be expected given the MP abundance in the water column, and gut evacuation

that can be expected given the food intake, we applied a mass-balance modelling. In this model,

using literature-derived parameters on clupeid feeding and food processing as well as ambient MP

concentrations, we estimated MP burden in the herring with the body size similar to those in our

collection. Finally, we evaluated whether HOC concentrations in the fish muscle were related to

the weight-specific MP gut content of the same individual.

Materials and Methods

Fish collection and sample characteristics

The Baltic herring used for our analyses were collected by the Swedish National Monitoring

Program for Contaminants in Marine Biota conducted by the Swedish Museum of Natural History

(Stockholm, Sweden). To avoid possible bias by known point sources, we randomly selected 130

specimens that had been collected at thirteen reference monitoring stations (Fig. 1), thus covering

a sufficiently large geographical area that would provide a representative range of HOC and MP

exposure for the analysis.

The sex ratio of the selected fish was approximately 50:50 and uniform across sampling sites. The

individuals were 3-7 years old, with a total length of 173 ± 18 mm and body weight 35 ± 12 g

(mean $\pm$ SD). The reproductive phase determined by gametocytic maturity was classified on a

five-degree scale according to Bucholtz et al. (2008) and included deformed gonads (stage 1), post

spawned individuals (stage 2), juveniles (stage 3), individuals with developing gonads (stage 4)

and fish with mature gonads (stage 5). Each fish was dissected, and the muscle tissues taken from

the middle dorsal muscle layer were used for HOC analysis, whereas the entire gastrointestinal

tract (GIT) was used for the MP analysis. All sampling was performed according to standard

procedures (TemaNord 1995). After dissecting, each individual GIT was packed in aluminum foil

to avoid cross-contamination. All GIT samples were immediately frozen at -20 °C and stored until

MP analysis at the Department of Environmental Science and Analytical Chemistry, Stockholm

University, Sweden.

MP quantification in the gastrointestinal tract of fish

Each GIT was placed in a glass Petri dish, opened with surgical scissors, and rinsed with deionized, particle-free water. Using a stereo microscope, the bolus was examined, and any items resembling MP were extracted by stainless steel pincers and transferred to clean Eppendorf tubes. The appearance of the putative MP was recorded and each particle was categorized according to its shape (fiber or fragment) and color. Hereafter, the number of MP per individual fish is referred to as *MP burden*.

To relate HOC concentration in the muscle tissue and MP intake by fish, the fish size must be taken into account when expressing MP counts. Moreover, the bolus size varied considerably among the individual fish and geographical areas (Supporting Information Table S1), indicating variability in feeding activity shortly before sampling and/or gut evacuation that might have been related to stress during the sampling. To account for this variability and the corresponding variation in the MP burden, we normalized the individual MP counts to its gut fullness; the latter was assessed by visual observation on a five-step semi-quantitative scale: 0 (empty, no food items), 0.25, 0.5 0.75 or 1 (full). The obtained values were further normalized to the individual body weight and termed *weight-specific MP burden* [number of MP / (gut fullness × body weight) (g wet weight)]. This allowed relating HOC concentrations in the fish to the expected MP burden in the GIT on a weight basis.

To identify whether the putative MP were synthetic polymers, we followed the recommendations of Norén (2007) and Hidalgo-Ruz et al. (2012). Particles 1-5 mm in diameter were recorded and classified as MP, if all the following criteria were met: (i) uniform, unnaturally bright or of an unnatural color, (ii) lack of organic structures, and (iii) uniform diameter over the entire length of a fiber. To test the accuracy of the visual identification, a random subset of 20 samples containing MP (i.e., the gut contents of 20 individual fish) was analyzed using the hot needle test (De Witte et al. 2014).

Controls and blanks

To prevent contamination by airborne particles during the examination, the dissections were performed under a Fumex local extractor (Wesch et al. 2016); each sample being analyzed for 10 min. A Petri dish filled with filtered deionized water was placed next to a test sample to serve as a

blank for the quantification and characterization of potential contamination during the analysis. When working with samples, a cotton lab coat and nitrile gloves were used: moreover, the type and color of clothing were recorded to enable contamination back-tracing. All procedural blanks contained plastic particles (mainly single fibers) of unknown origin. However, all these particles were < 1 mm and thus did not contribute to the MP counts used in the statistical analysis. If quantifiable amounts of blank contamination with particles > 1 mm were to be found, such samples would be excluded from any further analyses.

Chemical analysis

Following the guidelines of the Swedish National Monitoring Program for Contaminants in Marine Biota, the muscle samples were analyzed for polychlorinated biphenyls (PCB 28, 52, 101, 118, 138, 153 and 180), organochlorine pesticides (DDE, DDD, DDT, HCB, AHCH, BHCH, and Lindane) and polybrominated flame retardants (BDE 28, 47, 99, 100, 153, 154 and HBCD). For most compounds, 10 g of muscle tissue from individual fish were used, whereas 1 g samples of muscle tissue from 10 individuals were pooled for a few analytically challenging compounds. An overview of the analyzed contaminants and their average concentrations in herring muscle tissue are provided in Table 1, while details of the analytical procedures and quality assurance are provided elsewhere (Bignert et al. 2016).

Data analysis and statistics

Relationships between biological factors, geography and ingested microplastic

We used generalized additive models (GAM) in package *mgcv* to examine relationships between specific biological variables (*weight*, *gut fullness*, *age* and *reproductive phase*) and *MP burden*; *sea basin* was used as a random factor in the model since the inclusion of this term lowered the Akaike Information Criterion from 570 to 417. The model was specified as:

MP burden = β s(weight) + s(gut fullness) + s(age) + s(reproductive phase) + random(sea basin) + ε

The multicollinearity between the explanatory variables was evaluated as low (< 0.37) using concurvity measures (Amodio et al. 2014) calculated by the *mgcv*-package. Due to the

overrepresentation of zeros in the data (overdispersion) for the MP burden, the model was run using zero-inflated Poisson error structures. Model performance was assessed using residual plots. Differences in the MP burden between the basins were tested using Permanova with *station* nested within *basin* as a random factor (Anderson 2001). The significance level was set at $\alpha = 0.05$; all statistical analyses were conducted in R 3.5.0 (R Core Team 2014).

Relationships between HOCs and ingested microplastic

Maximum-likelihood Factor Analysis with Varimax rotation was used to assess the degree of association between the chemical variables and weight-specific MP burden in the GIT. Prior to the analysis, Bartlett's test of sphericity was performed to confirm patterned relationships between the variables and was statistically significant ($\chi^2_{15} = 175$, $p < 0.0001$). A scree plot was used to determine the number of factors to retain, and factor loadings > 0.7 were considered statistically significant (MacCallum et al. 2001). When measured values were below the limit of quantification (LOQ), they were imputed by LOQ divided by the square root of two (Succop et al. 2004). The analyzed chemical concentrations were summed and grouped into their respective contaminant groups (PCBs, PBDEs and organochlorine pesticides).

Modeling plastic ingestion by herring

To evaluate whether the observed MP burden could be predicted using ambient MP abundance data and food processing rates, we modeled the ingestion of MP using literature-derived parameters on food uptake, egestion, and MP abundance in the study area. The rationale is that observed MP abundance in the gut would reflect average exposure levels assuming that (1) MP concentrations are fairly homogeneous in the outer coastal areas (Gorokhova 2015, Gewert et al. 2017), which are the main feeding grounds of herring (Flinkman et al. 1998), (2) the MP abundance in the water column, where the fish feed, is similar to that at the surface, where the data on the relevant size fraction of MP (1- 5 mm) were collected; (3) MP ingestion by herring is non-selective and thus proportional to the MP abundance in the water, and (4) gut evacuation rates are non-discriminatory, i.e., MP are egested at the same rate as prey remains. Then, the MP burden (MP ind^{-1}) at any given time, t, can be written as the mass balance between the uptake and loss rates (Eq. 1):

$$MP_t = MP(t - dt) + (IR - Eg)\,dt, \qquad\qquad (1)$$

where *IR* and *ER* are the ingestion and egestion rates (MP h^{-1}), respectively. They can be calculated

as:

$IR = CMP \times CR$ (2)

and

$ER = GER \times MP_t$,   (3)

where *CMP* is the ambient MP concentration (number of MP L^{-1}), *CR* is the clearance rate (L h^{-1};

the volume of water swept clear of particles per individual and hour), and *GER* is the gut

evacuation rate (h^{-1}).

We used literature data to parameterize the model (Fig. S1, Table S2). The MP concentrations in

the target size range (1-5 mm) from surface waters in the outer Stockholm archipelago (Gewert

et al. 2017) were used as CMP values. Clearance rates were estimated using reported feeding

rates for North Sea herring on *Calanus finmarchicus,* a copepod of similar size as the

microplastics considered here, and the main prey for herring (Varpe and Fiksen 2010) (see

Supporting Information 1.1 for derivation of CR). As published gut evacuation rates for adult

herring were not available, we used experimental values reported for other clupeids of similar

size, European pilchard (*Sardina pilchardus*) (Costalago and Palomera 2014) and South

American pilchard (*Sardinops sagax*) (van der Lingen 1998), which have similar feeding

ecology and physiology as Baltic herring (Collard et al. 2017). The physiological rates used in

the model corresponded to the average size of our fish.

The model was implemented using STELLA® ver. 9.4.1 software (iSee systems, Inc. Lebanon,

NH, U.S.A.) to estimate MP burden (MP ind^{-1}) dynamics in a fish population at a given MP

abundance. The intrapopulation variability was simulated using a Monte Carlo generator with

1000 permutations (details on the simulation settings are provided in Supporting information 1.2).

To validate the model, we compared the simulated data distribution from the model to the field

data using descriptive statistics, χ^2 and the two-sample Cramér-von Mises tests.

Results

Observed MP burden

Particles identified by visual inspection as MP were found in 44 out of the 130 individuals (33.8%;

range: 0 to 51 pieces of plastic fiber and/or fragments ind^{-1}). In those 44 individuals, the mean

abundance was 7.8 ± 12.2 particles ind^{-1} (± SD). The dominant type of MP were fibers of various

colors (87.6%), while fragments were less frequent (12.4%). However, the black fibers were

identified as non-plastic by the hot needle test and were, therefore, excluded from further analyses.

When the black fibers were omitted, only 37 individuals contained MP (28.4 %; range: 0 to 51

MP) with a mean abundance of 8.4 MP ind^{-1} ± 12.8. The proportion of the fibers was 86.2%. When

all examined individual were considered, the population average was 2.4 MP ind^{-1} with the 95%

bootstrap confidence interval ranging 1.4 - 4.2 MP ind^{-1}. The variation in the MP burden between

the stations and basins was high (Fig. 2, Supporting Information Table S1), and we did not find

any significant differences in the MP burden between the basins (*station* nested within *basin* as a

random factor, pseudo $F_{4,117} = 1.1$, $p = 0.37$).

Predicted vs. observed MP burden and frequency of occurrence

The model predicted that 81% of fish contained MP, with a mean MP burden of 4.7 MP ind^{-1};

these values were about twice as high as the observed values. The ranges of the frequency

distributions for the simulated and observed values were overlapping, although the field

observations were more strongly skewed towards zero values compared to the model prediction

(Fig. 3 A, Supporting Information Table S3). The difference between the distributions was

statistically significant (Cramér-von Mises $T = 144$, $p < 0.0001$). When zeros were excluded, the

distributions, albeit still significantly different (Cramér-von Mises $T = 23.4$, $p < 0.0001$), became

more similar (Fig. 3 B, Supporting Information Table S 3), indicating that much of the difference

between the distributions was driven by the significantly higher proportion of zero observations

in the field data ($\chi^2 = 162.3$, $p < 0.0001$).

Linkage between MP intake and HOCs

We found no relationship between the weight-specific MP burden and the concentration of any of

the HOCs (Fig. 4). As a variable, weight-specific MP burden loaded weakly and negatively (-0.13)

on the first axis and moderately positive (0.62) on the second axis. In contrast, the organochlorine

pesticides and PBDEs loaded significantly and positively on the first axis, while the PCBs loaded

moderately positive (0.55) on the first and significantly positive (0.83) on the second axis.

Together, the two factors explained a cumulative variance of 85.1% (Supporting Information Table

S4). Hence, no contaminant group had loadings clustering with those for the weight-specific MP

burden.

Biological factors related to MP burden

The MP burden was positively and nearly linearly related to fish *body weight* (GAM, $\chi^2 = 26.5$, p
< 0.0001, Fig. 5 A). In contrast, a negative effect was found for *reproductive phase*, where MP
burden was significantly lower in fish that had reached sexual maturity (GAM $\chi^2 = 56.4$, p <
0.0001, Fig. 5 B). *Gut fullness* only had a negative effect on MP burden when the GIT was empty
of food items (GAM $\chi^2 = 53.6$, p < 0.0001, Fig. 5 C) while *Age* displayed a weak negative
relationship with MP burden (GAM $\chi^2 = 6.8$, p < 0.01, Fig. 5 D). However, albeit statistically
significant, the effect of *Age* was not particularly strong and most probably of low biological
importance.

Discussion

Microplastics are not very common in herring guts

Microplastics (mostly fibers) were found in about one third of the fish. While these values are in
good agreement with those reported for herring by Beer et al. (2018) for the central Baltic Sea
(20% containing MP, with 93% fibers), other studies found considerably lower MP frequency of
occurrence and fiber contribution to total MP in herring. Both Foekema et al. (2013) and Rummel
et al. (2016) found plastics in only 2% of herring samples from the North Sea and the Southern
Baltic Sea, with fibers accounting for less than 10% of MP. Having excluded fibers from their
analyses, Budimir et al. (2018) reported a frequency of occurrence as low as 1.8% in herring from
the northern Baltic Sea. These discrepancies between different studies could be related to
differences in fish size and gut fullness. For example, Foekema et al. (2013) used fish that were
considerably larger (>200 mm total length) which most likely already had switched from filter
feeding to raptorial feeding on larger prey (Huse and Toresen 1996). This change in feeding mode
would result in a lower ingestion rate of zooplankton-sized plastic particles and thus in a lower
overall MP burden. In the study of Rummel et al. (2016), many fish stomachs were empty, which
probably was related to arrested feeding in concert with spawning, and, possibly, stress-induced
gut evacuation caused by the fish sampling (Wilkins 1967, Vinson and Angradi 2011). This lends
further support to our findings that MP burden increases with fish size (cf. Beer et al. 2018) and
decreases with reproductive phase. In addition, one would expect the amount of ingested MP to
scale with the absolute size of gut bolus or gut fullness. However, since this relationship was weak

(Fig. 5 C), our findings only partly support this expectation. One possible explanation for this

could be slower egestion of MP compared to prey, similar to the selective retention of plastic fibers

in amphipods (Au et al. 2015) and fragments in cladocerans (Ogonowski et al. 2016), which would

result in a temporary accumulation of MP in the fish gut and obscure the expected positive

relationship between the gut fullness and MP burden. While fish size appears to be the strongest

covariable for standardizing internalized MP content, gut fullness was also influential, particularly

for fish with no food in the GIT, which may occur during periodic fasting (Darbyson et al. 2003).

The range of the MP burden predicted by our simple model was similar to that observed in the

field caught specimens, although the proportion of fish predicted to contain MP was more than

two-fold higher. This is, however, not surprising because, frequency of zero values was driven by

the CMP variability that was derived from surface collection of MP. However, contrary to the

model assumption of homogeneous MP distribution in the water column, it is likely that the MP

distribution is patchy, their concentrations vary with depth (Gorokhova 2015) and form aggregates

too large to be mistaken for food (Long et al. 2015, Lagarde et al. 2016). Therefore, the distribution

of MP concentrations originating from surface collections and used to model MP encounter rate

might not reflect the actual situation. The observed population MP burden was also more variable,

which is likely to be related to diel variations in feeding and gut evacuation under natural

conditions (Seyhan and Grove 2003) not accounted for by the model. Other biological factors,

such as maturity level, ontogenetic changes in feeding, and behavior, also may have affected the

probability of MP ingestion and thus contributed to intrapopulation variability in the MP burden.

Finally, fishing methods (which may induce gut evacuation) and time of capture (which may

reflect diurnal differences in feeding activity) may have contributed to the observed discrepancy

in the MP burden distribution. Nevertheless, given the simplicity of the model and the uncertainties

associated with its parameters, the predicted values were sufficiently close to those found in the

field, indicating that MP uptake can be predicted provided that we have reliable MP abundance

estimates.

No correlation between weight-specific MP burden and HOCs

The transfer of hydrophobic contaminants from ingested plastics to biota has been described as the

so-called "Trojan horse" effect (Cole et al. 2011). While this transfer has been demonstrated under

laboratory conditions (Besseling et al. 2013, Rochman et al. 2013, Browne et al. 2013, Batel et al.

2016), recent modelling studies indicate that natural sources play a much more important role than

MP in explaining HOC bioaccumulation patterns in aquatic organisms (Koelmans 2015, Koelmans

et al. 2016). We did not find any correlation between HOC concentrations in herring muscle and

MP burden, although it could be argued that omitting small MP (< 1 mm) from our analysis, could

have biased the results. Indeed, by focusing on the larger MP, we ignored potentially important

influence of a higher total surface area and thus higher HOC desorption rates (Hendriks et al. 2001,

Hartmann et al. 2017). However, the ingestion of such small particles by fish of this size is rather

unlikely, because filter-feeding herring have a relatively low capacity to retain small particles due

to their rather wide gill raker spacing (Gibson 1988, Collard et al. 2017) and actively avoid smaller

prey while feeding raptorially (Aro et al. 1989, Casini et al. 2004). In fact, this line of reasoning

has been supported by several other studies reporting a predominant retention of MP of > 1 mm

by similarly-sized herring (Lenz et al. 2016, Collard et al. 2017, Beer et al. 2018). Moreover, given

the short residence time (Grigorakis et al. 2017) of internalized plastics and the slow desorption

kinetics of many HOCs, the lack of correlation between the MP and organic contaminants is rather

expected and in line with other reports for fish and other aquatic animals (Herzke et al. 2016, Rehse

et al. 2018, Kleinteich et al. 2018).

Causality is difficult to prove using environmental samples, where many different parameters may

affect contaminant body burden of an organism (Hartmann et al. 2017), including various biotic

factors that have significant effects on both MP (this study) and HOC levels (Persson et al. 2013,

Silva Barni et al. 2014). However, our findings suggest that there is no tenable relationship

between the MP intake and tissue contaminant concentrations in the Baltic herring (Fig. 4).

Similarly, no correlation has been found between the amount of ingested plastic and HOC

concentrations in northern fulmars (*Fulmarus glacialis*) from the Norwegian coast (Herzke et al.

2016), even though the birds had ingested much larger amounts of plastic and their gut passage

time for plastic debris is several orders of magnitude longer than in herring (Ryan 2015). This lack

of relationship is also supported by the relatively constant MP burden observed in Baltic herring

over the past three decades (Beer et al. 2018), whereas muscle concentrations of HOCs decreased

significantly (Bignert et al. 2016). The mass-balance model indicates that our measurements of the

MP burden are ecologically plausible given the currently reported abundances of MP in the Baltic

surface water, thus supporting the reliability of the MP burden estimates in the Baltic herring

reported here and in other studies and providing confidence in the methods employed. Taken

together, these findings contrast the currently held paradigm that microplastics are an important

source of HOCs in aquatic organisms (Mato et al. 2001, Rochman et al. 2013).

Conclusions

Our findings suggest that microplastic ingestion by planktivorous fish is relatively low, even in a semi-enclosed sea like the Baltic, where the MP loading is expectedly high. In agreement with other studies, we also found that biological factors, such as fish size and reproductive state may affect both feeding in general and selectivity towards MP, and, hence, the MP burden. However, we found no indication that the muscle tissue HOC concentrations are related to the amount of ingested plastic. Thus, our study further strengthens the view that MP contribute extremely little to the herring diet and play a negligible role in explaining contaminant bioaccumulation in the fish.

Acknowledgements

This work was supported through the Joint Programming Initiative Healthy and Productive Seas and Oceans (JPI-Oceans) WEATHER-MIC project by the Swedish Research Council for Environment, Agricultural Sciences and Spatial Planning (FORMAS) [grant number 942-2015-1866], FORMAS project; ir-PLAST [grant number 2015-932], through the joint Baltic Sea research and development programme (BONUS) MICROPOLL project by the Swedish Innovation Agency VINNOVA [grant number 2017-00979] and the Baltic Ecosystem Adaptive Management (BEAM) project. The study was designed, analyzed and compiled solely at the responsibility of the authors without the involvement of the funding agencies. Preliminary results were presented at the 15[th] International Conference on Environmental Science and Technology in 2017, Rhodes, Greece, paper number: CEST2017_00690. A preprint of this manuscript has been deposited in BioRxiv; DOI: https://doi.org/10.1101/363127.

References

References

Ahlbeck, I., S. Hansson, O. Hjerne, and C. W. Ramcharan. 2012. Evaluating fish diet analysis methods by
individual-based modelling. Canadian Journal of Fisheries and Aquatic Sciences 69:1184–1201.

Amodio, S., M. Aria, and A. D'Ambrosio. 2014. On concurvity in nonlinear and nonparametric
regression models. Statistica 74:85–98.

Anderson, M. J. 2001. A new method for non-parametric multivariate analysis of variance. Austral
Ecology 26:32–46.

Aro, E., I. Vuorinen, and J. Flinkman. 1989. The prey preference of Baltic herring in the northern Baltic
Sea. International Counc. for the Exploration of the Sea, Copenhagen (Denmark).

Au, S. Y., T. F. Bruce, W. C. Bridges, and S. J. Klaine. 2015. Responses of Hyalella azteca to acute and
chronic microplastic exposures: Effects of Microplastic Exposure on Hyalella azteca.
Environmental Toxicology and Chemistry 34:2564–2572.

Batel, A., F. Linti, M. Scherer, L. Erdinger, and T. Braunbeck. 2016. Transfer of benzo[a]pyrene from
microplastics to Artemia nauplii and further to zebrafish via a trophic food web experiment:
CYP1A induction and visual tracking of persistent organic pollutants. Environmental Toxicology
and Chemistry 35:1656–1666.

Beer, S., A. Garm, B. Huwer, J. Dierking, and T. G. Nielsen. 2018. No increase in marine microplastic
concentration over the last three decades – A case study from the Baltic Sea. Science of The Total
Environment 621:1272–1279.

Besseling, E., A. Wegner, E. M. Foekema, M. J. van den Heuvel-Greve, and A. A. Koelmans. 2013.
Effects of Microplastic on Fitness and PCB Bioaccumulation by the Lugworm Arenicola marina
(L.). Environmental Science & Technology 47:593–600.

Bignert, A., S. Danielsson, S. Faxneld, and E. Nyberg. 2016. Comments Concerning the National
Swedish Contaminant Monitoring Programme in Marine Biota. Stockholm, Sweden.

Browne, M. A., S. J. Niven, T. S. Galloway, S. J. Rowland, and R. C. Thompson. 2013. Microplastic
Moves Pollutants and Additives to Worms, Reducing Functions Linked to Health and
Biodiversity. Current Biology 23:2388–2392.

Bucholtz, R. H., J. Tomkiewicz, and J. Dalskov. 2008. Manual to determine gonadal maturity of herring
(Clupea harengus L.). Pages 1–45. DTU Aqua-report 197-08, National Institute of Aquatic
Resources, Charlottenlund.

Budimir, S., O. Setälä, and M. Lehtiniemi. 2018. Effective and easy to use extraction method shows low
numbers of microplastics in offshore planktivorous fish from the northern Baltic Sea. Marine
Pollution Bulletin 127:586–592.

Casini, M., M. Cardinale, and F. Arrhenius. 2004. Feeding preferences of herring (Clupea harengus) and
sprat (Sprattus sprattus) in the southern Baltic Sea. ICES Journal of Marine Science 61:1267–
1277.

413 Cole, M., P. Lindeque, E. Fileman, C. Halsband, and T. S. Galloway. 2015. The Impact of Polystyrene
414    Microplastics on Feeding, Function and Fecundity in the Marine Copepod Calanus helgolandicus.
415    Environmental Science & Technology 49:1130–1137.

416 Cole, M., P. Lindeque, E. Fileman, C. Halsband, R. Goodhead, J. Moger, and T. S. Galloway. 2013.
417    Microplastic Ingestion by Zooplankton. Environmental Science & Technology 47:6646–6655.

418 Cole, M., P. Lindeque, C. Halsband, and T. S. Galloway. 2011. Microplastics as contaminants in the
419    marine environment: A review. Marine Pollution Bulletin 62:2588–2597.

420 Collard, F., B. Gilbert, G. Eppe, L. Roos, P. Compère, K. Das, and E. Parmentier. 2017. Morphology of
421    the filtration apparatus of three planktivorous fishes and relation with ingested anthropogenic
422    particles. Marine Pollution Bulletin 116:182–191.

423 Costalago, D., and I. Palomera. 2014. Feeding of European pilchard (Sardina pilchardus) in the
424    northwestern Mediterranean: from late larvae to adults. Scientia Marina 78:41–54.

425 Darbyson, E., D. P. Swain, D. Chabot, and M. Castonguay. 2003. Diel variation in feeding rate and prey
426    composition of herring and mackerel in the southern Gulf of St Lawrence. Journal of Fish
427    Biology 63:1235–1257.

428 De Witte, B., L. Devriese, K. Bekaert, S. Hoffman, G. Vandermeersch, K. Cooreman, and J. Robbens.
429    2014. Quality assessment of the blue mussel (Mytilus edulis): Comparison between commercial
430    and wild types. Marine Pollution Bulletin 85:146–155.

431 Dehaut, A., A.-L. Cassone, L. Frère, L. Hermabessiere, C. Himber, E. Rinnert, G. Rivière, C. Lambert, P.
432    Soudant, A. Huvet, G. Duflos, and I. Paul-Pont. 2016. Microplastics in seafood: Benchmark
433    protocol for their extraction and characterization. Environmental Pollution 215:223–233.

434 EFSA Panel on Contaminants in the Food Chain (CONTAM). 2016. Presence of microplastics and
435    nanoplastics in food, with particular focus on seafood. EFSA Journal 14:n/a-n/a.

436 Flinkman, J., E. Aro, I. Vuorinen, and M. Viitasalo. 1998. Changes in northern Baltic zooplankton and
437    herring nutrition from 1980s to 1990s: top-down and bottom-up processes at work. Mar Ecol
438    Prog Ser 165:127–136.

439 Foekema, E. M., C. De Gruijter, M. T. Mergia, J. A. van Franeker, A. J. Murk, and A. A. Koelmans.
440    2013. Plastic in North Sea Fish. Environmental Science & Technology 47:8818–8824.

441 Gewert, B., M. Ogonowski, A. Barth, and M. MacLeod. 2017. Abundance and composition of near
442    surface microplastics and plastic debris in the Stockholm Archipelago, Baltic Sea. Marine
443    Pollution Bulletin 120:292–302.

444 Gibson, R. N. 1988. Development, morphometry and particle retention capability of the gill rakers in the
445    herring, Clupea harengus L. Journal of Fish Biology 32:949–962.

446 Gorokhova, E. 2015. Screening for microplastic particles in plankton samples: How to integrate marine
447    litter assessment into existing monitoring programs? Marine Pollution Bulletin.

Gouin, T., N. Roche, R. Lohmann, and G. Hodges. 2011. A Thermodynamic Approach for Assessing the
Environmental Exposure of Chemicals Absorbed to Microplastic. Environmental Science &
Technology 45:1466–1472.

Grigorakis, S., S. A. Mason, and K. G. Drouillard. 2017. Determination of the gut retention of plastic
microbeads and microfibers in goldfish (Carassius auratus). Chemosphere 169:233–238.

Hartmann, N. B., S. Rist, J. Bodin, L. H. Jensen, S. N. Schmidt, P. Mayer, A. Meibom, and A. Baun.
2017. Microplastics as vectors for environmental contaminants: Exploring sorption, desorption,
and transfer to biota. Integrated Environmental Assessment and Management 13:488–493.

Hendriks, A. J., A. van der Linde, G. Cornelissen, and D. T. H. M. Sijm. 2001. The power of size. 1. Rate
constants and equilibrium ratios for accumulation of organic substances related to octanol-water
partition ratio and species weight. Environmental Toxicology and Chemistry 20:1399–1420.

Herzke, D., T. Anker-Nilssen, T. H. Nøst, A. Götsch, S. Christensen-Dalsgaard, M. Langset, K. Fangel,
and A. A. Koelmans. 2016. Negligible Impact of Ingested Microplastics on Tissue
Concentrations of Persistent Organic Pollutants in Northern Fulmars off Coastal Norway.
Environmental Science & Technology 50:1924–1933.

Hidalgo-Ruz, V., L. Gutow, R. C. Thompson, and M. Thiel. 2012. Microplastics in the Marine
Environment: A Review of the Methods Used for Identification and Quantification.
Environmental Science & Technology 46:3060–3075.

Huse, G., and R. Toresen. 1996. A comparative study of the feeding habits of herring (clupea harengus,
clupeidae, 1.) and capelin (mallotus villosus, osmeridae, müller) in the barents sea. Sarsia
81:143–153.

Jovanović, B. 2017. Ingestion of microplastics by fish and its potential consequences from a physical
perspective. Integrated Environmental Assessment and Management 13:510–515.

Kleinteich, J., S. Seidensticker, N. Marggrander, and C. Zarfl. 2018. Microplastics Reduce Short-Term
Effects of Environmental Contaminants. Part II: Polyethylene Particles Decrease the Effect of
Polycyclic Aromatic Hydrocarbons on Microorganisms. International Journal of Environmental
Research and Public Health 15:287.

Koelmans, A. A. 2015. Modeling the Role of Microplastics in Bioaccumulation of Organic Chemicals to
Marine Aquatic Organisms. A Critical Review. Pages 309–324 *in* M. Bergmann, L. Gutow, and
M. Klages, editors. Marine Anthropogenic Litter. Springer International Publishing, Cham.

Koelmans, A. A., A. Bakir, G. A. Burton, and C. R. Janssen. 2016. Microplastic as a Vector for
Chemicals in the Aquatic Environment: Critical Review and Model-Supported Reinterpretation
of Empirical Studies. Environmental Science & Technology:3315–3326.

Lagarde, F., O. Olivier, M. Zanella, P. Daniel, S. Hiard, and A. Caruso. 2016. Microplastic interactions
with freshwater microalgae: Hetero-aggregation and changes in plastic density appear strongly
dependent on polymer type. Environmental Pollution 215:331–339.

484 Lenz, R., K. Enders, T. Kirk Sørensen, and C. A. Stedmon. 2016. Analysis of microplastic in the
485   stomachs of herring and cod from the North Sea and Baltic Sea. Page 30. DTU Aqua-report, DTU
486   Aqua National Institute of Aquatic Resources, Copenhagen, Denmark.

487 van der Lingen, C. D. 1998. Gastric evacuation, feeding periodicity and daily ration of sardine *Sardinops*
488   *sagax* in the southern Benguela upwelling ecosystem. South African Journal of Marine Science
489   19:305–316.

490 Long, M., B. Moriceau, M. Gallinari, C. Lambert, A. Huvet, J. Raffray, and P. Soudant. 2015.
491   Interactions between microplastics and phytoplankton aggregates: Impact on their respective
492   fates. Marine Chemistry 175:39–46.

493 Lusher, A. L., G. Hernandez-Milian, J. O'Brien, S. Berrow, I. O'Connor, and R. Officer. 2015.
494   Microplastic and macroplastic ingestion by a deep diving, oceanic cetacean: The True's beaked
495   whale Mesoplodon mirus. Environmental Pollution 199:185–191.

496 Lusher, A. L., M. McHugh, and R. C. Thompson. 2013. Occurrence of microplastics in the
497   gastrointestinal tract of pelagic and demersal fish from the English Channel. Marine Pollution
498   Bulletin 67:94–99.

499 MacCallum, R. C., K. F. Widaman, K. J. Preacher, and S. Hong. 2001. Sample Size in Factor Analysis:
500   The Role of Model Error. Multivariate Behavioral Research 36:611–637.

501 Mato, Y., T. Isobe, H. Takada, H. Kanehiro, C. Ohtake, and T. Kaminuma. 2001. Plastic Resin Pellets as
502   a Transport Medium for Toxic Chemicals in the Marine Environment. Environmental Science &
503   Technology 35:318–324.

504 Norén, F. 2007. Small plastic particles in Swedish West Coast Waters. KIMO Sweden, Lysekil, Sweden.

505 Ogonowski, M., C. Schür, Å. Jarsén, and E. Gorokhova. 2016. The Effects of Natural and Anthropogenic
506   Microparticles on Individual Fitness in Daphnia magna. PLOS ONE 11:e0155063.

507 Oliveira, M., A. Ribeiro, and L. Guilhermino. 2012. Effects of short-term exposure to microplastics and
508   pyrene on Pomatoschistus microps (Teleostei, Gobiidae). Comparative Biochemistry and
509   Physiology Part A: Molecular & Integrative Physiology 163, Supplement:S20.

510 Persson, S., A. Rotander, B. van Bavel, B. Brunström, B.-M. Bäcklin, and U. Magnusson. 2013. Influence
511   of age, season, body condition and geographical area on concentrations of chlorinated and
512   brominated contaminants in wild mink (Neovison vison) in Sweden. Chemosphere 90:1664–
513   1671.

514 R Core Team. 2014. R: A language and environment for statistical computing. R Foundation for
515   Statistical Computing, Vienna, Austria.

516 Rehse, S., W. Kloas, and C. Zarfl. 2018. Microplastics Reduce Short-Term Effects of Environmental
517   Contaminants. Part I: Effects of Bisphenol A on Freshwater Zooplankton Are Lower in Presence
518   of Polyamide Particles. International Journal of Environmental Research and Public Health
519   15:280.

Rochman, C. M., E. Hoh, T. Kurobe, and S. J. Teh. 2013. Ingested plastic transfers hazardous chemicals
to fish and induces hepatic stress. Scientific Reports 3.

Rochman, C. M., R. L. Lewison, M. Eriksen, H. Allen, A.-M. Cook, and S. J. Teh. 2014. Polybrominated
diphenyl ethers (PBDEs) in fish tissue may be an indicator of plastic contamination in marine
habitats. Science of The Total Environment 476–477:622–633.

Rummel, C. D., M. G. J. Löder, N. F. Fricke, T. Lang, E.-M. Griebeler, M. Janke, and G. Gerdts. 2016.
Plastic ingestion by pelagic and demersal fish from the North Sea and Baltic Sea. Marine
Pollution Bulletin 102:134–141.

Rusina, T. P., F. Smedes, and J. Klanova. 2010. Diffusion coefficients of polychlorinated biphenyls and
polycyclic aromatic hydrocarbons in polydimethylsiloxane and low-density polyethylene
polymers. Journal of Applied Polymer Science 116:1803–1810.

Ryan, P. G. 2015. How quickly do albatrosses and petrels digest plastic particles? Environmental
Pollution 207:438–440.

Seyhan, K., and D. J. Grove. 2003. A New Approach in Modelling Gastric Emptying in Fish. Turkish
Journal of Veterinary and Animal Sciences 27:1043–1047.

Silva Barni, M. F., M. Gonzalez, and K. S. B. Miglioranza. 2014. Assessment of persistent organic
pollutants accumulation and lipid peroxidation in two reproductive stages of wild silverside
(Odontesthes bonariensis). Ecotoxicology and Environmental Safety 99:45–53.

Succop, P. A., S. Clark, M. Chen, and W. Galke. 2004. Imputation of data values that are less than a
detection limit. Journal of Occupational and Environmental Hygiene 1:436–441.

TemaNord. 1995. Nordic environmental specimen banking: Methods in use in ESB: Manual for Nordic
countries. TemaNord Scandinavia: Nordic Council of Ministers.

Varpe, Ø., and Ø. Fiksen. 2010. Seasonal plankton–fish interactions: light regime, prey phenology, and
herring foraging. Ecology 91:311–318.

Vinson, M. R., and T. R. Angradi. 2011. Stomach Emptiness in Fishes: Sources of Variation and Study
Design Implications. Reviews in Fisheries Science 19:63–73.

Wardrop, P., J. Shimeta, D. Nugegoda, P. D. Morrison, A. Miranda, M. Tang, and B. O. Clarke. 2016.
Chemical Pollutants Sorbed to Ingested Microbeads from Personal Care Products Accumulate in
Fish. Environmental Science & Technology 50:4037–4044.

Wesch, C., K. Bredimus, M. Paulus, and R. Klein. 2016. Towards the suitable monitoring of ingestion of
microplastics by marine biota: A review. Environmental Pollution 218:1200–1208.

Wilkins, N. P. 1967. Starvation of the herring, Clupea harengus L.: survival and some gross biochemical
changes. Comparative biochemistry and physiology 23:503–518.

 Figures and tables

Figure 1. Sampling sites within the Swedish National Monitoring Program for Contaminants in
Marine Biota included in this study, BotB Bothnian Bay, BS Bothnian Sea, NBP Northern Baltic
Proper, WGB Western Gotland Basin and BB Bornholm Basin. 1 Rånefjärden, 2 Harufjärden, 3
Holmöarna, 4 Gaviksfjärden, 5 Långvindsfjärden, 6 Bothnian Sea offshore site, 7
Ängsskärsklubb, 8 Lagnö, 9 Baltic proper offshore site, 10 Landsort, 11 Byxelkrok, 12 Utlängan
and 13 Western Hanö bight.

564 **Figure 2.** Boxplot of Log10-transformed MP abundance in the gastrointestinal tract (GIT) of

565 herring per basin ordered from north to south. Data are presented as medians (vertical lines), inter

566 quartile range, IQR (boxes), 1.5 IQR (whiskers) and outliers (points) being > 1.5 IQR. The black

567 slices of the pie charts indicate the proportion of examined herring with no MP in the GIT.

Figure 3. Frequency distribution of the MP burden based on the model simulations (dark grey bars) and field observations (light grey bars). Panel A shows the entire dataset and panel B presents only fish with MP in the GIT (i.e., the non-zero values). The dashed vertical lines indicate the mean values for the model simulations (long dash) and the observations (short dash).

Figure 4. Factor scores (axes) and loadings (arrows) of contaminants (HBCD, HCB and the sum of PCBs, BDEs and DDs) and weight-specific MP burden.

Figure 5. Generalized additive models (GAMs) showing partial response curves for the explanatory biological variables: *body weight* (A), *reproductive phase* (B), *gut fullness* (C) and *age* (D). The classes for *reproductive phase* correspond to: 1 = deformed gonads, 2 = post spawned, 3 = juvenile, 4 = developing gonads and 5 = mature gonads. The vertical axis shows the relative influence of the explanatory variable on the prediction of MP burden on the base of partial residuals. Grey bands indicate 95% confidence interval for each curve.

Table 1. Overview of the HOCs in herring muscle tissue and descriptive statistics of their concentrations ($\mu g\ g^{-1}$ fish muscle). SD – standard deviation.

Chemical group	Chemical species	Abbreviation	Mean concentration in fish muscle (µg/g)	SD	Median	Min	Max
Polychlorinated biphenyls (PCBs)	2,4,4′-PCB	PCB 28	0.0039	0.0019	0.0033	0.0018	0.0106
	2,2′,5,5′-PCB	PCB 52	0.0067	0.0044	0.0057	0.0025	0.0199
	2,2′,4,5,5′-PCB	PCB 101	0.0222	0.0149	0.0176	0.0069	0.0713
	2,3′,4,4′,5′-PCB	PCB 118	0.0204	0.0135	0.0155	0.0062	0.0694
	2,2′,3,4,4′,5′-PCB	PCB 138	0.0591	0.0408	0.0444	0.0152	0.1896
	2,2′,4,4′,5,5′-PCB	PCB 153	0.0417	0.0280	0.0345	0.0120	0.1320
	2,2′,3,4,4′,5,5′-PCB	PCB 180	0.0181	0.0114	0.0149	0.0029	0.0551
Organochlorine pesticides	4,4′-DDT	DDT	0.0205	0.0218	0.0134	0.0037	0.0966
	4,4'-DDE	DDE	0.1055	0.0960	0.0815	0.0160	0.4130
	4,4'-DDD	DDD	0.0284	0.0333	0.0163	0.0016	0.1391
	α-1,2,3,4,5,6-Hexachlorocyclohexane	AHCH	0.0029	0.0005	0.0030	0.0018	0.0039
	β-1,2,3,4,5,6-Hexachlorocyclohexane	BHCH	0.0056	0.0027	0.0057	0.0018	0.0099
	γ-1,2,3,4,5,6-Hexachlorocyclohexane	Lindane	0.0029	0.0005	0.0030	0.0018	0.0039
Brominated flame retardants (BDEs)	2,4,4′-TriBDE	BDE 28	0.0002	0.0001	0.0002	0.0001	0.0005
	2,2′,4,4′-TetraBDE	BDE 47	0.0051	0.0032	0.0041	0.0016	0.0160
	2,2′,4,4′,5-PentaBDE	BDE 99	0.0012	0.0009	0.0009	0.0005	0.0044
	2,2′,4,4′,6-PentaBDE	BDE 100	0.0012	0.0007	0.0011	0.0004	0.0035
	2,2′,4,4′,5,5′-HexaBDE	BDE 153	0.0002	0.0002	0.0002	0.0001	0.0009
	2,2′,4,4′,5,6′-HexaBDE	BDE 154	0.0006	0.0004	0.0005	0.0002	0.0018
	1,2,5,6,9,10-Hexabromocyclododecane	HBCD	0.0114	0.0096	0.0088	0.0025	0.0470
Other	Hexachlorobenzene	HCB	0.0289	0.0209	0.0246	0.0114	0.1013