Is the predation risk of mate-searching different between the sexes?

Viraj R. Torsekar*, Kavita Isvaran, Rohini Balakrishnan

Centre for Ecological Sciences, Indian Institute of Science, Bangalore, India

*Correspondence to virajt@iisc.ac.in +919590927391

1 Abstract

In animals that communicate for mate searching, typically one sex signals and the

other searches. Differential predation risk of mate searching between the sexes is

hypothesised to determine which sex invests more effort in mate searching. However,

relative levels of risk posed by the two mate searching strategies remain largely

unexplored. Furthermore, higher risks of signalling compared to responding to signals

have been hypothesised to drive reduction in asymmetry of mate searching effort

between the sexes in orthopterans, but rarely tested explicitly. Hence, we estimated

the relative predation risk experienced in natural populations by signalling males and

responding females. We did this by quantifying predation risk as a probability of

mortality in the context of acoustic communication in a tree cricket, *Oecanthus henryi*

from its ecologically relevant predator, a lynx spider, *Peucetia viridans*. Probability of

mortality was quantified by partitioning it into three spatial components at which

crickets and spiders interact, using a combination of extensive field observations and

manipulative experiments in a semi-natural setup. We found no differences in

predation risk faced by calling males and responding females, supporting the

hypothesis that similar sex-specific costs can explain shared mate searching

responsibilities. Our findings also suggest that direct benefits offered by males to

females upon pair formation may better explain females contributing towards mate

searching effort in orthopterans.

22 Keywords

23 Communication, sex roles, sex-specific costs, sex-biased predation, crickets

Introduction

Searching for mates typically involves some physical activity for pair formation. Mate searching effort is defined 'as a costly investment in traits that facilitate encounters with potential mates, including mobility, advertisement calls or displays, and pheromone production' (Fromhage et al. 2016). Several factors have been proposed to determine which of the two sexes contributes more towards mate searching (McCartney et al. 2012; Fromhage et al. 2016). However, sex differences in costs from predation risk, proposed as a potential determinant (Fromhage et al. 2016), has rarely been tested in natural populations (Heller 1992; Raghuram et al. 2015).

Sex differences in benefits of multiple matings, differential gametic and parental investments, and time- and predation-related costs have been proposed to determine which of the two sexes invests more effort in mate searching (Trivers 1972; Hammerstein and Parker 1987; Fromhage et al. 2016). Since males typically get higher maximum potential benefits from multiple matings than females (Bateman 1948; Rodríguez-Muñoz et al. 2010; Byers and Dunn 2012), and females are predicted to be the limiting resource to males by investing more in offspring (Andersson 1994), males are expected to be the searching sex. Contrary to this expectation, in a classic paper, Hammerstein and Parker (1987) showed that either sex can potentially search, arguing that sex differences in relative parental investment do not determine which sex searches for mates. But in most animals, males searching for mates is the fundamental mode of pair formation (Darwin 1871; Thornhill 1979). Among other factors, the cost of mate searching has been suggested as a potential determinant of this asymmetry (Fromhage et al. 2016). Fromhage et al. (2016) proposed that higher costs for females could explain the prevalence of male searching,

calling it the 'sex-specific hypothesis'. One of the ubiquitous costs experienced

during mate search is predation (Gwynne 1987; Magnhagen 1991; Heller and Arlettaz

1994; Zuk and Kolluru 1998; Zuk et al. 2006; Raghuram et al. 2015). Predation risk

as a cost of mate searching, however, has not been widely studied between the

participating sexes.

Although mate searching by males is prevalent in most animals, in orthopterans and

some other taxa females physically move to localise signalling males who are

predominantly sedentary (Darwin 1871; Thornhill 1979). Thus, females share mate

searching responsibilities, thereby reducing the asymmetry between the sexes

observed in most animal taxa. What factors explain this reduction in asymmetry and

its maintenance? Thornhill (1979) attributes this to two potential factors: direct

benefits provided by males to females on pair formation and/or the risks associated

with signalling. In many species of orthoptera, on pair formation, males provide direct

benefits such as a burrow for safe shelter (Gwynne 1995) or courtship nuptial gifts

(Arnqvist and Nilsson 2000). McCartney et al. (2012) examined the interspecific mate

searching differences across 32 taxa from a katydid genus, using theory and

observational data. Their findings provide comparative evidence for the hypothesis

that mate searching by females can be explained by the direct benefits offered by

males if these are substantial. The alternative hypothesis states that since males

benefit more from multiple mating, they should be selected to perform risky mate

searching activities; hence, males are expected to face higher risks while signalling

(Thornhill 1979; McCartney et al. 2012). The relative risk of signalling versus

responding as a potential determinant of reduction in sexual asymmetry in mate

searching effort between the sexes, has however rarely been tested.

Predation risk in the context of mate searching communication has predominantly

been studied from the signaller's perspective (Zuk and Kolluru 1998). Many studies

have demonstrated how sexual advertisement in the form of conspicuous calls makes

signallers vulnerable to predation (Walker 1964; Bell 1979; Ryan et al. 1982;

Belwood and Morris 1987; reviewed in Zuk and Kolluru 1998). However, studies

analysing predator diet found evidence for responding females being at an equal, if

not higher predation risk, in comparison with signalling males (Heller and Arlettaz

1994; Raghuram et al. 2015). Thus, both signalling and responding to signals entail

predation risk, since signalling attracts eavesdropping predators (Zuk and Kolluru

1998) and movement towards a signal increases exposure to predators (Gwynne 1987;

Heller and Arlettaz 1994; Raghuram et al. 2015). There is however, a paucity of

studies that attempt to estimate the relative predation risk of signallers and responders

(but see Heller 1992; Raghuram et al. 2015).

We examined the mate searching costs from predation risk in tree crickets by

estimating predation risk experienced by calling males and responding females. We

studied predation risk using an approach novel to the field of communication, though

commonly used in ecology (Holling 1959; Lima and Dill 1990; Hebblewhite et al.

2005; Brechbühl et al. 2011). We partitioned risk into constituent parts, each

characterised by a discrete spatial scale at which predator and prey interact: co-

occurrence, encounter and being eaten. At each scale, the binary response of prey

either succeeding or failing to avoid the predator helped estimate probabilities.

Predation risk was estimated as a product of these probabilities (represented in Fig. 1).

Such a comprehensive approach is critical to determine any trade-offs across different

scales that might not reflect in the total predation risk if studied only at the scale of

proportion of sexes in predator diets, predator visitations in playback experiments, or

predator preferences of certain prey behaviours in a controlled environment (Tuttle

and Ryan 1981, Heller and Arlettaz 1994, Alem et al. 2011).

To summarise, the sex-specific cost hypothesis attributes ubiquitous male mate

searching in most animal taxa to higher costs of searching in females (Fromhage et al.

2016). But unlike in most animal taxa, both sexes in orthopterans share mate

searching effort, with sedentary males signalling, and silent females localising these

signals. Hence, given these reduced sexual asymmetries of mate searching effort, we

expect costs to be similar between the two sexes in orthopterans. Whereas, explicitly

testing whether higher costs explain this reduced asymmetry in mate searching in

orthopterans, we expect the costs of male signalling to be higher (Thornhill 1979).

Methods

We carried out our study on a tree cricket species, *Oecanthus henryi* whose major

predator is the green lynx spider, *Peucetia viridans*. *Oecanthus henryi* is found

extensively in the dry scrubland of southern India, predominantly on bushes of *Hyptis*

suaveolens. *Oecanthus* species exhibit a mating system typical of true crickets

(Gryllidae), where the males produce a long-range species-specific call and females

do not call, but detect, recognize and locate males of their species (Walker 1957).

Males of *O. henryi* typically call from *H. suaveolens* leaves and the females negotiate

the complex architecture of the bushes and approach the calling males (Bhattacharya

2016). *O. henryi* male calls are made up of rhythmic chirps (Metrani and

Balakrishnan 2005). *P. viridans* (family Oxyopideae) is commonly observed on

Fig. 1 Graphical representation of experimental design employed in the study to

estimate predation risk. Each box represents a cricket interacting with its predator,

a spider, at a particular spatial scale. Each arrow represents transition to the next

spatial scale. Text above each box indicates what constituent of predation risk is

studied and text below indicates how it was estimated. Probabilities from each

spatial scale were multiplied to estimate predation risk.

Hyptis suaveolens bushes and has been observed preying upon tree crickets and

honeybees (VRT and RB, personal observations).

All field surveys, sampling sessions and experiments were carried out in a

homogenous patch of *H. suaveolens* on unused farmland, near Ullodu village

(13°38'27.2"N 77°42'01.1"E) in Chikkaballapur district of Karnataka state in southern

India. Laboratory experiments were performed on the campus of the Indian Institute

of Science, Bangalore. All experiments in semi-natural conditions were carried out in

an outdoor enclosure, constructed with a steel frame of dimensions 6m x 6m x 3m and

fastened with a fibre mosquito mesh (mesh size: 0.1cm x 0.2 cm), on the campus of

Indian Institute of Science, Bangalore.

We measured predation risk as a product of three probabilities: 1) probability of co-

occurrence of *O. henryi* and *P. viridans* on a bush (POC), 2) probability of *O. henryi*

encountering *P. viridans*, given their co-occurrence (POE), and 3) probability of *O.*

henryi being eaten by *P. viridans*, given an encounter (POBE).

150   Predation risk = POC x POE x POBE

The study was carried out in four parts. First, we performed extensive sampling to

investigate who the major predators of *O. henryi* were in the field. Second, we

sampled a population of *O. henryi* in the field to estimate the probability with which

they co-occur with their major predator species*,* considering a single *H. suaveolens*

bush as a unit (POC). Then we conducted two experiments in a semi-natural outdoor

enclosure to obtain the probability of encounter between *O. henryi* and *P. viridans*

when both co-occurred on the same bush (POE), and the probability with which *O.*

henryi are eaten by *P. viridans* upon encounter (POBE, represented in Fig. 1). The

sampling and experiments to measure the probabilities were carried out on two

different treatment regimes: calling males and responding females of *O. henryi*.

Predator sampling

Playback experiments were carried out in the field to determine acoustically orienting

aerial predators of *O. henryi*. Visitations by aerial predators were compared at

speakers playing back *O. henryi* calls with paired silent speakers for a total of 40

hours (details in supplementary information). For discovering arboreal predators of *O.*

henryi, relative abundance sampling sessions and predation experiments were carried

out. The extensive relative abundance sampling helped determine who the potential

predators of *O. henryi* were. To establish whether these potential predators are real

predators, detailed predation experiments were conducted in the lab and field.

Experiments were also conducted to understand how body size and starvation period

of the predator affects predation so as to better design further experiments (details in

supplementary information).

Probability of co-occurrence

To determine the probability of co-occurrence between *O. henryi* and its major

predator *P. viridans*, incidences of *O. henryi* co-occurring with large *P. viridans* on

the same bush were recorded in the field between November and May. Between

1900hrs and 2115hrs, calling males were located using their calls, whereas females

were localized using 5x5m quadrat sampling or opportunistically. Once localized,

these crickets were focally sampled for at least 30 minutes, allowing time for males to

call, to help distinguish callers from non-callers. The bush or bushes, on which these

crickets were observed while sampling, were thoroughly searched for the presence of

P. viridans at the end of the sampling session. If a spider was present, distances

between the cricket and the spider and whether they were present on the same or

different branches was recorded. Post sampling, crickets and spiders were collected

and brought back to the field station for marking and sizing, respectively. Crickets

were marked with a unique tricolour code using nontoxic paint markers (Edding

780, Edding, St Albans, U.K.), to avoid resampling; spiders were sized to confirm

their ability to predate on crickets, and both were released back into the field. Male *O.*

henryi were considered as callers based on whether or not they called more than 20%

of the time they were observed. 20% of the calling effort was chosen as a cut-off so as

to avoid choosing infrequent signallers as callers. Since there was no intuitive way to

categorise communicating and non-communicating females during field observations,

observed females were randomly classified as responding and non-responding based

on a supplementary experiment (that estimated the relative frequency of

communicating (phonotactic) females in the same population). This random sampling

involved computationally segregating (sampled for 10,000 iterations) the observed

females into responding and non-responding females, using the proportion of wild-

caught females known to be responsive (experiment details in supplementary

information).

Probability of encounter given co-occurrence

To determine the probability of encounter between *O. henryi* and *P. viridans* given

their co-occurrence on a bush, experiments were carried out in semi-natural

conditions, inside a large outdoor enclosure. Crickets and spiders were collected from

wild populations from in and around Peresandra, Karnataka, India (13°35'25.3"N

77°46'50.4"E), a few days before the experiment. Crickets were maintained on *H.*

suaveolens bushes and spiders were provided *Gryllus bimaculatus* nymphs 2-3 times

a week, both in the laboratory. Female crickets were collected as nymphs from the

field and fed on an apple diet till they eclosed into adults after which they were

maintained on *H. suaveolens* bushes. This exercise ensured virginity of all tested

females, which increases the propensity of females to perform phonotaxis. Male and

female crickets were maintained separately. Spiders were maintained in individual

plastic boxes (6 cm diameter, 4 cm height). Spiders were starved for 48 hours prior to

a trial and crickets were transferred to *H. suaveolens* bushes in outdoor cages at least a day before a trial to acclimatise them. No cricket was repeated across or within treatments and no spiders were repeated within treatments.

Each trial involved releasing one cricket and one spider on a *H. suaveolens* bush and observing the interaction for about 120 minutes. Crickets were released on bushes at least 4 hours before the trial was started. The spiders were released at 1900 hours, marking the beginning of a trial. They were released on the bush at a height and distance from its centre, picked randomly from the interquartile range of their respective uniform distributions that were obtained from field data (refer to earlier section). Spiders were released on either the same or on a different branch as the cricket at 1900 hours with the proportion with which they were observed in the field. On the same or different branch, the distance at which spiders were released from crickets was drawn randomly from the interquartile range of a uniform distribution of distances at which crickets and spiders were observed in the field. Once the spider was released at 1900 hours, the interaction between the cricket and spider was observed for about 120 minutes by scan sampling each individual alternatively every 30 seconds. An encounter was defined as any spatial proximity between the cricket and spider, within 4 cm of each other, on the same branch. This distance was the outer range from which *P. viridans* attacked *O. henryi* and also the distance at which the cricket could potentially antennate the spider (VRT, personal observations).

Any male that called for more than 20% of the time it was observed was classified as a calling male. Females were played back conspecific male calls from a speaker (X-mini Capsule Speaker V1.1, Xmi Pte Ltd, Singapore) placed 60 cm away, across the

bush from the female's position at 1900 hrs. A trial was considered only if the female

performed phonotaxis and reached within 20 cm of the speaker. The speaker was

fixed on a stand, which was adjusted to the median height at which calling males were

observed in the field (42 cm). The SPL of each call broadcast from the speaker was

adjusted to be 61 dB (r.m.s. re. 2 x 10^{-5} Nm^{-2}) at the female's location with the help of

a ½" microphone (Brüel and Kjær A/S, Denmark, Type 4189, 20 Hz to 20 kHz) fitted

on a Sound Level Meter Type 2250 (Brüel and Kjær A/S, Denmark). Since the call

carrier frequency changes with temperature in *O. henryi,* the choice of call to be

played back was based on the temperature recorded at 1900 hours every evening. The

call was chosen from among 3 representative calls that were recorded at 22°C, 24°C

and 26°C (calls recorded by Rittik Deb, Deb 2015), whichever was closest to the

recorded temperature. This call was played back in a loop using Audition software

(Adobe, Version 5.0.2) on a MacBook Pro (2011) laptop using X-mini (Capsule

Speaker V1.1, Xmi Pte Ltd, Singapore) speakers, for the entire duration of the

experiment.

Probability of being eaten given encounter

The probability of *O. henryi* being eaten by *P. viridans* once encounter occurs was

examined empirically in the same outdoor semi-natural setup. *O. henryi* and *P.*

viridans were collected and maintained using the same protocol mentioned in the

earlier section. The same two treatments were maintained for this experiment.

Crickets were released at least 4 hours before the commencement of the experiment at

1900 hours on the bush at a height and distance from its centre as explained in the last

experiment. From 1900 hours onwards, focal observations were made for at least 45

minutes to allow males to call and to allow females to perform phonotaxis. Following

these observations and based on whether the male called and female performed

phonotaxis, a spider was gently released within close proximity of the cricket, not

more than 6 cm away from it, using a *H. suaveolens* stick. A trial was considered only

if the spider attacked, which was confirmed by videotaping all interactions. A spider

capturing the cricket was scored as the cricket being eaten by the spider.

Analyses

Since comparing estimates of rare and non-normally distributed events can be

challenging, we employed non-parametric bootstrapping and permutation tests

(Manly 2006). These are robust methods for obtaining confidence intervals and *P*-

values, respectively, since they make few assumptions about the underlying

distributions (Manly 2006; Nakagawa and Cuthill 2007). Bootstrapping was used to

generate 95% confidence intervals around each probability. This process involved

sampling with replacement, for 10,000 iterations, from the original vector of success/

failures used to calculate the probability. Overlap in confidence intervals was used to

infer statistical significance for each relevant comparison (Cumming and Finch 2005).

Additionally, permutation tests were carried out to assess statistical significance

(Manly 2006). We used software R, Version 3.3.3 (R Core Team 2017) to run all

analyses, and the 'ggplot2' package (Wickham 2009) to plot all graphics.

Results

Not much is known about the identity of predators of *Oecanthus* species (Ponce-

Wainer and Del Castillo 2008, but see O'Neill and O'Neill 2003). We conducted a

series of experiments and sampling sessions to elucidate the predators of *O. henryi*.

Playback experiments were carried out in the field to observe if there are any

Fig. 2 Size distinction between *P. viridans* individuals that predated on *O. henryi* and

the ones that did not. All spiders that captured crickets were larger than 5.12mm

in body length.

acoustically orienting aerial predators of *O. henryi*. Bats (species unknown) flew past

the playback speaker on 4 separate occasions, and past the control speaker on 5

occasions out of a total of 40 hours of observation. Approaches to both speakers by

bats was very similar (4 and 5 times in 40 hours, respectively). Also, a praying mantis

approached a broadcast speaker once. In relative abundance sampling to investigate

arboreal predators, 15 5x5m plots were sampled, amounting to 1083 bushes. A total

of 127 *P. viridans* individuals, 129 spiders belonging to the web-building guild, and 1

praying mantis were observed, along with many beetles, roaches, termites and moths

which were not enumerated since they are not potential predators of tree crickets. Of

these, *P. viridans* and spiders belonging to the web-building guild were categorized as

potential predators. In the field predation experiment, 16 out of 30 *P. viridans* that

were offered *O. henryi*, captured and consumed them. All *P. viridans* that captured *O.*

henryi were larger than 5.12 mm in body length (n=14) (Fig. 2). In similar sets of

Fig. 3 Constituent probabilities of predation risk faced by *O. henryi* from its predator,

P. viridans. Predation risk experienced by communicating crickets was

partitioned into three spatial scales: probability of (a) co-occurrence, (b)

encounter and (c) being eaten. Pairwise comparisons between calling males and

responding females at the 3 spatial scales are represented with associated *P*-

values. Values in parentheses are sample sizes.

Treatments	Probability of co-occurrence	Probability of encounter	Probability of being eaten	Predation risk
Calling males	0.2 (35)	0.211 (38)	0.0 (18)	0.0 (91)
Responding females	0.232 (13)	0.356 (45)	0.04 (25)	0.0033 (83)

Table 1. Probability values of calling males and responding females listed for 3

different scales and the total predation risk. Values in parentheses are sample sizes.

Fig. 4 Predation risk faced by *O. henryi* from its predator, *P. viridans*, as calling

males and responding females. Probabilities are bootstrapped values represented

as 95% confidence intervals. Values in parentheses are sample sizes. Pairwise

comparison is represented with associated *P*-value.

experiments, spiders belonging to the web-building guild, were found not to be major

predators of *O. henryi* (details in supplementary information).

The probability with which crickets co-occur with large spiders was similar between

calling males and responding females ($P = 0.779$; Table 1, Fig. 3a). These results did

not qualitatively change even when all sampled females (n = 43) were considered

instead of only the randomly sampled segregate classified as responding females ($P =$

0.790). Once co-occurring on the same bush, both calling males and responding

females encountered spiders with the same probability ($P = 0.221$; Table 1, Fig. 3b).

When an encounter was forced between the spiders and crickets, relatively few

individuals, whether calling males or responding females were captured and eaten by

P. viridans (Table 1). Thus, on encounter, the probability of being eaten by the spider

is similar for both ($P = >0.999$; Table 1, Fig. 3c). Consequently, the product of these

probabilities, the predation risk, is also similar between calling males and responding

females ($P = 0.558$; Table 1, Fig. 4).

Discussion

Our study investigated the relative predation risk of communicating individuals,

signallers and responders, in natural populations on a within-night time scale.

Observing interactions at various spatial scales allowed a comprehensive

quantification of predation risk by taking into account predator-prey dynamics. Our

choice of predator was justified using extensive field surveys and acoustic sampling

sessions to determine all potential predators and carefully pruning that list using

predation experiments to select a predator that is ecologically relevant to our system

of choice. We found that the green lynx spider *P. viridans* was the primary predator of

O. henryi. Spiders have been observed to be predators of several species of crickets

(Hedrick and Kortet 2006; Dangles et al. 2006; Storm and Lima 2010). Spiders are

sensitive to both long-range acoustic (Shamble and Hoy 2016) and substrate-borne

vibratory cues (Barth 2002). Such multimodal sensitivity could allow them to

perceive both acoustic calls and locomotory cues, making spiders good model

predators for studying costs of communication that involves calling by males, and

movement by females.

Predation risk of calling in males and responding to calls in females was similar

across all spatial scales relevant to predator-prey dynamics. We examined and

compared the probability of communicating crickets co-occurring on bushes with

spiders. This probability was similar, indicating that risk faced from distribution of

spiders at a broad-scale is not influenced by whether the cricket is a signalling male or

a responding female. While co-occurring on the same bush, the probability of

encountering a spider was similar between the predominantly sedentary calling males

and mostly mobile responding females. A possible explanation for this result is that

communicating individuals are taking similar necessary evasive measures to spatially

avoid predators (reviewed in Sih 2005). It will be interesting to investigate whether

the two sexes spatially avoid predators while communicating. In addition, we

investigated the probability with which crickets were captured by spiders when

attacked and found no differences. This probability was extremely low for both sexes

in comparison with the earlier two probabilities. This result is not unexpected since

crickets are known to have high escape success against spider attacks, attributable to

their air-sensing systems (Dangles et al. 2006). Although probability of calling males

being eaten by spiders was zero, it was statistically similar to the non-zero probability

of responding females. Since almost certainly the probability of calling males being

captured by spiders in nature is not zero, this probability was interpreted as being very

low. Finally, the resultant product of the three probabilities, considered as predation

risk, was similar for calling males and responding females, suggesting the cost from

predation was comparable for mate searching crickets. Also, this probability was

extremely low for the two sexes, primarily due to high escape probabilities exhibited

by crickets when attacked by spiders.

The sex-specific cost hypothesis predicts sex differences in mate searching effort if

searching is more costly for females than for males. Since in orthopterans this

asymmetry in mate searching effort is reduced, we expected costs from predation risk

between the sexes to be similar. Our results, that costs of mate searching from

predation risk are similar for the signalling males and responding females, support the

sex-specific cost hypothesis as an explanation of shared mate searching

responsibilities in orthopterans. More such examinations of mate searching costs

between the sexes are important to determine whether searching costs indeed are

systematically higher for females than males. Furthermore, it will be interesting to test

whether the extent of sexual asymmetry in mate searching corresponds to the

differences in sex-specific costs, across different taxa.

We also tested one of the two factors proposed to explain females sharing mate

searching responsibilities with males in long distance signalling taxa: high risk of

signalling (Thornhill 1979). Risk associated with signalling has been shown in several

taxa in which males signal for mate searching (Walker 1964, Bell 1979, Ryan et al.

1982, Belwood and Morris 1987, reviewed in Zuk and Kolluru 1998). But, we suggest

when studies test this hypothesis, they should not only provide evidence that the risk

is high for signalling males, but also that this risk is higher than that for responding

females. Very few studies have estimated risk of signalling in comparison with

searching. Our findings, in corroboration with two other studies (Heller 1992,

Raghuram et al. 2014), refute the hypothesis of signalling males facing higher risks

than responding females to explain the reduction in mate searching effort (Thornhill

1979) in long distance signalling taxa. The hypothesis that direct benefits offered by

males drives evolution of female mate searching was however supported by

theoretical and observational results in katydids (McCartney et al. 2012). In addition,

courtship feeding as observed in *Oecanthus* species (Houghton 1909; Fulton 1915),

has been shown to be an ancestral trait in the orthopteran suborder Ensifera,

suggesting that males of most Ensiferan genera will offer direct benefits to their

female counterparts on pair formation (Gwynne 1997). Hence, reduction in sexual

asymmetry in mate searching efforts in orthopterans can perhaps be better explained

by provision of resources by males to females rather than invoking higher risk of

signalling.

In conclusion, although predation risk of signalling in males has been considered to

be high (Zuk and Kolluru 1998), when compared with risk of responding, our findings

show these risks to be similar. It is only by addressing predation risk between the

communicating sexes across several relevant spatial scales that we could compare

risks faced by the two mate searching strategies. More such detailed comparisons of

predation associated costs between the mate searching sexes would help update our

understanding of whether at all there are systematic cost differences in mate searching

strategies. Finally, we also show that overall predation costs of communication per

night are low and that a predation event is very rare, which raises questions on the

importance of predation as a major selection pressure on the evolution of

communication.

432 Author contributions

433 VRT participated in conceptualising and designing the study, carried out data

434 collection and analysis and wrote the manuscript. KI contributed to data analysis,

435 interpretation and writing the manuscript. RB contributed to conceptualising and

436 designing the study, interpretation of data and writing the manuscript.

438 Acknowledgements

We thank the DBT-IISc partnership programme (2012-2017) of the Department of

Biotechnology, Government of India for funding the fieldwork and Ministry of

Human Resource Development for funding the research fellowship of VRT. We thank

Diptarup Nandi, Rittik Deb and Monisha Bhattacharya for helpful discussions and

Diptarup Nandi for comments on the manuscript. We thank Mr. B. Sridhar (Garden

and Nursery Technical Officer) and Nursery staff for their support. We would like to

thank Ayan, Babu, Chaithra, Chirag, Dakshin, Girish, Himamshu, Harsha, Ismail,

Lakshmipriya, Manoj, Meenakshi, Meghana, Murthy, Shivaraju, Sidanth, Sonu,

Sriniketh, Srinivasan, Sunaina and Vinayaka for their help in making behavioural

observations and Manjunatha Reddy for his assistance in fieldwork.

References

References

Alem S, Koselj K, Siemers BM, Greenfield MD (2011) Bat predation and the

evolution of leks in acoustic moths. Behav Ecol Sociobiol 65:2105–2116 doi:

10.1007/s00265-011-1219-x

Andersson MB (1994) Sexual selection, Princeton University Press, Princeton

Arnqvist G, Nilsson T (2000) The evolution of polyandry: multiple mating and female

fitness in insects. Anim Behav 60:145–164. doi: 10.1006/anbe.2000.1446

Balakrishnan R (2016) Behavioral ecology of insect acoustic communication. In:

Pollack GS, Mason AC, Popper AN, Fay RR (eds) Insect hearing. Springer

International Publishing, Switzerland, pp 49–80

Barth FG, (2002) A spider's world: senses and behavior. Springer-Verlag, New York

Bateman AJ (1948) Intra-sexual selection in *Drosophila*. Heredity 2:349–368.

Bell PD (1979) Acoustic attraction of herons by crickets. J N Y Entomol Soc 87:126–

127

Belwood JJ, Morris GK (1987) Bat predation and its influence on calling behavior in

neotropical katydids. Science 238:64–67 doi: 10.1126/science.238.4823.64

Bhattacharya M (2016) Investigating pattern recognition and bi-coordinate sound

localization in the tree cricket species. PhD dissertation, Centre for Ecological

Sciences, Indian Institute of Science, Bangalore, India.

Brechbühl R, Casas J, Bacher S (2011) Diet choice of a predator in the wild:

overabundance of prey and missed opportunities along the prey capture sequence.

Ecosphere 2:1-15 doi: 10.1890/ES11-00323.1

Byers J, Dunn S (2012) Bateman in nature: predation on offspring reduces the

potential for sexual selection. Science 338:802–804 doi:10.1126/science.1224660

Core Team R (2017) A language and environment for statistical computing. R

Foundation for Statistical Computing. http://www.R- project.org.

Cumming G, Finch S (2005) Inference by eye: confidence intervals and how to read

pictures of data. Am Psychol 60:170–180 doi: 10.1037/0003-066X.60.2.170

Darwin C (1871) The descent of man and selection in relation to sex. John Murray,

London, UK

Dangles O, Ory N, Steinmann T, Christides JP, Casas J (2006) Spider's attack versus

cricket's escape: velocity modes determine success. Anim Behav 72:603–610.

doi: 10.1016/j.anbehav.2005.11.018

Deb R (2015) Mate choice, mate sampling and baffling behaviour in the tree cricket.

PhD dissertation, Centre for Ecological Sciences, Indian Institute of Science,

Bangalore, India.

Fromhage L, Jennions MD, Kokko H (2016) The evolution of sex roles in mate

searching. Evolution 70:617-624 doi: 10.1111/evo.12874

Fulton BB (1915) The tree crickets of New York: life history and bionomics. N Y

Agric Exp Station Tech Bull 42:1–47

Gwynne DT (1987) Sex-biased predation and the risky mate-locating behaviour of

male tick-tock cicadas (Homoptera: Cicadidae). Anim Behav 35:571–576 doi:

10.1016/S0003-3472(87)80283-X

Gwynne DT (1995) Phylogeny of the Ensifera (Orthoptera): A hypothesis supporting

493   multiple origins of acoustical signalling, complex spermatophores and maternal

494   care in crickets, katydids, and weta. J Orthop Res 4:203–218. doi:

495   10.2307/3503478

496   Gwynne DT (1997) The evolution of edible 'sperm sacs' and other forms of courtship

497   feeding in crickets, katydids and their kin (Orthoptera: Ensifera). In: Choe JC and

498   Crespi BJ (eds) The evolution of mating systems in insects and arachnids.

499   Cambridge University Press, Cambridge, UK, pp 110–129

500   Hammerstein P, Parker GA (1987) Sexual selection: games between the sexes. In:

501   Bradbury JW, Andersson MB (eds) Sexual selection: testing the alternatives. John

502   Wiley & Sons, Chichester, UK, pp 119–142

503   Hebblewhite M, Merrill EH, McDonald TL (2005) Spatial decomposition of

504   predation risk using resource selection functions: an example in a wolf–elk

505   predator–prey system. Oikos 111:101–111 doi: 10.1111/j.0030-

506   1299.2005.13858.x

507   Hedrick AV, Kortet R (2006) Hiding behaviour in two cricket populations that differ

508   in predation pressure. Anim Behav 72: 1111–1118 doi:

509   10.1016/j.anbehav.2006.03.018

510   Heller KG (1992) Risk shift between males and females in the pair-forming behavior

511   of bushcrickets. Naturwissenschaften 79:89–91

Heller KG, Arlettaz R (1994) Is there a sex ratio bias in the bushcricket prey of the

scops owl due to predation on calling males? J Orthop Res 2:41–42

Holling CS (1959) The components of predation as revealed by a study of small-

mammal predation of the European Pine Sawfly. Can Entomol 91:293–320 doi:

10.4039/Ent91293-5

Houghton CO (1909) Observations on the mating habits of *Oecanthus*. Entomol

News, 274–279

Lima SL, Dill LM (1990) Behavioral decisions made under the risk of predation: a

review and prospectus. Can J Zool 68:619–640 doi: 10.1139/z90-092

Magnhagen C (1991) Predation risk as a cost of reproduction. TREE 6:183–186.

Manly BFJ (2006) Randomization, Bootstrap and Monte Carlo Methods in Biology,

3rd edn. Chapman and Hall/CRC, New York

McCartney J, Kokko H, Heller KG, Gwynne DT (2012). The evolution of sex

differences in mate searching when females benefit: new theory and a

comparative test. - Proc R Soc Lond *B* Biol Sci 279:1225-1232 doi:

10.1098/rspb.2011.1505

Metrani S, Balakrishnan R (2005) The utility of song and morphological characters in

delineating species boundaries among sympatric tree crickets of the genus

Oecanthus (Orthoptera: Gryllidae: Oecanthinae): a numerical taxonomic

approach. J Orthop Res 14:1–16 doi: 10.1665/1082

6467(2005)14[1:TUOSAM]2.0.CO;2

Nakagawa S, Cuthill IC (2007) Effect size, confidence interval and statistical

significance: a practical guide for biologists. Biol Rev Camb Philos Soc 82:591–

605 doi: 10.1111/j.1469-185X.2007.00027.x

O'Neill KM, O'Neill RP (2003) Sex allocation, nests, and prey in the grass-carrying wasp *Isodontia mexicana* (Saussure) (Hymenoptera: Sphecidae) - J Kans Entomol Soc 76:447–454

Ponce-Wainer JX, Del Castillo RC (2008) Female mate choice and no detected predation risk in relation to the calling song of *Oecanthus niveus* (Gryllidae: Oecanthinae). Ann Entomol Soc Am 101:260–265 doi: 10.1603/0013-8746(2008)101[260:FMCAND]2.0.CO;2

Raghuram H, Deb R, Nandi D, Balakrishnan R (2015) Silent katydid females are at higher risk of bat predation than acoustically signalling katydid males. Proc R Soc Lond *B* Biol Sci 282:20142319 doi: 10.1098/rspb.2014.2319

Rodríguez-Muñoz R, Bretman A, Slate J, Walling CA, Tregenza T (2010) Natural and sexual selection in a wild insect population. Science 328:1269–1272 doi: 10.1126/science.1188102

Ryan MJ, Tuttle MD, Rand SA (1982) Bat predation and sexual advertisement in a neotropical anuran. Am Nat 119:136–139. doi: 10.1086/283899

Shamble PS, Menda G, Golden JR, Nitzany EI, WaldenK, Beatus T, Elias DO, Cohen I, Miles RN, Hoy RR (2016) Airborne acoustic perception by a jumping spider. Curr Biol 26:2913-2920 doi: 10.1016/j.cub.2016.08.041

Sih A (2005) Predator-prey space use as an emergent outcome of a behavioral response race. In: Barbosa P, Castellanos I (eds) Ecology of predator-prey interactions. Oxford University Press, New York, pp 240–255

Storm JJ, Lima SL (2010) Mothers forewarn offspring about predators: a

transgenerational maternal effect on behavior. Am Nat 175:382–390 doi:

10.1086/650443

Tuttle MD, Ryan MJ (1981) Bat predation and the evolution of frog vocalizations in

the neotropics. Science 214:677–678 doi: 10.1126/science.214.4521.677

Thornhill R (1979) Male and female sexual selection and the evolution of mating

strategies in insects. In: Blum M, Blum N (eds) Sexual selection and reproductive

competition in insects, Academic Press, New York, pp 81–121

Trivers RL (1972) Sexual selection and parental investment. In: Campbell BG (ed)

Sexual selection and the descent of man, Aldine Press, Chicago, pp 136–179

Walker TJJ (1957) Specificity in the response of female tree crickets (Orthoptera,

Gryllidae, Oecanthinae) to calling songs of the males. Ann Entomol Soc Am 50:

626-636 doi: 10.1093/aesa/50.6.626

Walker TJJ (1964) Experimental demonstration of a cat locating orthopteran prey by

the prey's calling song. The Florida Entomol 47:163–165 doi: 10.2307/3493290

Wickham W (2009) ggplot2: Elegant graphics for data analysis. Springer-Verlag,

New York

Zuk M, Kolluru GR (1998) Exploitation of sexual signals by predators and

parasitoids. Q Rev Biol 73:415–438. doi: 10.1086/420412

Zuk M, Rotenberry JT, Tinghitella RM (2006) Silent night: adaptive disappearance of

a sexual signal in a parasitized population of field crickets. Biol Lett 2:521-524

doi: 10.1098/rsbl.2006.0539