Kinetic analysis of yeast-yeast interactions in oenological conditions

Oliver Harlé[1], Judith Legrand[3], Catherine Tesnière[2], Martine Pradal[2], Jean-Roch Mouret[2], Thibault Nidelet[2*]

Affiliations :

1 : STLO, INRA, Agrocampus Ouest, Rennes, France

2 : SPO, Univ Montpellier, INRA, Montpellier SupAgro, Montpellier, France

3 : GQE-Le Moulon, INRA, Univ. Paris-Sud, CNRS, AgroParisTech, Université Paris-Saclay, 91190 Gif-sur-Yvette, France

[*] **Corresponding Author**: thibault.nidelet@inra.fr

keywords: winemaking fermentation, yeast-yeast interactions, population dynamics, non-*Saccharomyces* species, transgressive interactions, ecosystems

Abstract

Fermentation by microorganisms is a key step in the production of traditional food products such as bread, cheese, beer and wine. In these fermentative ecosystems, microorganisms interact in various ways (competition, predation, commensalism and mutualism). Traditional wine fermentation is a complex microbial process performed by different yeast species which can be classified in two groups: *Saccharomyces* and non-*Saccharomyces* species. To better

understand the different interactions occurring within wine fermentation, isolated cultures were

compared to cultures involving one reference strain of *S. cerevisiae* species and one strain of

five non-*Saccharomyces* yeast species out of five (*Metschnikowia pulcherrima, Metschnikowia*

fructicola, Torulaspora delbrueckii, Hanseniaspora opuntiae and Hanseniaspora uvarum). In

each case, we studied the population dynamics, the resources consumption and the production

of metabolites of central carbon metabolism. This deep phenotyping of competition kinetics

allowed us to identify the main mechanisms of interaction. *T. delbrueckii* and *S. cerevisiae*

compete for resources with comparable fitness in our experimental conditions. *S. cerevisiae*

and *H. uvarum* and *H. opuntiae* are also competing for resources although both *Hanseniaspora*

strains are characterized by a strong mortality in isolated and mixed fermentations. *M.*

pulcherrima and *M. fructicola* have a negative interaction with *S. cerevisiae*, provoking a

decrease in viability in co-culture, probably caused by the synthesis of a killer toxin. Overall,

this work highlights the interest of measuring the population and metabolites kinetics in order

to understand yeast-yeast interactions. These results are a first step for ecological engineering

and the intelligent design of optimal multi-starter consortiums using modeling tools.

Importance

Most oenological fermentations are using today fermentation starters composed of a

Saccharomyces cerevisiae single species. Recently the interest of other yeast species has

increased in order to diversify the organoleptic qualities of the wine. However, these other

species cannot finish the fermentation and are therefore most often associated with *S. cerevisiae*

strain in multi-species starters. Until now, there is a lack of knowledge on the interaction

mechanisms that support these multi-species starters. The goal of this work is to increase our

knowledge of yeast-yeast interaction in winemaking fermentation. Furthermore, acquired data

will be used in the future to build a mathematical model predicting the production dynamics of

mixed fermentations. This knowledge is of great importance to guide the design of optimized

yeast consortium to increase wine quality and therefore has a great economic potential.

53   Introduction

In natural or anthropized environments, microbial species are part of an ecosystem and interact

positively or negatively in a complex network. Until recently, process optimization in

agriculture or food processing was mostly based on the selection of single strains. However,

this paradigm is now being challenged and the scientific community is increasingly seeking to

exploit and optimize consortia consisting of several strains and/or species. Indeed, many studies

have shown that more diverse anthropized environments have many advantages in terms of

resilience, disease resistance or yield of production (1). Efforts are now being made to design

optimal consortia of various species and strains whose interactions will be exploited to

maximize a given criterion.

In oenology which is both an economical and societal important food ecosystem, the addition

of 'starters' composed of selected yeasts at the beginning of the fermentation process is

common. It is estimated that 80% of the oenological fermentations in the world are conducted

with starters (2, 3). Most often, these fermentation starters are composed of a single

Saccharomyces cerevisiae strain selected for its ability to complete the fermentation. Indeed,

numerous experiments have shown that *S. cerevisiae* species, which is rare at the beginning of

the fermentation, most often becomes predominant at the end of the fermentation,

demonstrating their superior fermentation abilities (4–7). However, in recent years, multi-

species starters have emerged with the aim to increase the aromatic complexity of wines. These

new starters most often combine one strain of *S. cerevisiae* allowing to complete the

fermentation and one other species strain, often from a different genus than *Saccharomyces*,

bringing a greater variety of flavors (4, 5, 7, 8).

There are numerous experiments and even industrial products that mix *Saccharomyces* and non-

Saccharomyces strains to improve the organoleptic qualities of wine (5, 9, 10). Non-

Saccharomyces strains used in these experiments are very diverse with more than 23 different

species (5) including among others *Torulaspora delbrueckii, Metschnikowia pulcherrima,*

Metschnikowia fructicola, Hanseniaspora opuntiae and *Hanseniaspora uvarum. T. delbrueckii*

is one of the few species that is capable to finish the fermentation and has shown to increase

wine quality in mixed cultures with *S. cerevisiae* (9, 11, 12). Bely et al (11) have shown that a

mixed culture *S. cerevisiae* and *T. delbrueckii* improves the analytical profile of sweet wines

notably by decreasing volatile acidity and acetaldehyde production (with 53% less in volatile

acidity and 60% less acetaldehyde than a pure culture of *S. cerevisiae*) (11). On the opposite,

species from the *Metschnikowia* genus ferments poorly in oenological conditions but also can

have interesting attributes (4). In conjunction with *S. cerevisiae*, the presence of *M. pulcherrima*

can reduce ethanol concentrations (4, 13), can increase 'citrus/grape fruit' and 'pear' attributes

(14), as well as the persistence of 'smocky' and 'flowery' attributes (15). *M. pulcherrima* also

has an amensalism effect on *S. cerevisiae* by depletion of iron via the production of

pulcherriminic acid (16). *M. fructiola* has been less studied and never in conjunction with *S.*

cerevisiae but still presents an interesting capacity to inhibit *Botrytis* growth (17). At last, the

Hanseniaspora genus also has also been studied in sequential or simultaneous fermentation

with *S. cerevisiae* and has shown to increase volatile compound production during winemaking

(4). It has been shown to increase 'tropical fruit', 'berry', 'floral', and 'nut aroma' character

(18) linked to higher concentrations of acetate esters, esters of MCFAs, isoamyl alcohol, 2-

phenylethanol and α-terpineol (19).

Despite these diverse studies, the composition and the protocol of inoculation of these multi-

strains starters are still very empirical and only based on input/output balance without

considering the dynamics of the microbial populations and their interactions. This lack of

knowledge about the yeast-yeast interactions prevents improving a rationale design of multi-

strain starters (20). To address this problem, we precisely followed population dynamics and

metabolite productions during oenological fermentations realized in unique or mixed cultures

of yeasts. This detailed analysis of the fermentation dynamics will enhance our understanding

of the interaction mechanisms and of the influence of yeasts-yeasts interactions on the

composition of the final product.

Results

In this work we compared the performances in winemaking conditions of isolated cultures of

six different yeast species (*Saccharomyces cerevisiae*, *Torulaspora delbrueckii*, *Metschnikowia*

pulcherrima, *Metschnikowia fructicola*, *Hanseniaspora opuntiae* and *Hanseniaspora uvarum,*

one strain by species) with paired co-cultures mixing each of the five non-*Saccharomyces*

strains with one GFP marked *Saccharomyces cerevisiae* strain. Obtained results will make it

possible to evaluate the intensity of yeast-yeast interactions. By comparing the output of

isolated and mixed cultures, we evaluated the intensity of yeast-yeast interactions and/or its

consequences on ecosystem services production.

We first investigated the consequences of co-culture onthe dynamic of CO_2 production which

is a good indicator of the fermentation progress. The values of the maximum rate of CO_2

production (***Vmax***, Fig. 1a, Table. 1) and of the maximum CO_2 produced were estimated

(***CO₂max***, Fig. 1b, Table. 1). The ***Vmax*** was highly dependent on the species (p.value < 0.001).

S. cerevisiae cultures (***Sc***) displayed the maximum ***Vmax*** (0.99 ± 0.02 g.L^{-1}.h^{-1}) followed by *T.*

delbrueckii (**Td**, 0.51 ± 0.05 g.L^{-1}.h^{-1}), followed by *Hanseniaspora* species (**Hu** 0.33 ± 0.04 g.L^{-1}.h^{-1},

and **Ho** 0.42 ± 0.02 g.L^{-1}.h^{-1}) and finally *Metschnikowia* species (**Mp** 0.165 ± 0.02 g.L^{-1}.h^{-1}

and **Mf** 0.17 ± 0.04 g.L^{-1}.h^{-1}). The five mixed cultures had an intermediate **Vmax** values

between those of **Sc** and **Td** isolated cultures (Fig. 1a). Mixed culture containing **Mp**, **Td** or **Mf**

species had significantly higher **Vmax** than those containing **Hu** or **Ho** species. we considered

two types of cultures, those where the fermentation was finished within xx hours (**CO2max** >

80g.L^{-1}) and those where it was not (Fig. 1b). All mixed cultures and the Sc cultures were able

to finish the fermentation while all non-*Saccharomyces* isolated cultures did not. **Td** isolated

cultures had the highest **CO₂max** among non-*Saccharomyces* species (65 ± 4 g.L^{-1}), followed

by *Hanseniaspora* species (**Hu** 30 ± 0.4 g.L^{-1}, and **Ho** 46 ± 0.6 g.L^{-1}) and finally *Metschnikowia*

species (**Mp** 22 ± 0.4 g.L^{-1} and **Mf** 20 ± 1 g.L^{-1}). For these two parameters, the mixed cultures

can be grouped with the **Sc** indicating a prevalent impact of the *S. cerevisiae* species on the

fermentation kinetics.

We then looked at the population dynamics for each fermentation (Fig. 2) and determined the

maximum growth rate of the population (**mu**), the maximum population size, also called the

carrying capacity (**K**) and the relative frequency of each species at the end of mixed cultures

(Table. 1). The fermentations conducted by *S. cerevisiae* alone goes through an exponential

growth rate (**mu$_{Sc}$** $= 5.6 \pm 0.8$ 10^6cells.mL^{-1}.h^{-1}) and reaches a maximum population around 1.5

10^8cells.mL^{-1} (**K$_{Sc}$** $= 1.56 \pm 0.16$ 10^8cells.mL^{-1}) that remains constant until the end of the

fermentation. **Td** fermentations have dynamics comparable to those of **Sc** with a similar growth

rate but a smaller carrying capacity (**K$_{Td}$** $= 1.23 \pm 0.05$ 10^8cells.mL^{-1}). The fermentations

mediated by *Hanseniaspora* species alone had as similar growth dynamics similar to **Sc** at the

beginning of the fermentation with though a higher growth rate (**mu$_{Ho}$** $= 7.7 \pm 2.3$ 10^6cells.mL^{-1}.h^{-1},

mu$_{Hu}$ $= 13.4 \pm 4.3$ 10^6cells.mL^{-1}.h^{-1}). On the opposite, their stationary phase is quite

different from that of *Sc*; characterized by a higher cell mortality with a population drop of

about 70% by the end of the fermentation. The fermentation mediated by *Metschnikowia*

species have growth dynamics globally similar to *Sc* fermentations: a similar growth rate (mu_{mp}

$= 2.3 \pm 0.4 \ 10^6$cells.mL^{-1}.h^{-1}, $mu_{Mf} = 3.4 \pm 0.7 \ 10^6$cells.mL^{-1}.h^{-1}), a much smaller carrying

capacity ($K_{Mp} = 0.6 \pm 0.01 \ 10^6$cells.mL^{-1}, $K_{Mf} = 0.8 \pm 0.05 \ 10^6$cells.mL^{-1}) and no mortality

during the stationary phase. In most cases, the mixed cultures display an intermediated pattern

between the two corresponding isolated cultures (Fig. 2). However, the mixed cultures with

Metschnikowia species harbor a cell mortality during the stationary phase contrary to their

respective isolated cultures. In the case of *ScvsMp* fermentation, only *S. cerevisiae* population

decreases significantly during the stationary phase, while in *ScvsMf* fermentation; both sub

populations significantly diminish. As a measure of fitness, we can also follow the variations

of *S. cerevisiae* frequency along the fermentation. In all mixed cultures, *S. cerevisiae*

significantly increased during the fermentation from 10% initially to frequencies varying

between 32% and 96% at the end of the culture depending on species (Table. 1). In all cases,

except the mix with the *T. delbrueckii* (prop$_{ScvsTd}$ = 0.32 $\pm$ 0.02), *S. cerevisiae* finishes by

dominating the mixed cultures (Table. 1).

We then looked at resources consumption (glucose, fructose and sources of assimilable

nitrogen) and estimated the final concentration of sugars (fructose and glucose) and amino-

acids (Fig. 3, Table 1). The *Sc* fermentations consumed all sugars leaving less than 0.1% of the

initial concentration (Fig. 3a). Among isolated non-sacc, *Td* fermentations was the one

consuming the highest fraction of the initial sugars leaving around 31% of sugars followed by

Ho (45%), *Hu* (67%), *Mf* (68%) and *Mp* (71%). Furthermore, all species but *Hanseniaspora*

opuntiae have consumed more glucose than fructose. The mixed cultures consumed more

sugars than all isolated non-*Saccharomyces* fermentations but less than *Sc* fermentations with

a preference for glucose, indicating the major impact of the *S. cerevisiae* strain in sugar

consumption (consistent with the CO_2 production).

The consumption of nitrogen assimilable sources (NAS, amino-acids and ammonium) harbours

the same pattern (Fig. 3b). NAS was consumed almost entirely in **Sc** isolated culture and in all

mixed populations whereas the fraction of NAS consumed varied between 84% and 94% in

non-*saccharomyces* isolated culture. However, the preference for the different sources of

nitrogen are different for each species (Fig. 3c). Glycine and histidine are the least consumed

by **Td**. **Ho** and **Hu** have similar behaviors: they both consume only half of the ammonium, 90%

of histidine and 89% and 79% of arginine respectively. In the same way, both *Metschnikowia*

species present a similar pattern; the non-exhausted NAS are identical for the two species:

ammonium, glycine, aspartate, arginine tyrosine, phenylalanine and alanine. Therefore, we can

classify NAS in function of their global preference by all tested yeasts. The obtained ranking

by order of preference is: glutamine, methionine, glutamate, valine, threonine, serine,

tryptophan, alanine, histidine, arginine, aspartate, glycine and surprisingly ammonium.

In parallel of the monitoring of the must resources consumption, we also investigated the

production of metabolites from the CCM (Central Carbon Metabolism): ethanol, glycerol,

succinate, pyruvate, acetate and alpha-ketoglutarate (Table 1). Theses productions has been

measured at the end of the cultures where sugar consumption being quite different from one

fermentation to another. Therefore, it appears more appropriate to first compare the production

yields (total production / sugar consumption) instead of the total productions. Moreover, we

decide to compare yields of production relatively to the **Sc** yield that is our reference (Fig. 4).

In the case of ethanol, only **Mf** fermentations has a yield of production statistically than the **Sc**

inferior (-32%). For glycerol, all non-*saccharomyces* grown alone but **Td** had a greater yield

than **Sc**. The mixed fermentations glycerol yield was not significantly different from the Sc one.

Regarding acetate, only *Hanseniaspora* species grown alone or in mixed culture harbored a greater yield than Sc (Fig. 4). For succinate, there could be a negative interaction between *Sc* and *Hanseniaspora* or *Sc* and *Metschnikowia*, as yields in mixed culture were inferior than in the corresponding isolated cultures. However, these differences were not significant after the correction for multiple tests. At last, we did not observe any significant variation of alpha-ketoglutarate yield of production, linked to its lower concentration and difficulty to be precisely measured.

The total production of each fermentation is the result of the combination of species yields, total sugar consumption and respective population dynamics during the fermentation. Therefore, differences observed in the total productions of mixed cultures result from additive or subtractive effects between these 3 parameters. Regarding ethanol, the total production is directly linked to the consumption of resources and all mixed cultures are equivalent to **Sc** fermentation. The case of glycerol is more interesting. Indeed, even if the sugar consumption for **ScvsHu** and **ScvsMp** mixed cultures are lower than *Sc* isolated culture, the total production of glycerol is superior to the corresponding isolated cultures (Glycerol$_{SCvsMp}$ = 6.1±0.1 g.L^{-1}, Glycerol$_{Sc}$ = 5.3±0.4 g.L^{-1}, Glycerol$_{Mp}$ = 3.7±0.2 g.L^{-1}). This result comes from the positive combination of the superior yield of glycerol production of *Hanseniaspora* and *Metschnikowia cells* and the population dynamics. In all other cases, even if the total production does not lie between the production bonds of the corresponding isolated cultures for some mixed cultures, these differences are never statistically significant.

Discussion

This manuscript presents one of the first works focusing on the kinetics of yeast-yeast

interactions between two species during the fermentative process in oenology. Indeed, so far,

most of the work presented in the literature focus on input-output balances (4, 21).

First, this work provides additional information about the population dynamics in winemaking

conditions of non-saccharomyces yeasts that are still poorly known. In agreement with previous

results (12, 22, 23), we found that *T. delbrueckii* has a very similar dynamic than *S. cerevisiae*

and is able to finish the fermentation on its own, while not being as efficient or as fit as *S.*

cerevisiae. However, other studies suggest that this result is not constant and depends on strains

and environmental conditions. The *T. delbrueckii* strain used in Belda et al (14) was less

performant than their *S. cerevisiae* strain. Our results also show that *T. delbrueckii* contrary to

S. cerevisiae does not consume all NAS (the limiting nutrient, (23), leaving half of the glycine

unconsumed. This difference in the consumption of nitrogen sources may explain the ability of

S. cerevisiae to invade mixed culture (12) despite its low inoculation rate. Despite differences

of carbon metabolites production between the isolated cultures of *T. delbrueckii* and *S.*

cerevisiae (linked to the differences in resources consumption), the yields are not statistically

different. The mixed cultures are identical to *S. cerevisiae* pure cultures because of *S. cerevisiae*

invasion. Globally the interaction mechanism between these two species seems to be a simple

competition for resources without much potential for optimization.

The two species, *H. uvarum* and *H. opuntiae* both present a very different pattern in

fermentation from *S. cerevisiae*. Their population dynamics are characterized by a faster growth

rate (reaching a maximum population similar to *S. cerevisiae*), followed by a long phase of

mortality where viability drops to 30% (24). Consequently, these two strains are not able to

finish the fermentation and leave most of the sugars (24). Nitrogen consumption is also specific

to this genus; only half of available ammonium is used. Unlike with *T. delbrueckii*, the yield of

production of carbon metabolites are very different than with *S. cerevisiae* and moreover the

mixed fermentations display cases of transgressive interactions: their glycerol production is

higher than both isolated cultures, despite a lower sugar consumption. These overproductions

of glycerol can be explained by the conjunction of the population dynamics, resource

consumptions (the ***Ho*** and ***Hu*** fermentation leaving ⅔ of sugars) and the individual species

yields of glycerol production which are from two to three times greater than that of *S. cerevisiae*.

However, this interaction it is not found in previous works (24) and, in this present work, seems

to be related to competition for resources without any change in the population dynamic and

metabolism of each strain between the mixed and pure cultures.

At last, the two *Metschnikowia* strains have unexpected fermentation dynamics. The ***Mp*** and

Mf isolated cultures are characterized by a short growth phase reaching a small carrying

capacity and followed by a long stationary phase without mortality. Nevertheless, in spite of

this very high viability, the measured resources (sugars and nitrogen) are not entirely consumed.

Why the cells stop growing remains unclear. A possible explanation could be linked to

unmeasured resources. The third most important resource of the medium is the lipids that are

present in the medium under the form of phytosterols. It has also been shown that some

Metschnikowia species are unable to consume lipids from the medium and therefore are limited

to their own production (see Fig S1). For all yeast species, lipid synthesis requires oxygen and

is therefore impossible in anaerobic conditions. From these elements, one can hypothesize that

Metschnikowia cells are producing their own lipids from oxygen at the beginning of the

fermentation and once the oxygen is exhausted they stop growing. Interestingly, despite the fact

that isolated cultures of *S. cerevisiae*, *M. pulcherrima* and *M. fructicola* did not present any

mortality, their corresponding mixed cultures (***ScvsMp*** and ***ScvsMf***) present 30% mortality.

Moreover, this mortality seems to impact differently both species. In the mixed **ScvsMp**

fermentation only the *S. cerevisiae* cells are dying; in **ScvsMf** both species are impacted. To

explain the survival of *M. pulcherrima* cells compared to *S. cerevisiae* cells, it can be

hypothesized that *Metschnikowia* cells produce toxins (a ferric salt of pulcherriminic acid, (25)

that kill *S. cerevisiae* cells (16, 26, 27). In **ScvsMf** cultures, the mortality of both species

suggests an interaction mechanism different from the one observed between *M. pulcherrima*

and *S. cerevisiae*. A two-phase interaction could explain such pattern : *M. fructicola* synthesizes

a toxin that kills only *S. cerevisiae* cells and thereafter *M. fructicola* cells dies for another

reason, like ethanol sensibility. Indeed, the production of ethanol is almost four time greater in

mixed cultures than in isolated *Metschnikowia* fermentations. The reason why no loss of

viability is observed for *M. pulcherrima* when placed in mixed culture with *S. cerevisiae* is

probably a better tolerance to ethanol stress compared to *M. fructicola*. As a conclusion, the

interaction mechanism between the *Metschnikowia* species and *S. cerevisiae* appears to be very

different from that between *S. cerevisiae* and the other studied non-saccharomyces species. It

is not a competition for resources in sugars or nitrogen but rather the possibility of a killer toxin

synthesized by the *Metschnikowia* strains or a competition for a non-measured resource (lipids,

vitamins, minerals).

If we look globally, we can first note that all mixed cultures finished the fermentation. No major

antagonistic phenomena were identified between the *S. cerevisiae* strain chosen and the "non-

Saccharomyces" yeasts studied. Moreover, despite the presence of differences in yield and

interactions between strains, the rapid dominance of *S. cerevisiae* (representing however only

10% of the population at t0) results at mixed cultures that are overall little different from *S.*

cerevisiae pure cultures. This result confirms the good adaptation of *S. cerevisiae* to the

winemaking conditions (4–6, 21). Nevertheless, interesting data appear in our research work.

Indeed, despite the pre-eminence of *S. cerevisiae*, we have already highlighted transgressive

interactions between *S. cerevisiae* and *H. uvarum* and *M. pulcherrima* leading to the

overproduction of glycerol in mixed cultures. This observation opens the door for potential

more interesting results: when *S. cerevisiae* is inoculated at lower frequencies (0.01, 0.001), the

fact that production yields are different between the species can lead to more important

transgressive interactions (lower ethanol production for example) in mixed cultures.

Furthermore, most of detected interactions can be explained by competition for resources or the

production of toxins without indication of an alteration of individual cell behavior linked to the

presence of other species in the same medium. This observation linked to our deep phenotyping

of population dynamics could allow the modeling of mixed oenological fermentations. Using

such model, transgressive interactions linked to consumption of resources could be investigated

and inform the choice of couple of species and the protocol of inoculations (specifically by

changing the *S. cerevisiae* starting frequency). It could be used to predict mixed cultures

dynamics from data on isolated cultures, potentially saving a lot of time and money and open

the possibility of ecological engineering. The development of such mathematical model will

only be possible thanks to a deep tracking of population dynamics to understand underlying

mechanisms of growth and mortality. Therefore, to better identify and understand yeast-yeast

interaction mechanisms, we should extend our deep phenotyping of mixed population dynamics

to additional species and investigate intra-specific variability as well as interactions with culture

environments.

319 **Mat & Met**

In this work, we used one strain of 6 different species (one strain by species): *Saccharomyces*

cerevisiae (***Sc***), *Torulaspora delbrueckii* (***Td***), *Metschnikowia pulcherrima* (***Mp***),

Metschnikowia fructicola (**Mf**)*, Hanseniaspora uvarum* (**Hu**) and *Hanseniaspora opuntiae*

(**Ho**). The *S. cerevisiae* strain is a haploid strain marked with GFP (59A-GFP, (2)). The 5 non-

Saccharomyces strains are natural strains isolated from viognier and sauvignon must in

different regions of France. For each species, 3 replicates of isolated cultures were performed

(except for *S. cerevisiae* that have a total of 8 replicates in different blocks). In addition, for

each non-*Saccharomyces*, 3 replicates of a mixed culture with the Sc strain was performed. In

all mixed fermentations, the starting proportion of *S.cerevisiae* cells was 10%. In this text, the

fermentations are referred by the species that perform them. The isolated cultures are referred

as: **Sc**, **Td**, **Mp**, **Mf**, **Hu** and **Ho** and the mixed cultures as **ScvsTd, ScvsMp, ScvsMf, ScvsHu**

and **ScvsHo**.

Initial cultures (12 h, in 50 ml YPD medium, 28 °C) were used to inoculate fermentation at a

density of 10^6 cells/ml therefore for mixed culture the density of *S. cerevisiae* cells was 0.1×10^6

cells/mL and the density of non-sacc cells was 0.9×10^6 cells/mL. Fermentations were carried

out in synthetic medium (SM) mimicking standard grape juice (28). The SM used in this study

contained 200 g/L of sugar (100 g/L of glucose and 100 g/L of fructose) and 200 mg/L of

assimilable nitrogen (as a mixed of ammonium chloride and amino acids). The concentrations

of weak acids, salts and vitamins were identical to those described by Seguinot et al. (29). The

pH of the medium was adjusted to 3.3 with 10M of NaOH. The SM medium was first saturated

with bubbling air during 40 minutes, then it was supplemented with 5 mg/L of phytosterols

(85451, Sigma Aldrich) to satisfy the lipid requirements of yeast cells during anaerobic growth.

Fermentation took place in 1.1-liter fermentors equipped with fermentation locks to maintain

anaerobiosis, at 20 °C, with continuous magnetic stirring (500 rpm) during approximately 200h.

CO_2 release was followed by automatic measurements of fermentor weight loss every 20 min.

The amount of CO_2 released allowed us to monitor the progress of the fermentation and evaluate

the maximum of released CO_2 (***CO2max***) as well as the maximum rate of CO_2 released (***Vmax***).

Samples were harvested after 6h, 12h and 24h, then every 12h during the first week and every

24h during the second week of fermentation. At each sample, the population density, the

viability and the proportion of *S. cerevisiae* cells were determined using a flow cytometer BD

Accuri™ C6 Plus as described in Delobel et al. (30). From these population data and by fitting

a growth population model (with the grofit package of R, (31), the carrying capacity (**K**) and

the maximum growth rate (**mu**) for each fermentation were estimated.

The final concentration of carbon metabolites in the medium (acetate, succinate, glycerol,

alpha-ketoglutarate, pyruvate, ethanol, glucose and fructose) with high-pressure liquid

chromatography (32). From these metabolite concentrations, we first calculated the consumed

sugar concentration as the difference between the final and the initial concentration of either

glucose or fructose. Then we calculated the yield of metabolite production by dividing the final

concentration by the corresponding consumed sugar concentration. Finally, we report these

yields to the yield of *S. cerevisiae* isolated cultures that we consider as our reference.

At last, the ammonium concentration after 100h of fermentation was determined enzymatically

with R-Biopharm (Darmstadt, Germany) and the free amino acid content of the must was

estimated with cation exchange chromatography with post-column ninhydrin derivatization

(33).

The experimental work was performed in 5 different blocks. Each block was composed of three

replicates of non-*Saccharomyces* fermentations (for example **Td**), three replicates of the

corresponding mixed fermentations with *S. cerevisiae* (for example **ScvsTd**) and one or two

fermentations of isolated *S. cerevisiae* cells (**Sc**). The block effect was evaluated on the

parameters of the **Sc** fermentation. For most of the studied parameters the block effect was not

significant. For the parameters where there was a block effect (mu and K), a statistical

correction for the block effect did not modify our results. Therefore, for simplification, we

consider that there was no block effect and compare all fermentations without any correction

for block effect. For each measured parameters, an anova was performed to evaluated the type

of fermentation (***Sc***, ***Td***, ***Mp***, ***Mf***, ***Hu***, ***Ho***, ***ScvsTd***, ***ScvsMp***, ***ScvsMf***, ***ScvsHu*** and ***ScvsHo***)

effect and then a Tukey-test was performed to determine statistical groups and two by two

statistical differences. All statistical analyses were performed using R (34) and Rstudio (35).

All data, analysis and figures scripts can be found in this github adresse:

https://github.com/tnidelet/Harle-et-al-2018.

Acknowledgments.

380　We thank Christian Picou, Marc Perez, Faiza Macna for technical assistance and Delphine

381　Sicard for her advice.

References

1.　Barot S, Allard V, Cantarel A, Enjalbert J, Gauffreteau A, Goldringer I, Lata J-C, Roux

X, Niboyet A, Porcher E. 2017. Designing mixtures of varieties for multifunctional

agriculture with the help of ecology. A review. Agron Sustain Dev 37:13.

2.　Marsit S, Dequin S. 2015. Diversity and adaptive evolution of Saccharomyces wine yeast:

a review. FEMS Yeast Research 15:fov067.

3.　Sablayrolles J. 2008. Fermented beverages: the example of winemaking. Advances in

Fermentation Technology New Delhi: Asiatech Publishers 322–47.

4.　Varela C. 2016. The impact of non-Saccharomyces yeasts in the production of alcoholic

beverages. Appl Microbiol Biot 100:9861–9874.

5.　Jolly NP, Varela C, Pretorius IS. 2014. Not your ordinary yeast: non-Saccharomyces

yeasts in wine production uncovered. Fems Yeast Res 14:215–237.

6.　Pinto C, Pinho D, Cardoso R, Custódio V, Fernandes J, Sousa S, Pinheiro M, Egas C,

Gomes AC. 2015. Wine fermentation microbiome: a landscape from different Portuguese

wine appellations 6:905.

7.　García-Ríos E, Gutiérrez A, Salvadó Z, Arroyo-López FN, Guillamon JM. 2013. Fitness

advantage of commercial wine yeasts according to nitrogen concentration, temperature

and ethanol content under microvinification conditions. Appl Environ Microbiol

AEM.03405-13.

8.　Ivey M, Massel M, Phister TG. 2013. Microbial Interactions in Food Fermentations.

Annual Review of Food Science and Technology 4:141–162.

9.　Sadineni V, Kondapalli N, Obulam V. 2012. Effect of co-fermentation with

Saccharomyces cerevisiae and Torulaspora delbrueckii or Metschnikowia pulcherrima on

the aroma and sensory properties of mango wine. Ann Microbiol 62:1353–1360.

10.　Hernández-Orte P, Cersosimo M, Loscos N, Cacho J, Garcia-Moruno E, Ferreira V. 2008.

The development of varietal aroma from non-floral grapes by yeasts of different genera.

Food Chemistry 107:1064–1077.

11.　Bely M, Stoeckle P, Isabelle M-P, Dubourdieu D. 2008. Impact of mixed Torulaspora

delbrueckii–Saccharomyces cerevisiae culture on high-sugar fermentation. Int J Food

Microbiol 122:312–320.

12. Azzolini M, Fedrizzi B, Tosi E, Finato F, Vagnoli P, Scrinzi C, Zapparoli G. 2012. Effects

of Torulaspora delbrueckii and Saccharomyces cerevisiae mixed cultures on fermentation

and aroma of Amarone wine 235:303–313.

13. Contreras A, Hidalgo C, Henschke PA, Chambers PJ, Curtin C, Varela C. 2014.

Evaluation of Non-Saccharomyces Yeasts for the Reduction of Alcohol Content in Wine.

Appl Environ Microbiol 80:1670–1678.

14. Belda I, Navascués E, Marquina D, Santos A, Calderon F, Benito S. 2015. Dynamic

analysis of physiological properties of Torulaspora delbrueckii in wine fermentations and

its incidence on wine quality. Appl Microbiol Biot 99:1911–1922.

15. González-Royo E, Pascual O, Kontoudakis N, Esteruelas M, Esteve-Zarzoso B, Mas A,

Canals JM, Zamora F. 2015. Oenological consequences of sequential inoculation with

non-<Emphasis Type="Italic">Saccharomyces</Emphasis> yeasts (<Emphasis

Type="Italic">Torulaspora delbrueckii</Emphasis> or <Emphasis

Type="Italic">Metschnikowia pulcherrima</Emphasis>) and <Emphasis

Type="Italic">Saccharomyces cerevisiae</Emphasis> in base wine for sparkling wine

production. Eur Food Res Technol 240:999–1012.

16. Oro L, Ciani M, Comitini F. 2014. Antimicrobial activity of Metschnikowia pulcherrima

on wine yeasts. J Appl Microbiol 116:1209–1217.

17. Kurtzman CP, Droby S. 2001. Metschnikowia fructicola, a New Ascosporic Yeast with

Potential for Biocontrol of Postharvest Fruit Rots. Systematic and Applied Microbiology

24:395–399.

18.   Hu K, Qin Y, Tao Y-S, Zhu X-L, Peng C-T, Ullah N. Potential of Glycosidase from Non-

434   Saccharomyces Isolates for Enhancement of Wine Aroma. Journal of Food Science

435   81:M935–M943.

436   19.   Tristezza M, Tufariello M, Capozzi V, Spano G, Mita G, Grieco F. 2016. The Oenological

437   Potential of Hanseniaspora uvarum in Simultaneous and Sequential Co-fermentation with

438   Saccharomyces cerevisiae for Industrial Wine Production. Front Microbiol 7.

439   20.   Song H-S, Cannon WR, Beliaev AS, Konopka A. 2014. Mathematical Modeling of

440   Microbial Community Dynamics: A Methodological Review. Processes 2:711–752.

441   21.   García M, Braulio E-Z, Arroyo T. 2016. Non-Saccharomyces Yeasts: Biotechnological

442   Role for Wine Production.

443   22.   van Breda V, Jolly N, van Wyk J. 2013. Characterisation of commercial and natural

444   Torulaspora delbrueckii wine yeast strains. Int J Food Microbiol 163:80–88.

445   23.   Taillandier P, Lai Q, Anne J-O, Brandam C. 2014. Interactions between Torulaspora

446   delbrueckii and Saccharomyces cerevisiae in wine fermentation: influence of inoculation

447   and nitrogen content 30:1959–1967.

448   24.   Moreira N, Mendes F, de Pinho P, journal of … H-T. 2008. Heavy sulphur compounds,

449   higher alcohols and esters production profile of Hanseniaspora uvarum and Hanseniaspora

450   guilliermondii grown as pure and …. International journal of ….

451   25.   MacDonald. 1965. Biosynthesis of pulcherriminic acid. Biochem J 96:533–8.

452   26.   Sipiczki M. 2006. Metschnikowia Strains Isolated from Botrytized Grapes Antagonize

453   Fungal and Bacterial Growth by Iron Depletion. Appl Environ Microb 72:6716–6724.

27. Türkel S, Ener B. 2009. Isolation and Characterization of New Metschnikowia

pulcherrima Strains as Producers of the Antimicrobial Pigment Pulcherrimin. Zeitschrift

Für Naturforschung C 64:405–410.

28. Bely M, Sablayrolles JM, Barre P. 1990. Description of Alcoholic Fermentation Kinetics:

Its Variability and Significance. Am J Enol Vitic 41:319–324.

29. Seguinot P, SANCHEZ I, Bloem A, Ortiz-Julien A, Camarasa C. 2017. Impact of nitrogen

sources on the fermentative kinetic of non - Saccharomyces yeasts.

30. Delobel P, Pradal M, Blondin B, Tesniere C. A 'fragile cell' sub-population revealed

during cytometric assessment of Saccharomyces cerevisiae viability in lipid-limited

alcoholic fermentation. Letters in Applied Microbiology 55:338–344.

31. grofit: Fitting Biological Growth Curves with R | Kahm | Journal of Statistical Software.

32. Camarasa C, Sanchez I, Brial P, Bigey F, Dequin S. 2011. Phenotypic landscape of

Saccharomyces cerevisiae during wine fermentation: evidence for origin-dependent

metabolic traits. PLoS ONE 6:e25147.

33. Crépin L, Nidelet T, Sanchez I, Dequin S, Camarasa C. 2012. Sequential Use of Nitrogen

Compounds by Saccharomyces cerevisiae during Wine Fermentation: a Model Based on

Kinetic and Regulation Characteristics of Nitrogen Permeases 78:8102–8111.

34. R Core Team (2016). R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

35. RStudio (2012). RStudio: Integrated development environment for R (Version 0.96.122)

[Computer software]. Boston, MA. Retrieved May 20, 2012. Available from

http://www.rstudio.org/.

Tables

Table 1

Parameter values (mean ± standard deviation) for each type of fermentation.

	Mu	K	prop	Vmax	MaxCO2	Sugar	NAS	Ethanol	Glycerol	Succinate	Acetate	Pyruvate	Alpha
Sc	5.6±0.8^{cd}	1.6±0.2^{bc}	NA	0.99±0.03^{a}	88.2±2.2^{a}	0.2±0.2^{f}	10.4±8.6^{d}	95.2±8.4^{a}	5.3±0.4^{c}	1.05±0.28^{a}	0.63±0.06^{d}	0.12±0.08^{b}	0.08±0.04^{ab}
Ho	7.7±2.3^{bc}	1.6±0.1^{bc}	NA	0.42±0.02^{d}	46.4±0.6^{d}	92.3±1.4^{b}	122.2±3.5^{bc}	50.1±0.2^{c}	5.2±0.0^{c}	0.35±0.00^{b}	0.99±0.02^{b}	0.13±0.05^{ab}	0.09±0.1^{ab}
ScvsHo	5.7±0.3^{bcd}	**2.3±0.1^{a}**	0.83±0.02^{b}	0.57±0.01^{c}	82.7±0.6^{b}	23.3±1.3^{de}	**6.4±2.4^{d}**	87.2±0.8^{d}	**6.0±0.0^{abc}**	0.42±0.04^{b}	0.84±0.01^{c}	**0.08±0.00^{b}**	**0.03±0.0^{b}**
Hu	15.6±2.9^{a}	1.4±0.1^{cd}	NA	0.34±0.04^{d}	30.7±0.5^{e}	134.3±11.2^{a}	173.3±24.8^{b}	34.9±1.1^{d}	5.5±0.2^{bc}	0.66±0.03^{b}	1.15±0.03^{a}	0.17±0.04^{ab}	0.14±0.09^{ab}
ScvsHu	8.7±0.6^{b}	**1.7±0.1^{b}**	0.76±0.01^{c}	0.53±0.04^{c}	82.1±1.5^{b}	16.5±6.5^{e}	**4.6±4.2^{d}**	89.5±3.8^{d}	**6.6±0.3^{a}**	**0.60±0.06^{b}**	0.86±0.07^{bc}	0.16±0.04^{ab}	0.09±0.06^{ab}
Mf	3.4±0.8^{de}	0.8±0.05^{e}	NA	0.17±0.04^{e}	20.6±1.2^{f}	135.8±9.7^{a}	256.6±2.9^{a}	20.6±0.3^{d}	2.8±0.1^{e}	0.29±0.01^{b}	0.13±0.01^{g}	0.10±0.01^{b}	0.10±0.00^{ab}
ScvsMf	**3.1±0.2^{de}**	1.3±0.03^{d}	0.96±0.02^{a}	0.76±0.03^{b}	86.8±0.7^{ab}	9.9±2.1^{ef}	**1.8±0.1^{d}**	92.1±3.2^{d}	**6.0±0.1^{abc}**	0.58±0.02^{b}	0.46±0.02^{e}	0.12±0.01^{b}	0.08±0.01^{ab}
Mp	2.4±0.4^{e}	0.6±0.01^{e}	NA	0.16±0.02^{e}	21.8±0.4^{f}	142±3.7^{a}	305.5±5.9^{a}	23.4±1.8^{d}	3.7±0.2^{d}	0.26±0.03^{b}	0.07±0.06^{g}	0.29±0.04^{ab}	0.13±0.01^{ab}
ScvsMp	2.8±0.1^{de}	1.3±0.01^{d}	0.50±0.02^{d}	0.68±0.04^{b}	83.3±2.2^{b}	20.8±6.0^{de}	**2.1±0.2^{d}**	86.5±3.4^{ab}	**6.1±0.1^{ab}**	0.52±0.04^{b}	0.46±0.02^{e}	0.24±0.01^{ab}	**0.19±0.01^{ab}**
Td	5.3±1.0^{cde}	1.2±0.05^{d}	NA	0.51±0.06^{c}	65.7±4.0^{c}	63.5±11.5^{c}	112.6±21.1^{c}	56±5.3^{c}	4.0±0.3^{d}	0.42±0.04^{b}	0.17±0.03^{fg}	0.14±0.06^{ab}	0.02±0.01^{b}
ScvsTd	**4.2±0.6^{de}**	1.5±0.01^{bcd}	0.32±0.02^{e}	0.72±0.02^{b}	83.9±3.2^{ab}	35.8±5.2^{d}	**1.8±0.1^{d}**	76.8±3.7^{b}	5.2±0.2^{c}	0.63±0.06^{b}	0.27±0.06^{f}	**0.20±0.11^{ab}**	0.05±0.03^{b}

Mu : 10^6cells.mL^{-1}.h^{-1}; K : 10^8 cells.mL^{-1}, prop : no unity, Vmax : g.L^{-1}.h^{-1}, MaxCO2 : g.L^{-1}, Final Sugar : g.L^{-1}, Final NAS : mg.L^{-1}, Ethanol : g.L^{-1}, Glycerol : g.L^{-1}, Succinate : g.L^{-1}, Acetate : g.L^{-1}, Pyruvate : g.L^{-1}, Alpha : g.L^{-1}

Bold : values of mixed culture, that are not between the two values of corresponding isolated cultures, sign of transgressive interactions.

Superscript : Statistical groups obtain from a Tukey test

501 # Figures

503 **FIG 1**

504 Representation of a) the maximum rate of CO_2 production (***Vmax***) and b) the total CO_2

505 produced in function of the species driving each fermentation (average ± standard deviation).

506 The small letters indicate the statistical groups from a Tukey analysis.

FIG2

Global monitoring of the kinetics of the total living population (left), and sub-population in the mixed cultures (right) across the fermentation. All points represent a sample (average ± standard error). The full lines are for total population and the dashed lines for the two sub populations into mixed cultures. At the end of dashed lines, the final proportion of both sub-populations in mixed cultures is indicated. The light colors represent pure culture of 'non-

Saccharomyces' strains and the darker to the corresponding culture in competition with *S. cerevisiae*. The isolated cultures of *S. cerevisiae* are represented in black. **Sc, Mp, Mf, Ho, Hu, Td, ScvsMp, ScvsMf, ScvsHo, ScvsHu, ScvsTd** correspond respectively to cultures with: isolated *S. cerevisiae*, isolated *M. pulcherrima*, isolated *M. fructicola*, isolated *H. opuntiae*, isolated *H.uvarum*, isolated *T. delbrueckii*, *M. pulcherrima* mixed with *S. cerevisiae*, *M. fructicola* mixed with *S. cerevisiae*, *H. opuntiae* mixed with *S. cerevisiae*, *H. uvarum* mixed with *S. cerevisiae* and *T. delbrueckii* mixed with *S. cerevisiae*.

C

	Mp	Mf	Td	Hu	Ho	Sc vs Ho	Sc vs Hu	Sc	Sc vs Mp	Sc vs Td	Sc vs Mf
Glutamine	99%	100%	99%	100%	100%	100%	100%	100%	100%	100%	100%
Methionine	90%	100%	100%	100%	100%	100%	100%	100%	100%	100%	100%
Glutamate	98%	97%	99%	99%	99%	99%	99%	99%	100%	100%	99%
Valine	86%	97%	97%	100%	100%	100%	100%	100%	100%	100%	100%
Threonine	87%	94%	100%	100%	100%	100%	100%	100%	100%	100%	100%
Serine	87%	93%	99%	99%	99%	99%	100%	100%	100%	100%	100%
Tryptophan	82%	85%	85%	99%	100%	100%	100%	100%	100%	100%	100%
Alanine	72%	77%	97%	100%	100%	99%	100%	100%	100%	100%	100%
Histidine	91%	98%	73%	91%	91%	100%	100%	99%	100%	100%	100%
Phenylalanine	69%	71%	99%	100%	100%	100%	100%	100%	100%	100%	100%
Tyrosine	66%	66%	96%	100%	100%	100%	100%	100%	100%	100%	100%
Arginine	80%	82%	96%	79%	89%	100%	100%	98%	100%	100%	100%
Aspartate	65%	72%	98%	97%	97%	97%	97%	98%	97%	98%	98%
Glycine	65%	69%	44%	87%	95%	92%	94%	98%	98%	98%	99%
Ammonium	60%	59%	86%	51%	53%	100%	100%	100%	100%	100%	100%

FIG3

Consumption of sugars and nitrogen assimilable sources (NAS) for each type of

fermentation. A) Final concentration of sugar (average ± standard deviation). B) Final

concentration of NAS (average ± standard deviation). C) Percentage of consumption of each

NAS in each type of fermentation represented as a color gradient from green (<75 %) to red (>

75%). *Sc, Mp, Mf, Ho, Hu, Td, ScvsMp, ScvsMf, ScvsHo, ScvsHu, ScvsTd* correspond

respectively to cultures with: isolated *S. cerevisiae*, isolated *M. pulcherrima*, isolated *M.*

fructicola, isolated *H. opuntiae*, isolated *H.uvarum*, isolated *T. delbrueckii*, *M. pulcherrima*

mixed with *S. cerevisiae*, *M. fructicola* mixed with *S. cerevisiae*, *H. opuntiae* mixed with *S.*

cerevisiae, *H. uvarum* mixed with *S. cerevisiae* and *T. delbrueckii* mixed with *S. cerevisiae*.

FIG4

Yield (average ± standard deviation) of carbon metabolite production relative to the yield of

production of isolated *S. cerevisiae* cultures for acetate, alpha-ketoglutarate, ethanol, glycerol,

pyruvate and succinate and each type of fermentation. *Sc, Mp, Mf, Ho, Hu, Td, ScvsMp,*

ScvsMf, ScvsHo, ScvsHu, ScvsTd correspond respectively to cultures with: isolated *S.*

cerevisiae, isolated *M. pulcherrima*, isolated *M. fructicola*, isolated *H. opuntiae*, isolated

H.uvarum, isolated *T. delbrueckii*, *M. pulcherrima* mixed with *S. cerevisiae*, *M. fructicola*

mixed with *S. cerevisiae*, *H. opuntiae* mixed with *S. cerevisiae*, *H. uvarum* mixed with *S.*

cerevisiae and *T. delbrueckii* mixed with *S. cerevisiae*.