Kinetic analysis of yeast-yeast interactions in oenological conditions

Oliver Harlé[1], Judith Legrand[3], Catherine Tesnière[2], Martine Pradal[2], Jean-Roch Mouret[2], Thibault Nidelet[2*]

Affiliations :

1 : STLO, INRA, Agrocampus Ouest, Rennes, France

2 : SPO, Univ Montpellier, INRA, Montpellier SupAgro, Montpellier, France

3 : GQE-Le Moulon, INRA, Univ. Paris-Sud, CNRS, AgroParisTech, Université Paris-Saclay, 91190 Gif-sur-Yvette, France

[*] **Corresponding Author**: thibault.nidelet@inra.fr

keywords: winemaking fermentation, population dynamics, non-*Saccharomyces* species, transgressive interactions, ecosystems

Abstract

Fermentation by microorganisms is a key step in the production of traditional food products such as bread, cheese, beer and wine. In these fermentative ecosystems, microorganisms interact in various ways, namely competition, predation, commensalism and mutualism. Traditional wine fermentation is a complex microbial process performed by *Saccharomyces* and non-*Saccharomyces* yeast species. To better understand the different interactions occurring

within wine fermentation, isolated yeast cultures were compared with mixed co-cultures of one reference strain of *S. cerevisiae* with one strain of four non-*Saccharomyces* yeast species (*Metschnikowia pulcherrima, M. fructicola, Hanseniaspora opuntiae and H. uvarum*). In each case, we studied population dynamics, resource consumption and the production of metabolites from central carbon metabolism. This deep phenotyping of competition kinetics allowed us to identify the main mechanisms of interaction between species. *S. cerevisiae* competed with *H. uvarum* and *H. opuntiae* for resources although both *Hanseniaspora* species were characterized by a strong mortality either in isolated or mixed fermentations. *M. pulcherrima* and *M. fructicola* displayed a negative interaction with the *S. cerevisiae* strain tested, with a decrease in viability in co-culture, probably due to iron depletion via the production of pulcherriminic acid. Overall, this work highlights the importance of measuring cell populations and their metabolite kinetics to understand yeast-yeast interactions. These results are a first step towards ecological engineering and the rational design of optimal multi-species starter consortia using modeling tools.

Introduction

In natural or anthropized environments, microbial species are part of an ecosystem and interact positively or negatively, forming a complex network. Until recently, process optimization in agriculture or food processing was mostly based on the selection of single strains. However, this paradigm is now being challenged and the scientific community is increasingly seeking to exploit and optimize consortia of several strains and/or species. Indeed, many studies have shown that more diverse anthropized environments have many advantages in terms of resilience, disease resistance or yield (Barot et al., 2017). Efforts are now being made to design

optimal consortia of various species and strains whose interactions will be exploited to

maximize given criteria such as fermentation quality, aromatic complexity or other organoleptic

characteristics.

Wine fermentation is both an economically and societally important food ecosystem, where the

addition of fermentation 'starters' composed of selected yeasts at the beginning of the

fermentation process is common. In fact, around 80% of oenological fermentations worldwide

are conducted with starters (Marsit and Dequin, 2015; Sablayrolles, 2008). Most often, these

"starters" are only composed of a single *Saccharomyces cerevisiae* strain selected for its ability

to complete fermentation. Indeed, numerous experiments have shown that *S. cerevisiae*, with

an initially low population, most often becomes the predominant species at the end of the

fermentation, demonstrating its superior fermentative abilities (García-Ríos et al., 2014; Jolly

et al., 2014; Pinto et al., 2015; Varela, 2016). However, in recent years, multi-species starters

have emerged, aiming at increasing the aromatic complexity of wines. They most often combine

one strain of *S. cerevisiae* allowing to complete fermentation and another species, often from a

different genus, contributing to a greater variety of flavors (García-Ríos et al., 2014; Ivey et al.,

2013; Jolly et al., 2014; Varela, 2016).

Indeed, there are numerous experiments and even industrial products making use of such mixed

starters to improve wine's organoleptic qualities (Jolly et al., 2014; P et al., 2008; Sadineni et

al., 2012). The non-*Saccharomyces* strains used in these experiments are very diverse, with

more than 23 different species (Jolly et al., 2014) including *Torulaspora delbrueckii*,

Metschnikowia pulcherrima, *Metschnikowia fructicola*, *Hanseniaspora opuntiae* and

Hanseniaspora uvarum. Species in the *Metschnikowia* genus ferment poorly in oenological

conditions but can have interesting attributes (Varela, 2016): in conjunction with *S. cerevisiae*,

M. pulcherrima can reduce ethanol concentrations (Contreras et al., 2014; Varela, 2016),

increase 'citrus/grape fruit' and 'pear' attributes (Belda et al., 2015), as well as allow the

persistence of 'smocky' and 'flowery' characteristics (Elena et al., 2015). *M. pulcherrima* also

has an amensalism effect on *S. cerevisiae* through iron depletion via the production of

pulcherriminic acid (Oro et al., 2014). *M. fructicola* has been less studied and never in

conjunction with *S. cerevisiae* although it presents the interesting ability to inhibit *Botrytis*

growth (Kurtzman and Droby, 2001). Last, the *Hanseniaspora* genus, studied in sequential or

simultaneous fermentation with *S. cerevisiae,* has been shown to increase volatile compound

production during winemaking (Varela, 2016). It notably increased the 'tropical fruit', 'berry',

'floral' and 'nut aroma' characters (Hu et al., n.d.), that were linked to higher concentrations of

acetate esters, esters of MCFAs, isoamyl alcohol, 2-phenylethanol and α-terpineol (Tristezza et

al., 2016).

Despite these various studies, the composition and protocol of inoculation of these multi-strains

starters are still very empirical and only based on the input/output balance, without considering

the dynamics of the microbial populations or their interactions. This lack of knowledge about

yeast-yeast interactions prevents implementing a rational design of multi-strain starters (Song

et al., 2014). To address this problem, we decided to focus our study on population dynamics

and metabolite production during oenological fermentations performed in isolated or mixed

yeast cultures. Since our goal was not to obtain optimal mixes but to understand the mechanism

of microbial interaction, we chose to compare the population dynamics and yields between

isolated cultures of strains from five species (one *S. cerevisiae* and four non-*Saccharomyces*)

and four corresponding mixed cultures always including the *S. cerevisiae* strain as reference.

We were thus able to identify key microbial interaction mechanisms that are further discussed.

Results

In this work, we compared in winemaking conditions the performance of isolated cultures of

five different strains from five yeast species (*Saccharomyces cerevisiae*, *Metschnikowia*

pulcherrima, *Metschnikowia fructicola*, *Hanseniaspora opuntiae* and *Hanseniaspora uvarum*)

and mixed co-cultures combining each of the four non-*Saccharomyces* species with one GFP-

labelled *S. cerevisiae* strain representing 10% of the initial inoculate. As our research objective

was not to evaluate the final product but to study interaction dynamics between yeast species,

it was decided to stop the monitoring of fermentations at a given time, even if the sugar supply

was not completely exhausted. For all fermentation with the *S. cerevisiae* reference strain,

sugars were exhausted after around 200-220 h while in fermentations with isolated non-

Saccharomyces strains, the sugar supply was still not exhausted after 400h. Here, we focused

on the first 300 hours of fermentation.

By comparing the output of isolated and mixed cultures, we evaluated the intensity of yeast-

yeast interactions and/or their consequences on ecosystem service production.

CO$_2$ kinetics

We first investigated the influence of species and co-culture on the dynamics of CO_2 production

(proportional to sugar consumption), which is a good indicator of the fermentation progress.

Indeed, CO_2 production is easy to monitor (based on weight measurement) and is directly

proportional to ethanol synthesis and sugar consumption. The values of the maximum rate of

CO_2 production (***Vmax***, Fig. 1a, Table. 1) and of the maximum CO_2 produced were estimated

(***CO₂max***, Fig. 1b, Table. 1). ***Vmax*** was highly dependent on the species (p.value < 0.001): *S.*

cerevisiae cultures (***Sc***) displayed the highest value (***Vmax$_{Sc}$*** = 0.99 ± 0.02 g.L^{-1}.h^{-1}), followed

by both *Hanseniaspora* species (***Vmax$_{Hu}$*** = 0.33 ± 0.04 g.L^{-1}.h^{-1}, and ***Vmax$_{Ho}$*** = 0.42 ± 0.02

g.L^{-1}.h^{-1}) and finally both *Metschnikowia* species (***Vmax$_{Mp}$*** = 0.165 ± 0.02 g.L^{-1}.h^{-1} and ***Vmax$_{Mf}$***

= 0.17 ± 0.04 g.L^{-1}.h^{-1}). The four mixed cultures had intermediate ***Vmax*** values between those

of ***Sc*** and the highest **Vmax** of all *non-sacc* cultures (Fig. 1a). Mixed cultures containing

Metschnikowia species had significantly higher ***Vmax*** values than those containing

Hanseniaspora species (Fig 1a). Although we did not monitor all cultures until the exhaustion

of glucose and fructose, it was however possible to estimate the capacity of a given species to

complete fermentation by estimating the amount of CO_2 produced during the first 300 hours.

Sc fermentations finished after around 220 hours with a ***CO2max$_{Sc}$*** = 88.2 ± 2.2 g.L^{-1}. We

therefore can make the hypothesis that all cultures that produced more than 80g CO_2.L^{-1} (90%

of ***Sc*** maximum) within 300 hours will be able to complete fermentation. Under this

assumption, all mixed cultures, but not isolated non-*Saccharomyces* cultures, would eventually

complete fermentation. Among the latter cultures , both *Hanseniaspora* species had the highest

CO₂max (***Hu*** 30 ± 0.4 g.L^{-1}, and ***Ho*** 46 ± 0.6 g.L^{-1}) followed by *Metschnikowia* species (***Mp*** 22

± 0.4 g.L^{-1} and ***Mf*** 20 ± 1 g.L^{-1}).

Population Kinetics

We also looked at population dynamics in each culture (Fig. 2) and determined the maximum

growth rate of the population (***µ***), the maximum population size, also termed carrying capacity

(***K***) and the relative abundance of each species after 300 hours of mixed culture, corresponding

in our case to the end of the monitoring period (Table. 1). Fermentations with *S. cerevisiae*

alone went through an exponential growth rate (***µ$_{Sc}$*** = 0.15 ± 0.02 h^{-1}) and reached a maximum

population of around 1.5×10^8cells.mL^{-1} (***K$_{Sc}$*** = 1.55 ± 0.15 10^8cells.mL^{-1}) that remained constant

until the end of the fermentation. Fermentations with either *Hanseniaspora* species alone had a

growth dynamic similar to ***Sc*** at the beginning of the fermentation but a higher growth rate (***µ$_{Ho}$***

= 0.19 ± 0.03 h^{-1}, ***µ$_{Hu}$*** = 0.62 ± 0.18 h^{-1}). On the opposite, their stationary phase was quite

different from that of ***Sc*** and characterized by a higher cell mortality with a population drop of

about 70% by the end of the process. Fermentations performed by *Metschnikowia* species in

isolation had growth dynamics mostly similar to **Sc** fermentations: a similar growth rate (μ_{mp} =

0.18 ± 0.03 h^{-1}, μ_{Mf} = 0.17 ± 0.2 h^{-1}), no mortality during the stationary phase but a much

reduced maximum population (K_{Mp} = 0.57 ± 0.01 10^6cells.mL^{-1}, K_{Mf} = 0.8 ± 0.05 10^6cells.mL^{-}

^{1}). In most cases, mixed cultures displayed an intermediate pattern between the two

corresponding isolated cultures (Fig. 2). However, mixed or isolated cultures with

Metschnikowia displayed different cell mortality rates during the stationary phase: in the case

of **ScvsMp** fermentations, only the *S. cerevisiae* population decreased significantly during the

stationary phase, while in **ScvsMf** fermentations, both subpopulations significantly decreased.

As a measure of fitness, we also followed the variations of *S. cerevisiae* frequency along the

fermentation. In all mixed cultures, *S. cerevisiae* was found dominant (frequency > 50%) in the

end, increasing significantly during fermentation from 10% initially to frequencies varying

between 50% (**ScvsMp**) and 96% (**ScvsMf**) (Table. 1)

Sugar and NAS consumption

We then looked at the final concentration of resources: sugars (fructose and glucose) and NAS

(sum of all assimilable nitrogen sources, i.e. ammonium and amino-acids) (Fig. 3, Table 1). In

Sc fermentations, less than 0.1% of the initial concentration of both sugars remained (Fig. 3a).

As seen in the paragraph concerning CO_2 production, non-*Saccharomyces* species in

monocultures did not complete fermentation in the 300h period and left respectively 45% of

sugars for **Ho**, 67% for **Hu**, 68% for **Mf** and 71% for **Mp**. Furthermore, all species except *H.*

opuntiae preferentially consumed glucose (see supplementary figure 1). Sugar consumption

was higher in mixed cultures than in single non-*Saccharomyces* species cultures (Table 1).

However, it was still lower than in **Sc** species cultures, also with a preference for glucose. This

indicates the major impact of *S. cerevisiae* on sugar consumption (consistent with the CO_2

production observed), compared to the other species studied.

The consumption of nitrogen assimilable sources (NAS, amino-acids and ammonium)

displayed the same pattern (Fig. 3b). NAS were almost entirely consumed both in *Sc* isolated

cultures and in all co-cultures, whereas in non-*Saccharomyces* isolated cultures the fraction of

NAS consumed varied between 84% and 94%. However, the preference for different nitrogen

sources varied with each species (Fig. 3c). Both *Hanseniaspora* species had similar behaviors,

consuming only half of the available ammonium, 90% of histidine and 89% or 79% of arginine

(Fig. 3c). *Metschnikowia* species presented a similar pattern. It was possible to classify these

non-*Saccharomyces* species preferences for the various NAS. The resulting ranking by order

of preference was glutamine, methionine, glutamate, valine, threonine, serine, tryptophan,

alanine, histidine, arginine, aspartate, glycine and, surprisingly last, ammonium.

Metabolite production

In parallel with must resources consumption monitoring, we also investigated the production

of metabolites from CCM (Central Carbon Metabolism): ethanol, glycerol, succinate, pyruvate,

acetate and alpha-ketoglutarate (Table 1). These measurements of metabolite production were

taken after 300 hours when sugars consumptions were quite different from one culture to

another depending on their dynamics. To allow figures comparison, we computed the

production yield (total production / sugar consumption) for each culture and, from these data,

we then estimated this yield relatively to that of *Sc* in isolated culture (Fig. 4).

In the case of ethanol, only *Mf* fermentations had a relative yield significantly inferior (-32%)

to 0 (0 being *Sc* yield). For glycerol, all isolated non-*Saccharomyces* fermentations had a

greater yield than *Sc* and mixed fermentations were intermediate between (but not significantly

different from) the corresponding isolated cultures. For acetate, only *Hanseniaspora* strains

displayed a higher yield (Fig. 4).

Finally, all mixed cultures seemed to have a lower succinate yield than both corresponding

isolated cultures (but not significantly after correction for multiple tests).

For each fermentation, the total production of metabolites resulted from the combination of

species yields, total sugar consumption and respective population dynamics during

fermentation. Therefore, differences observed in the total productions of mixed cultures were

the consequences of additive or subtractive effects observed for these 3 components.

Considering ethanol, its total production was directly linked to the consumption of resources

and all mixed cultures were equivalent to *Sc* fermentations (Table 1, supplementary figure S2).

The case of glycerol was more interesting. Indeed, even if the average sugar consumption was

lower in *ScvsHu* and *ScvsMp* mixed cultures than in isolated *Sc* culture, the total production of

glycerol was significantly higher than that of the corresponding isolated cultures

(Glycerol$_{SCvsMp}$ = 6.1±0.1 g.L^{-1}, Glycerol$_{Sc}$ = 5.3±0.4 g.L^{-1}, Glycerol$_{Mp}$ = 3.7±0.2 g.L^{-1}). This

resulted from the positive combination of the greater glycerol yield by *Hanseniaspora* and

Metschnikowia and their population dynamics. For all other metabolites, the total production

of mixed cultures was not significantly different from the corresponding isolated cultures

(Table 1).

Discussion
216

217　This study presents one of the first works focusing on the population dynamics and kinetics of

218　yeast-yeast interactions between two species during the alcoholic fermentative process. By

219　contrast, to our knowledge, most of the previous studies were concerned only with input-output

220　balances (García et al., 2016; Varela, 2016). Firstly, it was possible to group the non-

Saccharomyces strains performance according to their genus (see PCA in supplementary figures S3); *H. uvarum* and *H. opuntiae* both presented similar phenotypes. Their population dynamics were characterized by a similarly high growth rate (reaching a maximum population similar to *S. cerevisiae*), followed by a long mortality phase where viability dropped to 30% (Moreira et al., 2008; Tristezza et al., 2016). Their NAS consumption was also similar, with half of the available ammonium not being consumed. Even their yield and total production of CCM metabolites were very similar.

For almost all these characteristics, *Hanseniaspora* mixed cultures presented intermediate phenotypes compared to the corresponding isolated cultures. The only exception was the total production of glycerol that seemed to be superior in mixed cultures whereas their sugar consumption was inferior; this is characteristic of a transgressive interaction (often referred to as over-yielding), i.e. a situation in which the ecosystem performance is higher than that of the best-yielding species present (when cultivated alone). This glycerol overproduction in mixed cultures has already been observed in previous works (Ciani and Ferraro, 1996; Tristezza et al., 2016) but is not systematic depending on the strains and experimental conditions (Moreira et al., 2008). In the present study, the glycerol overproduction and sugar consumption observed in mixed cultures could be explained by the joint effect of population dynamics (*S. cerevisiae* slowly dominating the population), resource consumptions (the ***Ho*** and ***Hu*** fermentation leaving ⅔ of sugars) and the glycerol yields of *Hanseniaspora* strains that were two to three times higher that of *S. cerevisiae*.

Regarding *Hanseniaspora* species, ***Mp*** and ***Mf*** isolated cultures were characterized by a short growth phase reaching a small carrying capacity and followed by a long stationary phase with no mortality. Nevertheless, despite this very high viability, the resources measured (sugars and nitrogen) were not entirely consumed. The reason why the cells stopped growing remains

unclear. A possible explanation could be linked to unmonitored resources. For instance, the

third most important resource in the synthetic medium was lipids, available as phytosterols.

Concurrently, it has been shown that some *Metschnikowia* species were unable to import lipids

from the medium, depending instead on their own lipid synthesis (see Fig S1). For all yeast

species, lipid synthesis requires oxygen and is therefore impossible in anaerobic conditions.

From these data, one can hypothesize that *Metschnikowia* cells were able to produce their own

lipids in the aerobic conditions prevailing at the beginning of the fermentation. Then, with the

progress of fermentation and ensuing oxygen limitation, *Metschnikowia* cells became unable to

synthetize the necessary lipids and stopped multiplying.

Interestingly, even though *S. cerevisiae*, *M. pulcherrima* and *M. fructicola* mortality rates were

low in isolated cultures, the corresponding mixed cultures (***ScvsMp*** and ***ScvsMf***) presented 30%

mortality. Moreover, this mortality seemed to impact both species differently. In the mixed

ScvsMp culture, only *S. cerevisiae* cells eventually died, whereas in ***ScvsMf*** both species were

negatively affected. The survival of *M. pulcherrima* cells compared to *S. cerevisiae* cells could

be explained by the production of pulcherriminic acid by *M. pulcherrima* (MacDonald, 1965).

Indeed, pulcherriminic acid is known to deplete iron from the medium, which has a lethal effect

on *S. cerevisiae* cells (Oro et al., 2014; Sipiczki, 2006; Türkel and C, 2009). In ***ScvsMf*** cultures,

the mortality observed in both species suggests a more complex mechanism of interaction

(although it is not clear whether *M. fructicola* also produces pulcherriminic acid; Kurtzman and

Droby, 2001). To explain these results, we could hypothesize the conjunction of two different

mechanisms of interaction. It could be that *M. fructicola* synthesized a metabolite

(pulcherriminic acid?) impacting the viability of *S. cerevisiae* cells (through iron depletion?),

with *M. fructicola* cells dying thereafter for another reason such as sensitivity to ethanol.

Indeed, the production of ethanol was almost four times higher in mixed cultures than in

isolated *Metschnikowia* fermentations. Under such hypothesis, the reason why no loss of

viability was observed for *M. pulcherrima* in mixed culture with *S. cerevisiae* could probably

be related to a better tolerance of *M. pulcherrima* to ethanol stress compared to *M. fructicola*.

As in *Hanseniaspora*, we also observed an overproduction of glycerol in *Metschnikowia* mixed

cultures. This also could be explained by the same conjunction of yield, sugar consumption and

population dynamics. These overproductions of glycerol open the way for potentially more

interesting results. In fact, one can hypothesize that, by exploiting this specific yield and

population dynamics, for example by inoculating the S. *cerevisiae* strain at lower relative

frequencies (0.01, 0.001, 0.0001), more important transgressive interactions (such as lower

ethanol production) could be achieved in such mixed cultures.

For all *Metschnikowia* and *Hanseniaspora* mixed cultures, we only discussed transgressive

interactions when mixed cultures over-produced (or under-produced) a given metabolite.

Indeed, it was very difficult to identify interactions when the productions of mixed

fermentations were within the range of isolated cultures productions. As the relative frequency

of both strains in mixed cultures evolved during fermentation, it was difficult to link the final

mix to the contributions of each species. It was even more difficult to assess whether these

contributions combined additively or with interaction. New statistical developments will be

needed to answer this question (Barot et al., 2017; Finckh and Mundt, 1992).

If we look more generally at *Metschnikowia* and *Hanseniaspora* mixed cultures with *S.*

cerevisiae, we could not evidence any major antagonistic phenomena. For almost all assays,

mixed cultures performance stood always between that of the corresponding isolated cultures

(Fig 1, 2 & 3), with the exception of the total production of glycerol. Moreover, despite the

differences in yield and interactions between species, the rapid dominance of *S. cerevisiae*

(increasing from 10% to at least 50% during the fermentation) resulted in mixed cultures that

were overall not different from *S. cerevisiae* pure cultures. This result is in agreement with the

good adaptation of *S. cerevisiae* to winemaking conditions (García et al., 2016; Jolly et al.,

2014; Pinto et al., 2015; Varela, 2016) but could be different with a different set of strains. This

result is important in the context of ecological engineering. In fact, our results confirmed that

S. cerevisiae has a much better fitness than the non-*Saccharomyces* strains studied in this paper.

Therefore, if we want mixed culture behavior to deviate from that of *S. cerevisiae* monoculture,

it must be ensured that non-*Saccharomyces* cells dominate the culture as soon as possible. To

achieve this, two conceivable options are currently tested: either to reduce the proportion of *S.*

cerevisiae at t_0 or to successively inoculate (Tristezza et al., 2016) first the non-*Saccharomyces*

species , and only then the *S. cerevisiae* strain. These two options could be equivalent

depending on the type of interaction(s) that occurs. If strain behaviors in isolated or mixed

cultures are identical, then all interactions are mediated by the medium through the competition

for resources and the production of constitutive toxins such as ethanol (producing a toxin only

in mixed fermentation would be a behavior change) and could be qualified as "indirect". Indeed,

all the interactions observed in the present study could be explained by these types of

interactions.

Mathematical models could be designed from data on isolated cultures to predict the impact of

indirect interactions in mixed cultures. This would allow simulating numerous mixes of species

with various initial conditions and identify optimal strategies depending on one or several given

criteria. Using these approaches could limit the number of necessary tests, potentially saving a

lot of time and money and opening the way to a more methodical ecological engineering. The

development of such mathematical models will only be possible thanks to a deep tracking of

population dynamics to understand underlying mechanisms of growth and mortality.

Obviously, it is also critical to validate this approach by i) first extending the number of species

co-cultured with *S. cerevisiae*, ii) investigating intra-specific variability and strain-strain

interactions between species, iii) investigating the impact of the environment of culture

(temperature, grape variety, etc.).

Mat & Met

Strains

In this work, we used one strain of 5 different species (one strain per species): *Saccharomyces*

cerevisiae (***Sc***)*, Metschnikowia pulcherrima* (***Mp***)*, M. fructicola* (***Mf***)*, Hanseniaspora uvarum*

(***Hu***) and *H. opuntiae* (***Ho***). The *S. cerevisiae* strain is a haploid strain from EC1118 labelled

with GFP (59A-GFP, Marsit and Dequin, 2015). The *Hanseniaspora uvarum* (CLIB 3221) *H.*

opuntiae (CLIB 3093) and *Metschnikowia pulcherrima* (CLIB 3235) strains originated from

the yeast CIRM (https://www6.inra.fr/cirm_eng/Yeasts/Strain-catalogue) and were isolated

from grape musts. The *Metschnikowia fructicola strain was* from the Lallemand collection.

For each species, 3 replicates of isolated cultures were performed (except for *S. cerevisiae* that

had a total of 8 replicates in different blocks). In addition, for each non-*Saccharomyces*, 3

replicates of a mixed culture with the **Sc** strain were performed. In all mixed fermentations, the

starting proportion of *S. cerevisiae* cells was set at 10%. In this text, fermentations were referred

to by the species that performed them, i.e. isolated cultures were referred to as: ***Sc***, ***Mp***, ***Mf***, ***Hu***

and ***Ho*** and mixed cultures as ***ScvsMp***, ***ScvsMf***, ***ScvsHu*** and ***ScvsHo***.

Medium

Initial cultures (12 h, in 50 ml YPD medium, 28 °C) were used to inoculate fermentation media

at a total density of 10^6 cells/mL; therefore, for mixed culture the *S. cerevisiae* cells density was

0.1×10^6 /mL and the non-*Saccharomyces* cells density was 0.9×10^6 /mL. Fermentations were

carried out in a synthetic medium (SM) mimicking standard grape juice (Bely et al., 1990). The

SM used in this study contained 200 g/L of sugar (100 g glucose and 100 g fructose per liter)

and 200 mg/L of assimilable nitrogen (as a mix of ammonium chloride and amino acids). The

concentrations of weak acids, salts and vitamins were identical to those described by Seguinot

et al. (Seguinot et al., 2017). The pH of the medium was adjusted to 3.3 with 10M NaOH. The

SM medium was first saturated with bubbling air during 40 minutes, then it was supplemented

with 5 mg/L phytosterols (85451, Sigma Aldrich) to fulfill the lipid requirements of yeast cells

during anaerobic growth.

Measurements

Fermentation took place in 1.1-liter fermentors equipped with fermentation locks to maintain

anaerobiosis, at 20 °C, with continuous magnetic stirring (500 rpm) during approximately 300h.

CO_2 release was followed by automatic measurements of fermentor weight loss every 20 min.

The amount of CO_2 released allowed us to monitor the progress of the fermentation and evaluate

the maximum of released CO_2 (***CO2max***) as well as the maximum rate of CO_2 released (***Vmax***).

Samples were harvested after 6h, 12h and 24h, then every 12h during the first week and every

24h during the second week of fermentation. For each sample, the population density, viability

and proportion of *S. cerevisiae* cells were determined using a BD Accuri™ C6 Plus flow

cytometer as described in Delobel et al., (2012. From these population densities (without taking

into account viability), we fitted a growth population model (with the growthcurver package

in R, Sprouffske and Wagner, 2016), and determined the carrying capacity (***K***) and maximum

growth rate (***mu***) for each fermentation.

The final concentrations of carbon metabolites in the medium (acetate, succinate, glycerol,

alpha-ketoglutarate, pyruvate, ethanol, glucose and fructose) were determined with high-

pressure liquid chromatography (Camarasa et al., 2011). From these metabolite concentrations,

we first calculated the consumed sugar concentration as the difference between the final and

the initial concentration of either glucose or fructose. Then we calculated the yield of metabolite

production by dividing the final concentration by the corresponding consumed sugar

concentration. Finally, we compared these yields to the yield of *S. cerevisiae* isolated cultures

considered as reference.

Finally, the ammonium concentration after 100h of fermentation was determined enzymatically

with R-Biopharm (Darmstadt, Germany) and the free amino acid content of the must was

estimated through cation exchange chromatography with post-column ninhydrin derivatization

(Crépin et al., 2012).

Statistical Analysis

The experimental work was performed in 5 different blocks. Each block was composed of three

replicates of non-*Saccharomyces* fermentations (for example ***Hu***), three replicates of the

corresponding mixed fermentations with *S. cerevisiae* (for example ***ScvsHu***) and one or two

fermentations of isolated *S. cerevisiae* cells (***Sc***). The block effect was evaluated on the

parameters of the ***Sc*** fermentation. For most studied parameters, the block effect was not

significant. For those parameters where a block effect was observed (***mu*** and ***K***), a statistical

correction for block effect did not modify our results. Therefore, for simplification purposes,

we compared all fermentations without any correction for the block effect parameter. For each

measured parameter, an ANOVA was performed to evaluate the type of fermentation (***Sc***, ***Mp***,

Mf, ***Hu***, ***Ho***, ***ScvsTd***, ***ScvsMp***, ***ScvsMf***, ***ScvsHu*** and ***ScvsHo***) effect and then a Tukey t-test was

performed to determine statistical groups and two-by-two statistical differences. All statistical

analyses were performed using R ("R Core Team (2016). R: A language and environment for

statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL

https://www.R-project.org/.," n.d.) and Rstudio ("RStudio (2012). RStudio: Integrated

development environment for R (Version 0.96.122) [Computer software]. Boston, MA.

Retrieved May 20, 2012. Available from http://www.rstudio.org/," n.d.). All data, analysis and

figures scripts can be found in this github adress: https://github.com/tnidelet/Harle-et-al-2018.

Acknowledgments.

We thank the CIRM, the Lallemand company, Jean-Luc Legras, Virginie Galeote and Jean-

Nicolas Jasmin for providing the strains used in this study. We thank also Christian Picou, Marc

Perez, Faiza Macna for technical assistance and Delphine Sicard for advice.

References

Barot, S., Allard, V., Cantarel, A., Enjalbert, J., Gauffreteau, A., Goldringer, I., Lata, J.-C., Roux, X., Niboyet, A., Porcher, E., 2017. Designing mixtures of varieties for multifunctional agriculture with the help of ecology. A review. Agron Sustain Dev 37, 13. https://doi.org/10.1007/s13593-017-0418-x

Belda, I., Navascués, E., Marquina, D., Santos, A., Calderon, F., Benito, S., 2015. Dynamic analysis of physiological properties of Torulaspora delbrueckii in wine fermentations and its incidence on wine quality. Appl Microbiol Biot 99, 1911–1922. https://doi.org/10.1007/s00253-014-6197-2

Bely, M., Sablayrolles, J.M., Barre, P., 1990. Description of Alcoholic Fermentation Kinetics: Its Variability and Significance. Am. J. Enol. Vitic. 41, 319–324.

Camarasa, C., Sanchez, I., Brial, P., Bigey, F., Dequin, S., 2011. Phenotypic landscape of Saccharomyces cerevisiae during wine fermentation: evidence for origin-dependent metabolic traits. PLoS ONE 6, e25147. https://doi.org/10.1371/journal.pone.0025147

Ciani, M., Ferraro, L., 1996. Enhanced Glycerol Content in Wines Made with Immobilized Candida stellata Cells. Appl. Environ. Microbiol. 62, 128–132.

Contreras, A., Hidalgo, C., Henschke, P.A., Chambers, P.J., Curtin, C., Varela, C., 2014. Evaluation of Non-Saccharomyces Yeasts for the Reduction of Alcohol Content in Wine. Appl. Environ. Microbiol. 80, 1670–1678. https://doi.org/10.1128/AEM.03780-13

Crépin, L., Nidelet, T., Sanchez, I., Dequin, S., Camarasa, C., 2012. Sequential Use of Nitrogen Compounds by Saccharomyces cerevisiae during Wine Fermentation: a Model Based on Kinetic and Regulation Characteristics of Nitrogen Permeases 78, 8102–8111. https://doi.org/10.1128/AEM.02294-12

Delobel, P., Pradal, M., Blondin, B., Tesniere, C., n.d. A 'fragile cell' sub-population revealed during cytometric assessment of Saccharomyces cerevisiae viability in lipid-limited alcoholic

fermentation. Letters in Applied Microbiology 55, 338–344. https://doi.org/10.1111/j.1472-765X.2012.03301.x

Elena, G.-R., Pascual, O., Kontoudakis, N., Esteruelas, M., Braulio, E.-Z., Mas, A., Canals, J., Zamora, F., 2015. Oenological consequences of sequential inoculation with non-Saccharomyces yeasts (Torulaspora delbrueckii or Metschnikowia pulcherrima) and Saccharomyces cerevisiae in base wine for sparkling wine production. Eur Food Res Technol 240, 999–1012. https://doi.org/10.1007/s00217-014-2404-8

Finckh, M.R. (International R.R.I., Mundt, C.C., 1992. Stripe rust, yield, and plant competition in wheat cultivar mixtures. Phytopathology (USA).

García, M., Braulio, E.-Z., Arroyo, T., 2016. Non-Saccharomyces Yeasts: Biotechnological Role for Wine Production. https://doi.org/10.5772/64957

García-Ríos, E., Gutiérrez, A., Salvadó, Z., Arroyo-López, F.N., Guillamon, J.M., 2014. The Fitness Advantage of Commercial Wine Yeasts in Relation to the Nitrogen Concentration, Temperature, and Ethanol Content under Microvinification Conditions. Appl. Environ. Microbiol. 80, 704–713. https://doi.org/10.1128/AEM.03405-13

Hu, K., Qin, Y., Tao, Y.-S., Zhu, X.-L., Peng, C.-T., Ullah, N., n.d. Potential of Glycosidase from Non-Saccharomyces Isolates for Enhancement of Wine Aroma. Journal of Food Science 81, M935–M943. https://doi.org/10.1111/1750-3841.13253

Ivey, M., Massel, M., Phister, T.G., 2013. Microbial Interactions in Food Fermentations. Annual Review of Food Science and Technology 4, 141–162. https://doi.org/10.1146/annurev-food-022811-101219

Jolly, N.P., Varela, C., Pretorius, I.S., 2014. Not your ordinary yeast: non-Saccharomyces yeasts in wine production uncovered. Fems Yeast Res 14, 215–237. https://doi.org/10.1111/1567-1364.12111

Kurtzman, C.P., Droby, S., 2001. Metschnikowia fructicola, a New Ascosporic Yeast with Potential for Biocontrol of Postharvest Fruit Rots. Systematic and Applied Microbiology 24, 395–399. https://doi.org/10.1078/0723-2020-00045

MacDonald, 1965. Biosynthesis of pulcherriminic acid. Biochem. J. 96, 533–8.

Marsit, S., Dequin, S., 2015. Diversity and adaptive evolution of Saccharomyces wine yeast: a review. FEMS Yeast Research 15, fov067. https://doi.org/10.1093/femsyr/fov067

Moreira, N., Mendes, F., de Pinho, P., journal of … H.-T., 2008. Heavy sulphur compounds, higher alcohols and esters production profile of Hanseniaspora uvarum and Hanseniaspora guilliermondii grown as pure and …. International journal of ….

Oro, L., Ciani, M., Comitini, F., 2014. Antimicrobial activity of Metschnikowia pulcherrima on wine yeasts. J Appl Microbiol 116, 1209–1217. https://doi.org/10.1111/jam.12446

P, H.-O., Cersosimo, M., Loscos, N., Cacho, J., E, G.-M., Ferreira, V., 2008. The development of varietal aroma from non-floral grapes by yeasts of different genera. Food Chem 107, 1064–1077. https://doi.org/10.1016/j.foodchem.2007.09.032

Pinto, C., Pinho, D., Cardoso, R., Custódio, V., Fernandes, J., Sousa, S., Pinheiro, M., Egas, C., Gomes, A.C., 2015. Wine fermentation microbiome: a landscape from different Portuguese wine appellations 6, 905.

R Core Team (2016). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL   https://www.R-project.org/., n.d.

RStudio (2012). RStudio: Integrated development environment for R (Version 0.96.122) [Computer software]. Boston, MA. Retrieved May 20, 2012. Available from http://www.rstudio.org/, n.d.

Sablayrolles, J., 2008. Fermented beverages: the example of winemaking. Advances in Fermentation Technology. New Delhi: Asiatech Publishers 322–47.

Sadineni, V., Kondapalli, N., Obulam, V., 2012. Effect of co-fermentation with Saccharomyces cerevisiae and Torulaspora delbrueckii or Metschnikowia pulcherrima on the

aroma and sensory properties of mango wine. Ann Microbiol 62, 1353–1360.

https://doi.org/10.1007/s13213-011-0383-6

Seguinot, P., SANCHEZ, I., Bloem, A., Ortiz-Julien, A., Camarasa, C., 2017. Impact of

nitrogen sources on the fermentative kinetic of non - Saccharomyces yeasts.

Sipiczki, M., 2006. Metschnikowia Strains Isolated from Botrytized Grapes Antagonize

Fungal and Bacterial Growth by Iron Depletion. Appl Environ Microb 72, 6716–6724.

https://doi.org/10.1128/aem.01275-06

Song, H.-S., Cannon, W.R., Beliaev, A.S., Konopka, A., 2014. Mathematical Modeling of

Microbial Community Dynamics: A Methodological Review. Processes 2, 711–752.

https://doi.org/10.3390/pr2040711

Sprouffske, K., Wagner, A., 2016. Growthcurver: an R package for obtaining interpretable

metrics from microbial growth curves. BMC Bioinformatics 17, 172.

https://doi.org/10.1186/s12859-016-1016-7

Tristezza, M., Tufariello, M., Capozzi, V., Spano, G., Mita, G., Grieco, F., 2016. The

Oenological Potential of Hanseniaspora uvarum in Simultaneous and Sequential Co-

fermentation with Saccharomyces cerevisiae for Industrial Wine Production. Front.

Microbiol. 7. https://doi.org/10.3389/fmicb.2016.00670

Türkel, S., C, E.-B., 2009. Isolation and characterization of new Metschnikowia pulcherrima

strains as producers of the antimicrobial pigment pulcherrimin. Zeitschrift für Naturforschung

C.

Varela, C., 2016. The impact of non-Saccharomyces yeasts in the production of alcoholic

beverages. Appl Microbiol Biot 100, 9861–9874. https://doi.org/10.1007/s00253-016-7941-6

Tables

Table 1

Parameter values (mean ± standard deviation) for each type of fermentation.

	μ	K	prop	Vmax	MaxCO2	Sugar	NAS	Ethanol	Glycerol	Succinate	Acetate	Pyruvate	Alpha
Sc	0.15±0.02[bc]	1.6±0.2[bcd]	NA	0.99±0.03[a]	88.2±2.2[a]	0.2±0.2[d]	10.4±8.6[c]	95.2±8.4[a]	5.3±0.4[c]	1.05±0.28[a]	0.63±0.06[d]	0.12±0.08[bc]	0.08±0.04[ab]
Ho	0.19±0.03[bc]	1.6±0.1[bc]	NA	0.42±0.02[d]	46.4±0.6[d]	92.3±1.4[b]	122.2±3.5[b]	50.1±0.2[b]	5.2±0.0[c]	0.35±0.00[b]	0.99±0.02[b]	0.13±0.05[bc]	0.09±0.1[ab]
ScvsHo	**0.11±0.00[bc]**	**2.3±0.1[a]**	0.83±0.02[b]	0.57±0.01[c]	82.7±0.6[bc]	23.3±1.3[c]	**6.4±2.4[c]**	87.2±0.8[a]	**6.0±0.0[abc]**	0.42±0.04[b]	0.84±0.01[c]	**0.08±0.00[c]**	**0.03±0.0[b]**
Hu	0.62±0.18[a]	1.4±0.1[cde]	NA	0.34±0.04[d]	30.7±0.5[e]	134.3±11.2[a]	173.3±24.8[b]	34.9±1.1[c]	5.5±0.2[bc]	0.66±0.03[b]	1.15±0.03[a]	0.17±0.04[abc]	0.14±0.09[ab]
ScvsHu	0.24±0.01[b]	**1.7±0.1[b]**	0.76±0.01[c]	0.53±0.04[c]	82.1±1.5[c]	16.5±6.5[c]	**4.6±4.2[c]**	89.5±3.8[a]	**6.6±0.3[a]**	**0.60±0.06[b]**	0.86±0.07[bc]	0.16±0.04[abc]	0.09±0.06[ab]
Mf	0.17±0.02[bc]	0.8±0.05[f]	NA	0.17±0.04[e]	20.6±1.2[f]	135.8±9.7[a]	256.6±2.9[a]	20.6±0.3[c]	2.8±0.1[e]	0.29±0.01[b]	0.13±0.01[f]	0.10±0.01[bc]	0.10±0.00[ab]
ScvsMf	**0.09±0.01[bc]**	1.3±0.03[de]	0.96±0.02[a]	0.76±0.03[b]	86.8±0.7[ab]	9.9±2.1[cd]	**1.8±0.1[c]**	92.1±3.2[a]	**6.0±0.1[abc]**	0.58±0.02[b]	0.46±0.02[e]	0.12±0.01[bc]	0.08±0.01[ab]
Mp	0.18±0.03[bc]	0.6±0.01[f]	NA	0.16±0.02[f]	21.8±0.4[f]	142±3.7[a]	305.5±5.9[a]	23.4±1.8[c]	3.7±0.2[d]	0.26±0.03[b]	0.07±0.06[f]	0.29±0.04[a]	0.13±0.01[ab]
ScvsMp	**0.09±0.00[c]**	1.3±0.01[e]	0.50±0.02[d]	0.68±0.04[b]	83.3±2.2[bc]	20.8±6.0[c]	**2.1±0.2[c]**	86.5±3.4[a]	**6.1±0.1[ab]**	0.52±0.04[b]	0.46±0.02[e]	0.24±0.01[ab]	**0.19±0.01[a]**

Mu : h[-1]; K : 10[8] cells.mL[-1], prop : no unity, Vmax : g.L[-1].h[-1], MaxCO2 : g.L[-1], Final Sugar : g.L[-1], Final NAS : mg.L[-1], Ethanol : g.L[-1], Glycerol : g.L[-1], Succinate : g.L[-1], Acetate : g.L[-1], Pyruvate : g.L[-1], Alpha : g.L[-1]

Bold: values of mixed culture that are not intermediate between the two values of the corresponding isolated cultures, a sign of transgressive interactions.

Superscript: Statistical groups obtained from a Tukey test

Figures
523

FIG 1
525

Representation of a) the maximum rate of CO_2 production (***Vmax***) and b) the total CO_2
526

produced in function of the species driving each fermentation (average ± standard deviation).
527

The small letters indicate the statistical groups from a Tukey analysis.
528

FIG2

Global monitoring of the kinetics of the total living population (left), and sub-population in the mixed cultures (right) across fermentation. Each point represents a sample (average ± standard error). full lines are for total population and dashed lines for the two sub populations in mixed cultures. At the end of dashed lines, the final proportion of both sub-populations in mixed cultures is indicated. The light colors represent pure cultures of 'non-*Saccharomyces*'

strains and dark ones to the corresponding culture in competition with *S. cerevisiae*. The isolated cultures of *S. cerevisiae* are represented in black. ***Sc, Mp, Mf, Ho, Hu, ScvsMp, ScvsMf, ScvsHo*** and ***ScvsHu*** correspond respectively to cultures with: isolated *S. cerevisiae*, isolated *M. pulcherrima*, isolated *M. fructicola*, isolated *H. opuntiae*, isolated *H.uvarum*, *M. pulcherrima* mixed with *S. cerevisiae*, *M. fructicola* mixed with *S. cerevisiae*, *H. opuntiae* mixed with *S. cerevisiae* and *H. uvarum* mixed with *S. cerevisiae*.

FIG3

Consumption of sugars and nitrogen assimilable sources (NAS) for each type of fermentation. A) Final concentration of sugar (average ± standard deviation). B) Final concentration of NAS (average ± standard deviation). C) Percentage of consumption of each NAS in each type of fermentation represented as a color gradient from green (<75 %) to red (> 75%). ***Sc, Mp, Mf, Ho, Hu, ScvsMp, ScvsMf, ScvsHo*** and ***ScvsHu*** correspond respectively to cultures with: isolated *S. cerevisiae*, isolated *M. pulcherrima*, isolated *M. fructicola*, isolated *H. opuntiae*, isolated *H.uvarum*, *M. pulcherrima* mixed with *S. cerevisiae*, *M. fructicola* mixed with *S. cerevisiae*, *H. opuntiae* mixed with *S. cerevisiae* and *H. uvarum* mixed with *S. cerevisiae*.

FIG4

Yield (average ± standard deviation) of carbon metabolite production relative to the yield of production of isolated *S. cerevisiae* cultures for acetate, alpha-ketoglutarate, ethanol, glycerol, pyruvate and succinate and each type of fermentation. ***Sc, Mp, Mf, Ho, Hu, Td, ScvsMp,***

561 ***ScvsMf, ScvsHo*** and ***ScvsHu, ScvsTd*** correspond respectively to cultures with: isolated *S.*

562 *cerevisiae*, isolated *M. pulcherrima*, isolated *M. fructicola*, isolated *H. opuntiae*, isolated

563 *H.uvarum*, *M. pulcherrima* mixed with *S. cerevisiae*, *M. fructicola* mixed with *S. cerevisiae*, *H.*

564 *opuntiae* mixed with *S. cerevisiae* and *H. uvarum* mixed with *S. cerevisiae*.