Investigations of the mechanisms of interactions between four non-conventional species with *Saccharomyces cerevisiae* in oenological conditions

Oliver Harlé[1], Judith Legrand[3], Catherine Tesnière[2], Martine Pradal[2], Jean-Roch Mouret[2], Thibault Nidelet[2*]

Affiliations :

1 : STLO, INRA, Agrocampus Ouest, Rennes, France

2 : SPO, Univ. Montpellier, INRA, Montpellier SupAgro, Montpellier, France

3 : GQE-Le Moulon, INRA, Univ. Paris-Sud, CNRS, AgroParisTech, Université Paris-Saclay, Gif-sur-Yvette, France

[*] **Corresponding Author**: thibault.nidelet@inra.fr

keywords: winemaking fermentation, population dynamics, non-*Saccharomyces* species, interactions, ecosystems

Abstract

Fermentation by microorganisms is a key step in the production of traditional food

products such as bread, cheese, beer and wine. In these fermentative ecosystems,

microorganisms interact in various ways, namely competition, predation, commensalism and

mutualism. Traditional wine fermentation is a complex microbial process performed by

Saccharomyces and non-*Saccharomyces* (NS) yeast species. To better understand the different

interactions occurring within wine fermentation, isolated yeast cultures were compared with

mixed co-cultures of one reference strain of *S. cerevisiae* with one strain of four NS yeast

species (*Metschnikowia pulcherrima, M. fructicola, Hanseniaspora opuntiae and H. uvarum*).

In each case, we studied population dynamics, resource consumed and metabolites produced

from central carbon metabolism. This phenotyping of competition kinetics allowed us to

confirm the main mechanisms of interaction between strains of four NS species. *S. cerevisiae*

competed with *H. uvarum* and *H. opuntiae* for resources although both *Hanseniaspora* species

were characterized by a strong mortality either in isolated or mixed fermentations. *M.*

pulcherrima and *M. fructicola* displayed a negative interaction with the *S. cerevisiae* strain

tested, with a decrease in viability in co-culture, probably due to iron depletion via the

production of pulcherriminic acid. Overall, this work highlights the importance of measuring

specific cell populations in mixed cultures and their metabolite kinetics to understand yeast-

yeast interactions. These results are a first step towards ecological engineering and the rational

design of optimal multi-species starter consortia using modeling tools. In particular the

originality of this paper is for the first times to highlight the joint-effect of different species

population dynamics on glycerol production and also to discuss on the putative role of lipid

uptake on the limitation of some non-conventional species growth although interaction

processes.

1. Introduction

In natural or anthropized environments, microbial species are part of an ecosystem and interact positively or negatively, forming a complex network. Until recently, process optimization in agriculture or food processing was mostly based on the selection of single strains. However, this paradigm is now being challenged and the scientific community is increasingly seeking to exploit and optimize consortia of several strains and/or species. Indeed, many studies have shown that more diverse anthropized environments have many advantages in terms of resilience, disease resistance or yield . Efforts are now being made to design optimal consortia of various species and strains whose interactions will be exploited to maximize given criteria such as fermentation quality, aromatic complexity or other organoleptic characteristics.

Wine fermentation is both an economically and societally important food ecosystem, where the addition of fermentation 'starters' composed of selected yeasts at the beginning of the fermentation process is common. In fact, around 80% of oenological fermentations worldwide are conducted with starters (1,2). Most often, these "starters" are only composed of a single *Saccharomyces cerevisiae* (S. c.) strain selected for its ability to complete fermentation. Indeed, numerous experiments have shown that *S. cerevisiae*, with an initially low population, most often becomes the predominant species at the end of the fermentation, demonstrating its superior fermentative abilities (3–6). However, in recent years, multi-species starters have emerged, aiming at increasing the aromatic complexity of wines. They most often combine one strain of *S. cerevisiae* allowing to complete fermentation and another species, often from a different genus, contributing to a greater variety of flavors (3,4,6,7).

Indeed, there are numerous experiments and even industrial products making use of such mixed starters to improve wine's organoleptic qualities (4,8,9). The non-*Saccharomyces* (NS) strains used in these experiments are very diverse, with more than 23 different species including

Torulaspora delbrueckii, *Metschnikowia pulcherrima*, *Metschnikowia fructicola*, *Hanseniaspora opuntiae* and *Hanseniaspora uvarum*. Species in the *Metschnikowia* genus ferment poorly in oenological conditions but can have interesting attributes: in conjunction with *S. cerevisiae*, a strain of *M. pulcherrima* could reduce ethanol concentrations (6,10), increase 'citrus/grape fruit' and 'pear' attributes (11), as well as allow the persistence of 'smocky' and 'flowery' characteristics (12). *M. pulcherrima* also has an amensalism effect on *S. cerevisiae* through iron depletion via the production of pulcherriminic acid (13). *M. fructicola* has been less studied and never in conjunction with *S. cerevisiae* although it presents the interesting ability to inhibit *Botrytis* growth (14). Last, the *Hanseniaspora* genus, studied in sequential or simultaneous fermentation with *S. cerevisiae,* has been shown to increase volatile compound production during winemaking (6). It notably increased the 'tropical fruit', 'berry', 'floral' and 'nut aroma' characters (15), that were linked to higher concentrations of acetate esters, esters of MCFAs, isoamyl alcohol, 2-phenylethanol and α-terpineol (16).

Despite these various studies, the composition and protocol of inoculation of these multi-strains starters are still very empirical and only based on the input/output balance, without considering the dynamics of the microbial populations or their interactions. This lack of knowledge about yeast-yeast interactions prevents implementing a rational design of multi-strain starters (17). To address this problem, we decided to focus our study on population dynamics and metabolites produced during oenological fermentations performed in isolated or mixed yeast cultures. Since our goal was not to obtain optimal mixes but to understand the mechanism of microbial interaction, we chose to compare the population dynamics and yields between isolated cultures of strains from five species (one *S. cerevisiae* and four NS) and four corresponding mixed cultures always including the *S. cerevisiae* strain as reference. We were thus able to identify key microbial interaction mechanisms that are further discussed.

2. Results

In this work, we compared in winemaking conditions the performance of single cell cultures of five different strains from five yeast species (*Saccharomyces cerevisiae*, *Metschnikowia pulcherrima*, *Metschnikowia fructicola*, *Hanseniaspora opuntiae* and *Hanseniaspora uvarum*) and mixed co-cultures combining each of the four NS species with one GFP-labelled *S. cerevisiae* strain representing 10% of the initial inoculate. We choose to stop the monitoring of fermentation at a given time, even if the sugar supply was not completely exhausted. Thus, for all fermentation with the *S. cerevisiae* reference strain, sugars were exhausted after around 200-220 h while in fermentations with single NS strains, the sugar supply was still not exhausted after 400h. Here, we focused on the first 300 hours of fermentation.

By comparing the output of single strain and mixed strain cultures, we evaluated the intensity of yeast-yeast interactions and/or their consequences on ecosystem service production.

2.1. CO$_2$ kinetics

We first investigated the influence of species and co-culture on the dynamics of CO$_2$ production (proportional to sugar consumption), which is a good indicator of the fermentation progress. Indeed, CO$_2$ production is easy to monitor (based on weight measurement) and is directly proportional to ethanol synthesis and sugar consumption. The values of the maximum rate of CO$_2$ production (***Vmax***, Fig 1a) and of the maximum CO$_2$ produced were estimated (***CO$_2$max***, Fig 1b). ***Vmax*** was highly dependent on the species (p.value < 0.001): *S. cerevisiae* cultures (***Sc***) displayed the highest value (***Vmax$_{Sc}$*** = 0.99 ± 0.02 g.L^{-1}.h^{-1}), followed by both *Hanseniaspora* species (***Vmax$_{Hu}$*** = 0.33 ± 0.04 g.L^{-1}.h^{-1}, and ***Vmax$_{Ho}$*** = 0.42 ± 0.02 g.L^{-1}.h^{-1}) and finally both *Metschnikowia* species (***Vmax$_{Mp}$*** = 0.165 ± 0.02 g.L^{-1}.h^{-1} and ***Vmax$_{Mf}$*** = 0.17 ± 0.04 g.L^{-1}.h^{-1}). The four mixed cultures had intermediate ***Vmax*** values between those of ***Sc*** and

the highest **Vmax** of all *non-sacc* cultures (Fig 1a). Mixed cultures containing *Metschnikowia*

species had significantly higher **Vmax** values than those containing *Hanseniaspora* species (Fig

1a). Although we did not monitor all cultures until the exhaustion of glucose and fructose, it

was however possible to estimate the capacity of a given species to complete fermentation by

estimating the amount of CO_2 produced during the first 300 hours. **Sc** fermentations finished

after around 220 hours with a $CO2max_{Sc} = 88.2 \pm 2.2$ g.L^{-1}. We therefore can make the

hypothesis that all cultures that produced more than 80g CO_2.L^{-1} (90% of **Sc** maximum) within

300 hours will be able to complete fermentation. Under this assumption, all mixed cultures, but

no NS single strain cultures, would eventually complete fermentation. Among the latter

cultures, both *Hanseniaspora* species had the highest **CO_2max** (**Hu** 30 ± 0.4 g.L^{-1}, and **Ho** 46

± 0.6 g.L^{-1}) followed by *Metschnikowia* species (**Mp** 22 ± 0.4 g.L^{-1} and **Mf** 20 ± 1 g.L^{-1}).

FIG 1

Maximum rate of CO_2 production, *Vmax* (A) and total CO_2 produced (B) in function of

the single or mixed strain species driving each fermentation. Values corresponds to average

$\pm$ standard deviation. The small letters indicate the statistical groups from a Tukey analysis.

2.2. Population Kinetics

We also looked at population dynamics in each culture (Fig 2) and determined the maximum growth rate of the population (μ), the maximum population size, also termed carrying capacity (K) and the relative abundance (by cytometry) of each species after 300 hours of mixed culture, corresponding in our case to the end of the monitoring period (S1 Table). Fermentations with *S. cerevisiae* alone went through an exponential growth rate ($\mu_{Sc} = 0.15 \pm 0.02$ h^{-1}) and reached a maximum population of around $1.5*10^{8}$cells.mL^{-1} ($K_{Sc} = 1.55 \pm 0.15$ 10^{8}cells.mL^{-1}) that remained constant until the end of the fermentation. Fermentations with either *Hanseniaspora* species alone had a growth dynamic like *Sc* at the beginning of the fermentation but a higher growth rate ($\mu_{Ho} = 0.19 \pm 0.03$ h^{-1}, $\mu_{Hu} = 0.62 \pm 0.18$ h^{-1}). On the opposite, their stationary phase was quite different from that of *Sc* and characterized by a higher cell mortality with a population drop of about 70% by the end of the process. Fermentations performed by *Metschnikowia* species in single strain culture had growth dynamics mostly similar to *Sc* fermentations: a similar growth rate ($\mu_{mp} = 0.18 \pm 0.03$ h^{-1}, $\mu_{Mf} = 0.17 \pm 0.2$ h^{-1}), no mortality during the stationary phase but a much reduced maximum population ($K_{Mp} = 0.57 \pm 0.01$ 10^{6}cells.mL^{-1}, $K_{Mf} = 0.8 \pm 0.05$ 10^{6}cells.mL^{-1}). In most cases, mixed cultures displayed an intermediate pattern between the two corresponding single strain cultures (Fig 2). However, mixed or single strain cultures with *Metschnikowia* displayed different cell mortality rates during the stationary phase: in the case of *ScvsMp* fermentations, only the *S. cerevisiae* population decreased significantly during the stationary phase, while in *ScvsMf* fermentations, both subpopulations significantly decreased. As a measure of fitness, we also followed the variations of *S. cerevisiae* frequency along the fermentation. In all mixed cultures, *S. cerevisiae* was found dominant (frequency > 50%) in the end, increasing significantly during fermentation from 10% initially to frequencies varying between 50% (*ScvsMp*) and 96% (*ScvsMf*) (Fig 2).

FIG 2

Global monitoring of the kinetics of the total living population (left), and sub-population in the mixed cultures (right) across fermentation. Each population was detected by flow cytometry as indicated in the Material and Methods section. Each point represents a sample (average ± standard error). Full lines are for total population and dashed lines for the two sub populations in mixed cultures. At the end of dashed lines, the final proportion of both sub-populations in mixed cultures is indicated. The light colors represent single straind cultures of 'non-*Saccharomyces*' strains and dark ones to the corresponding culture in competition with *S. cerevisiae*. The single strand cultures of *S. cerevisiae* are represented in black.

2.3. Sugar and nitrogen assimilable source consumption

We then looked at the final concentration of resources: sugars (fructose and glucose) and nitrogen assimilable source (NAS) i.e. ammonium and amino-acids (Fig 3). In *Sc* fermentations, less than 0.1% of the initial concentration of both sugars remained (Fig 3a). As seen in the paragraph concerning CO_2 production, NS species in single strain cultures did not complete fermentation in the 300h period and left respectively 45% of sugars for *Ho*, 67% for *Hu*, 68% for *Mf* and 71% for *Mp*. Furthermore, all species except *H. opuntiae* preferentially

consumed glucose (S1 Fig). Sugar consumption was higher in mixed cultures than in single NS

species cultures (Fig 3a). However, it was still lower than in *Sc* species cultures, also with a

preference for glucose. This indicates the major impact of *S. cerevisiae* on sugar consumption

(consistent with the CO_2 production observed), compared to the other NS species studied.

The consumption of NAS displayed the same pattern (Fig 3b). NAS were almost entirely

consumed both in *Sc* single strain cultures and in all co-cultures, whereas in NS single strain

cultures the fraction of NAS consumed varied between 84% and 94%. However, the preference

for different nitrogen sources varied with each NS species (Fig 3c). Both *Hanseniaspora*

species had similar behaviors, consuming only half of the available ammonium, 90% of

histidine and 89% or 79% of arginine (Fig 3c). *Metschnikowia* species presented a similar

pattern. It was possible to classify these NS species preferences for the various NAS. The

resulting ranking by order of preference was glutamine, methionine, glutamate, valine,

threonine, serine, tryptophan, alanine, histidine, arginine, aspartate, glycine and, surprisingly

last, ammonium.

FIG3

Consumption of sugars and nitrogen assimilable sources (NAS) for each type of

fermentation. A) Final concentration of sugar (average ± standard deviation). B) Final

concentration of NAS (average ± standard deviation). C) Percentage of consumption of each

NAS in each type of fermentation represented as a color gradient from green (<75 %) to red (>

75%).

2.4. Metabolite production

In parallel with must resources consumption monitoring, we also investigated the production of metabolites from Central Carbon Metabolism (CCM): ethanol, glycerol, succinate, pyruvate, acetate and alpha-ketoglutarate (Fig 4). These measurements of metabolite production were taken after 300 hours when sugars consumptions were quite different from one culture to another depending on their dynamics. To allow figures comparison, we computed the

production yield (total production / sugar consumption) for each culture and, from these data,

we then estimated this yield relatively to that of *Sc* in single strain culture (Fig 4).

FIG 4

Yield of carbon metabolite production relative to the yield of production of single

S. cerevisiae. Average Yield are given with standard deviations for acetate, alpha-ketoglutarate,

ethanol, glycerol, pyruvate and succinate and each type of fermentation.

In the case of ethanol, only *Mf* fermentations had a relative yield significantly inferior (-

32%) to 0 (0 being *Sc* yield). For glycerol, all isolated NS fermentations had a greater yield

than *Sc* and mixed fermentations were intermediate between (but not significantly different

from) the corresponding single strain cultures. For acetate, only *Hanseniaspora* strains

displayed a higher yield (Fig 4).

Finally, all mixed cultures seemed to have a lower succinate yield than both

corresponding isolated cultures (but not significantly after correction for multiple tests).

For each fermentation, the total production of metabolites resulted from the combination

of species yields, total sugar consumption and respective population dynamics during

fermentation. Therefore, differences observed in the total productions of mixed cultures were

the consequences of additive or subtractive effects observed for these 3 components.

Considering ethanol, its total production was directly linked to the consumption of resources

and all mixed cultures were equivalent to *Sc* fermentations (S2 Fig). The case of glycerol was

more interesting. Indeed, even if the average sugar consumption was lower in *ScvsHu* and

ScvsMp mixed cultures than in single strain *Sc* culture, the total production of glycerol was

significantly higher than that of the corresponding single strain cultures (Glycerol$_{ScvsMp}$ =

6.1±0.1 g.L^{-1}, Glycerol$_{Sc}$ = 5.3±0.4 g.L^{-1}, Glycerol$_{Mp}$ = 3.7±0.2 g.L^{-1}). This resulted from the

positive combination of the greater glycerol yield by *Hanseniaspora* and *Metschnikowia* and

their population dynamics. For all other metabolites, the total production of mixed cultures was

not significantly different from the corresponding single strain cultures (S1 Table).

3. Discussion

244   This study presents one of the first works focusing on the population dynamics and

245   kinetics of yeast-yeast interactions between two species (S.C. and NS) during the alcoholic

246   fermentative process (18), but dynamics are distinct according to the *M. pulcherrima* and *H.*

247   *uvarum* strains tested in this study. Firstly, all strains grew in single strain and mixed strain

248   cultures. *Metschnikowia* single strain cultures present lower Vmax values than *Hanseniaspora*

single strain cultures, while mixed cultures containing *Metschnikowia* species had significantly higher Vmax values than those containing *Hanseniaspora* species. It appears that the better growth behavior of *Hanseniaspora* strains limited more *S. cerevisiae*'s growth than *Metschnikowia* in mixed culture. The growth characteristics of the competing strains directly impact the mixed fermentation characteristics.

Mixed population dynamics were characterized by similar growth rate (reaching a maximum population like *S. cerevisiae* single strain culture), followed by a long phase with decreasing mortality. NS viability dropped to 30% in accordance with Moreira et al., 2008 and Tristezza et al., 2016. Their NAS consumption was also similar, with half of the available ammonium not being consumed. Even their yield and total production of CCM metabolites were very similar. For almost all these characteristics, mixed cultures presented intermediate phenotypes compared to the corresponding single strain cultures.

However, there is a remarkable exception considering the total production of glycerol that is superior in mixed cultures whereas their sugar consumption was inferior (Figure 4). This point is characteristic of a transgressive interaction (often referred to as over-yielding), i.e. a situation in which the ecosystem performance is higher than that of the best-yielding species present (when cultivated alone). This glycerol overproduction in mixed cultures has already been observed in previous works (16,19) but is not systematic depending on the strains and experimental conditions (20). In the present study, the glycerol and sugar consumption observed in mixed cultures is for the first time explained by the joint effect of population dynamics (*S. cerevisiae* slowly dominating the population), resource consumptions (the **NS** fermentation leaving ⅔ of sugars) and the glycerol yields of NS strains that were two to three times higher that of *S. cerevisiae*.

These overproductions of glycerol open the way for potentially more interesting results. Our results point out that inoculation conditions in mixed cultures must be considered according

to fermentation aims and strains characteristics. Exploiting this specific yield and population

dynamics, for example by inoculating the S. *cerevisiae* strain at lower relative frequencies

(0.01, 0.001, 0.0001), more important transgressive interactions (such as lower ethanol

production) could be achieved in mixed cultures.

Meanwhile NS fermentations present similarities, it was possible to group the NS strains

performance according to their genus (see PCA in figures 5). While *H. uvarum* and *H. opuntiae*

both presented similar phenotypes, *M. pulcherrima* and *M. fructicola* presented a more distinct

phenotype. Regarding *Metschnikowia* species, **Mp** and **Mf** single strain cultures were

characterized by a short growth phase reaching a small carrying capacity and followed by a

long stationary phase, maintaining viability at a high level.

Nevertheless, despite *Metschnikowia* high viability, the resources measured (sugars and

nitrogen) were not entirely consumed. The reason why the cells stopped growing remains

unclear. A possible explanation could be linked to resources that have not been considered in

the present study. For instance, the third most important resource to be considered during

fermentation, even using synthetic medium, is lipid content, available as phytosterols and fatty

acids. In fact, for all yeast species, yeast lipid synthesis requires oxygen and is therefore

impossible in anaerobic conditions. With the progress of fermentation and ensuing oxygen

limitation yeast should import lipids to survive. If not, it stops to multiply. It is the lack of lipid

uptake under anaerobiosis that we can hypothesized for some *Metschnikowia* species. In fact,

we have shown that some of them were unable to import lipids from the medium, depending

instead on their own (internal) lipid synthesis for their growth (see Fig S1).

Interestingly, even though *S. cerevisiae*, *M. pulcherrima* and *M. fructicola* mortality rates

were low in single strain cultures, the corresponding mixed cultures (**ScvsMp** and **ScvsMf**)

presented 30% mortality (after 200 hours of fermentation). Moreover, this mortality seemed to

impact both species differently. In the mixed **ScvsMp** culture, only *S. cerevisiae* cells eventually

died, whereas in **ScvsMf** both species were negatively affected. The survival of *M. pulcherrima*

cells compared to *S. cerevisiae* cells could be explained by the production of pulcherriminic

acid by *M. pulcherrima* (21). Indeed, pulcherriminic acid is known to deplete iron from the

medium, which has a lethal effect on *S. cerevisiae* cells (13,22,23). In **ScvsMf** cultures, the

mortality observed in both species suggests a more complex mechanism of interaction

(although it is not clear whether *M. fructicola* also produces pulcherriminic acid; Kurtzman and

Droby, 2001). To explain these results, we could hypothesize the conjunction of two different

mechanisms of interaction. It could be that *M. fructicola* synthesized a metabolite

(pulcherriminic acid?) impacting the viability of *S. cerevisiae* cells (through iron depletion?),

with *M. fructicola* cells dying thereafter for another reason such as sensitivity to ethanol.

Indeed, the production of ethanol was almost four times higher in mixed cultures than in single

strain *Metschnikowia* fermentations. Under such hypothesis, the reason why no loss of viability

was observed for *M. pulcherrima* in mixed culture with *S. cerevisiae* could probably be related

to a better tolerance of *M. pulcherrima* to ethanol stress compared to *M. fructicola*.

For all *Metschnikowia* and *Hanseniaspora* mixed cultures, we only discussed

transgressive interactions when mixed cultures over-produced (or under-produced) a given

metabolite. Indeed, it was very difficult to identify interactions when the productions of mixed

fermentations were within the range of isolated cultures productions. As the relative frequency

of both strains in mixed cultures evolved during fermentation, it was difficult to link the final

mix to the contributions of each species. It was even more difficult to assess whether these

contributions combined additively or with interaction. New statistical developments or dual

transcriptomics will be needed to answer this question (24–26).

To discuss more generally the mechanisms of interactions of *Metschnikowia* and

Hanseniaspora cultures mixed with *S. cerevisiae*, we could not evidence any major antagonistic

phenomena. For almost all assays, mixed cultures performance stood always between that of

the corresponding isolated cultures (Fig 1, 2 & 3), except for the total production of glycerol.

Moreover, despite the differences in yield and interactions between species, the rapid

dominance of *S. cerevisiae* (increasing from 10% to at least 50% during the fermentation)

resulted in mixed cultures that were overall not different from *S. cerevisiae* single strain

cultures. This result is in agreement with the good adaptation of *S. cerevisiae* to winemaking

conditions (4–6,27) but could be different with a different set of strains. This result is important

in the context of ecological engineering. In fact, our results confirmed that *S. cerevisiae* has a

much better fitness than the NS strains studied in this paper. Therefore, if we want mixed culture

behavior to deviate from that of *S. cerevisiae* single strain culture, it must be ensured that NS

cells dominate the culture as soon as possible. To achieve this, two conceivable options are

currently tested: either to reduce the proportion of *S. cerevisiae* at t_0 or to successively inoculate

(16) first the NS species , and only then the *S. cerevisiae* strain. These two options could be

equivalent depending on the type of interaction(s) that occurs. If strain behaviors in single

strain or mixed cultures are identical, then all interactions are mediated by the medium through

the competition for resources and the production of constitutive toxins such as ethanol

(producing a toxin only in mixed fermentation would be a behavior change) and could be

qualified as "indirect".

Mathematical models could be designed from data on isolated cultures to predict the

impact of indirect interactions in mixed cultures. This would allow simulating numerous mixes

of species with various initial conditions and identify optimal strategies depending on one or

several given criteria. Using these approaches could limit the number of necessary tests,

potentially saving a lot of time and money and opening the way to a more methodical ecological

engineering. The development of such mathematical models will only be possible thanks to a deep tracking of population dynamics to understand underlying mechanisms of growth and mortality. Obviously, it is also critical to validate this approach by i) first extending the number of species co-cultured with *S. cerevisiae*, ii) investigating intra-specific variability and strain-strain interactions between species, iii) investigating the impact of the environment of culture (temperature, grape variety, etc.).

4. Mat & Met

4.1. Strains

In this work, we used one strain of 5 different species (one strain per species): *Saccharomyces cerevisiae* (**Sc**), *Metschnikowia pulcherrima* (**Mp**), *M. fructicola* (**Mf**), *Hanseniaspora uvarum* (**Hu**) and *H. opuntiae* (**Ho**). The *S. cerevisiae* strain is a haploid strain from EC1118 labelled with GFP (59A-GFP, Marsit and Dequin, 2015). The *Hanseniaspora uvarum* (CLIB 3221) *H. opuntiae* (CLIB 3093) and *Metschnikowia pulcherrima* (CLIB 3235) strains originated from the yeast CIRM (https://www6.inra.fr/cirm_eng/Yeasts/Strain-catalogue) and were isolated from grape musts. The *Metschnikowia fructicola strain was* from the Lallemand collection.

For each strain, 3 replicates of single strain cultures were performed (except for *S. cerevisiae* that had a total of 8 replicates in different blocks). In addition, for each NS strain, 3 replicates of a mixed culture with the **Sc** strain were performed. In all mixed fermentations, the starting proportion of *S. cerevisiae* cells was set at 10%. In this text, fermentations were referred

to by the species that performed them, i.e. single strain cultures were referred to as: **Sc**, **Mp**,

Mf, **Hu** and **Ho** and mixed strain cultures as **ScvsMp**, **ScvsMf**, **ScvsHu** and **ScvsHo**.

4.2. Medium

Initial cultures (12 h, in 50 ml YPD medium, 28 °C) were used to inoculate fermentation

media at a total density of 10^6 cells/mL; therefore, for mixed culture the *S. cerevisiae* cells

density was 0.1×10^6 /mL and the NS cells density was 0.9×10^6 /mL. Fermentations were carried

out in a synthetic medium (SM) mimicking standard grape juice (28). The SM used in this study

contained 200 g/L of sugar (100 g glucose and 100 g fructose per liter) and 200 mg/L of

assimilable nitrogen (as a mix of ammonium chloride and amino acids). The concentrations of

weak acids, salts and vitamins were identical to those described by Seguinot et al. (29). The pH

of the medium was adjusted to 3.3 with 10M NaOH. The SM medium was first saturated with

bubbling air for 40 minutes, then it was supplemented with 5 mg/L phytosterols (85451, Sigma

Aldrich) solubilized in Tween 80 to fulfill the lipid requirements (sterols and fatty acids) of

yeast cells during anaerobic growth.

4.3. Measurements

Fermentation took place in 1.1-liter fermentors equipped with fermentation locks to

maintain anaerobiosis, at 20 °C, with continuous magnetic stirring (500 rpm) during

approximately 300h. CO_2 release was followed by automatic measurements of fermentor

weight loss every 20 min. The amount of CO_2 released allowed us to monitor the progress of

the fermentation and evaluate the maximum of released CO_2 (**CO2max**) as well as the

maximum rate of CO_2 released (**Vmax**). Samples were harvested after 6h, 12h and 24h, then

every 12h during the first week and every 24h during the second week of fermentation. For

each sample, the population density cells were determined using a BD Accuri™ C6 Plus flow

cytometer as described in (30). Viability was determined using propidium iodide staining and BD Accuri™ C6 Plus flow cytometer adapted from (31). Proportions of *S. cerevisiae* in mixed-culture was established thanks to the green fluorescence produced by the *S. cerevisiae* 59A-GFP, the forward and side scatters from the BD Accuri™ C6 Plus flow cytometer and machine learning using the caret package in R (32). From these population densities (without taking into account viability), we fitted a growth population model (with the growthcurver package in R, (33), and determined the carrying capacity (***K***) and maximum growth rate (***mu***) for each fermentation.

The final concentrations of carbon metabolites in the medium (acetate, succinate, glycerol, alpha-ketoglutarate, pyruvate, ethanol, glucose and fructose) were determined with high-pressure liquid chromatography (34). From these metabolite concentrations, we first calculated the consumed sugar concentration as the difference between the final and the initial concentration of either glucose or fructose. Then we calculated the yield of metabolite production by dividing the final concentration by the corresponding consumed sugar concentration. Finally, we compared these yields to the yield of *S. cerevisiae* single strain cultures considered as reference.

Finally, the ammonium concentration after 100h of fermentation was determined enzymatically with R-Biopharm (Darmstadt, Germany) and the free amino acid content of the must was estimated through cation exchange chromatography with post-column ninhydrin derivatization (35).

4.4. Statistical Analysis

The experimental work was performed in 5 different blocks. Each block was composed of three replicates of NS fermentations (for example ***Hu***), three replicates of the corresponding mixed fermentations with *S. cerevisiae* (for example ***ScvsHu***) and one or two fermentations of

single strain *S. cerevisiae* cells (***Sc***). The block effect was evaluated on the parameters of the

Sc fermentation. For most studied parameters, the block effect was not significant. For those

parameters where a block effect was observed (***mu*** and ***K***), a statistical correction for block

effect did not modify our results. Therefore, for simplification purposes, we compared all

fermentations without any correction for the block effect parameter. For each measured

parameter, an ANOVA was performed to evaluate the type of fermentation (***Sc***, ***Mp***, ***Mf***, ***Hu***,

Ho, ***ScvsTd***, ***ScvsMp***, ***ScvsMf***, ***ScvsHu*** and ***ScvsHo***) effect and then a Tukey t-test was

performed to determine statistical groups and two-by-two statistical differences. All statistical

analyses were performed using R (36) and Rstudio (37). All data, analysis and figures scripts

can be found in this github address: https://github.com/tnidelet/Git-Harle-et-al-2019.

Acknowledgments.

434 We thank the CIRM, the Lallemand company, Jean-Luc Legras, Virginie Galeote and Jean-

435 Nicolas Jasmin for providing the strains used in this study. We thank also Christian Picou, Marc

436 Perez, Faiza Macna for technical assistance and Delphine Sicard for advices.

References

1. Marsit S, Dequin S. Diversity and adaptive evolution of Saccharomyces wine yeast: a
review. FEMS Yeast Research. 1 nov 2015;15(7):fov067.

2. Sablayrolles J. Fermented beverages: the example of winemaking. Advances in
Fermentation Technology New Delhi: Asiatech Publishers. 2008;322–47.

3. García-Ríos E, Gutiérrez A, Salvadó Z, Arroyo-López FN, Guillamon JM. The Fitness
Advantage of Commercial Wine Yeasts in Relation to the Nitrogen Concentration,
Temperature, and Ethanol Content under Microvinification Conditions. Appl Environ
Microbiol. 15 janv 2014;80(2):704-13.

4. Jolly NP, Varela C, Pretorius IS. Not your ordinary yeast: non-Saccharomyces yeasts in
wine production uncovered. Fems Yeast Res. 2014;14(2):215-37.

5.   Pinto C, Pinho D, Cardoso R, Custódio V, Fernandes J, Sousa S, et al. Wine
fermentation microbiome: a landscape from different Portuguese wine appellations.
2015;6:905.

6.   Varela C. The impact of non-Saccharomyces yeasts in the production of alcoholic
beverages. Appl Microbiol Biot. 2016;100(23):9861-74.

7.   Ivey M, Massel M, Phister TG. Microbial Interactions in Food Fermentations. Annual
Review of Food Science and Technology. 2013;4(1):141-62.

8.   Sadineni V, Kondapalli N, Obulam V. Effect of co-fermentation with Saccharomyces
cerevisiae and Torulaspora delbrueckii or Metschnikowia pulcherrima on the aroma and
sensory properties of mango wine. Ann Microbiol. 2012;62(4):1353-60.

9.   P H-O, Cersosimo M, Loscos N, Cacho J, E G-M, Ferreira V. The development of
varietal aroma from non-floral grapes by yeasts of different genera. Food Chem.
2008;107(3):1064-77.

10. Contreras A, Hidalgo C, Henschke PA, Chambers PJ, Curtin C, Varela C. Evaluation of
Non-Saccharomyces Yeasts for the Reduction of Alcohol Content in Wine. Appl
Environ Microbiol. 1 mars 2014;80(5):1670-8.

11. Belda I, Navascués E, Marquina D, Santos A, Calderon F, Benito S. Dynamic analysis
of physiological properties of Torulaspora delbrueckii in wine fermentations and its
incidence on wine quality. Appl Microbiol Biot. 2015;99(4):1911-22.

12. Elena G-R, Pascual O, Kontoudakis N, Esteruelas M, Braulio E-Z, Mas A, et al.
Oenological consequences of sequential inoculation with non-Saccharomyces yeasts
(Torulaspora delbrueckii or Metschnikowia pulcherrima) and Saccharomyces cerevisiae
in base wine for sparkling wine production. Eur Food Res Technol.
2015;240(5):999-1012.

13. Oro L, Ciani M, Comitini F. Antimicrobial activity of Metschnikowia pulcherrima on
wine yeasts. J Appl Microbiol. 2014;116(5):1209-17.

14. Kurtzman CP, Droby S. Metschnikowia fructicola, a New Ascosporic Yeast with
Potential for Biocontrol of Postharvest Fruit Rots. Systematic and Applied
Microbiology. janv 2001;24(3):395-9.

15. Hu K, Qin Y, Tao Y-S, Zhu X-L, Peng C-T, Ullah N. Potential of Glycosidase from
Non-Saccharomyces Isolates for Enhancement of Wine Aroma. Journal of Food Science.
81(4):M935-43.

16. Tristezza M, Tufariello M, Capozzi V, Spano G, Mita G, Grieco F. The Oenological
Potential of Hanseniaspora uvarum in Simultaneous and Sequential Co-fermentation
with Saccharomyces cerevisiae for Industrial Wine Production. Front Microbiol
[Internet]. 2016 [cité 2 juill 2018];7. Disponible sur:
https://www.frontiersin.org/articles/10.3389/fmicb.2016.00670/full

17. Song H-S, Cannon WR, Beliaev AS, Konopka A. Mathematical Modeling of Microbial
Community Dynamics: A Methodological Review. Processes. 17 oct 2014;2(4):711-52.

18. Wang C, Mas A, Esteve-Zarzoso B. The Interaction between Saccharomyces cerevisiae
and Non-Saccharomyces Yeast during Alcoholic Fermentation Is Species and Strain
Specific. Front Microbiol [Internet]. 2016 [cité 8 oct 2019];7. Disponible sur:
https://www.frontiersin.org/articles/10.3389/fmicb.2016.00502/full

19. Ciani M, Ferraro L. Enhanced Glycerol Content in Wines Made with Immobilized
Candida stellata Cells. Appl Environ Microbiol. 1 janv 1996;62(1):128-32.

20. Moreira N, Mendes F, de Pinho P, journal of … H-T. Heavy sulphur compounds, higher
alcohols and esters production profile of Hanseniaspora uvarum and Hanseniaspora
guilliermondii grown as pure and …. International journal of …. 2008;

21. MacDonald. Biosynthesis of pulcherriminic acid. Biochem J. 1965;96(2):533-8.

22. Sipiczki M. Metschnikowia Strains Isolated from Botrytized Grapes Antagonize Fungal
and Bacterial Growth by Iron Depletion. Appl Environ Microb. 2006;72(10):6716-24.

23. Türkel S, C E-B. Isolation and characterization of new Metschnikowia pulcherrima
strains as producers of the antimicrobial pigment pulcherrimin. Zeitschrift für
Naturforschung C. 2009;

24. Barot S, Allard V, Cantarel A, Enjalbert J, Gauffreteau A, Goldringer I, et al. Designing
mixtures of varieties for multifunctional agriculture with the help of ecology. A review.
Agron Sustain Dev. 2017;37(2):13.

25. Finckh MR (International RRI, Mundt CC. Stripe rust, yield, and plant competition in
wheat cultivar mixtures. Phytopathology (USA) [Internet]. 1992 [cité 17 janv 2019];
Disponible sur: http://agris.fao.org/agris-search/search.do?recordID=US9307396

26. Wolf T, Kämmer P, Brunke S, Linde J. Two's company: studying interspecies
relationships with dual RNA-seq. Current Opinion in Microbiology. 1 avr 2018;42:7-12.

27. García M, Braulio E-Z, Arroyo T. Non-Saccharomyces Yeasts: Biotechnological Role
for Wine Production. 2016.

28. Bely M, Sablayrolles JM, Barre P. Description of Alcoholic Fermentation Kinetics: Its
Variability and Significance. Am J Enol Vitic. 1 janv 1990;41(4):319-24.

29. Seguinot P, SANCHEZ I, Bloem A, Ortiz-Julien A, Camarasa C. Impact of nitrogen
sources on the fermentative kinetic of non - Saccharomyces yeasts [Internet]. 2017 [cité
2 juill 2018]. np. Disponible sur: https://hal.archives-ouvertes.fr/hal-01604146

30. Delobel P, Pradal M, Blondin B, Tesniere C. A 'fragile cell' sub-population revealed
during cytometric assessment of Saccharomyces cerevisiae viability in lipid-limited
alcoholic fermentation. Letters in Applied Microbiology. 55(5):338-44.

31. Riccardi C, Nicoletti I. Analysis of apoptosis by propidium iodide staining and flow
cytometry. Nat Protoc. août 2006;1(3):1458-61.

32. Kuhn M. Building Predictive Models in R Using the caret Package. Journal of Statistical
Software. 10 nov 2008;28(1):1-26.

33. Sprouffske K, Wagner A. Growthcurver: an R package for obtaining interpretable
metrics from microbial growth curves. BMC Bioinformatics. 19 avr 2016;17(1):172.

34. Camarasa C, Sanchez I, Brial P, Bigey F, Dequin S. Phenotypic landscape of
Saccharomyces cerevisiae during wine fermentation: evidence for origin-dependent
metabolic traits. PLoS ONE. 2011;6(9):e25147.

35. Crépin L, Nidelet T, Sanchez I, Dequin S, Camarasa C. Sequential Use of Nitrogen
Compounds by Saccharomyces cerevisiae during Wine Fermentation: a Model Based on
Kinetic and Regulation Characteristics of Nitrogen Permeases. 2012;78(22):8102-11.

36. R Core Team (2016). R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria. URL   https://www.R-
project.org/.

37. RStudio (2012). RStudio: Integrated development environment for R (Version 0.96.122)
[Computer software]. Boston, MA. Retrieved May 20, 2012. Available from
http://www.rstudio.org/.

Supporting information

S1 FIG

Glucose and Fructose consumption kinetics in function of different strains. Each point represents a sample (average ± standard deviation).

S2 FIG

Total production of carbon metabolite in function of the strains driving the fermentation. Average production are given with standard deviations for acetate, alpha-ketoglutarate, ethanol, glycerol, pyruvate and succinate.

551
552
553 **S3 FIG**

554 **Principal component analysis of carbon metabolites and growth parameter of isolated**
555 **cultures.** The mixed cultures are a second time projected on the plan determiner by only
556 isolated cultures. In the top right is represented the circle of variables.
557

559
S4 FIG

Serial tenfold dilutions of two Metschnikovia pulcherrima strains (A and B) spotted
onto various synthetic standard agar media (SM425, 425 mg/l assimilable nitrogen) with
Tween 80, (Tw, 0.06%), supplemented or not with phytosterol (Phyto, 20 mg/L), in the
presence or not of fluconazole (FLC, 256 μg/mL). Plates were incubated at 28°C for five
days in air or in anaerobiosis.
566

569 **Growth parameter values for each type of fermentation.**

	μ	K	prop	Vmax	MaxCO2	Sugar	NAS	Ethanol	Glycerol	Succinate	Acetate	Pyruvate	Alpha
Sc	0.15±0.02[bc]	1.6±0.2[bcd]	NA	0.99±0.03[a]	88.2±2.2[a]	0.2±0.2[d]	10.4±8.6[c]	95.2±8.4[a]	5.3±0.4[c]	1.05±0.28[a]	0.63±0.06[d]	0.12±0.08[bc]	0.08±0.04[ab]
Ho	0.19±0.03[bc]	1.6±0.1[bc]	NA	0.42±0.02[d]	46.4±0.6[d]	92.3±1.4[b]	122.2±3.5[b]	50.1±0.2[b]	5.2±0.0[c]	0.35±0.00[b]	0.99±0.02[b]	0.13±0.05[bc]	0.09±0.1[ab]
ScvsHo	**0.11±0.00[bc]**	**2.3±0.1[a]**	0.83±0.02[b]	0.57±0.01[c]	82.7±0.6[bc]	23.3±1.3[c]	**6.4±2.4[c]**	87.2±0.8[a]	**6.0±0.0[abc]**	0.42±0.04[b]	0.84±0.01[c]	**0.08±0.00[c]**	**0.03±0.0[b]**
Hu	0.62±0.18[a]	1.4±0.1[cde]	NA	0.34±0.04[d]	30.7±0.5[e]	134.3±11.2[a]	173.3±24.8[b]	34.9±1.1[c]	5.5±0.2[bc]	0.66±0.03[b]	1.15±0.03[a]	0.17±0.04[abc]	0.14±0.09[ab]
ScvsHu	0.24±0.01[b]	**1.7±0.1[b]**	0.76±0.01[c]	0.53±0.04[c]	82.1±1.5[c]	16.5±6.5[c]	**4.6±4.2[c]**	89.5±3.8[a]	**6.6±0.3[a]**	**0.60±0.06[b]**	0.86±0.07[bc]	0.16±0.04[abc]	0.09±0.06[ab]
Mf	0.17±0.02[bc]	0.8±0.05[f]	NA	0.17±0.04[e]	20.6±1.2[f]	135.8±9.7[a]	256.6±2.9[a]	20.6±0.3[c]	2.8±0.1[e]	0.29±0.01[b]	0.13±0.01[f]	0.10±0.01[bc]	0.10±0.00[ab]
ScvsMf	**0.09±0.01[bc]**	1.3±0.03[de]	0.96±0.02[a]	0.76±0.03[b]	86.8±0.7[ab]	9.9±2.1[cd]	**1.8±0.1[c]**	92.1±3.2[a]	**6.0±0.1[abc]**	0.58±0.02[b]	0.46±0.02[e]	0.12±0.01[bc]	0.08±0.01[ab]
Mp	0.18±0.03[bc]	0.6±0.01[f]	NA	0.16±0.02[e]	21.8±0.4[f]	142±3.7[a]	305.5±5.9[a]	23.4±1.8[c]	3.7±0.2[d]	0.26±0.03[b]	0.07±0.06[f]	0.29±0.04[a]	0.13±0.01[ab]
ScvsMp	**0.09±0.00[c]**	1.3±0.01[e]	0.50±0.02[d]	0.68±0.04[b]	83.3±2.2[bc]	20.8±6.0[c]	**2.1±0.2[c]**	86.5±3.4[a]	**6.1±0.1[ab]**	0.52±0.04[b]	0.46±0.02[e]	0.24±0.01[ab]	**0.19±0.01[a]**

581 Mu : h-1; K : 10^8 cells.mL-1, prop : no unity, Vmax : g.L-1.h-1, MaxCO2 : g.L-1, Final Sugar : g.L-1, Final NAS : mg.L-1, Ethanol : g.L-1,

582 Glycerol : g.L-1, Succinate : g.L-1, Acetate : g.L-1, Pyruvate : g.L-1, Alpha : g.L-1

583 **Bold:** values of mixed culture that are not intermediate between the two values of the corresponding isolated cultures, a sign of transgressive

584 interactions.

585 Superscript: Statistical groups obtained from a Tukey test.