Genetic evidence for two carbon fixation pathways in symbiotic and free-living bacteria: The
Calvin-Benson-Bassham cycle and the reverse tricarboxylic acid cycle

Maxim Rubin-Blum[1,2*], Nicole Dubilier[1,3], Manuel Kleiner[4*]

[1]Max-Planck Institute for Marine Microbiology, Celsiusstrasse 1, 28359 Bremen, Germany

[2]Israel Limnology and Oceanography Research, Tel Shikmona, 3108000, Haifa, Israel

[3]MARUM, University of Bremen, 28359 Bremen, Germany

[4]Department of Plant & Microbial Biology, North Carolina State University, Raleigh, NC, USA

*Corresponding authors

mrubin@ocean.org.il

manuel_kleiner@ncsu.edu

Abstract: Very few bacteria are able to fix carbon via both the reverse tricarboxylic acid (rTCA)

and the Calvin-Benson-Bassham (CBB) cycles, such as symbiotic, sulfur-oxidizing bacteria that

are the sole carbon source for the marine tubeworm *Riftia pachyptila,* the fastest growing

invertebrate. To date, this co-existence of two carbon fixation pathways had not been found in a

cultured bacterium and could thus not be studied in detail. Moreover, it was not clear if these two

pathways were encoded in the same symbiont individual, or if two symbiont populations, each

with one of the pathways, co-existed within tubeworms. With comparative genomics, we show

that *Thioflavicoccus mobilis,* a cultured, free-living gammaproteobacterial sulfur oxidizer,

possesses the genes for both carbon fixation pathways. Here, we also show that both the CBB and

rTCA pathways are likely encoded in the genome of the sulfur-oxidizing symbiont of the

tubeworm *Escarpia laminata* from deep-sea asphalt volcanoes in the Gulf of Mexico. Finally, we

provide genomic and transcriptomic data suggesting a potential electron flow towards the rTCA

cycle carboxylase 2-oxoglutarate:ferredoxin oxidoreductase, via a rare variant of NADH

dehydrogenase/heterodisulfide reductase. This electron bifurcating complex, together with

NAD(P)+ transhydrogenase and Na+ translocating Rnf membrane complexes may improve the

efficiency of the rTCA cycle in both the symbiotic and the free-living sulfur oxidizer.

Importance: Primary production on Earth is dependent on autotrophic carbon fixation, which

leads to the incorporation of carbon dioxide into biomass. Multiple metabolic pathways have

been described for autotrophic carbon fixation, but most autotrophic organisms were assumed to

have the genes for only one of these pathways. Our finding of a cultivable bacterium with two

carbon fixation pathways in its genome opens the possibility to study the potential benefits of

having two pathways and the interplay between these pathways. Additionally, this will allow the

investigation of the unusual, and potentially very efficient mechanism of electron flow that could

drive the rTCA cycle in these autotrophs. Such studies will deepen our understanding of carbon

fixation pathways and could provide new avenues for optimizing carbon fixation in

biotechnological applications.

Observation

Primary production by autotrophic organisms drives the global carbon cycle. Currently, seven naturally occurring pathways for inorganic carbon fixation are known in autotrophic organisms (1). The dominant carbon fixation pathway used by plants, algae, and many bacteria is the Calvin-Benson-Bassham (CBB) cycle. The six more efficient, alternative pathways are limited to autotrophic microbes that live in reducing habitats, due to the oxygen sensitivity of these alternative pathways (2, 3). Only a handful of autotrophic organisms have more than one carbon fixation pathway: The sulfur-oxidizing symbionts of marine tubeworms such as *Riftia, Escarpia, Tevnia and Lamellibrachia* (the symbionts of these hosts are closely related to each other) have and express both the oxygen-tolerant CBB cycle and the oxygen-sensitive reverse tricarboxylic acid (rTCA) cycle (4–8). Only a few free-living bacteria may have the genes for both cycles, such as the large sulfur bacteria, *Beggiatoa* and *Thiomargarita* spp., in which all CBB cycle genes and some rTCA cycle genes were found to co-exist in their genomes (9–11). The CBB cycle in the symbionts and the large sulfur bacteria is potentially more energy efficient than the classical version of the CBB cycle based on the replacement of the fructose-1,6-bisphosphatase with a pyrophosphate dependent enzyme (9, 10, 12, 13). In addition, it is likely that the interplay between the CBB and rTCA cycle under fluctuating redox conditions contributes to the high efficiency of carbon fixation in tubeworm symbioses (4, 5, 14), and consequently to the extremely high growth rates of tubeworms, which grow faster than any other known invertebrate (15).

Given that tubeworm symbionts and large sulfur bacteria could not yet be cultivated, it was not possible to investigate the co-occurrence of two carbon fixation cycles in detail to better understand the biochemical and physiological mechanisms that enable the interplay between these two pathways. In this study, we sequenced the genome and transcriptome of the symbiont from the tubeworm *Escarpia laminata* and compared its genome to those of other tubeworm symbionts and free-living microbes. These comparisons led us to discover the presence of co-occurring CBB and rTCA cycles in the genome of a cultured bacterium.

Co-occurrence of rTCA cycle genes with RuBisCo in symbiotic and free-living

gammaproteobacteria. Genes for enzymes that are specific to the rTCA pathway, that is the ATP

citrate lyase (*aclAB* genes), 2-oxoglutarate:ferredoxin oxidoreductase (OGOR, *korABCD* genes),

and a putative fumarate reductase (*tfrAB* genes, homologs of genes encoding a thiol:fumarate

reductase from *Methanobacterium thermoautotrophicum* (16)), were assumed to occur in only a

few symbiotic Gammaproteobacteria. We discovered, using comparative genomics, that these

rTCA cycle enzymes also occur in some Chromatiaceae, including the cultivated sulfur oxidizer

Thioflavicoccus mobilis and a gammaproteobacterial genome from an environmental

metagenome(17), (**Fig. 1**). The type II ATP citrate lyases of tubeworm symbionts and *T. mobilis*

were likely acquired via horizontal gene transfer from other bacterial clades (**Suppl. Fig. S1**), (6).

These gammaproteobacteria also encode either Form I or II RuBisCO, or both (**Suppl. Note 1**).

Presence of the rTCA and the CBB pathways in the genome of a single bacterium: Due to the

fragmented nature of the previously available genomes of tubeworm symbionts, past studies

could not determine whether the genes for both pathways are present in a single genome or if the

two pathways are distributed in a strain-specific manner, i.e. only one of the two pathways is

present in the genome of a single cell (3). Here, we provide two lines of evidence that both

pathways can co-occur in the genome of a single organism. First, sequencing coverage for the

genes of both pathways in the *E. laminata* symbiont was similar to that of single-copy marker

genes (**Suppl. Table 1**). Since genes that are strain specific are expected to have lower coverage

than the rest of the genome (18), the similar coverage of genes encoding the two pathways and

single-copy genes suggests that in the *E. laminata* symbiont both pathways are present in all cells.

Second, in the closed genome of the cultured *T. mobilis*, both the genes encoding the rTCA and

the CBB cycle co-occur, providing evidence that these genes co-exist in a single genome.

Our transcriptomic analyses of *E. laminata* tubeworm symbionts revealed high expression levels

of both the rTCA and the CBB cycle genes (Fig. 2). This observation is consistent with previous

proteomic analyses of the *Riftia* symbiont (4, 5). The high expression levels of genes from the

rTCA and the CBB cycle suggests that both pathways play an important metabolic role in these

symbionts. It is, however, not clear whether these cycles function simultaneously within single

symbiont cells, or are differentially expressed within the symbiont population (3).

The rTCA gene clusters are conserved among the tubeworm symbionts and some

Chromatiaceae bacteria. In the tubeworm symbionts, the cultivated *T. mobilis* and the

gammaproteobacterial genome from an environmental metagenome, there was a considerable

level of conservation of the rTCA gene clusters, at the sequence and synteny levels (**Fig. 2**). The

aclAB genes that encode the two subunits of the ATP citrate lyase were accompanied by those

that encode bidirectional TCA cycle enzymes, including *acn* (aconitase)*, idh (*isocitrate

dehydrogenase), and *mdh* (malate dehydrogenase). The other rTCA specific genes *korABCD*

(four-subunit OGOR) and *tfrAB* (putative thiol:fumarate reductase), were also present in the

rTCA gene cluster. Similar to the ATP citrate lyase, the four-subunit OGOR, as well as the thiol:

fumarate reductase are very rare among Gammaproteobacteria, and were probably acquired via a

single horizontal gene transfer event from a distant bacterial clade (**Fig. 1**, **Suppl. Fig. S2 and**

S3). A dimeric OGOR (*korAB* genes), more common than the four-subunit enzyme among

gammaproteobacterial autotrophs, yet absent in *T. mobilis,* was located elsewhere in the genome

of the *E. laminata* symbiont. The *korAB* genes were co-localized with genes that encode

additional rTCA cycle enzymes (**Suppl. Note 2, Suppl. Fig. S4**).

An array of genes that encode several electron translocating complexes were integrated into the

rTCA cycle gene clusters. These complexes included an electron-bifurcating NADH

dehydrogenase/heterodisulfide reductase complex (*flxABCD-hdrABC* genes, **Suppl. Note 3**), a

NAD(P)+ transhydrogenase and Na+ translocating Rnf membrane complex (*pntAB* and

rnfABCDGE genes, **Suppl. Note 4**). Most interestingly, the conserved interspersing of the

korABCD and *tfrAB* genes with the *flxABCD-hdrABC* genes hints at the possibility that these

proteins form a complex that efficiently shuttles electrons directly to the OGOR and the thiol:

fumarate reductase (**Suppl. Figure S5**). If this is the case, the carbon fixation efficiency of the

rTCA cycle would be most likely considerably higher than the canonical rTCA cycle.

Conclusions. Until now, the only bacteria known to possess two carbon fixation pathways were

sulfur-oxidizing, tubeworm symbionts, and possibly also large sulfur bacteria, all of which are

currently not amenable to cultivation-based studies. With the discovery of the co-existence of the

CBB and rTCA cycle in the cultivable *T. mobilis*, experimental studies are now feasible. Such

studies would reveal if these pathways are expressed under different physicochemical conditions,

and potentially allow the biotechnological optimization of efficiency and yield in production

processes that rely on autotrophic carbon fixers. To our knowledge, the use of organisms with

multiple carbon fixation pathways has not been used as a design principle for these applications.

Methods: Comparative genomics and transcriptomics. Publically available genomes from

NCBI and JGI-IMG collections, as well as de-novo assembled genomes of *Escarpia laminata*

symbionts (estimated completeness 99.5%), were used for genomic comparison (see

Supplementary Methods). To verify presence/absence of target gene homologs in sequenced

organisms we used NCBI's BLAST against the nucleotide collection and non-redundant protein

database (19). *E. laminata* symbiont genomes were used as a template for genome-centered

transcriptomics (sequences available under the BioProject accession number PRJNA471406).

Phylogenetic and phylogenomic analyses. Phylogenomic treeing was performed using scripts

available at phylogenomics-tools (DOI:10.5281/zenodo.46122). Twenty-three marker proteins

that are universally conserved across the bacterial domain were extracted from genomes using

the AMPHORA2 pipeline (20). Twenty-three single-copy markers were used for alignment with

MUSCLE (21). The marker alignments were concatenated into a single partitioned alignment,

and poorly aligned regions were removed. Functional protein sequences were aligned with

MAFFT (22). Maximum Likelihood trees were calculated with IQ-tree (23) and MEGA7 (24),

using the best-fitting model.

Author Contributions

MRB, ND and MK conceived the study. MRB and MK analyzed the samples. MRB, ND

and MK wrote the manuscript.

Acknowledgments

The authors thank all individuals who helped during the R/V Meteor research cruise

M114, including onboard technical and scientific personnel, the captain and crew, and the ROV

MARUM-Quest team. We thank the Max Planck-Genome-Centre Cologne

(http://mpgc.mpipz.mpg.de/home/) for generating the metagenomic and the metatranscriptomic

data used in this study. We thank Julie Reveillaud for providing genomes of Mid-Cayman Rise

tubeworm symbionts, and Matthias Winkel, Marc Mußmann and Jake V. Bailey for helpful and

supportive comments on a bioRxiv preprint version of this manuscript. The Campeche Knoll

cruise was funded by the German Research Foundation (DFG – Deutsche

Forschungsgemeinschaft). We are grateful to the Mexican authorities for granting permission to

conduct this research in the southern Gulf of Mexico (permission of DGOPA: 02540/14 from 5

November 2014). This study was funded by the Max Planck Society, the MARUM DFG-Research

Center / Excellence Cluster "The Ocean in the Earth System" at the University of Bremen, an

ERC Advanced Grant (BathyBiome, 340535), a Gordon and Betty Moore Foundation Marine

Microbial Initiative Investigator Award to ND (Grant GBMF3811) and the NC State Chancellor's

Faculty Excellence Program Cluster on Microbiomes and Complex Microbial Communities

(MK).

Conflict of interest

The authors declare no conflict of interest.

References

1.   Figueroa IA, Barnum TP, Somasekhar PY, Carlström CI, Engelbrektson AL, Coates JD.

2018. Metagenomics-guided analysis of microbial chemolithoautotrophic phosphite

oxidation yields evidence of a seventh natural CO_2 fixation pathway. Proc Natl Acad Sci

115:E92–E101.

2.   Erb TJ. 2011. Carboxylases in natural and synthetic microbial pathways. Appl Environ

Microbiol 77:8466–8477.

3.   Hügler M, Sievert SM. 2011. Beyond the Calvin cycle: Autotrophic carbon fxation in the

Ocean. Ann Rev Mar Sci 3:261–289.

4.   Gardebrecht A, Markert S, Sievert SM, Felbeck H, Thürmer A, Albrecht D, Wollherr A,

Kabisch J, Le Bris N, Lehmann R, Daniel R, Liesegang H, Hecker M, Schweder T. 2012.

Physiological homogeneity among the endosymbionts of Riftia pachyptila and *Tevnia*

jerichonana revealed by proteogenomics. ISME J 6:766–76.

5.   Markert S, Arndt C, Felbeck H, Becher D, Sievert SM, Hugler M, Albrecht D, Robidart J,

Bench S, Feldman RA, Hecker M, Schweder T. 2007. Physiological Proteomics of the

Uncultured Endosymbiont of *Riftia pachyptila*. Science 315:247–250.

6.   Markert S, Gardebrecht A, Felbeck H, Sievert SM, Klose J, Becher D, Albrecht D, Thürmer

178   A, Daniel R, Kleiner M, Hecker M, Schweder T. 2011. Status quo in physiological

179   proteomics of the uncultured *Riftia pachyptila* endosymbiont. Proteomics 11:3106–3117.

180 7. Robidart JC, Bench SR, Feldman RA, Novoradovsky A, Podell SB, Gaasterland T, Allen

181   EE, Felbeck H. 2008. Metabolic versatility of the *Riftia pachyptila* endosymbiont revealed

182   through metagenomics. Environ Microbiol 10:727–737.

183 8. Reveillaud J, Anderson R, Reves-Sohn S, Cavanaugh C, Huber JA. 2018. Metagenomic

184   investigation of vestimentiferan tubeworm endosymbionts from Mid-Cayman Rise reveals

185   new insights into metabolism and diversity. Microbiome 6:19.

186 9. Winkel M, Carvalho V, Woyke T, Richter M, Schulz-Vogt HN, Flood B, Bailey J,

187   Mußmann M. 2016. Single-cell sequencing of *Thiomargarita* reveals genomic flexibility for

188   adaptation to dynamic redox conditions. Front Microbiol 7:964.

189 10. MacGregor BJ, Biddle JF, Harbort C, Matthysse AG, Teske A. 2013. Sulfide oxidation,

190   nitrate respiration, carbon acquisition, and electron transport pathways suggested by the

191   draft genome of a single orange Guaymas Basin *Beggiatoa* (*Cand.* Maribeggiatoa) sp.

192   filament. Mar Genomics 11:53–65.

193 11. Flood BE, Fliss P, Jones DS, Dick GJ, Jain S, Kaster A-K, Winkel M, Mußmann M, Bailey J.

194   2016. Single-Cell (Meta-)Genomics of a dimorphic *Candidatus* Thiomargarita nelsonii

195   reveals genomic plasticity. Front Microbiol 7:1–17.

196   12.   Kleiner M, Wentrup C, Lott C, Teeling H, Wetzel S, Young J, Chang Y-JY-J, Shah M,

197   VerBerkmoes NC, Zarzycki J, Fuchs G, Markert S, Hempel K, Voigt B, Becher D, Liebeke

198   M, Lalk M, Albrecht D, Hecker M, Schweder T, Dubilier N. 2012. Metaproteomics of a

199   gutless marine worm and its symbiotic microbial community reveal unusual pathways for

200   carbon and energy use. Proc Natl Acad Sci USA 109:E1173-82.

201   13.   Kleiner M, Petersen JM, Dubilier N. 2012. Convergent and divergent evolution of

202   metabolism in sulfur-oxidizing symbionts and the role of horizontal gene transfer. Curr

203   Opin Microbiol 15:621–31.

204   14.   Klatt JM, Polerecky L. 2015. Assessment of the stoichiometry and efficiency of CO_2

205   fixation coupled to reduced sulfur oxidation. Front Microbiol 6:484.

206   15.   Bright M, Klose J, Nussbaumer AD. 2013. Giant tubeworms. Curr Biol 23:R224–R225.

207   16.   Heim S, Kunkel A, Thauer RK, Hedderich R. 1998. Thiol : fumarate reductase (Tfr) from

208   *Methanobacterium thermoautotrophicum.* Identification of the catalytic sites for fumarate

209   reduction and thiol oxidation. Eur J Biochem 253:292–299.

210   17.   Anantharaman K, Brown CT, Hug LA, Sharon I, Castelle CJ, Probst AJ, Thomas BC,

211   Singh A, Wilkins MJ, Karaoz U, Brodie EL, Williams KH, Hubbard SS, Banfield JF. 2016.

212   Thousands of microbial genomes shed light on interconnected biogeochemical processes

213   in an aquifer system. Nat Commun 7:13219.

18.   Pasić L, Rodriguez-Mueller B, Martin-Cuadrado A-B, Mira A, Rohwer F, Rodriguez-

Valera F. 2009. Metagenomic islands of hyperhalophiles: the case of *Salinibacter ruber*.

BMC Genomics 10:570.

19.   Johnson M, Zaretskaya I, Raytselis Y, Merezhuk Y, McGinnis S, Madden TL. 2008. NCBI

BLAST: a better web interface. Nucleic Acids Res 36:W5–W9.

20.   Wu M, Scott AJ. 2012. Phylogenomic analysis of bacterial and archaeal sequences with

AMPHORA2. Bioinformatics 28:1033–1034.

21.   Edgar RC. 2004. MUSCLE: Multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res 32:1792–1797.

22.   Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:

Improvements in performance and usability. Mol Biol Evol 30:772–780.

23.   Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ. 2015. IQ-TREE: A fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol

32:268–274.

24.   Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular evolutionary genetics analysis

version 7.0 for bigger datasets. Mol Biol Evol 33:1870–1874.

Figure 1: Phylogenomic tree showing occurrence of RuBisCO (CbbM/CbbL), ATP citrate lyase (AclAB), 4-subunit 2-oxoglutarate:ferredoxin oxidoreductase (KorABCD), putative thiol:fumarate reductase (TfrAB) and 2-subunit 2-oxoglutarate:ferredoxin oxidoreductase (KorAB) in the genomes of tubeworm symbionts (green), purple sulfur bacteria (purple) and other related bacteria (58 organisms total, alignment of 2526 amino-acid sites from 23 single-copy markers). The maximum likelihood tree was built with IQ-tree using the LG+R6 model of substitution (20). The tree is unrooted although outgroup "thiotrophic symbionts of bathymodiolin mussels and clams" is drawn at root. Branch labels are SH-aLRT support (%) / ultrafast bootstrap support (%). Accession numbers are provided in Supplementary Table 2.
* Was not included in the tree due to several missing single-copy marker genes or multiple versions of these genes, making an accurate phylogenomic placement challenging.
** Only the *aclB* gene was present.

Figure 2: The rTCA cycle gene clusters in symbiotic and free-living bacteria, and the respective transcriptomic gene expression levels in the symbionts of *Escarpia laminata* tubeworm (*aclA*, log (TPM)=3.6; *korA*, log (TPM)=3.3; *hdrA*, log (TPM)=2.9; for comparison - *atpB*, log (TPM)=2.0; *cbbM*, log (TPM)=5.0. TPM, transcripts per kilobase million. *rbr*, rubrerythrin. *dsr**, oxidoreductase related to the NADPH-dependent glutamate synthase small chain, clustered with sulfite reductase. The dotted line is the median expression value for *E. laminata* genes.