Genetic evidence for two carbon fixation pathways in symbiotic and free-living bacteria: The
Calvin-Bassham cycle and the reverse tricarboxylic acid cycle

Maxim Rubin-Blum[1,2*], Nicole Dubilier[1,3], Manuel Kleiner[4*]

[1]Max-Planck Institute for Marine Microbiology, Celsiusstrasse 1, 28359 Bremen, Germany

[2]Israel Limnology and Oceanography Research, Tel Shikmona, 3108000, Haifa, Israel

[3]MARUM, University of Bremen, 28359 Bremen, Germany

[4]Department of Plant & Microbial Biology, North Carolina State University, Raleigh, NC, USA

*Corresponding authors

mrubin@ocean.org.il

manuel_kleiner@ncsu.edu

Abstract: Sulfide-oxidizing bacterial endosymbionts of marine tubeworms such as *Riftia pachyptila* are able to fix carbon via both the reverse tricarboxylic acid (rTCA) and the Calvin-Benson-Bassham (CBB) cycles. Until recently, this co-existence of two carbon fixation pathways was thought to be unique to tubeworm symbionts. However, it was not clear if these two pathways were encoded in the same symbiont individual, or if two symbiont populations, each with one of the pathways, co-existed within tubeworms. We show here that both the CBB and rTCA pathways are likely encoded in the genome of the sulfur-oxidizing symbiont of the tubeworm *Escarpia laminata* from deep-sea asphalt volcanoes in the Gulf of Mexico. Furthermore, with comparative genomics, we show that *Thioflavicoccus mobilis,* a cultured, free-living gammaproteobacterial sulfur oxidizer, also possesses the genes for both carbon fixation pathways. Finally, we provide genomic and transcriptomic data suggesting a potential electron flow towards the rTCA cycle carboxylase 2-oxoglutarate:ferredoxin oxidoreductase, via a rare variant of NADH dehydrogenase/heterodisulfide reductase. This electron bifurcating complex, together with NAD(P)+ transhydrogenase and Na+ translocating Rnf membrane complexes may drive the rTCA cycle in both the symbiotic and the free-living sulfur oxidizer.

Importance: Primary production on Earth is dependent on autotrophic carbon fixation, which leads to the incorporation of carbon dioxide into biomass. Multiple metabolic pathways have been described for autotrophic carbon fixation, but all known autotrophic organisms were assumed to have the genes for only one of these pathways. The only possible exception to this paradigm was the sulfur-oxidizing symbiont of tubeworms, but this could have been an artifact of strain diversity within the symbiont population. Our results revealed that two pathways for carbon fixation co-occur in both symbiotic and free-living bacteria. Our finding of a cultivable bacterium with two carbon fixation pathways in its genome opens the possibility to study the potential benefits of having two pathways and the interplay between these cycles. This will deepen our understanding of carbon fixation pathways and could provide new avenues for optimizing carbon fixation in biotechnological applications.

Observation

Primary production by autotrophic organisms drives the global carbon cycle. Currently, seven naturally occurring pathways for inorganic carbon fixation are known in autotrophic organisms (1). The dominant carbon fixation pathway used by plants, algae, and many bacteria is the Calvin-Benson-Bassham (CBB) cycle. The six more efficient, alternative pathways are limited to autotrophic microbes that live in reducing habitats, due to the oxygen sensitivity of these alternative pathways (2, 3). All investigated autotrophic organisms have only one carbon fixation pathway. The only known exception are intracellular, sulfur-oxidizing symbionts of marine tubeworms such as *Riftia, Escarpia, Tevnia and Lamellibrachia* (the symbionts of these hosts are closely related to each other). These symbionts employ both the oxygen-tolerant CBB cycle and the oxygen-sensitive reverse tricarboxylic acid (rTCA) cycle. However, it was not clear if both pathways were encoded in the genome of a single symbiont individual, or if multiple strains with very similar genomes that encoded either one or the other pathway co-occurred within tubeworms (4–8). The CBB cycle in these symbionts is potentially more energy efficient than the classical version of the CBB cycle based on the replacement of the fructose-1,6-bisphosphatase with a pyrophosphate dependent enzyme (9, 10). In addition, it is likely that the interplay between the CBB and rTCA cycle under fluctuating redox conditions contributes to the high efficiency of carbon fixation in these symbioses (4, 5, 11), and consequently to the extremely high growth rates of tubeworms, which grow faster than any other known invertebrate (12).

Given that tubeworm symbionts could not yet be cultivated, it was not possible to investigate the co-occurrence of two carbon fixation cycles in detail to better understand the biochemical and physiological mechanisms that enable the interplay between these two pathways. In this study, we sequenced the genome and transcriptome of the symbiont from the tubeworm *Escarpia laminata* and compared its genome to those of other tubeworm symbionts and free-living microbes. These comparisons led us to discover the presence of co-occurring CBB and rTCA cycles in the genome of a cultured bacterium.

Co-occurrence of rTCA cycle genes with RuBisCo in symbiotic and free-living gammaproteobacteria. Genes for enzymes that are specific to the rTCA pathway, that is the ATP citrate lyase (*aclAB* genes), 2-oxoglutarate:ferredoxin oxidoreductase (OGOR, *korABCD* genes), and a putative fumarate reductase (*tfrAB* genes, homologs of genes encoding a thiol:fumarate reductase from *Methanobacterium thermoautotrophicum*, (13)), were assumed to occur in only a few symbiotic Gammaproteobacteria. We discovered, using comparative genomics, that these rTCA cycle enzymes also occur in some Chromatiaceae, including the cultivated sulfur oxidizer *Thioflacicoccus mobilis* and a gammaproteobacterial genome from an environmental metagenome(14), (**Fig. 1**). The type II ATP citrate lyases of tubeworm symbionts and *T. mobilis* were likely acquired via horizontal gene transfer from other bacterial clades (**Suppl. Fig. S1**), (6). These gammaproteobacteria also encode either Form I or II RuBisCO, or both (**Suppl. Note 1**).

Presence of the rTCA and the CBB pathways in the genome of a single bacterium: Due to the fragmented nature of the previously available genomes of tubeworm symbionts, past studies could not determine whether the genes for both pathways are present in a single genome or if the two pathways are distributed in a strain-specific manner, i.e. only one of the two pathways is present in the genome of a single cell (3). Here, we provide two lines of evidence that both pathways can co-occur in the genome of a single organism. First, sequencing coverage for the genes of both pathways in the *E. laminata* symbiont was similar to that of single-copy marker genes (**Suppl. Table 1**). Since genes that are strain specific are expected to have lower coverage than the rest of the genome (15), the similar coverage of genes encoding the two pathways and single-copy genes suggests that in the *E. laminata* symbiont both pathways are present in all cells. Second, in the closed genome of the cultured *T. mobilis*, both the genes encoding the rTCA and the CBB cycle co-occur, providing evidence that these genes co-exist in a single genome.

Our transcriptomic analyses of *E. laminata* tubeworm symbionts revealed high expression levels of both the rTCA and the CBB cycle genes (Fig. 2). This observation is consistent with previous proteomic analyses of the *Riftia* symbiont (4, 5). The high expression levels of genes from the

rTCA and the CBB cycle suggests that both pathways play an important metabolic role in these

symbionts. It is, however, not clear whether these cycles function simultaneously within single

symbiont cells, or are differentially expressed within the symbiont population (3).

The rTCA gene clusters are conserved among the tubeworm symbionts and some

Chromatiaceae bacteria. In the tubeworm symbionts, the cultivated *T. mobilis* and the

gammaproteobacterial genome from an environmental metagenome, there was a considerable

level of conservation of the rTCA gene clusters, at the sequence and synteny levels (**Fig. 2**). The

aclAB genes that encode the two subunits of the ATP citrate lyase were accompanied by those

that encode bidirectional TCA cycle enzymes, including *acn* (aconitase), *idh* (isocitrate

dehydrogenase), and *mdh* (malate dehydrogenase). The other rTCA specific genes *korABCD*

(four-subunit OGOR) and *tfrAB* (putative thiol:fumarate reductase), were also present in the

rTCA gene cluster. Similar to the ATP citrate lyase, the four-subunit OGOR, as well as the thiol:

fumarate reductase are very rare among Gammaproteobacteria, and were probably acquired via a

single horizontal gene transfer event from a distant bacterial clade (**Fig. 1**, **Suppl. Fig. S2 and**

S3). A dimeric OGOR (*korAB* genes), more common than the four-subunit enzyme among

gammaproteobacterial autotrophs, yet absent in *T. mobilis,* was located elsewhere in the genome

of the *E. laminata* symbiont. The *korAB* genes were co-localized with genes that encode

additional rTCA cycle enzymes (**Suppl. Note 2, Suppl. Fig. S4**).

An array of genes that encode several electron translocating complexes were integrated into the

rTCA cycle gene clusters. These complexes included an electron-bifurcating NADH

dehydrogenase/heterodisulfide reductase complex (*flxABCD-hdrABC* genes, **Suppl. Note 3**), a

NAD(P)+ transhydrogenase and Na+ translocating Rnf membrane complex (*pntAB* and

rnfABCDGE genes, **Suppl. Note 4**). Most interestingly, the conserved interspersing of the

korABCD and *tfrAB* genes with the *flxABCD-hdrABC* genes hints at the possibility that these

proteins form a complex that efficiently shuttles electrons directly to the OGOR and the thiol:

fumarate reductase (**Suppl. Figure S5**).

Conclusions. Until now, the only bacteria known to possess two carbon fixation pathways were

sulfur-oxidizing, tubeworm symbionts, which are currently not amenable to cultivation-based

studies. With the discovery of the co-existence of the CBB and rTCA cycle in the cultivable *T.*

mobilis, experimental studies are now feasible. Such studies would reveal if these pathways are

expressed under different physicochemical conditions, and potentially allow the biotechnological

optimization of efficiency and yield in production processes that rely on autotrophic carbon

fixers. To our knowledge, the use of organisms with multiple carbon fixation pathways has not

been used as a design principle for these applications.

Methods: Comparative genomics and transcriptomics. Publically available genomes from

NCBI and JGI-IMG collections, as well as de-novo assembled genomes of *Escarpia laminata*

symbionts (estimated completeness 99.5%), were used for genomic comparison (see

Supplementary Methods). To verify presence/absence of target gene homologs in sequenced

organisms we used NCBI's BLAST against the nucleotide collection and non-redundant protein

database (16). *E. laminata* symbiont genomes were used as a template for genome-centered

transcriptomics (sequences available under the BioProject accession number PRJNA471406).

Phylogenetic and phylogenomic analyses. Phylogenomic treeing was performed using scripts

available at phylogenomics-tools (DOI:10.5281/zenodo.46122). Twenty-three marker proteins

that are universally conserved across the bacterial domain were extracted from genomes using

the AMPHORA2 pipeline (17). Twenty-three single-copy markers were used for alignment with

MUSCLE (18). The marker alignments were concatenated into a single partitioned alignment,

and poorly aligned regions were removed. Functional protein sequences were aligned with

MAFFT (19). Maximum Likelihood trees were calculated with IQ-tree (20) and MEGA7 (21),

using the best-fitting model.

Author Contributions

MRB, ND and MK conceived the study. MRB and MK analyzed the samples. MRB, ND

and MK wrote the manuscript.

Acknowledgments

The authors thank all individuals who helped during the R/V Meteor research cruise

M114, including onboard technical and scientific personnel, the captain and crew, and the ROV

MARUM-Quest team. We thank the Max Planck-Genome-Centre Cologne

(http://mpgc.mpipz.mpg.de/home/) for generating the metagenomic and the metatranscriptomic

data used in this study. We thank Julie Reveillaud for providing genomes of Mid-Cayman Rise

tubeworm symbionts. The Campeche Knoll cruise was funded by the German Research

Foundation (DFG – Deutsche Forschungsgemeinschaft). We are grateful to the Mexican

authorities for granting permission to conduct this research in the southern Gulf of Mexico

(permission of DGOPA: 02540/14 from 5 November 2014). This study was funded by the Max

Planck Society, the MARUM DFG-Research Center / Excellence Cluster "The Ocean in the Earth

System" at the University of Bremen, an ERC Advanced Grant (BathyBiome, 340535), a Gordon

and Betty Moore Foundation Marine Microbial Initiative Investigator Award to ND (Grant

GBMF3811) and the NC State Chancellor's Faculty Excellence Program Cluster on Microbiomes

and Complex Microbial Communities (MK).

Conflict of interest

The authors declare no conflict of interest.

References

1. Figueroa IA, Barnum TP, Somasekhar PY, Carlström CI, Engelbrektson AL, Coates JD.

2018. Metagenomics-guided analysis of microbial chemolithoautotrophic phosphite

oxidation yields evidence of a seventh natural CO_2 fixation pathway. Proc Natl Acad Sci

115:E92–E101.

2. Erb TJ. 2011. Carboxylases in natural and synthetic microbial pathways. Appl Environ

Microbiol 77:8466–8477.

3. Hügler M, Sievert SM. 2011. Beyond the Calvin cycle: autotrophic carbon fixation in the

ocean. Ann Rev Mar Sci 3:261–289.

4. Gardebrecht A, Markert S, Sievert SM, Felbeck H, Thürmer A, Albrecht D, Wollherr A,

Kabisch J, Le Bris N, Lehmann R, Daniel R, Liesegang H, Hecker M, Schweder T. 2012.

Physiological homogeneity among the endosymbionts of *Riftia pachyptila* and *Tevnia*

jerichonana revealed by proteogenomics. ISME J 6:766–76.

5. Markert S, Arndt C, Felbeck H, Becher D, Sievert SM, Hugler M, Albrecht D, Robidart J,

Bench S, Feldman RA, Hecker M, Schweder T. 2007. Physiological Proteomics of the

Uncultured Endosymbiont of *Riftia pachyptila*. Science 315:247–250.

6. Markert S, Gardebrecht A, Felbeck H, Sievert SM, Klose J, Becher D, Albrecht D, Thürmer

A, Daniel R, Kleiner M, Hecker M, Schweder T. 2011. Status quo in physiological

proteomics of the uncultured *Riftia pachyptila* endosymbiont. Proteomics 11:3106–3117.

7.   Robidart JC, Bench SR, Feldman RA, Novoradovsky A, Podell SB, Gaasterland T, Allen

EE, Felbeck H. 2008. Metabolic versatility of the *Riftia pachyptila* endosymbiont revealed

through metagenomics. Environ Microbiol 10:727–737.

8.   Reveillaud J, Anderson R, Reves-Sohn S, Cavanaugh C, Huber JA. 2018. Metagenomic

investigation of vestimentiferan tubeworm endosymbionts from Mid-Cayman Rise reveals

new insights into metabolism and diversity. Microbiome 6:19.

9.   Kleiner M, Wentrup C, Lott C, Teeling H, Wetzel S, Young J, Chang Y-JY-J, Shah M,

VerBerkmoes NC, Zarzycki J, Fuchs G, Markert S, Hempel K, Voigt B, Becher D, Liebeke

M, Lalk M, Albrecht D, Hecker M, Schweder T, Dubilier N. 2012. Metaproteomics of a

gutless marine worm and its symbiotic microbial community reveal unusual pathways for

carbon and energy use. Proc Natl Acad Sci USA 109:E1173-1182.

10.   Kleiner M, Petersen JM, Dubilier N. 2012. Convergent and divergent evolution of

metabolism in sulfur-oxidizing symbionts and the role of horizontal gene transfer. Curr

Opin Microbiol 15:621–631.

11.   Klatt JM, Polerecky L. 2015. Assessment of the stoichiometry and efficiency of CO_2

fixation coupled to reduced sulfur oxidation. Front Microbiol 6.

12.   Bright M, Klose J, Nussbaumer AD. 2013. Giant tubeworms. Curr Biol 23:R224–R225.

13.   Heim S, Kunkel A, Thauer RK, Hedderich R. 1998. Thiol : fumarate reductase (Tfr) from

196   *Methanobacterium thermoautotrophicum.* Identification of the catalytic sites for fumarate

197   reduction and thiol oxidation. Eur J Biochem 253:292–299.

198   14.   Anantharaman K, Brown CT, Hug LA, Sharon I, Castelle CJ, Probst AJ, Thomas BC,

199   Singh A, Wilkins MJ, Karaoz U, Brodie EL, Williams KH, Hubbard SS, Banfield JF. 2016.

200   Thousands of microbial genomes shed light on interconnected biogeochemical processes

201   in an aquifer system. Nat Commun 7:13219.

202   15.   Pasić L, Rodriguez-Mueller B, Martin-Cuadrado A-B, Mira A, Rohwer F, Rodriguez-

203   Valera F. 2009. Metagenomic islands of hyperhalophiles: the case of *Salinibacter ruber.*

204   BMC Genomics 10:570.

205   16.   Johnson M, Zaretskaya I, Raytselis Y, Merezhuk Y, McGinnis S, Madden TL. 2008. NCBI

206   BLAST: a better web interface. Nucleic Acids Res 36:W5–W9.

207   17.   Wu M, Scott AJ. 2012. Phylogenomic analysis of bacterial and archaeal sequences with

208   AMPHORA2. Bioinformatics 28:1033–1034.

209   18.   Edgar RC. 2004. MUSCLE: Multiple sequence alignment with high accuracy and high

210   throughput. Nucleic Acids Res 32:1792–1797.

211   19.   Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:

212   Improvements in performance and usability. Mol Biol Evol 30:772–780.

213   20.   Nguyen LT, Schmidt HA, Von Haeseler A, Minh BQ. 2015. IQ-TREE: A fast and effective

214   stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol

215   32:268–274.

216   21.   Kumar S, Stecher G, Tamura K. 2016. MEGA7: Molecular evolutionary genetics analysis

217   version 7.0 for bigger datasets. Mol Biol Evol 33:1870–1874.

Figures

Figure 1: Phylogenomic tree showing occurrence of RuBisCO (CbbM/CbbL), ATP citrate lyase (AclAB), 4-subunit 2-oxoglutarate:ferredoxin oxidoreductase (KorABCD), putative thiol:fumarate reductase (TfrAB) and 2-subunit 2-oxoglutarate:ferredoxin oxidoreductase (KorAB) in the genomes of tubeworm symbionts (green), purple sulfur bacteria (purple) and other related bacteria (58 organisms total, alignment of 2526 amino-acid sites from 23 single-copy markers). The maximum likelihood tree was built with IQ-tree using the LG+R6 model of substitution (20). The tree is unrooted although outgroup "thiotrophic symbionts of bathymodiolin mussels and clams" is drawn at root. Branch labels are SH-aLRT support (%) / ultrafast bootstrap support (%). Accession numbers are provided in Supplementary Table 2.

Figure 2: The rTCA cycle gene clusters in symbiotic and free-living bacteria, and the respective transcriptomic gene expression levels in the symbionts of *Escarpia laminata* tubeworm (*aclA*, log (TPM)=3.6; *korA*, log (TPM)=3.3; *hdrA*, log (TPM)=2.9; for comparison - *atpB*, log (TPM)=2.0); *cbbM*, log (TPM)=5.0. TPM, transcripts per kilobase million.