New tools for diet analysis: nanopore sequencing of metagenomic DNA from rat

stomach contents to quantify diet

Nikki E. Freed, William S. Pearman, Adam N. H. Smith, Georgia Breckell, James Dale,

Olin K. Silander

Addresses of all authors: Institute of Natural and Mathematical Sciences, Massey

University, Auckland 0745, New Zealand

4. Corresponding authors: Olin K. Silander, Institute of Natural and Mathematical

Sciences, Massey University, Auckland 0745, New Zealand, olinsilander@gmail.com,

+64 9 213 6618; Nikki E. Freed, Institute of Natural and Mathematical Sciences, Massey

University, Auckland 0745, New Zealand, freednikki@gmail.com, +64 9 213 6639

Abstract

Background: Using metagenomics to determine animal diet offers a new and promising

alternative to current methods. Here we show that rapid and inexpensive diet

quantification is possible through metagenomic sequencing with the portable Oxford

Nanopore Technologies (ONT) MinION. Using an amplification-free approach, we

profiled the stomach contents from wild-caught rats.

Results: We conservatively identified diet items from over 50 taxonomic orders, ranging

across nine phyla that include plants, vertebrates, invertebrates, and fungi. This

highlights the wide range of taxa that can be identified using this simple approach. We

calibrate the accuracy of this method by comparing the characteristics of reads

matching the ground-truth host genome (rat) to those matching diet items, and show

that at the family-level, false positive taxon assignments are approximately 97.5%

accurate. We also suggest a way to mitigate for database biases in metagenomic

approaches. Finally, we implement a constrained ordination analysis and show that we

can identify the sampling location of an individual rat within tens of kilometres based on

diet content alone.

Conclusions: This work establishes proof-of-principle for long-read metagenomic

methods in quantitative diet analysis. We show that diet content can be quantified even

with limited expertise, using a simple, amplification free workflow and a relatively

inexpensive and accessible next generation sequencing method. Continued increases

in the accuracy and throughput of ONT sequencing, along with improved genomic

databases, suggests that a metagenomic approach to quantification of animal diets will

become an important method in the future.

Keywords: Metagenomics, nanopore, diet analysis, rat, next generation sequencing

Background

Accurate quantification of animal diets yields critical insight into ecosystem and food web dynamics. However, unbiased and sensitive assessment of diet content is difficult to achieve. This is largely due to the limited accuracy of many current methods. Such methods include visual inspection of gut contents (1,2), which presents bias against items are most easily degraded (for example, soft-bodied species); stable isotope analysis (3,4), which yields only broad information on diet, such as whether diet items are terrestrial or marine in origin (5,6); and time-lapse video (7,8), for which species identification is difficult for small prey items or in low-light conditions.

To circumvent these issues, DNA-based methods (9,10) have become popular. Perhaps the most widely applied DNA-based method is metabarcoding. This approach relies on PCR amplification and sequencing of conserved regions from nuclear, mitochondrial, or plastid genomes (9). With adequate primer selection, this method can detect a wide range of species, and does not require specific expertise, which is often necessary for other methods.

However, DNA metabarcoding is not free from bias. PCR primers must be specifically tailored to particular sets of taxa or species (11). Although "universal" PCR primer pairs have been developed (for example targeting all bilaterians or even all eukaryotes (12), all primer sets exhibit bias towards certain taxa. Five-fold differences in fungal operational taxonomic units (OTU) estimates have been found when using different sets of fungal-specific PCR primer pairs (13). It has also been shown that published

universal primer pairs were capable of amplifying only between 57% and 91% of

tested metazoan species, with as few as 33% of species in some phyla being amplified

at all (e.g. cnidarians)(14). Different genomic loci from the same species can exhibit up

to 2,000-fold differences in DNA concentration as inferred using qPCR (15). The choice

of polymerase can also bias diversity metrics when using metabarcoding (16). For

these reasons, an approach that circumvents PCR and thus avoids these biases is

desirable.

Metagenomic sequencing aims to directly sequence all of the DNA in a sample without

introducing bias. Although there are still biases with this approach, for example due to

nucleotide content affecting the likelihood of a molecule being sequenced, these are

inherently less than those introduced by metabarcoding. Metagenomic approaches

have most frequently been used to yield insights into microbial diversity and function

(17–24). Metagenomic applications aimed at eukaryotic taxa identification are less

common. Several metagenomic diet studies have implemented filtering steps to select

only mitogenomic sequence or metabarcode regions, or have used abridged

databases before data analysis in order to mitigate database biases (25–27). However,

to our knowledge, very few studies have used unfiltered metagenomic sequence

analysis to infer diet (30,31).

Here, we establish a proof-of-principle methodology to accurately classify

metagenomic sequences from eukaryotic taxa and determine diet content using low-

accuracy, long-read sequencing, Oxford Nanopore (ONT).

We quantified rat diets from several locations in the North Island of New Zealand using

stomach samples. Using these samples and methodology provides three distinct

advantages.

First, rats are extremely omnivorous. As such, they serve as an excellent means to

quantify the breadth of taxa that can be detected using a metagenomic long read

approach.

Second, the use of stomach samples means that a significant number of reads will be

host reads. This allows us to assess the characteristics of true positive sequence reads

(rat-derived reads that match rat database sequences), as well as false positive reads

(rat-derived reads that match non-rat database sequences). We can then determine

whether reads matching diet items have similar characteristics to known true positive

reads. This use of host reads is exactly analogous to feeding the rats a diet of known

content (i.e. rat) and testing whether the contents of the known diet can be accurately

identified using this method.

Third, quantifying rat diets has important ecological implications. It is well-established

that the relatively recent introduction of mammalian predators to New Zealand has had

significant negative effects on many of the native animal populations. This ranges from

insects (34), to reptiles (35), to molluscs (36), to birds (37,38), with downstream effects

on terrestrial and aquatic ecosystems (39). To counteract the effects of mammalian

predators, an ambitious plan is currently being put into place that aims for the

eradication of all mammalian predators from New Zealand (including possums, rats,

stoats, and hedgehogs), by 2050 (http://www.doc.govt.nz/predator-free-2050; (40). A

useful step toward this goal would be to prioritise the management of predators and

establish in which locations native species experience the highest levels of predation.

To do so requires establishing the diet content of local mammalian predators.

Importantly, using DNA sequencing methods to profile diet does not require high depth

to accurately profile the breadth of diet items consumed by an animal. Here we aim to

quantify all diet items present in the diet at 1% or more. At this fraction, if we assume

that read counts are Poisson distributed, with only 2,000 reads, 99% of the time we will

quantify such items within 2-fold of their true amount. Thus, diet quantification

presents a clear example of a situation in which sequencing depth is not a critical

factor.

Results
DNA sequencing and assignment of reads to taxa

We selected eight rats from each of three locations near Auckland, New Zealand for diet

quantification. Each location comprised a different type of habitat: undisturbed inland

native forest (Waitakere Regional Parklands, WP); native bush surrounding an estuary

(Okura Bush Walkway, OB); and restored coastal wetland (Long Bay Regional Park,

LB). We isolated DNA from whole stomach contents from each rat (see Methods). We

obtained DNA from entire homogenised stomach samples. We sequenced these DNA

samples using the Oxford Nanopore ligation sequencing kit (LSK-108) and native

barcoding kit to multiplex the samples within a sequencing run. We performed two

separate runs, obtaining a total of 82,977 reads (January 2017) and 96,150 reads

(March 2017) from each. These numbers are not far below expectations given the flow

cell and kit chemistry and MinKNOW software versions available at that time (47).

However, these read numbers are considerably below those expected for current ONT

flow cells and software, which has improved per flow cell output more than 100-fold

above these numbers.

We found large variation in the numbers of reads per barcode: approximately 10-fold for

the January samples, and up to 40-fold in March (**Fig. 1A** and **1B**). We hypothesise that

this is due to the highly variable quality of DNA in each sample. This did not appear to

have strong effects on read accuracy, as the median phred quality scores per read

ranged from 7-12 (0.80 - 0.94 accuracy) for both runs.

The degradation of the DNA during digestion in the stomach, as well as fragmentation

during DNA isolation (48) and sequencing library preparation led to relatively short

median read lengths of 606 bp and 527 bp for the January and March datasets,

respectively (**Fig. 1C**). Within runs, read length and quality were similar. Although these

median read lengths are considerably shorter than other nanopore sequencing results

from both our and others work (49), we found wide variation in length, with almost 10%

of all reads being longer than 1200 bp. Notably, reads of this length can allow more

precise taxonomic identification than accurate short reads (32).

To quantify diet contents we first BLASTed all sequences against a combined database

of the NCBI nt database (the partially non-redundant nucleotide sequences from all

traditional divisions of GenBank excluding genome survey sequence, EST, high-

throughput genome, and whole genome shotgun

(ftp://ftp.ncbi.nlm.nih.gov/blast/db/README)) and the NCBI other_genomic database

(RefSeq chromosome records for non-human organisms

(ftp://ftp.ncbi.nlm.nih.gov/blast/db/README)). We used BLAST as it is generally viewed

as the gold standard method in metagenomic analyses (50). Of the 133,022 barcoded

reads, 30,535 (23%) hit a sequence in the combined nt and other_genomic database at

an e-value cut-off of 1e-2.

Fig. 1. Results of nanopore metagenomic sequencing of rat stomach contents. (A)
and (B) Barcode distributions for January and March runs, respectively. We
multiplexed the samples on the flow cells, using 12 barcodes per flow cell. The
distribution of read numbers across barcodes varied by up to 40-fold. 20% (January)
and 30% (March) of all reads could not be assigned to a barcode ("None"). The inability
to assign these reads to a barcode is due primarily to their lower quality. **(C) Read**
length distribution for January and March nanopore runs. 90% of the reads were
between 350 bp and 1,580 bp in length, with only 0.55% being longer than 4,000 bp.

We first aimed to assess the quality of these hits. We found a bimodal distribution of

alignment lengths and e-values (**Fig. 2A**). We also noticed that mean read quality had

substantial effects on the likelihood of a read yielding a BLAST hit, with almost 40% of

high accuracy reads having hits in the March dataset, as compared to 1% of low

accuracy reads (**Fig. 2B**). We hypothesized that many of the short alignments with high

e-values were false positives. We thus first filtered this hit set, only retaining BLAST hits

with e-values less than 1e-20 and alignments greater than 100 bp. To give some

intuition for this length cut-off, a 100bp read with five 1 bp indels and seven mismatches

(88% identity) would have an identical raw score and e-value (given the default match,

mismatch, and gap parameters) to a single-end 70 bp Illumina read with a 100%

identical database match. Similar or quality filters based on length and identity have

been imposed previously (30). A total of 22,154 hits passed this e-value filter.

Fig. 2. BLAST hits of metagenomic reads. (a) Alignment lengths and e-values were
bimodally distributed. The y-axis is plotted on a log scale, with zero e-values
suppressed by adding a small number (1e-190) to each e-value. The horizontal red
dotted line indicates the e-value cut-off we implemented and the vertical red dotted line
indicates the length cut-off (e-value < 1e-20 and alignment length of 100, respectively)
to decrease false positive hits. **(b) The fraction of reads with high quality BLAST hits
(e-value < 1e-20) increased as a function of read accuracy.** We binned the data
according to mean read accuracy (bin width = 0.02) and calculated the fraction of reads
within each bin that have a high quality BLAST hit (alignment length greater than 100bp
and e-value less than 1e-20) for the January and March runs separately (blue and
orange points, respectively). The number of reads in each bin is indicated above each
point (in thousands). There is a clear positive correlation between mean accuracy and
the likelihood of a high quality BLAST hit, reaching almost 40% for high accuracy reads
(>92.5%) for the March dataset.

We next used MEGAN6 (41) to assign reads to specific taxa. MEGAN6 employs an

LCA algorithm to assign reads to a taxon. For example, if a read has BLAST hits to five

different species, three of which have bit scores within 20% of the best hit, the read will

be assigned to the genus, family, order, or higher taxon level that is the LCA of those

best-hit three species (51). If a read matches one species far better than any other, by

definition, the LCA is that species.

16,820 reads (76%) were assigned to a taxon by MEGAN. Of these, 31% were

assigned by MEGAN as being bacterial, and 55% of these were *Lactobacillus spp*.

These results match previous studies on rat stomach microbiomes, which have found

lactobacilli to be the dominant taxa (52–55). Plant-associated *Pseudomonas* and

Lactococcus taxa were also common, at 7% and 6%, respectively.

MEGAN assigned reads to a wide range of eukaryotic taxa. To conservatively infer

taxon presence, we first reclassified MEGAN species-level assignments to the level of

genus. Even so, many clear false positive assignments remained (e.g. hippo and naked

mole rat). These matches were generally short and of low identity. To reduce such false

positive taxon inferences, we used information from reads assigned to the genera

Rattus (rat) and *Mus* (mouse).

We inferred that the reads assigned to *Rattus* (2,696 reads in total) were true positive

genus-level assignments and that the reads assigned to *Mus* (2,798 reads in total) were

false positive genus-level assignments (and not true positive *Mus*-derived reads). If

these were true positive *Mus* reads, then they would be due to mouse predation by

some of the rats. Although rats are known to prey on mice (56), if this had occurred, we

would expect that (1) if a rat predated a mouse the ratio of mouse to rat reads would be

higher than in rats that had not predated mice; (2) in those same rats, the percent

identity of the reads assigned to *Mus* would be higher than in rats that had not predated

mice. However, we found that the ratio of mouse to rat reads and percent identity was

similar for all rats. This suggested that these *Mus* hits were indeed false positives and

could thus be used to determine the rate of false positive genus-level assignment.

The mean percent identity values of the best BLAST hits for *Rattus* and *Mus* reads

differed substantially, with *Rattus* reads having a median identity of 86.4%, and *Mus*

81.0% (**Fig. 3A**). The mean percent identity for *Rattus* reads corresponds very well to

that expected given the mean quality scores of the reads (86.4% identity corresponds

to a mean quality score of 8.7, similar to what we observed; **Fig. S1A-C**).

To further investigate alignment characteristics differentiating true positive (*Rattus*) and

false positive (*Mus*) assignments, we considered the ratio of alignment length to read

length. If the full read aligns, this ratio is one. More generally, higher quality alignments

should have higher ratios. Indeed, we found this ratio differed substantially between

the *Rattus*- and *Mus*-assigned reads: the median ratio of alignment length to read

length was 0.57 for *Rattus* and 0.52 for *Mus* (**Fig. 3B**).

We used this analysis to select cut-off values for assigning reads as having true

positive or false positive taxon assignments. For genus-level assignments, we required

at least 82.5% alignment identity, and an alignment length to read length ratio of at

least 0.55. These cut-offs exclude 88% of the reads falsely assigned to *Mus*, instead

correctly assigning them one taxon level higher, to the Family *Muridae*.

However, this rate of false positive assignments is still relatively high. We thus also

quantified the rate of false positive assignment at the level of family. We first identified

all reads classified as being from the order *Rodentia.* We then implemented specific

cut-offs, requiring reads and database matches to have 77.5% identity, an alignment

length to read length ratio of at least 0.1, and a total alignment length of at least 150

bp. With these cut-offs, 97.3% of all reads assigned to the Order *Rodentia* were

classified in the likely true positive Family *Muridae* (which contains the genus *Rattus*).

The remaining 2.7% were assigned to the family *Cricetidae* (voles and lemmings),

except or four reads assigned to *Spalacidae* (mole-rats) (**Fig. S2**). These are clear false

positive assignments, as it is highly unlikely that these families were predated.

However, these results establish that by implementing specific cut-off values for

Fig. 3. Distributions of percent identity and length for alignments of reads
matching *Rattus* (rat), *Mus* (mouse), and diet items. (a) The percent identity for
alignments of rat (*Rattus*) and diet items is much higher than for mouse (*Mus*).
Histograms of the percent identity of the alignment of the top BLAST hit with the read.
Mus matches have substantially lower percent identity compared to both *Rattus* and diet
items. The dotted lines indicate the cut-offs that we implemented for inferring reads as
belonging to a specific genus (above 82.5% identity) or family (above 77.5% identity).
(b) Ratios of alignment lengths to read lengths of rat (*Rattus*) and diet items are

higher than for mouse (*Mus*). This plot is analogous to that in (a). The dotted line
indicates the cut-off that we implemented for inferring reads as belonging to a specific
genus (above 0.55).

alignments, we can ensure a low rate of false positive assignments at the family level,

especially for taxa with genomic sequence well-represented in the database.

However, it is possible that for taxa that are not well-represented in databases, read

classification will be less accurate. If this occurred, we would expect that reads would

frequently be assigned to the wrong taxon, resulting in lower percent identities and low

alignment lengths for those reads. We thus checked whether diet items had percent

identities and alignment lengths similar to the true positive *Rattus* alignments, or

instead whether they were more similar to the false positive *Mus* alignments. We found

that the majority of diet items had alignment percent identities that overlapped with the

Rattus reads. Furthermore, the alignment length to read length ratios often exceed

those for the *Rattus* reads. This suggests that the diet taxa assignments are often

correct down to the level of genus (as the *Rattus*-assigned reads are correct to the

level of genus) and are not false positive assignments. However, here we

conservatively report diet items at the level of family.

For reads that did not pass the above cut-offs, we placed taxon assignments at the

level of order, or used the taxon level assigned by MEGAN. Using these cut-offs, 16%

of all reads were classified at the genus level; 71% were classified at the family-level or

below; 89% were classified at the order-level or below; and 98% were classified at the

phylum-level or below.

Beyond the small fraction of false positive family assignments within *Rodentia*

(discussed above), a small fraction of clear false positives remained in other orders.

Again, most had short alignments (**Table S4**). The exception to this were three reads

from two rats matching *Buthidae* (scorpions), which had alignment lengths of 762 bp,

664 bp, and 298 bp with identities of 83%, 88%, and 79%, respectively. It is unlikely

these are true positives, and instead we hypothesise that these rats predated

harvestmen (*Opiliones*), a closely related sister taxon within *Arachnida*, but lacking

significant amounts of genomic data. Despite the presence of these false positive taxa,

we did not further increase the stringency of our filters, suggesting that we have a small

rate of false positive inference at the family level (2.7% in the case of *Muridae*), and

most likely similar rates for other family assignments.

Identification of diet

Within each rat, we identified a wide variety of plant, animal, and fungal orders, ranging

from two to 25 orders per rat (mean 8.7; **Fig. 4**). In total, we identified taxa from 68

different Families, 55 different Orders, 15 different Classes, and eight different Phyla

(**Fig. 5**). In this analysis, we consider all non-host, non-bacterial reads as potential diet

items, although not all are necessarily so. Plants were the primary diet item, with the

largest fraction of rats consuming four predominant orders: *Poales* (grasses), *Fabales*

(legumes), *Arecales* (palms), and *Araucariales* (podocarps). The dominance of plant

matter (fruits and seeds) in rat diets has been established previously (57,58). Animal

taxa made up a smaller component of each rat's diet, with *Insecta* dominating:

Hymenoptera (bees, wasps, and ants), *Coleoptera* (beetles), *Lepidoptera* (moths and

butterflies), *Blattodea* (cockroaches), *Diptera* (flies), and *Phasmatodea* (stick insects). In

addition, *Stylommatophora* (slugs and snails) were present in substantial numbers (**Fig.**

5A and **5B**). Fungi were only a small component of the rats' diet, although several

orders were present: *Sclerotiniales* (plant pathogens)*, Saccharomycetales* (budding

yeasts)*, Mucorales* (pin molds)*, Russulales* (brittlegills and milk-caps), and

Chytotheriales (black yeasts)*.* Finally, for many rats, a substantial proportion of the

stomach contents were parasitic worms (primarily *Spirurida* (nematodes) and

Hymenolepididae (tapeworms)).

Due to our metagenomic approach, the fraction of each element of the rats' diets is

distorted by biases in genomic databases: whole genome data exists for only a few

taxa, while mtDNA, rDNA, metabarcode loci, and microsatellite sequence data are

present in the database for many animal and plant genera. To quantify database-driven

bias, for each taxon we determined the fraction of hits that mapped to mtDNA, rDNA,

microsatellites, or EST libraries (we refer to this as *non-genomic*), versus the fraction of

hits that mapped to DNA sequences annotated as arising from genome sequencing

projects). We expect that for animals with sequenced genomes, these two fractions

should be approximately constant, largely determined by the relative amounts of mtDNA

and nuclear DNA in the cell. For animals without sequenced genomes, there should be

considerably more hits to mtDNA, plastid, rDNA, and microsatellites (non-genomic

sequence), and few (if any) genomic hits. By comparing these fractions for taxa with

complete genome sequences in the database to taxa without complete genomes we

aimed to assess the effects of this bias.

Fig. 4. Numbers of taxa in individual rats. Each boxplot indicates the range of families (left boxes) or orders (right boxes) consumed by each rat in each location (OB: Okura Bush, native bush; LB: Long Bay Park, restored wetland; WP: Waitakere Park, native forest). The numbers for individual rats (eight per location) are plotted in grey.

Fig. 5. Proportions of taxa in the diets of individual rats. (a) Reads assigned to taxa at the family and (b) order level. The rows correspond to a single rat, with the proportions of reads for that rat assigned to each family or order indicated in shades of blue and yellow. Reads that were not assigned to a specific family or order are indicated at the right side of the figure. The families and orders have been sorted so that the most common diet components appear on the left. Only the 55 most common families are shown. Note that the color gradations presented on the scale are not linear.

We first considered only animals with sequenced genomes in the database. For these, we found that the fraction of reads that mapped to non-genomic sequence (mtDNA, rDNA, microsatellite, plastid, and EST library) ranged from 0% (*Coturnix* and *Numida*) to 27% (*Rattus*) to 39% (*Gallus*) (**Fig. 6**). This is a considerable range, and we hypothesise that variation in the fraction of non-genomic reads is due to the type of tissue sequenced. For *Rattus* the sequenced tissue was primarily stomach muscle, which has a relatively high fraction of mtDNA, explaining the large fraction of reads mapping to mtDNA.

For plants with sequenced genomes, the fraction of reads matching non-genomic

sequence (mostly mtDNA, plastid, and rDNA) was generally lower: between 2%

(*Cenchrus*) and 12% (*Zea*). Thus, on average, for animals with sequenced genomes

present in the database, approximately 30% of all reads mapped to non-genomic

sequences; for plants, approximately 5% mapped to non-genomic sequences.

In contrast, for taxa with little or no genomic sequence in the database, the vast majority

of matches were non-genomic (mtDNA, plastid, rDNA, or microsatellite loci): 90% of

Phoenix (date palm) hits; all *Helix* (snail); and all *Rhaphidophorae* (cave weta) hits. All

Arthurdendyus (the endemic New Zealand flatworm) hits were solely to rDNA loci.

This overrepresentation of non-genomic sequence is in strong contrast to animals that

do have sequenced genomes. In that case we expect only 30% of all reads should map

to non-genomic sequence. Thus, these results indicate that for animal taxa with little or

no genomic sequence data, we have underestimated the actual number of sequences

from that taxon by approximately two- to three-fold. For plant taxa with little or no

genomic sequence data, we have likely underestimated read abundance by

approximately 20-fold.

Figure 6. Fractions of reads matching genomic and non-genomic sequence for the best BLAST hit of each read. For the species with complete genomes, the fraction of reads matching genomic sequence ranges from 40% (*Solanum*) to 100%. This large range is likely due to the tissue from which the DNA was isolated. For example, muscle tissue has a higher fraction of mtDNA to nuclear DNA than egg. For species without fully sequenced genomes, this fraction ranges from 0% to 20% (*Phoenix*, which has a small amount of genomic data present in the database).

After classification of the reads at the genus, family, and order level, further examination

of the data suggested that specific taxa were overrepresented in the diets of rats from

particular locations. For example, six out of eight rats from the native estuarine bush

habitat (OB) consumed *Arecaceae*, while only one in the restored wetland area (LB) did.

All three rats that consumed *Phaseanidae* were from the native estuarine habitat (OB).

All five rats that consumed *Solanales* were from the restored wetland area. These

372   patterns suggested that it might be possible to use diet components alone to pinpoint

373   the habitat from which each rat was sampled.

nMDS and CAP analysis by location

In order to determine if diet composition of the rats differed consistently between

locations, we first performed an unconstrained analysis using non-metric

multidimensional scaling (nMDS) on taxa assigned at the family level. The input for the

nMDS was the dissimilarity matrix (Bray-Curtis distance of diet at the family level).

NMDS uses rank-based distances to cluster samples that are most similar.

The family-level unconstrained ordination (nMDS) showed no obvious grouping of rats

with respect to the locations (**Fig. 7A**), indicating that locations did not correspond to the

predominant axes of variation among the diets. We next performed a constrained

ordination method, Canonical Analysis of Principal coordinates (CAP, see Methods).

CAP identified axes of variation, if any, that distinguish the diets of rats from different

locations (**Fig. 7B**). We found that the CAP axes correctly classified the locations of 19

out of 24 (79%) rats using a leave-one-out procedure. The families having the largest

correlations with the first two principal coordinates, and thus most responsible for the

separation between groups, were primarily plants: *Arecaceae*, *Podocarpaceae*,

Piperaceae, and *Pinaceae*. In addition, insect groups (*Cerambycids* and *Formicids*) and

birds (*Phaseanidae* and *Numididae*) played a role (**Fig. 7C**).

The families driving similarity within the three locations (i.e., those that had the greatest

within-location SIMPER scores, see Methods) varied among locations. LB had average

Bray-Curtis within-location similarity of 13%; mostly attributable to *Hymenolepidae*

(accounting for 51% of the within-group similarity), *Solanaceae* (11%), and *Fabaceae*

(11%). The average similarity for OB was 21%, with the greatest contributing taxa

being *Arecaceae* (33%), *Poaceae* (23%), *Fabaceae* (9%), and *Phasianidae* (8%). The

average similarity for WP was 24%, with the greatest contributing taxa being *Poaceae*

(72%) (**Table S4**).

Fig. 7. Unconstrained nMDS (a) and constrained CAP (b) ordinations of the diets
of rats from three locations. Both ordinations were based on Bray-Curtis
dissimilarities of square root transformed proportions of reads attributed to each
family. The locations were a native estuarine bush (OB, orange); a restored marine
wetland (LB, purple); and a native forest (WP, light blue). The CAP ordination is
repeated in panel **(c)** as a biplot with the rats omitted to show the Pearson correlations
between families and the first two CAP axes. The eight families with the strongest
correlations are shown, indicating the taxa associated with each location.

Discussion

Accuracy and sensitivity

Here we have shown that using a simple metagenomic approach with error-prone long

reads allows rapid and accurate classification of rat diet components (approximately

2.7% error in taxon assignments at the family-level). We expect that this technique can

be used to infer diet for a wide variety of animal and sample types, including samples

that use less invasive collection methods, such as fecal matter. The sensitivity of this

approach will likely improve as the accuracy and yield of ONT sequencing continues to

increase. The analysis here is based on less than 200,000 reads from two flow cells.

Current yields for similar read length distributions are in excess of ten million reads per

flow cell. As ONT sequencing accuracy improves (currently just above 96% modal

accuracy), this will also increase the applicability of this method. This is illustrated by

the fact that the fraction of reads yielding BLAST hits increased substantially for higher

quality reads, approaching 40% for high quality reads in our dataset (reads with greater

than 92.5% accuracy, **Fig. 2B)**.

As the species sampling of genomic databases increases (59), the taxon-level

precision of this method will improve. Given the current rate of genomic sequencing,

with careful sampling, the vast majority of multicellular plant and animal families (and

even genera) will likely have at least one type species with a sequenced genome within

the next decade. Continued advancement in sequence database search algorithms as

compared to current methods (23,24,60) should considerably decrease the

computational workload necessary to find matching sequences.

Methodological advantages

As genomic databases become more complete, metagenomic approaches will offer

significant advantages due to decreased bias. We found that rats consumed many soft-

bodied species (e.g. mushrooms, flat worms, slugs, and lepidopterans) that would be

difficult to identify using visual inspection of stomach contents. Achieving data on such

a wide variety of taxa (across multiple phyla) would be difficult to quantify using

metabarcoding, as there are no universal 18S or COI universal primers capable of

amplifying sequences in all these taxa. While it might be possible to use several

different primer sets targeted at different phyla or orders, quantitatively comparing diet

components across these using sequences amplified with different primer sets is

extremely difficult due to differences in primer binding and PCR efficiency. A second

advantage of metagenomic sequencing (as has been pointed out previously), is that it

allows for "holistic" views of ecology, as it simultaneously profiles parasites,

microbiome, and prey species (30).

The ONT-based sequencing method has several unique advantages, and perhaps the

most obvious is the accessibility of the platform. Compared to other high throughput

sequencing technologies (e.g. Illumina, IonTorrent, or PacBio), there is no initial capital

investment required to use the platform. On a per-sample basis, data generation is

inexpensive (assuming 12 multiplexed samples, approximately $150 USD per sample,

and half this price if reagents are purchased in bulk). Library preparation and

sequencing can be extremely rapid, going from DNA sample to sequence in less than

two hours (61). Furthermore, the sequencing platform itself is highly portable. Given (1)

that ONT-based methods are now similar in cost-per-read (approximately $650 for 10

million reads) than the most accessible Illumina method (MiSeq, approximately $1300

for 20 million reads); and (2) that even marginal increases in read length are likely to

significantly improve species identification, we expect that ONT-based methods should

soon become useful for routine ecological monitoring of species (62).

Some modifications to our approach might further increase the precision of our ability to

infer community composition. Any error-prone long read dataset (i.e. PacBio or ONT)

has both short (e.g. 500 bp) and long (e.g. 5000 bp) reads, as well as high quality (e.g.

mean accuracy greater than 90%) and low quality (e.g. mean accuracy less than 80%)

reads. When inferring community composition, a null expectation is that taxa should be

equally represented by long, high quality reads as they are by short, low quality reads. If

some taxa are represented only by short, low quality reads, this suggests that these

taxa may be false positive inferences. Similarly, the difficulty in correctly mapping short

inaccurate reads could be mitigated by weighting the probability of taxon mapping by

the number of long, accurate reads that map to certain taxa. Thus, the fact that not all

reads are extremely long and accurate does not mean that they cannot all be used to

infer taxon presence in metagenomic analyses.

For many diet studies, the aim is to ideally resolve the biomass, or nutritional content, or

prey numbers. However, as a caveat to the methods utilised here, and to all

sequencing-based methods, the accuracy of estimating these numbers is constrained

by the fact that different tissues and different taxa have different amounts of DNA (both

nuclear and mitochondrial) per gram of biomass. It is nearly to impossible to fully

account for this variation. Regardless, there is considerable utility in using sequencing

based approaches for diet assessment, not least because it is one of the few methods

that allows the full breadth of the diet to be observed, as illustrated here by the number

of different orders we find that rats predate.

Conclusions

Here we have shown that a rapid error-prone long read metagenomic approach is able to accurately characterise diet taxa at the family level and distinguish between the diets of rats according to the locations from which they were sourced. This information may be used to guide conservation efforts toward specific areas and habitats in which native species are most at risk from this highly destructive introduced predator.

Methods
Study Areas

We trapped rats from three locations near Auckland, New Zealand. Each location comprised a different type of habitat: undisturbed inland native forest (Waitakere Regional Parklands, WP); native bush surrounding an estuary (Okura Bush Walkway, OB); and restored coastal wetland (Long Bay Regional Park, LB) (**Fig. 8**). Snap traps in OB and LB were baited with peanut butter, apple, and cinnamon wax pellets; or bacon fat and flax pellets.

Fig. 8. Location of rat sampling sites in the greater Auckland area in the North

Island of New Zealand. Each point indicates a trap where one rat was captured, with the colour of the points indicating the three broad locations: the native estuarine bush habitat of Okura Bush (OB), the restored wetland of Long Bay (LB), and the native forest of Waitakere Park (WP). The two insets show the three locations in higher resolution with topographical details. Green indicates park areas. Precise geographical coordinates were only available for five out of eight rats in WP.

Traps in WP were baited with chicken eggs, rabbit meat, or cinnamon scented poison pellets. From 16 November to 16 December 2016, traps were surveyed by established conservation groups at each site every 48 hours. A total of 36 rats were collected from these locations. Three of the rats from WP had poison in their stomachs. However, all three of these were killed by the snap traps, and as such are unlikely to have swallowed any bait. In addition, none were identified as having chicken or rabbit in their diet. The majority of rats collected (34/36) were determined to be male *Rattus rattus* by visual inspection. These 34 rats were selected for further analysis.

DNA Isolation

Within 48 hours of trapping, rats were stored at either -20°C or -80°C until dissection. We dissected out intact stomachs from each animal and removed the contents. After snap freezing in liquid nitrogen, we homogenised the stomach contents using a sterile mini blender to ensure sampling was representative of the entire stomach.

We purified DNA from 20 mg of homogenised stomach contents using the Promega Wizard Genomic DNA Purification Kit, with the following modifications to the Animal Tissue protocol: after protein precipitation, we transferred the supernatant to a new tube and centrifuged a second time to minimise protein carryover. The DNA pellet was washed twice with ethanol. These modifications were performed to improved DNA

purity. We rehydrated precipitated DNA by incubating overnight in molecular biology

grade water at 4°C and stored the DNA at -20°C. DNA quantity, purity, and quality was

ascertained by nanodrop and agarose gel electrophoresis. The DNA samples were

ranked according quantity and purity (based on A260/A280 and secondarily, A230/A280

ratios). The eight highest quality DNA samples from each of the three locations were

selected for sequencing. We did no size selection on the purified DNA.

DNA Sequencing

Sequencing was performed on two different dates (24 January 2017 and 17 March

2017) using a MinION Mk1B device and R9.4 chemistry. For each sequencing run, DNA

from each rat was barcoded using the 1D Native Barcoding Kit (Barcode expansion kit

EXP-NBD103 with sequencing kit SQK-LSK108) following the manufacturer's

instructions. This included an AMPure bead purification step to remove adaptors, which

also likely removed very short reads (less than 200 bp; see **Fig. 1A**). Twelve samples

were pooled and run on each flow cell, for a total of 24 individual rats. The flow cells had

1373 active pores (January 2017) and 1439 active pores (March 2017). Sequencing

was performed using local base calling in MinKnow v1.3.25 (January) or MinKnow

v1.5.5 (March), but for the analysis present here both runs were re-basecalled after data

collection using Albacore 2.2.7 with demultiplexing performed in Albacore and filtering

disable*d (optio*ns *--barcoding --disable_filtering*).

Sequence classification

All sequences were BLASTed (blastn v2.6.0+) against a locally compiled database

consisting of the combined NCBI other_genomic and nt databases (downloaded on 13[th]

June 2018 from NCBI). Default blastn parameters were used (match 2, mismatch -3,

gapopen -5, gapextend -2), and only hits with an e-value of 1e-2 or less were saved.

Due to the predominance of short indels present in nanopore sequence data, we used

an initial set of basecalled data to test whether changing these default penalties affected

the results (gapopen -1, gapextend -1). We found that these adjusted parameters did

not qualitatively change our results.

We assigned sequence reads to specific taxon levels using MEGAN6 (v.6.11.7 June

2018) (41). We only used reads with BLAST hits having an e-value of 1×10^{-20} or lower

(corresponding to a bit score of 115 or higher given the databases we used) and an

alignment length of 100 base pairs or more. To assign reads to taxon levels, we

considered all hits having bit scores within 20% of the bit score of the best hit (MEGAN

parameter Top Percent).

Multivariate analyses

Multivariate analyses were done using the software PRIMER v7 (42). The data used in

the multivariate analyses were in the form of a sample- (i.e. individual rat) by-family

matrix of read counts. All bacteria, rodent, and primate families were removed. The

majority of rodent hits were to rat and mouse, resulting from the rats' own DNA (see

below). The majority of the primate hits (32 in total) were assigned to Hominidae (19),

which likely resulted from sample contamination (**Table S3**).

The read counts were converted to proportions per individual rat by dividing by the

total count for each rat, to account for the fact that the number of reads varied

substantially among rats (43). The proportions were then square-root transformed so

that subsequent analyses were informed by the full range of taxa, rather than just the

most abundant families (44). We then calculated a matrix of Bray-Curtis dissimilarities,

which quantified the difference in the gut DNA of each pair of rats based on the

square-root transformed proportions of read counts across families (43).

We used unconstrained ordination, non-metric multidimensional scaling (nMDS)

applied to the dissimilarity matrix to examine the overall patterns in the diet

composition among rats. To assess the degree to which the diet compositions of rats

were distinguishable among the three locations, we applied canonical analysis of

principal coordinates (CAP) (45) to the dissimilarity matrix. CAP is a constrained

ordination which aims to find axes through multivariate data that best separates *a priori*

groups of samples (in this case, the groups are the locations from which the rats were

sampled); CAP is akin to linear discriminant analysis but it can be used with any

resemblance matrix. The out-of-sample classification success was evaluated using a

leave-one-out cross-validation procedure (45).

We used Similarity Percentage (SIMPER; (46)) to characterise and distinguish between

the locations. This allowed us to identify the families with the greatest percentage

contributions to (1) the Bray-Curtis similarities of diets within each location (**Table S5**)

and (2) the Bray-Curtis dissimilarities between each pair of locations (**Table S6**).

Declarations

Ethics approval

Sample collection was performed under (Auckland Council Permit to Undertake Research WS1064).

Availability of data and materials

Sequence data are available in the SRA archive (accession number PRJEB27647).

Competing interests

WP received funding from Oxford Nanopore Technologies (1000$USD) to present this work at a conference (Ecological Society of Australia 2018).

Funding

This work was supported by a Massey University Research Fund to NF, a Marsden Fund Grant (15-MAU-136) to JD and Marsden Fund Grant (MAU1703) to OS.

Authors' contributions

WP, JD, NF, and OS conceived the project. WP performed the stomach dissections. WP and NF optimised the genomic DNA isolation and library preparation. NF performed the nanopore sequencing. GB and OS processed and performed quality control on the sequencing data. WP and OS performed the sequence classification. WP, AS, NF, and OS analysed the data. WP, NF, AS, and OS wrote the paper, with input from all authors.

Acknowledgements

Thanks to Friends of Okura Bush, Mary Stewart from Auckland Council, and Gillian

Wadams and the volunteers at the Waitakere Ranges for collecting rat samples and

aiding in rat species identification.

References

1. Daniel MJ. Seasonal Diet Of The Ship Rat (Rattus Rattus) In Lowland Forest In New Zealand. Proc. 1973;20:21–30.

2. Pierce GJ, Boyle. A review of methods for diet analysis in piscivorous marine mammals. Oceanogr Mar Biol Annu Rev. 1991;29:409–86.

3. Major HL, Jones IL, Charette MR, Diamond AW. Variations in the diet of introduced Norway rats (Rattus norvegicus) inferred using stable isotope analysis. J Zool. 2007 Apr 1;271(4):463–8.

4. Carreon-Martinez L, Heath DD. Revolution in food web analysis and trophic ecology: diet analysis by DNA and stable isotope analysis. Mol Ecol. 2010 Jan;19(1):25–7.

5. Hobson KA. Use of stable-carbon isotope analysis to estimate marine and terrestrial protein content in gull diets. Can J Zool. 1987 May 1;65(5):1210–3.

6. Basha WA, Chamberlain AT, Zaki ME, Kandeel WA, Fares NH. Diet reconstruction through stable isotope analysis of ancient mummified soft tissues from Kulubnarti (Sudanese Nubia). Journal of Archaeological Science: Reports. 2016 Feb 1;5:71–9.

7. Dunlap M, Pawlik JR. Video-monitored predation by Caribbean reef fishes on an array of mangrove and reef sponges. Mar Biol. 1996 Mar 1;126(1):117–23.

8. Brown KP, Moller H, Innes J, Jansen P. Identifying predators at nests of small birds in a New Zealand forest. Ibis . 2008;140(2):274–9.

9. King RA, Read DS, Traugott M, Symondson WOC. Molecular analysis of predation: a review of best practice for DNA-based approaches. Mol Ecol. 2008 Feb;17(4):947–63.

10. Soininen EM, Valentini A, Coissac E, Miquel C, Gielly L, Brochmann C, et al. Analysing diet of small herbivores: the efficiency of DNA barcoding coupled with high-throughput pyrosequencing for deciphering the composition of complex plant mixtures. Front Zool. 2009 Aug 20;6:16.

11. Jarman SN, Gales NJ, Tierney M, Gill PC, Elliott NG. A DNA-based method for
identification of krill species and its application to analysing the diet of marine
vertebrate predators. Mol Ecol. 2002 Dec;11(12):2679–90.

12. Jarman SN, Deagle BE, Gales NJ. Group-specific polymerase chain reaction for
DNA-based analysis of species diversity and identity in dietary samples. Mol Ecol.
2004 May;13(5):1313–22.

13. Tedersoo L, Anslan S, Bahram M, Põlme S, Riit T, Liiv I, et al. Shotgun
metagenomes and multiple primer pair-barcode combinations of amplicons reveal
biases in metabarcoding analyses of fungi. MycoKeys. 2015;10:1.

14. Leray M, Yang JY, Meyer CP, Mills SC, Agudelo N, Ranwez V, et al. A new
versatile primer set targeting a short fragment of the mitochondrial COI region for
metabarcoding metazoan diversity: application for characterizing coral reef fish gut
contents. Front Zool. 2013 Jun 14;10:34.

15. Pawluczyk M, Weiss J, Links MG, Egaña Aranguren M, Wilkinson MD, Egea-
Cortines M. Quantitative evaluation of bias in PCR amplification and next-
generation sequencing derived from metabarcoding samples. Anal Bioanal Chem.
2015 Mar;407(7):1841–8.

16. Pereira RPA, Peplies J, Brettar I, Hoefle MG. Impact of DNA polymerase choice on
assessment of bacterial communities by a Legionella genus-specific next-
generation sequencing approach [Internet]. bioRxiv. 2018 [cited 2018 Jan 31]. p.
247445. Available from:
https://www.biorxiv.org/content/early/2018/01/12/247445.abstract

17. Hover BM, Kim S-H, Katz M, Charlop-Powers Z, Owen JG, Ternei MA, et al.
Culture-independent discovery of the malacidins as calcium-dependent antibiotics
with activity against multidrug-resistant Gram-positive pathogens. Nat Microbiol.
2018 Apr;3(4):415–22.

18. Anantharaman K, Brown CT, Hug LA, Sharon I, Castelle CJ, Probst AJ, et al.
Thousands of microbial genomes shed light on interconnected biogeochemical
processes in an aquifer system. Nat Commun. 2016 Oct 24;7:13219.

19. Xu Z, Knight R. Dietary effects on human gut microbiome diversity. Br J Nutr. 2015
Jan;113 Suppl:S1–5.

20. Fierer N, Leff JW, Adams BJ, Nielsen UN, Bates ST, Lauber CL, et al. Cross-biome
metagenomic analyses of soil microbial communities and their functional
attributes. Proc Natl Acad Sci U S A. 2012 Dec 26;109(52):21390–5.

21. Breitwieser FP, Salzberg SL. KrakenHLL: Confident and fast metagenomics
classification using unique k-mer counts. bioRxiv [Internet]. 2018; Available from:

668  https://www.biorxiv.org/content/early/2018/06/06/262956.abstract

669 22. Huson DH, Mitra S, Ruscheweyh H-J, Weber N, Schuster SC. Integrative analysis
670   of environmental sequences using MEGAN4. Genome Res. 2011 Sep;21(9):1552–
671   60.

672 23. Kim D, Song L, Breitwieser FP, Salzberg SL. Centrifuge: rapid and sensitive
673   classification of metagenomic sequences. Genome Res. 2016 Dec;26(12):1721–9.

674 24. Wood DE, Salzberg SL. Kraken: ultrafast metagenomic sequence classification
675   using exact alignments. Genome Biol. 2014 Mar 3;15(3):R46.

676 25. Paula DP, Linard B, Crampton-Platt A, Srivathsan A, Timmermans MJTN, Sujii ER,
677   et al. Uncovering Trophic Interactions in Arthropod Predators through DNA
678   Shotgun-Sequencing of Gut Contents. PLoS One. 2016 Sep 13;11(9):e0161841.

679 26. Srivathsan A, Ang A, Vogler AP, Meier R. Fecal metagenomics for the
680   simultaneous assessment of diet, parasites, and population genetics of an
681   understudied primate. Front Zool. 2016 Apr 21;13:17.

682 27. Arribas P, Andújar C, Hopkins K, Shepherd M, Vogler AP. Metabarcoding and
683   mitochondrial metagenomics of endogean arthropods to unveil the mesofauna of
684   the soil. Yu D, editor. Methods Ecol Evol. 2016 Sep 7;7(9):1071–81.

685 28. Linard B, Crampton-Platt A, Gillett CPDT, Timmermans MJTN, Vogler AP.
686   Metagenome Skimming of Insect Specimen Pools: Potential for Comparative
687   Genomics. Genome Biol Evol. 2015 May 14;7(6):1474–89.

688 29. Meiser A, Otte J, Schmitt I, Grande FD. Sequencing genomes from mixed DNA
689   samples - evaluating the metagenome skimming approach in lichenized fungi. Sci
690   Rep. 2017 Nov 2;7(1):14881.

691 30. Srivathsan A, Sha JCM, Vogler AP, Meier R. Comparing the effectiveness of
692   metagenomics and metabarcoding for diet analysis of a leaf-feeding monkey
693   (Pygathrix nemaeus). Mol Ecol Resour. 2015 Mar;15(2):250–61.

694 31. Søe MJ, Nejsum P, Seersholm FV, Fredensborg BL, Habraken R, Haase K, et al.
695   Ancient DNA from latrines in Northern Europe and the Middle East (500 BC–1700
696   AD) reveals past parasites and diet. PLoS One. 2018 Apr 25;13(4):e0195481.

697 32. Pearman WS, Freed NE, Silander OK. The advantages and disadvantages of
698   short- and long-read metagenomics to infer bacterial and eukaryotic community
699   composition [Internet]. bioRxiv. 2019 [cited 2019 Aug 13]. p. 650788. Available
700   from: https://www.biorxiv.org/content/10.1101/650788v2.abstract

701 33. Hillmann B, Al-Ghalith GA, Shields-Cutler RR, Zhu Q, Gohl DM, Beckman KB, et
702   al. Evaluating the Information Content of Shallow Shotgun Metagenomics.

mSystems [Internet]. 2018 Nov;3(6). Available from:
http://dx.doi.org/10.1128/mSystems.00069-18

34. Gibbs GW. Why are some weta (Orthoptera: Stenopelmatidae) vulnerable yet
others are common? J Insect Conserv. 1998 Dec 1;2(3-4):161–6.

35. Towns DR, Daugherty CH, Cree A. Raising the prospects for a forgotten fauna: a
review of 10 years of conservation effort for New Zealand reptiles. Biol Conserv.
2001 May 1;99(1):3–16.

36. Stringer IAN, Bassett SM, McLean MJ, McCartney J, Parrish GR. Biology and
conservation of the rare New Zealand land snail Paryphanta busbyi watti
(Mollusca, Pulmonata). Invertebr Biol. 2003 Sep 1;122(3):241–51.

37. Diamond JM, Veitch CR. Extinctions and introductions in the new zealand
avifauna: cause and effect? Science. 1981 Jan 30;211(4481):499–501.

38. Dowding JE, Murphy EC. The impact of predation by introduced mammals on
endemic shorebirds in New Zealand: a conservation perspective. Biol Conserv.
2001 May 1;99(1):47–64.

39. Graham NAJ, Wilson SK, Carr P, Hoey AS, Jennings S, MacNeil MA. Seabirds
enhance coral reef productivity and functioning in the absence of invasive rats.
Nature. 2018 Jul;559(7713):250–3.

40. Russell JC, Innes JG, Brown PH, Byrom AE. Predator-Free New Zealand:
Conservation Country. Bioscience. 2015 May 1;65(5):520–5.

41. Huson DH, Beier S, Flade I, Górska A, El-Hadidi M, Mitra S, et al. MEGAN
Community Edition - Interactive Exploration and Analysis of Large-Scale
Microbiome Sequencing Data. PLoS Comput Biol. 2016 Jun;12(6):e1004957.

42. Clarke KR, Gorley RN. PRIMER v7: User Manual/Tutorial. PRIMER-E, Plymouth;
2015 p. 296.

43. Clarke KR, Robert Clarke K, Somerfield PJ, Gee Chapman M. On resemblance
measures for ecological studies, including taxonomic dissimilarities and a zero-
adjusted Bray–Curtis coefficient for denuded assemblages. J Exp Mar Bio Ecol.
2006;330(1):55–80.

44. Clarke KR, Green RH. Statistical Design and Analysis for a "biological Effects"
Study. Mar Ecol Prog Ser. 1988;46:213–26.

45. Anderson MJ, Willis TJ. Canonical Analysis Of Principal Coordinates: A Useful
Method Of Constrained Ordination For Ecology. Ecology. 2003 Feb 1;84(2):511–
25.

46. Clarke KR. Non-parametric multivariate analyses of changes in community
structure. Austral Ecol. 1993;18(1):117–43.

47. Jain M, Koren S, Miga KH, Quick J, Rand AC, Sasani TA, et al. Nanopore
sequencing and assembly of a human genome with ultra-long reads. Nat
Biotechnol. 2018 Apr;36(4):338–45.

48. Deagle BE, Eveson JP, Jarman SN. Quantification of damage in DNA recovered
from highly degraded samples--a case study on DNA in faeces. Front Zool. 2006
Aug 16;3:11.

49. Jain M, Olsen HE, Paten B, Akeson M. The Oxford Nanopore MinION: delivery of
nanopore sequencing to the genomics community. Genome Biol. 2016 Nov
25;17(1):239.

50. McIntyre ABR, Ounit R, Afshinnekoo E, Prill RJ, Hénaff E, Alexander N, et al.
Comprehensive benchmarking and ensemble approaches for metagenomic
classifiers. Genome Biol. 2017 Dec 21;18(1):182.

51. Huson DH, Auch AF, Qi J, Schuster SC. MEGAN analysis of metagenomic data.
Genome Res. 2007 Mar;17(3):377–86.

52. Maurice CF, Knowles SCL, Ladau J, Pollard KS, Fenton A, Pedersen AB, et al.
Marked seasonal variation in the wild mouse gut microbiota. ISME J. 2015
Nov;9(11):2423–34.

53. Brownlee A, Moss W. The influence of diet on lactobacilli in the stomach of the rat.
J Pathol. 1961 Oct 1;82(2):513–6.

54. Li D, Chen H, Mao B, Yang Q, Zhao J, Gu Z, et al. Microbial Biogeography and
Core Microbiota of the Rat Digestive Tract. Sci Rep. 2017 Apr 4;8:45840.

55. Horáková Z, Zierdt CH, Beaven MA. Identification of lactobacillus as the source of
bacterial histidine decarboxylase in rat stomach. Eur J Pharmacol. 1971
Sep;16(1):67–77.

56. Bridgman LJ, Innes J, Gillies C, Fitzgerald N, King CM. Do ship rats display
predatory behaviour towards house mice? Anim Behav. 2013 Aug 1;86(2)):257–68.

57. Sweetapple PJ, Nugent G. Ship rat demography and diet following possum control
in a mixed podocarp—hardwood forest. N Z J Ecol. 2007;31(2):186–201.

58. Riofrío-Lazo M, Páez-Rosas D. Feeding Habits of Introduced Black Rats, Rattus
rattus, in Nesting Colonies of Galapagos Petrel on San Cristóbal Island,
Galapagos. PLoS One. 2015 May 18;10(5):e0127901.

59. Lewin HA, Robinson GE, Kress WJ, Baker WJ, Coddington J, Crandall KA, et al.

Earth BioGenome Project: Sequencing life for the future of life. Proc Natl Acad Sci
U S A. 2018 Apr 24;115(17):4325–33.

60. Nasko DJ, Koren S, Phillippy AM, Treangen TJ. RefSeq database growth
influences the accuracy of k-mer-based species identification. 2018;1–21.

61. Zaaijer S, Gordon A, Speyer D, Piccone R, Groen SC, Erlich Y. Rapid re-
identification of human samples using portable DNA sequencing. Elife [Internet].
2017 Nov 28;6. Available from: http://dx.doi.org/10.7554/eLife.27798

62. Kamenova S, Bartley TJ, Bohan DA, Boutain JR, Colautti RI, Domaizon I, et al.
Chapter Three - Invasions Toolkit: Current Methods for Tracking the Spread and
Impact of Invasive Species. In: Bohan DA, Dumbrell AJ, Massol F, editors.
Advances in Ecological Research. Academic Press; 2017. p. 85–182.

 Supplemental Tables

784 **Table S1**. Read numbers and total base pairs for each barcode in the January 2017
785 sequencing run.

Sample	Total reads	Total Mbp	Mean length
OB2	19907	14.62	734
WP11	10164	9.63	947
WP5	8237	6.78	823
LB7	7548	7.04	933
OB13	3644	3.63	995
WP9	2954	2.4	814
OB5	2850	2.06	721
WP8	2801	2.32	827
LB6	2531	1.6	632
OB7	2473	1.87	756
LB5	1641	1.16	705
LB3	1554	0.99	636
None	16673	13.01	781
Total	82977	67.1	809

 Table S2. Read numbers and total base pairs for each barcode in the March 2017
 sequencing run.

Sample	Total reads	Total Mbp	Mean length
LB1	17820	9.21	517
LB8	16923	13.13	776
WP2	10511	7.00	666
LB4	8684	4.92	567
OB11	5689	3.40	598
WP10	1563	0.99	633
OB12	1479	0.89	604
WP12	1309	0.78	596
LB2	1127	0.76	676
WP3	637	0.73	1141
OB14	541	0.37	683
OB10	435	0.24	555
None	29432	21.33	725
Total	96150	63.75	663

Table S3. Characteristics of alignments for reads assigned to the Primate family.
Many are both long and have high identity, suggesting that they are not false positive
assignments, but contamination.

Rat	Read length	Mean read accuracy	% ID	Alignment length	Genus
WP10	428	0.91	94.0	314	Homo
WP10	782	0.92	93.8	657	Homo
OB5	365	0.90	93.2	249	Homo
LB3	515	0.95	92.6	462	Homo
WP10	510	0.90	92.0	460	Homo
WP10	704	0.90	91.1	587	Homo
WP10	467	0.89	90.3	402	Homo
WP10	494	0.89	89.7	388	Homo
WP10	339	0.88	89.6	269	Homo
WP10	446	0.88	89.6	326	Homo
WP10	327	0.91	89.1	210	Homo
OB5	415	0.89	88.8	277	Homo
OB5	561	0.86	88.7	257	Homo
WP11	434	0.86	88.7	301	Homo
WP10	486	0.88	88.2	365	Homo
WP10	613	0.90	88.2	526	Homo
WP10	563	0.87	88.2	457	Homo
WP10	1373	0.91	87.7	1337	Homo
WP11	526	0.91	87.5	473	Homo
OB14	478	0.89	86.9	373	Homo
OB5	715	0.89	86.8	673	Homo
WP10	475	0.87	86.7	362	Homo
WP10	398	0.86	86.5	259	Homo
WP10	377	0.88	85.3	251	Homo
WP9	558	0.87	85.2	508	Homo
WP10	429	0.86	84.8	276	Homo
WP10	322	0.81	84.5	174	Homo
WP8	723	0.84	83.1	438	Homo
LB8	965	0.86	80.4	245	Rhinopithecus

LB5	3042	0.94	79.4	1018	Cebus
WP2	464	0.93	77.3	216	Homo
LB8	671	0.90	73.9	406	Aotus

Table S4. Characteristics of alignments for reads that are likely false positive assignments. Most are short long or have low identity, suggesting that they are false positive assignments. The exception are the reads matching Buthidae, which we hypothesize are due to the rats predation of the sister taxa, harvestmen.

Rat	Read length	Mean read accuracy	% ID	Alignment length	Genus	Megan family
WP5	1285	0.90	82.9	298	Centruroides	Buthidae
OB11	1874	0.90	88.0	664	Centruroides	Buthidae
WP5	1711	0.93	79.3	762	Centruroides	Buthidae
WP11	859	0.93	86.1	151	Octopus	Octopodidae
WP2	516	0.86	81.4	172	Oncorhynchus	Salmonidae
OB12	424	0.90	85.7	140	Xiphophorus	Poeciliidae
OB12	643	0.84	89.3	177	Xiphophorus	Poeciliidae

Table S5. SIMPER analysis of family contributions to group similarities.

Family	Average	Average	Similarity/SD	Percentage	Group
Hymenolepididae	3.37	6.87	0.34	51.2	LB
Solanaceae	1.57	1.48	0.34	11.1	LB
Fabaceae	1.74	1.41	0.44	10.5	LB
Arecaceae	2.86	7.11	1	33.4	OB
Poaceae	2.87	4.82	0.55	22.7	OB
Fabaceae	1.17	1.98	0.51	9.3	OB
Phasianidae	1.79	1.67	0.34	7.9	OB
Poaceae	5.08	17.61	0.62	72.1	WP

805   **Table S6.** SIMPER analysis of family contributions to group dissimilarities.

Species	Av Abund Group1	Avg Abund Group2	Avg. Diss	Diss/SD	% contrib	Group 1	Group2
Poaceae	1.95	5.08	15.15	1.04	16.74	LB	WP
Poaceae	2.87	5.08	11.29	1.26	13.78	OB	WP
Hymenolepididae	3.37	0.48	10.8	0.73	11.93	LB	WP
Hymenolepididae	3.37	0.29	9.37	0.79	10.32	LB	OB
Poaceae	1.95	2.87	8.37	1.1	9.22	LB	OB
Arecaceae	0.05	2.86	6.99	1.41	7.7	LB	OB
Arecaceae	2.86	1.31	5.92	1.29	7.23	OB	WP
Fabaceae	1.74	1.05	6.14	0.67	6.78	LB	WP
Podocarpaceae	0	2.38	5.34	0.83	5.9	LB	WP
Podocarpaceae	0.71	2.38	4.82	0.99	5.88	OB	WP
Fabaceae	1.74	1.17	4.87	0.81	5.37	LB	OB
Fabaceae	1.17	1.05	4.31	0.84	5.26	OB	WP

Datafile S1. Table of read BLAST hits and assigned MEGAN taxa with with reads
reclassified at the family or order level by filtering on read length to alignment length
ratio and percent identity.

Datafile S2. Table of read BLAST hits and assigned MEGAN taxa with no filters
applied.

Supplemental Figures

Fig S1. Correlation of read accuracy with alignment characteristics. Only rat reads
exhibit a clear positive relationship between accuracy and percent identity. (A) indicates
the relationship for reads assigned to *Rattus* and (B) for *Mus*. The orange line indicates
a running median; the red dotted line is the y=x line, which is expected if accuracy
corresponds exactly to percent identity.

Fig S2. Alignment characteristics of true positive and false positive taxon
assignments at the family level. False positive taxon assignments (*Cricetideaa*,
orange, and *Spalacidae*, green) have lower percent identity and shorter alignment
lengths than true positive taxon assignments (*Muridae*, black). Only a single false
positive taxon assignment has a read length to alignment length ratio greater than 0.5
and a percent identity greater than 85%. This suggests that with further filters or
methodologies (e.g. decision tree analysis using different read and alignment
characteristics) could, if necessary, decrease false positive rates even further.