1. Title: New tools for diet analyses: nanopore sequencing of metagenomic DNA from

stomach contents to quantify diet in an invasive population of rats

2. Authors: William Pearman, Adam N. H. Smith, Georgia Breckell, James Dale, Nikki E.

Freed, Olin K. Silander

3. Addresses of all authors: Institute of Natural and Mathematical Sciences, Massey

University, Auckland 0745, New Zealand

4. Keywords: Nanopore sequencing, diet analysis, rat, invasive

5. Communicating authors: Olin K. Silander, Institute of Natural and Mathematical

Sciences, Massey University, Auckland 0745, New Zealand, olinsilander@gmail.com,

+64 9 213 6618; Nikki E. Freed, Institute of Natural and Mathematical Sciences, Massey

University, Auckland 0745, New Zealand, freednikki@gmail.com, +64 9 213 6639

6. Running title: Quantifying rat diets by direct nanopore sequencing

Abstract

Introduced species of mammals in New Zealand have had catastrophic effects on

populations of diverse native species. Quantifying the diets of these omnivorous and

predatory species is critical for understanding which native species are most impacted,

and to prioritize which mammal species and locations should be targeted with control

programmes. A variety of methods have been applied to quantify diet components in

animals, including visual inspection of gut contents (Daniel 1973; Pierce and Boyle

1991), stable isotope analysis (Major et al. 2007; Carreon-Martinez and Heath 2010),

and time-lapse video (Brown and Brown 1997; Dunlap and Pawlik 1996). Increasingly,

DNA-based metabarcoding methods are being used (King et al. 2008; Soininen et al.

2009). These metabarcoding methods require a PCR step using primers that bind to

highly conserved genomic regions (e.g. mitochondrial COI) to amplify specific regions

for sequencing. This step introduces significant bias, primarily due to the lack of a

universal primer set (King et al. 2008). Here we show that direct metagenomic

sequencing using the Oxford Nanopore Minion allows rapid quantification of rat diets.

Using a sample of rats collected from within 100km of Auckland, NZ, we show that

these rats consume a wide variety of plant, invertebrate, vertebrate, and fungal taxa,

with substantial differences in diet content between locales. We then show that, based

on diet content alone, it is possible to pinpoint the sampling location of an individual rat

within tens of kilometres. We expect that the rapidly increasing accuracy and

throughput of nanopore-based sequencing, as well as increases in the species

diversity of genomic databases, will soon allow rapid and unbiased assessments of

animal diets in field settings.

Introduction

The relatively recent introduction of mammalian species to New Zealand has had

significant negative effects on many of the native animal populations, including insects

(Gibbs 1998), reptiles (Towns, Daugherty, and Cree 2001), molluscs (Stringer et al.

2003), and birds (Diamond and Veitch 1981; Dowding and Murphy 2001). These

mammalian predators include possums, rats, stoats, and hedgehogs, as well as

domestic pets such as cats (Gillies and Clout 2003). Currently, an ambitious plan is

being put into place that aims for the eradication of all mammalian predators by 2050

(http://www.doc.govt.nz/predator-free-2050; Russell et al. 2015). A useful step toward

this goal would be to prioritise the management of predators and geographical areas in

which native species are experiencing the highest levels of predation. To do so requires

establishing the diet content of these predators.

Unbiased and sensitive assessment of diet content is extremely difficult to achieve due

to the limited accuracy of available methods. Identification of prey items using visual

examination of stomach contents is strongly affected by which items are most easily

degraded (for example, soft-bodied species). Stable isotope analysis is imprecise,

yielding only broad information on diet such as relative consumption of protein and plant

matter, as well as information on whether prey items are terrestrial or marine in origin

(Hobson 1987; Basha et al. 2016). Time-lapse video (Brown et al. 2008) requires

identification of the specific prey item, often difficult or impossible for small prey items or

in low-light conditions.

Perhaps the most widely applied current molecular method is DNA metabarcoding. This

approach relies on the PCR amplification and DNA sequencing of conserved regions

from nuclear, mitochondrial, and/or plastid genomes (King et al. 2008). With adequate

primer selection, this method can detect a wide range of species, and does not require

specific expertise necessary for other methods (for example identifying degraded prey

items).

However, DNA metabarcoding is not free from bias. PCR primers must be specifically

tailored to particular sets of taxa or species (Jarman et al. 2002). Although more

"universal" PCR primer pairs have been developed (for example targeting all bilaterians

or even all eukaryotes; (Jarman, Deagle, and Gales 2004), all primer sets exhibit bias

towards certain taxa. Tedersoo et al. (Tedersoo et al. 2015) found five-fold differences

in fungal operational taxonomic units (OTU) estimates when using different sets of

fungal-specific PCR primer pairs. Leray et al. (Leray et al. 2013) found that published

universal primer pairs (i.e. those that do not target specific taxa) were capable of

amplifying only between 57% and 91% of tested metazoan species, with as few as 33%

of species in some phyla being amplified at all (e.g. cnidarians). Deagle et al. argued

that in general, COI regions are simply not sufficiently conserved, and thus should not

be used for metabarcoding studies at all (Deagle et al. 2014). Finally, Pawluczyk et al.

showed that different loci from the same species exhibit up to 2,000-fold differences in

qPCR-estimated DNA quantity within samples (Pawluczyk et al. 2015). It has even been

shown that the polymerase itself can bias diversity metrics when using metabarcoding

methods (Pereira et al. 2018). For these reasons, a less biased molecular-based

method is desirable.

Metagenomic approaches, in which all DNA in the sample is directly sequenced, offer a

promising alternative. Such methods have been used for diet analysis in only very few

and recent instances (Srivathsan et al. 2015; Paula et al. 2015; Srivathsan et al. 2016).

Although metagenomic approaches decrease the bias arising from PCR amplification of

specific DNA regions, additional biases can arise, as the presence or absence of

species and genera can only be inferred for those species or genera present in genomic

databases. Although this is similarly true for metabarcoding approaches, metabarcode

databases are considerably more comprehensive in terms of species representation as

compared to genomic databases. Importantly, genomic sequence databases are rapidly

increasing in species diversity, as are the methods to query these large databases

(Wood and Salzberg 2014; Kim et al. 2016).

Here, we quantify rat diet composition using a metagenomic approach based on

nanopore sequencing (Oxford Nanopore Technologies). The data suggest that rat diets

around Auckland, New Zealand, comprise a wide range of plant, animal, and fungal

families. We show using multivariate analyses that rat diets are distinguishable by

habitat and location. This work establishes long-read metagenomic methods as a

straightforward approach for diet quantification.

Materials and Methods

Study Areas

We trapped rats from three locations near Auckland, New Zealand. Each location

comprised a different type of habitat: undisturbed inland native forest (Waitakere

Regional Parklands, WP); native bush surrounding an estuary (Okura Bush Walkway,

OB); and restored coastal wetland (Long Bay Regional Park, LB) (**Fig. 1**). Traps in OB

and LB were baited with peanut butter, apple, and cinnamon wax pellets; or bacon fat

and flax pellets. Traps in WP were baited with chicken eggs, rabbit meat, or cinnamon

scented poison pellets. From 16 November to 16 December 2016, traps were surveyed

by established volunteer conservation groups at each site every 48 hours. A total of 36

rats were collected from these locations. The majority of rats collected (34/36) were

determined to be male by visual inspection, and identified as *Rattus norvegicus*. These

34 rats were selected for further analysis.

Fig. 1. Location of rat sampling sites in the greater Auckland area in the North Island of New Zealand. Each point indicates a trap where one rat was captured, with the colour of the points indicating the three broad locations: the native estuarine bush habitat of Okura Bush (OB), the restored wetland of Long Bay (LB), and the native forest of Waitakere Park (WP). The two insets show the three locations in higher resolution with topographical details. Green indicates park areas. Precise geographical coordinates were only available for five out of eight rats in WP.

DNA Isolation

Within 48 hours of trapping, rats were stored at either -20°C or -80°C until dissection. We removed intact stomachs from each animal, rinsed them with 70% ethanol, and removed the contents. After snap freezing in liquid nitrogen, we homogenised the stomach contents using a sterile mini blender to ensure sampling was representative of the entire stomach.

We purified DNA from 10-20 mg of homogenised stomach contents using the Promega

Wizard Genomic DNA Purification Kit, with the following modifications to the Animal

Tissue protocol: after protein precipitation, we transferred the supernatant to a new tube

and centrifuged a second time to minimise protein carryover. The DNA pellet was

washed twice with ethanol. These modifications were performed to improved DNA

purity. We rehydrated precipitated DNA by incubating overnight in molecular biology

grade water at 4°C and stored the DNA at -20°C. DNA quantity, purity, and quality was

ascertained by nanodrop and agarose gel. The DNA samples were ranked according

quantity and purity (based on A260/A280 and secondarily, A230/A280 ratios). The eight

highest quality DNA samples from each location were selected for DNA sequencing.

DNA Sequencing

Sequencing was performed on two different dates (24 January 2017 and 17 March

2017) using a MinION Mk1B device and R9.4 chemistry. For each sequencing run, DNA

from each rat was barcoded using the 1D Native Barcoding Kit (Barcode expansion kit

EXP-NBD103 with sequencing kit SQK-LSK108) following the manufacturer's

instructions. Twelve samples were pooled and run on each flow cell, for a total of 24

individual rats. The flow cells had 1373 active pores (January) and 1439 active pores

(March). Sequencing was performed using local base calling in MinKnow v1.3.25

(January) or MinKnow v1.5.5 (March), but both runs were re-basecalled after data

collection using Albacore 2.2.7 with the following command: *read_fast5_basecaller.py -i*

./fast5_input -t 48 -s basecalled_output -f FLO-MIN106 -k SQK-LSK108 --barcoding -r -

o fastq,fast5 -q 0 --disable_filtering.

Sequence classification

All sequences were BLASTed (blastn v2.6.0+) against a locally compiled database

consisting of the combined NCBI other_genomic and nt databases (downloaded on 13[th]

June 2018 from NCBI). Default blastn parameters were used (gapopen 5, gapextend 2),

and only hits with an e-value of 0.01 or less were retained. Due to the predominance of

short indels present in nanopore sequence data, we used the datasets basecalled in

January and March of 2017 to test whether changing these default penalties affected

the results (gapopen 1, gapextend 1). We found that these adjusted parameters did not

qualitatively change our results.

We assigned sequence reads to specific taxon levels using MEGAN6 (v.6.11.7 June

2018) (Huson et al. 2016). We only used reads with BLAST hits having an e-value of

1×10^{-20} or lower (corresponding to a bit score of 115 or higher) and an alignment length

of 100 bp or more. To assign reads to taxon levels, we considered all hits having bit

scores within 20% of the bit score of the best hit (MEGAN parameter Top Percent).

Statistical analyses

Multivariate analyses were done using the software PRIMER v7 (Clarke and Gorley

2015). The data used in the multivariate analyses were in the form of a sample- (i.e.

individual rat) by-family matrix of read counts. All bacteria, rodent, and primate families

were removed. The majority of rodent hits were to rat and mouse, resulting from the

rats' own DNA (see below). The majority of the primate hits were to human sequences,

which probably resulted from sample contamination.

The read counts were converted to proportions per individual rat, by dividing by the total

count for each rat, to account for the fact that the number of reads varied substantially

among rats (Clarke et al. 2006). The proportions were then square-root transformed so

that subsequent analyses were informed by the full range of taxa, rather than just the

most abundant families (Clarke and Green 1988). We then calculated a matrix of Bray-

Curtis dissimilarities, which quantified the difference in the gut DNA of each pair of rats

based on the square-root transformed proportions of read counts across families

(Clarke et al. 2006).

We used unconstrained ordination (specifically, non-metric multidimensional scaling

(nMDS)) applied to the dissimilarity matrix to examine the overall patterns in the diet

composition among rats. To assess the degree to which the diet compositions of rats

were distinguishable among the three locations, we applied canonical analysis of

principal coordinates (CAP) (Anderson and Willis 2003) to the dissimilarity matrix. CAP

is a constrained ordination which aims to find axes through multivariate data that best

separates *a priori* groups of samples (in this case, the groups are the locations from

which the rats were sampled); it is akin to linear discriminant analysis but CAP can be

used with any resemblance matrix. The out-of-sample classification success was

evaluated using a leave-one-out cross-validation procedure (Anderson and Willis 2003).

We used Similarity Percentage (SIMPER) (Clarke 1993)) to characterise taxa that

distinguished between the locations. This allowed us to identify the families with the

greatest percentage contributions to (1) the Bray-Curtis similarities of diets within each

location and (2) the Bray-Curtis dissimilarities between each pair of locations.

Results

DNA sequencing and assignment of reads to taxa

After DNA isolation and sequencing, we obtained a total of 82,977 reads from the

January sequencing run and 96,150 reads from the March run. Median read lengths

were 606 bp and 527 bp for the January and March datasets, respectively (**Fig. 2A**).

These lengths are considerably shorter than other nanopore sequencing results from

both our and others work (Jain et al. 2016). This is most likely due to degradation of the

DNA during digestion in the stomach as well as fragmentation during DNA isolation

(Deagle, Eveson, and Jarman 2006) and sequencing library preparation. The median

quality scores per read ranged from 7-12 (0.80 - 0.94 accuracy) for both runs (**Fig. S1**).

The number of reads per barcoded rat sample varied by 10-fold for January and up to

40-fold in March(**Fig. 2B** and **2C**). This is primarily due to the highly variable quality of

DNA in each sample. However, read length and quality were similar for all samples

(**Fig. S1**).

Fig. 2. Results of nanopore metagenomic sequencing of rat stomach contents. (a)

Read length distribution for January and March nanopore runs. Read lengths

varied between 300 and 3,000 bp. **(b) and (c) Barcode distributions for January and**

March runs, respectively. We multiplexed the samples on the flow cells, using 12

barcodes per flow cell. The distribution of read numbers across barcodes was quite

uneven, varying by up to 40-fold in some cases. 20% (January) and 30% (March) of all

reads could not be assigned to a barcode ("None"). The inability to assign these reads

to a barcode is due primarily to their lower quality.

To quantify diet contents we first BLASTed all sequences against a combined database

of the NCBI nt database (the partially non-redundant nucleotide sequences from all

traditional divisions of GenBank excluding genome survey sequence, EST, high-

throughput genome, and whole genome shotgun

(ftp://ftp.ncbi.nlm.nih.gov/blast/db/README)) and the NCBI other_genomic database

(RefSeq chromosome records for non-human organisms

(ftp://ftp.ncbi.nlm.nih.gov/blast/db/README)). We used BLAST as it is generally viewed

as the gold standard method in metagenomic analyses (McIntyre et al. 2017). Of the

133,022 barcoded reads, 30,535 (23%) hit a sequence in the combined nt and

other_genomic database at an e-value cut-off of 0.01.

As an initial assessment of the quality of these hits, we examined the alignment lengths

and e-values. We found a bimodal distribution of alignment lengths and a highly skewed

distribution of e-values (**Fig. 3A**). We hypothesized that many of the short alignments

with high e-values were false positives. We thus first filtered this hit set, only retaining

BLAST hits with e-values less than 1e-20. Similar quality filters have been imposed

previously (Srivathsan et al. 2015). A total of 22,331 hits passed this filter. Mean read

quality had substantial effects on the rates of BLAST success, with almost 40% of high

accuracy read having hits in the March dataset, as compared to 1% of low accuracy hits

(**Fig. 3B**).

To specifically assign each sequence read to a taxon, we analysed the BLAST results in

MEGAN6 (Huson et al. 2016). The algorithm employed in MEGAN6 assigns reads to a

most recent common ancestor (MRCA) taxon level. For example, if a read has BLAST

hits to five species, three of which have bit scores within 20% of the best hit, the read

will be assigned to the genus, family, order, or higher taxon level that is the MRCA of

those best-hit three species (Huson et al. 2007). If a read matches one species far

better than to any other, by definition, the MRCA is that species.

Of the reads from the rat stomachs with BLAST hits, 31% were assigned by MEGAN as

being bacterial. 55% of these were *Lactobacillus spp*. These results match previous

studies on rat stomach microbiomes, which have found lactobacilli to be the dominant

taxa (Maurice et al. 2015; Brownlee and Moss 1961; Li et al. 2017; Horáková, Zierdt,

and Beaven 1971). Plant-associated *Pseudomonas* and *Lactococcus* taxa were also

common, at 7% and 6%, respectively.

Fig. 3. BLAST hits of metagenomic reads. (a) Biplot showing the e-value and alignment length of the top BLAST hit for each read. We observed a bimodal distribution of alignment lengths and a skewed distribution of e-values. The y-axis is plotted on a log scale, with zero e-values suppressed by adding a small number (1e-190) to each e-value. The horizontal red dotted line indicates the e-value cut-off we implemented and the vertical red dotted line indicates the length cut-off (e-value < 1e-20 and alignment length of 100, respectively) to decrease false positive hits. **(b) The fraction of reads with high quality BLAST hits (e-value < 1e-20) increases as a function of read accuracy.** We binned the data according to mean read accuracy (bin width of 0.02) and calculated the fraction of reads within each bin that had a high quality BLAST hit for the January and March runs separately (blue and orange points, respectively). The number of reads in each bin is indicated above or below each point (in thousands). There is a clear positive correlation between mean accuracy and the likelihood of a high-quality BLAST hit, reaching almost 40% for very high quality reads (accuracy greater than 92.5%).

MEGAN assigned reads to a wide range of eukaryotic taxa. To conservatively infer

taxon presence, we first reclassified MEGAN species-level assignments to the level of

genus. However, after this, many clear false positive assignments remained (e.g. hippo

and naked mole rat). These matches were generally short and of low identity. To reduce

such false positive taxon inferences, we used information from reads assigned to the

genera *Rattus* (rat) and *Mus* (mouse). We inferred that the reads assigned to *Rattus*

(2,696 reads in total) were true positive genus-level assignments and that the reads

assigned to *Mus* (2,798 reads in total) were false positive genus-level assignments (and

not true positive *Mus*-derived reads). Although rats are known to prey on mice

(Bridgman et al. 2013), if this had occurred, we would expect that (1) the ratio of mouse

to rat reads would be higher in the subset of rats that had predated mice; (2) in those

same rats, the percent identity of the reads assigned to *Mus* would be higher than in

rats that had not predated mice. However, we found that the ratio of mouse to rat reads

was similar for all rats, and there was no evidence of higher percent identities for *Mus*

reads from rats that had higher ratios.

Notably, the mean percent identity values of the best BLAST hits for these two groups

of reads differed substantially, with *Rattus* reads having a median identity of 86.4%, and

Mus 81.0% (**Fig. 4A**). The mean percent identity for *Rattus* reads corresponds very well

to that expected given the mean quality scores of the reads (assuming the true

sequence of the read is 100% identical to *Rattus*, 86.4% identity corresponds to a mean

quality score of 8.7).

There was also a clear difference in the alignment lengths: the median ratio of

alignment length to read length was 0.57 for *Rattus* and 0.52 for *Mus* (**Fig. 4B**).

Importantly, the majority of diet items have percent identities that overlap with the

Rattus reads, and alignment length to read length ratios that exceed the *Rattus* reads,

suggesting that many diet taxa assignments are correct down to the level of genus.

However, to minimise the rates of false positive taxon assignments of diet items, we

implemented cut-offs based on the characteristics of the *Mus*- and *Rattus*-assigned

reads. For genus-level assignment, we required at least 82.5% identity and an

alignment length to read length ratio of at least 0.55. These cut-offs exclude 88% of the

reads falsely assigned to *Mus*, instead assigning them correctly to one taxon level

higher, the Family *Muridae*.

Fig. 4. Distributions of percent identity and length for alignments of reads

matching *Rattus* (rat), *Mus* (mouse), and diet items. (a) Percent identity for

alignments of rat (*Rattus*) and diet items is much higher than for mouse (*Mus*).

Histograms are shown for the percent identity of the alignment of the top BLAST hit with

the read. *Mus* matches show a clear shift to the left (lower percent identity) as

compared to *Rattus* and diet items. Although different genera, *Mus* and *Rattus* are in

the same family (*Muridae*). The dotted lines indicate the cut-offs that we implemented

for inferring reads as belonging to a specific genus (above 82.5% identity) or family

(above 77.5% identity). **(b) Ratios of alignment lengths to read lengths of rat**

(*Rattus*) and diet items are higher than for mouse (*Mus*). This plot is analogous to

that in (a). The dotted line indicates the cut-off that we implemented for inferring reads

as belonging to a specific genus (above 0.55).

For family-level assignments, we required 77.5% identity, an alignment length to read

length ratio of at least 0.1, and a total alignment length of at least 150 bp. Using higher

cut-offs for the ratio of alignment length to read length excluded a large number of likely

true positive taxa for which only short mtDNA or rDNA database sequences were

present in the databases. For all other read-to-taxon assignments, we placed the read

at the level of Order, or used the taxon level assigned by MEGAN. Using these cut-offs,

16% of all reads were classified at the Genus level; 71% were classified at the Family-

level or below; 89% were classified at the Order-level or below; and 98% were classified

at the Phylum-level or below.

Even though we imposed these relatively stringent filters, we observed that a small

number of likely false positive taxa remained. Most were single reads with short

alignments: *Poeciliidae* (177 bp); *Salmonidae* (172 bp); *Cyprinodontiformes* (140 bp and

177 bp); and *Octopodidae* (151 bp). The exception to this were three reads from two

rats matching *Buthidae* (scorpions), which had alignment lengths of 762 bp, 664 bp, and

298 bp. It is unlikely these are true positives, and instead that these rats predated

harvestmen, a closely related sister taxon within *Arachnida*. Despite the presence of

these false positive taxa, to ensure we could identify the majority of taxa at high

resolution (i.e. family-level), we did not further increase the stringency of our filters.

Identification of stomach contents

Within each rat, a wide variety of plant, animal, and fungal families and orders were

discernible, ranging from two to 25 orders per rat (mean 8.7) (**Fig. 5**). Plants were the

primary diet item, with the largest fraction of rats consuming four predominant orders:

Poales (grasses), *Fabales* (legumes), *Arecales* (palms), and *Araucariales* (podocarps).

The dominance of plant matter (fruits and seeds) in rat diets has been established

previously (Sweetapple and Nugent 2007; Riofrío-Lazo and Páez-Rosas 2015). Animal

taxa made up a smaller component of each rat's diet, with *Insecta* dominating:

Hymenoptera, *Coleoptera*, *Lepidoptera* (moths and butterflies), *Blattodea*

(cockroaches), *Diptera* (flies), and *Phasmatodea* (stick insects). In addition

Stylommatophora (slugs and snails) were present in substantial numbers (**Fig. 6A** and

6B). Fungi were only a small component, although several families and orders were

present: *Sclerotiniaceae* (plant pathogens), *Saccharomycetaceae* (budding yeasts),

Mucorales (pin molds), *Russulales* (brittlegills and milk-caps), and *Chytotheriales* (black

yeasts). Finally, for many rats, a substantial proportion of the stomach contents were

parasitic worms (primarily *Spirurida* (nematodes) and *Hymenolepididae* (tapeworms)).

Close examination of this data suggested that specific families and orders were

overrepresented in the diets of rats from particular locations. For example, six out of

eight rats from the native estuarine bush habitat (OB) consumed *Arecaceae*, while only

one in the restored wetland area (LB) did. All three rats that consumed *Phaseanidae*

were from the native estuarine habitat (OB). All five rats that consumed *Solanales* were

from the restored wetland area. These patterns suggested that it might be possible to

use diet components alone to pinpoint the habitat from which each rat was sampled.

Fig. 5. Numbers of taxa in individual rats. Each boxplot indicates the range of

families (left boxes) or orders (right boxes) consumed by each rat in each location (OB:

Okura Bush; LB: Long Bay Park; WP: Waitakere Park). The numbers for individual rats

(eight per location) are plotted in grey.

Fig. 6. Proportions of taxa in the diets of individual rats. Reads assigned to taxa at the family and order level are shown in the top and bottom panels, respectively. Each row corresponds to a single rat, with the proportions of reads for that rat assigned to each family or order indicated in shades of blue and yellow. Reads that were not assigned to a specific family or order are on the right end of the figure. The families and orders have been sorted so that the most common diet components appear on the left. Only the 55 most common families are shown. Note that the colour gradations presented on the scale are not linear.

nMDS and CAP analysis by location

In order to determine if diet composition of the rats differed consistently between locations, we first performed an unconstrained analysis using nMDS on taxa assigned at the family level. Using family rather than order or genus provides a balance between how precisely we identify the taxon of diet item (genus, family, order), and whether we assign a taxon at all. While family-level assignments are less precise than genus-level, only 16% of all reads were classified at the genus level, while 71% were classified at the family level.

The family-level unconstrained ordination (nMDS; **Fig. 7a**) showed no obvious grouping of rats with respect to the locations, indicating that locations did not correspond to the predominant axes of variation among the diets. However, a constrained ordination analysis (CAP; **Fig. 7b**) identified axes of variation that distinguished the diets of rats from different locations. The CAP axes correctly classified the locations of 19 out of 24 (79%) rats using a leave-one-out procedure.

The families that best characterised the three locations (i.e., had the greatest within-location SIMPER scores) varied among locations. WP had average Bray-Curtis within-location similarity of 13%; mostly attributable to *Hymenolepidae* (accounting for 51% of the within-group similarity), *Solanaceae* (11%), and *Fabaceae* (11%). The average similarity for OB was 21%, with the greatest contributing taxa being *Arecaceae* (33%), *Poaceae* (23%), *Fabaceae* (9%), and *Phasianidae* (8%). The average similarity for WP was 24%, with the greatest contributing taxa being *Poaceae* (72%) (**Table S4**). The families with the largest correlations with the first two principal coordinates and thus responsible for the separation between groups) were *Arecaceae*, *Podocarpaceae*, *Phasianidae*, and *Pinaceae* (**Fig. 7c**).

Fig. 7. Unconstrained nMDS (a) and constrained CAP (b) ordinations of the diets of rats from three locations. Both ordinations were based on Bray-Curtis dissimilarities of square root transformed proportions of reads attributed to each family. The locations were a native estuarine bush (OB, orange); a restored marine wetland (LB, purple); and a native forest (WP, light blue). The CAP ordination is repeated in panel **(c) with the rats omitted to show the Pearson correlations between families and the two cap axes.** The eight families with the strongest correlations are shown, indicating the taxa most strongly associated with each location.

Discussion

Accuracy and sensitivity

Here we have shown that using a simple long-read metagenomic DNA sequencing approach allows rapid classification of rat diet components. We expect that this technique for dietary analysis could be applied to a variety of animals and sample types, including samples that require less invasive collection (e.g. fecal matter). The sensitivity of this approach will likely improve as the accuracy and yield of Oxford Nanopore sequencing increases, and as the diversity of taxa in genomic sequence databases increases. Several aspects of the data support this.

405 First, we note that we did not find BLAST hits for the majority of reads. This is partially

406 due the relatively low accuracy of the Oxford Nanopore sequencing platform at the time

407 we preformed this analysis (approximately 87%). However, the fraction of reads yielding

408 hits in the database increased substantially for higher quality reads, approaching 40%

409 for very high-quality reads (**Fig. 3b)**. Other factors also likely reduced the numbers of

410 BLAST hits, such as the paucity of genome sequence data for many taxa. This is

411 convincingly illustrated by comparing across taxa the fraction of genomic hits to

412 mitochondrial or rDNA sequence hits: although genome data exists for only a few taxa,

413 mtDNA and rDNA sequence data are present in the database for the vast majority of

414 animal and plant genera.

415 For animals with sequenced genomes in the database, we found that only a minority of

416 reads mapped to mitochondrial or rDNA sequences (e.g. 32% of *Rattus* hits and 22% of

417 *Anoplophora* hits). For plants with sequenced genomes, the fraction of mtDNA or rDNA-

418 matching reads was even lower: between 4% (*Aegilops*) and 6% (*Triticum* and

419 *Solanum*). For genera with little or no genomic sequence, the vast majority of matches

420 were solely to mtDNA, rDNA, or microsatellite loci: 90% of *Phoenix* (date palm) hits; all

421 *Helix* (snail) and *Rhaphidophora* **(**cave weta) hits. All *Artioposthia* (New Zealand

422 flatworm) hits were to rDNA. These results indicate that for genera with no genomic

423 sequence data, we have underestimated the actual number of sequences from that

424 taxon by approximately three- to twenty-fold (for animals and plants, respectively).

425 As the species sampling of genomic databases increases (Lewin et al. 2018), the taxon-

426 level precision of this method will improve. Given the current rate of genomic

427 sequencing, with careful sampling, the vast majority of multicellular plant and animal

428 families (and even genera) will likely have at least one type species with a sequenced

429 genome within the next decade. Continued advancement in sequence database search

430 algorithms as compared to current methods (Wood and Salzberg 2014; Kim et al. 2016;

431 Nasko et al. 2018) should considerably decrease the computational workload necessary

432 to find matching sequences.

To decrease biases in genomic databases, some previous studies have performed

metagenomic classification using mitogenome data alone (Srivathsan et al. 2016; Paula

et al. 2016). Using such methods, they found between 0.004% and 0.008% of all

metagenomic reads matched mitogenomes from diet taxa. Limiting database searches

to mitogenomes partially ameliorates biases in terms of taxon field in terms of taxon

representation (i.e. most taxa will have similar levels of genomic representation in the

databases). However, it considerably decreases diet resolution given that for some

taxa, only a small percentage of sequence reads derive from the mitochondria as

opposed to the nuclear genome.

It is important to note that our interest in diet also includes resolving relative biomass

and relative numbers of each prey species, neither of which necessarily correlate well

with the amount of DNA (either mitochondrial or nuclear) purified from a sample. Even a

simple correction for the fraction of reads matching mitochondrial versus nuclear

genomes is difficult, as different plant and animal tissues differ considerably in the

relative amounts of mitochondrial versus nuclear DNA (e.g. leaf versus fruit).

Methodological advantages

We found that rats consumed many soft-bodied species (e.g. mushrooms, flat worms,

slugs, and lepidopterans) that would be difficult to identify using visual inspection of

stomach contents. Capturing such a wide variety of taxa would be difficult to quantify

using metabarcoding methods, as there are no universal 18S or COI universal primers

capable of amplifying sequences in all these taxa. While it might be possible to use

primer sets targeted at different phyla or orders, quantitatively comparing diet

components across these using sequences amplified with different primer sets is

extremely difficult due to differences in primer binding and PCR efficiency.

The nanopore MinION-based sequencing method used in this simple metagenomic

approach has several advantages. Compared to other high throughput sequencing

technologies (e.g. Illumina, IonTorrent, or PacBio), there is no initial capital investment

required to use the platform. On a per-sample basis, data generation is inexpensive

(approximately $150 USD per barcoded sample, and approximately half this price if

reagents are purchased in bulk). Library preparation and sequencing can be extremely

rapid, going from DNA sample to sequence in less than two hours (Zaaijer et al. 2017).

Furthermore, the sequencing platform itself is highly portable. As the cost of nanopore-

based sequencing continues to decrease (both per sample and per base pair), it should

become possible to use molecular methods for routine ecological monitoring of species

presence or absence in field settings, without significant investment in infrastructure

(Kamenova et al. 2017). Finally, we suggest that our approach of standardising the read

counts by sample, followed by an optional transformation such as square root and

dissimilarity-based multivariate ordination, offers a useful analytical pipeline for

analysing metagenomic diet-composition data.

Conclusion

Here we have shown that a rapid long-read metagenomic approach is able to accurately

characterise diet taxa at the family-level and distinguish between the diets of rats

according to the locations from which they were sourced, with almost 80% out-of-

sample classification success. This approach also identifies the taxa responsible for

such patterns. This information may be used to guide conservation efforts toward

specific areas and habitats in which native species are most at risk from this highly

destructive introduced predator.

Acknowledgements

This work was supported by a Massey University Research Fund to NF, a Marsden

Fund Grant (15-MAU-136) to JD and Marsden Fund Grant (MAU1703) to OS. Thanks to

Friends of Okura Bush, Mary Stewart from Auckland Council, and Gillian Wadams and

the volunteers at the Waitakere Ranges for collecting rat samples. Sample collection

was performed under (Auckland Council Permit to Undertake Research WS1064).

Data Accessibility

Sequence data are available in ENA (accession number PRJEB27647)

Author Contributions

WP, JD, NF, and OS conceived the project. WP performed the stomach dissections.

WP and NF optimised the genomic DNA isolation and library preparation. NF performed

the nanopore sequencing. GB and OS processed and performed quality control on the

sequencing data. WP and OS performed the sequence classification. WP, AS, NF, and

OS analysed the data. WP, NF, AS, and OS wrote the paper, with input from all authors.

Supplemental Tables

Table S1. Read numbers and total base pairs for each barcode in the January
sequencing run.

Rat	Total reads	Total Mbp	Mean length
OB2	19.907	14.62	734
WP11	10,164	9.63	947
WP5	8237	6.78	823
LB7	7548	7.04	933
OB13	3644	3.63	995
WP9	2954	2.4	814
OB5	2850	2.06	721
WP8	2801	2.32	827
LB6	2531	1.6	632
OB7	2473	1.87	756
LB5	1641	1.16	705
LB3	1554	0.99	636
None	16673	13.01	781
Total	82,977	67.1	

501   **Table S2**. Read numbers and total base pairs for each barcode in the March

502   sequencing run.

Barcode	Total	Total Mbp	Mean
LB1	17.820	9.21	517
LB8	16,923	13.13	776
WP2	10,511	7.00	666
LB4	8684	4.92	567
OB11	5689	3.40	598
WP10	1563	0.99	633
OB12	1479	0.89	604
WP12	1309	0.78	596
LB2	1127	0.76	676
WP3	637	0.73	1141
OB14	541	0.37	683
OB10	435	0.24	555
None	29,432	21.33	725
Total	96,150	63.75	

504 **Table S3.** SIMPER analysis of family contributions to group similarities. Abbreviations:

505 Av.Abund: average abundance; Av.Si: average similarity; Sim/SD: similarity over

506 standard deviation; Percent.contr: percent contribution.

Family	Av.Abun	Av.Si	Sim/SD	Percent.cont	Group
Hymenolepidida	3.37	6.87	0.34	51.2	LB
Solanaceae	1.57	1.48	0.34	11.1	LB
Fabaceae	1.74	1.41	0.44	10.5	LB
Arecaceae	2.86	7.11	1	33.4	OB
Poaceae	2.87	4.82	0.55	22.7	OB
Fabaceae	1.17	1.98	0.51	9.3	OB
Phasianidae	1.79	1.67	0.34	7.9	OB
Poaceae	5.08	17.61	0.62	72.1	WP

509 **Table S4.** SIMPER analysis of family contributions to group dissimilarities. The families
510 are sorted by percent contribution to group dissimilarity. Abbreviations: Abund.Group1
511 and Abund.Group2: abundance of each family in groups 1 and 2 (see Group 1 and 2
512 columns). Av.Diss: average dissimilarity; Diss/SD: Dissimilarity over standard deviation;
513 Percent.contr: percent contribution.

Family	Abund.Group1	Abund.Group2	Av.Diss	Diss/SD	Percent.contr	Group1	Group2
Poaceae	1.95	5.08	15.15	1.04	16.74	LB	WP
Poaceae	2.87	5.08	11.29	1.26	13.78	OB	WP
Hymenolepididae	3.37	0.48	10.8	0.73	11.93	LB	WP
Hymenolepididae	3.37	0.29	9.37	0.79	10.32	LB	OB
Poaceae	1.95	2.87	8.37	1.1	9.22	LB	OB
Arecaceae	0.05	2.86	6.99	1.41	7.7	LB	OB
Arecaceae	2.86	1.31	5.92	1.29	7.23	OB	WP
Fabaceae	1.74	1.05	6.14	0.67	6.78	LB	WP
Podocarpaceae	0	2.38	5.34	0.83	5.9	LB	WP
Podocarpaceae	0.71	2.38	4.82	0.99	5.88	OB	WP
Fabaceae	1.74	1.17	4.87	0.81	5.37	LB	OB
Fabaceae	1.17	1.05	4.31	0.84	5.26	OB	WP
Phasianidae	1.79	0	4.03	0.7	4.92	OB	WP
Phasianidae	0	1.79	4.28	0.72	4.71	LB	OB
Physalopteridae	1.63	0.18	3.79	0.59	4.63	OB	WP
Physalopteridae	0.46	1.63	4.15	0.65	4.57	LB	OB
Solanaceae	1.57	0	3.95	0.65	4.37	LB	WP
Solanaceae	1.57	0	3.56	0.67	3.92	LB	OB
Helicidae	0.87	0.47	3.46	0.52	3.82	LB	WP
Oleaceae	1.06	0.25	3.09	0.44	3.77	OB	WP
Arecaceae	0.05	1.31	3.4	0.65	3.75	LB	WP
Piperaceae	1.42	0	3.36	0.65	3.71	LB	WP
Helicidae	0.87	0.82	3.33	0.58	3.66	LB	OB
Cerambycidae	1.41	0.18	2.99	0.77	3.65	OB	WP
Cerambycidae	0	1.41	3.07	0.74	3.38	LB	OB
Oleaceae	0	1.06	3.06	0.41	3.37	LB	OB
Piperaceae	1.42	0	3.04	0.67	3.35	LB	OB
Onchocercidae	0.39	1.04	3.02	0.77	3.33	LB	OB
Geoplanidae	0.8	0.48	2.64	0.49	3.22	OB	WP
Thelaziidae	0.9	0.64	2.85	0.77	3.14	LB	OB
Helicidae	0.82	0.47	2.54	0.61	3.1	OB	WP
Onchocercidae	1.04	0	2.49	0.66	3.04	OB	WP
Formicidae	0.44	1.17	2.39	1.04	2.92	OB	WP

Sclerotiniaceae	0	1.28	2.63	0.57	2.91	LB	WP
Sclerotiniaceae	0	1.28	2.3	0.57	2.81	OB	WP
Formicidae	0	1.17	2.46	0.94	2.72	LB	WP
Rosaceae	0.92	0.3	2.13	0.58	2.61	OB	WP
Ectobiidae	0.63	0.34	2.24	0.5	2.48	LB	WP
Thelaziidae	0.9	0	2.13	0.56	2.36	LB	WP
Geoplanidae	0	0.8	2.04	0.37	2.25	LB	OB
Rosaceae	0	0.92	1.99	0.5	2.19	LB	OB
Numididae	0.78	0	1.68	0.56	2.05	OB	WP
Myrtaceae	0.66	0	1.8	0.38	1.99	LB	WP
Thelaziidae	0.64	0	1.53	0.51	1.86	OB	WP
Ranunculaceae	0.66	0	1.36	0.37	1.5	LB	WP

Datafile S1. Table of read BLAST hits and assigned MEGAN taxa with reads

reclassified at the family or order level by filtering on read length to alignment length

ratio and percent identity.

Datafile S2. Table of read BLAST hits and assigned MEGAN taxa with no filters

applied.

Supplemental Figures

Fig S1. Biplots of read lengths and qualities for each barcode in the January and

March runs.

Fig S2. Correlation of read accuracy with alignment characteristics. (a) Read

accuracy is positively correlated with the percent identity of the top BLAST hit. Points

show a subsample of reads; orange line indicates a running median; red dotted line is

the y=x line, which is expected if accuracy corresponds exactly to percent identity. (b)

Read accuracy and alignment length show no relationship. (c) Read accuracy and the

ratio of read length to alignment length are positively correlated.

References

1. Anderson, Marti J., and Trevor J. Willis. 2003. "Canonical Analysis Of Principal Coordinates: A Useful Method Of Constrained Ordination For Ecology." *Ecology* 84 (2): 511–25.

2. Basha, Walaa A., Andrew T. Chamberlain, Moushira E. Zaki, Wafaa A. Kandeel, and Nagui H. Fares. 2016. "Diet Reconstruction through Stable Isotope Analysis of Ancient Mummified Soft Tissues from Kulubnarti (Sudanese Nubia)." *Journal of Archaeological Science: Reports* 5 (February): 71–79.

3. Bridgman, L. J., J. Innes, Craig Gillies, Neil Fitzgerald, and Carolyn M. King. 2013. "Do Ship Rats Display Predatory Behaviour towards House Mice?" *Animal Behaviour* 86 (2)): 257–68.

4. Brown, K. P., H. Moller, J. Innes, and P. Jansen. 2008. "Identifying Predators at Nests of Small Birds in a New Zealand Forest." *The Ibis* 140 (2): 274–79.

5. Brownlee, A., and W. Moss. 1961. "The Influence of Diet on Lactobacilli in the Stomach of the Rat." *The Journal of Pathology* 82 (2): 513–16.

6. Carreon-Martinez, L., and D. D. Heath. 2010. "Revolution in Food Web Analysis

553   and Trophic Ecology: Diet Analysis by DNA and Stable Isotope Analysis."
554   *Molecular Ecology* 19 (1): 25–27.
555

556   7. Clarke, K. R. 1993. "Non-Parametric Multivariate Analyses of Changes in
557   Community Structure." *Austral Ecology* 18 (1): 117–43.
558

559   8. Clarke, K. R., and R. N. Gorley. 2015. "PRIMER v7: User Manual/Tutorial."
560   PRIMER-E, Plymouth.
561

562   9. Clarke, K. R., and R. H. Green. 1988. "Statistical Design and Analysis for a
563   'Biological Effects' Study." *Marine Ecology Progress Series* 46: 213–26.
564

565 10. Clarke, K. Robert, K. Robert Clarke, Paul J. Somerfield, and M. Gee Chapman.
566   2006. "On Resemblance Measures for Ecological Studies, Including Taxonomic
567   Dissimilarities and a Zero-Adjusted Bray–Curtis Coefficient for Denuded
568   Assemblages." *Journal of Experimental Marine Biology and Ecology* 330 (1): 55–
569   80.
570

571 11. Daniel, M. J. 1973. "Seasonal Diet Of The Ship Rat (Rattus Rattus) In Lowland
572   Forest In New Zealand." *Proceedings of the New Zealand Ecological Society* 20:
573   21–30.
574

575 12. Deagle, Bruce E., J. Paige Eveson, and Simon N. Jarman. 2006. "Quantification
576   of Damage in DNA Recovered from Highly Degraded Samples--a Case Study on
577   DNA in Faeces." *Frontiers in Zoology* 3 (August): 11.
578

579 13. Deagle, Bruce E., Simon N. Jarman, Eric Coissac, François Pompanon, and
580   Pierre Taberlet. 2014. "DNA Metabarcoding and the Cytochrome c Oxidase
581   Subunit I Marker: Not a Perfect Match." *Biology Letters* 10 (9).
582   https://doi.org/10.1098/rsbl.2014.0562.
583

584 14. Diamond, J. M., and C. R. Veitch. 1981. "Extinctions and Introductions in the
585   New Zealand Avifauna: Cause and Effect?" *Science* 211 (4481): 499–501.
586

587 15. Dowding, John E., and Elaine C. Murphy. 2001. "The Impact of Predation by
588   Introduced Mammals on Endemic Shorebirds in New Zealand: A Conservation
589   Perspective." *Biological Conservation* 99 (1): 47–64.
590

591 16. Dunlap, M., and J. R. Pawlik. 1996. "Video-Monitored Predation by Caribbean
592   Reef Fishes on an Array of Mangrove and Reef Sponges." *Marine Biology* 126
593   (1): 117–23.
594

595 17. Gibbs, George W. 1998. "Why Are Some Weta (Orthoptera: Stenopelmatidae)
596   Vulnerable yet Others Are Common?" *Journal of Insect Conservation* 2 (3-4):

161–66.

18. Gillies, Craig, and Mick Clout. 2003. "The Prey of Domestic Cats (Felis Catus) in Two Suburbs of Auckland City, New Zealand." *Journal of Zoology* 259 (3): 309–15.

19. Hobson, Keith A. 1987. "Use of Stable-Carbon Isotope Analysis to Estimate Marine and Terrestrial Protein Content in Gull Diets." *Canadian Journal of Zoology* 65 (5): 1210–13.

20. Horáková, Z., C. H. Zierdt, and M. A. Beaven. 1971. "Identification of Lactobacillus as the Source of Bacterial Histidine Decarboxylase in Rat Stomach." *European Journal of Pharmacology* 16 (1): 67–77.

21. Huson, Daniel H., Alexander F. Auch, Ji Qi, and Stephan C. Schuster. 2007. "MEGAN Analysis of Metagenomic Data." *Genome Research* 17 (3): 377–86.

22. Huson, Daniel H., Sina Beier, Isabell Flade, Anna Górska, Mohamed El-Hadidi, Suparna Mitra, Hans-Joachim Ruscheweyh, and Rewati Tappu. 2016. "MEGAN Community Edition - Interactive Exploration and Analysis of Large-Scale Microbiome Sequencing Data." *PLoS Computational Biology* 12 (6): e1004957.

23. Jain, Miten, Hugh E. Olsen, Benedict Paten, and Mark Akeson. 2016. "The Oxford Nanopore MinION: Delivery of Nanopore Sequencing to the Genomics Community." *Genome Biology* 17 (1): 239.

24. Jarman, S. N., B. E. Deagle, and N. J. Gales. 2004. "Group-Specific Polymerase Chain Reaction for DNA-Based Analysis of Species Diversity and Identity in Dietary Samples." *Molecular Ecology* 13 (5): 1313–22.

25. Jarman, S. N., N. J. Gales, M. Tierney, P. C. Gill, and N. G. Elliott. 2002. "A DNA-Based Method for Identification of Krill Species and Its Application to Analysing the Diet of Marine Vertebrate Predators." *Molecular Ecology* 11 (12): 2679–90.

26. Kamenova, S., T. J. Bartley, D. A. Bohan, J. R. Boutain, R. I. Colautti, I. Domaizon, C. Fontaine, et al. 2017. "Chapter Three - Invasions Toolkit: Current Methods for Tracking the Spread and Impact of Invasive Species." In *Advances in Ecological Research*, edited by David A. Bohan, Alex J. Dumbrell, and François Massol, 56:85–182. Academic Press.

27. Kim, Daehwan, Li Song, Florian P. Breitwieser, and Steven L. Salzberg. 2016. "Centrifuge: Rapid and Sensitive Classification of Metagenomic Sequences." *Genome Research* 26 (12): 1721–29.

28. King, R. A., D. S. Read, M. Traugott, and W. O. C. Symondson. 2008. "Molecular Analysis of Predation: A Review of Best Practice for DNA-Based Approaches." *Molecular Ecology* 17 (4): 947–63.

29. Leray, Matthieu, Joy Y. Yang, Christopher P. Meyer, Suzanne C. Mills, Natalia Agudelo, Vincent Ranwez, Joel T. Boehm, and Ryuji J. Machida. 2013. "A New Versatile Primer Set Targeting a Short Fragment of the Mitochondrial COI Region for Metabarcoding Metazoan Diversity: Application for Characterizing Coral Reef Fish Gut Contents." *Frontiers in Zoology* 10 (June): 34.

30. Lewin, Harris A., Gene E. Robinson, W. John Kress, William J. Baker, Jonathan Coddington, Keith A. Crandall, Richard Durbin, et al. 2018. "Earth BioGenome Project: Sequencing Life for the Future of Life." *Proceedings of the National Academy of Sciences of the United States of America* 115 (17): 4325–33.

31. Li, Dongyao, Haiqin Chen, Bingyong Mao, Qin Yang, Jianxin Zhao, Zhennan Gu, Hao Zhang, Yong Q. Chen, and Wei Chen. 2017. "Microbial Biogeography and Core Microbiota of the Rat Digestive Tract." *Scientific Reports* 8 (April): 45840.

32. Major, H. L., I. L. Jones, M. R. Charette, and A. W. Diamond. 2007. "Variations in the Diet of Introduced Norway Rats (*Rattus norvegicus*) Inferred Using Stable Isotope Analysis." *Journal of Zoology* 271 (4): 463–68.

33. Maurice, Corinne F., Sarah C. L. Knowles, Joshua Ladau, Katherine S. Pollard, Andy Fenton, Amy B. Pedersen, and Peter J. Turnbaugh. 2015. "Marked Seasonal Variation in the Wild Mouse Gut Microbiota." *The ISME Journal* 9 (11): 2423–34.

34. McIntyre, Alexa B. R., Rachid Ounit, Ebrahim Afshinnekoo, Robert J. Prill, Elizabeth Hénaff, Noah Alexander, Samuel S. Minot, et al. 2017. "Comprehensive Benchmarking and Ensemble Approaches for Metagenomic Classifiers." *Genome Biology* 18 (1): 182.

35. Nasko, Daniel J., Sergey Koren, Adam M. Phillippy, and Todd J. Treangen. 2018. "RefSeq Database Growth Influences the Accuracy of K-Mer-Based Species Identification," 1–21.

36. Paula, Débora P., Benjamin Linard, David A. Andow, Edison R. Sujii, Carmen S. S. Pires, and Alfried P. Vogler. 2015. "Detection and Decay Rates of Prey and Prey Symbionts in the Gut of a Predator through Metagenomics." *Molecular Ecology Resources* 15 (4): 880–92.

37. Paula, Débora P., Benjamin Linard, Alex Crampton-Platt, Amrita Srivathsan,

Martijn J. T. N. Timmermans, Edison R. Sujii, Carmen S. S. Pires, Lucas M. Souza, David A. Andow, and Alfried P. Vogler. 2016. "Uncovering Trophic Interactions in Arthropod Predators through DNA Shotgun-Sequencing of Gut Contents." *PloS One* 11 (9): e0161841.

38. Pawluczyk, Marta, Julia Weiss, Matthew G. Links, Mikel Egaña Aranguren, Mark D. Wilkinson, and Marcos Egea-Cortines. 2015. "Quantitative Evaluation of Bias in PCR Amplification and next-Generation Sequencing Derived from Metabarcoding Samples." *Analytical and Bioanalytical Chemistry* 407 (7): 1841–48.

39. Kerry P. Brown. 1997. "Predation at Nests of Two New Zealand Endemic Passerines; Implications for Bird Community Restoration." *Pacific Conservation Biology: A Journal Devoted to Conservation and Land Management in the Pacific Region* 3 (2): 91.

40. Pereira, Rui P. A., Jorg Peplies, Ingrid Brettar, and Manfred G. Hoefle. 2018. "Impact of DNA Polymerase Choice on Assessment of Bacterial Communities by a Legionella Genus-Specific next-Generation Sequencing Approach." *bioRxiv*. https://doi.org/10.1101/247445.

41. Pierce, Graham J., and Peter R. Boyle. 1991. "A Review of Methods for Diet Analysis in Piscivorous Marine Mammals." *Oceanography and Marine Biology: An Annual Review* 29: 409–86.

42. Riofrío-Lazo, Marjorie, and Diego Páez-Rosas. 2015. "Feeding Habits of Introduced Black Rats, *Rattus rattus*, in Nesting Colonies of Galapagos Petrel on San Cristóbal Island, Galapagos." *PloS One* 10 (5): e0127901.

43. Russell, James C., John G. Innes, Philip H. Brown, and Andrea E. Byrom. 2015. "Predator-Free New Zealand: Conservation Country." *Bioscience* 65 (5): 520–25.

44. Soininen, Eeva M., Alice Valentini, Eric Coissac, Christian Miquel, Ludovic Gielly, Christian Brochmann, Anne K. Brysting, et al. 2009. "Analysing Diet of Small Herbivores: The Efficiency of DNA Barcoding Coupled with High-Throughput Pyrosequencing for Deciphering the Composition of Complex Plant Mixtures." *Frontiers in Zoology* 6 (August): 16.

45. Srivathsan, Amrita, Andie Ang, Alfried P. Vogler, and Rudolf Meier. 2016. "Fecal Metagenomics for the Simultaneous Assessment of Diet, Parasites, and Population Genetics of an Understudied Primate." *Frontiers in Zoology* 13 (April): 17.

46. Srivathsan, Amrita, John C. M. Sha, Alfried P. Vogler, and Rudolf Meier. 2015.

"Comparing the Effectiveness of Metagenomics and Metabarcoding for Diet Analysis of a Leaf-Feeding Monkey (Pygathrix Nemaeus)." *Molecular Ecology Resources* 15 (2): 250–61.

47. Stringer, Ian A. N., Suzanne M. Bassett, Megan J. McLean, Jay McCartney, and G. Richard Parrish. 2003. "Biology and Conservation of the Rare New Zealand Land Snail Paryphanta Busbyi Watti (Mollusca, Pulmonata)." *Invertebrate Biology: A Quarterly Journal of the American Microscopical Society and the Division of Invertebrate Zoology/ASZ* 122 (3): 241–51.

48. Sweetapple, Peter J., and Graham Nugent. 2007. "Ship Rat Demography and Diet Following Possum Control in a Mixed Podocarp—hardwood Forest." *New Zealand Journal of Ecology* 31 (2): 186–201.

49. Tedersoo, Leho, Sten Anslan, Mohammad Bahram, Sergei Põlme, Taavi Riit, Ingrid Liiv, Urmas Kõljalg, et al. 2015. "Shotgun Metagenomes and Multiple Primer Pair-Barcode Combinations of Amplicons Reveal Biases in Metabarcoding Analyses of Fungi." *MycoKeys* 10: 1.

50. Towns, David R., Charles H. Daugherty, and Alison Cree. 2001. "Raising the Prospects for a Forgotten Fauna: A Review of 10 Years of Conservation Effort for New Zealand Reptiles." *Biological Conservation* 99 (1): 3–16.

51. Wood, Derrick E., and Steven L. Salzberg. 2014. "Kraken: Ultrafast Metagenomic Sequence Classification Using Exact Alignments." *Genome Biology* 15 (3): R46.

52. Zaaijer, Sophie, Assaf Gordon, Daniel Speyer, Robert Piccone, Simon Cornelis Groen, and Yaniv Erlich. 2017. "Rapid Re-Identification of Human Samples Using Portable DNA Sequencing." *eLife* 6 (November). https://doi.org/10.7554/eLife.27798.