Genetic screening for mutants with altered seminal root numbers in hexaploid wheat

using a high-throughput root phenotyping platform

Oluwaseyi Shorinola[*, †], Ryan Kaye [§], Guy Golan[‡], Zvi Peleg[‡], Stefan Kepinski [§] and Cristobal

Uauy[†].

*Bioscience Eastern and Central Africa - International Livestock Research Institute, Nairobi,

PO Box 30709, Kenya.

[§] Centre for Plant Sciences, Faculty of Biological Sciences, University of Leeds, Leeds,

LS2 9JT, UK.

[‡]The Robert H. Smith Institute of Plant Sciences and Genetics in Agriculture, The Hebrew

University of Jerusalem, Rehovot 7610001, Israel.

[†] John Innes Centre, Norwich Research Park, Norwich, NR4 7UH, UK.

Running title: Identification of wheat root mutants.

Keyword: Wheat, Root, TILLING, Mutations, Forward Genetics

Corresponding Authors:

Oluwaseyi Shorinola, Bioscience Eastern and Central Africa - International Livestock

Research Institute, Nairobi, P O Box 30709, Kenya, +254204223881, o.shorinola@cgiar.org.

Cristobal Uauy, John Innes Centre, Norwich Research Park, Norwich, NR4 7UH, UK, +44-

(0)1603-450195, cristobal.uauy@jic.ac.uk.

ABSTRACT

Roots are the main channel for water and nutrient uptake in plants. Optimisation of root architecture provides a viable strategy to improve nutrient and water uptake efficiency and maintain crop productivity under water-limiting and nutrient-poor conditions. We know little, however, about the genetic control of root development in wheat, a crop supplying 20% of global calorie and protein intake. To improve our understanding of the genetic control of seminal root development in wheat, we conducted a high-throughput screen for variation in seminal root number using an exome-sequenced mutant population derived from the hexaploid wheat cultivar Cadenza. The screen identified seven independent mutants with homozygous and stably altered seminal root number phenotypes. One mutant, Cadenza0900, displays a recessive extra seminal root number phenotype, while six mutants (Cadenza0062, Cadenza0369, Cadenza0393, Cadenza0465, Cadenza0818 and Cadenza1273) show lower seminal root number phenotypes most likely originating from defects in the formation and activation of seminal root primordia. Segregation analysis in F_2 populations suggest that the phenotype of Cadenza0900 is controlled by multiple loci whereas the Cadenza0062 phenotype fits a 3:1 mutant:wild-type segregation ratio characteristic of dominant single gene action. This work highlights the potential to use the sequenced wheat mutant population as a forward genetic resource to uncover novel variation in agronomic traits, such as seminal root architecture.

INTRODUCTION

The 1960's "Green Revolution" demonstrated the impact that changes to plant architecture in major crops like wheat and rice can have on increasing food production (Hedden 2003). While the Green Revolution focused on improving shoot architecture, it did not optimise root architecture, in part because selection was primarily for performance under management regimes involving high rates of fertiliser application (Lynch 2007). In addition to providing anchorage, the root is the main channel for water and nutrient uptake in crops and serves as an interface for symbiotic interaction with the soil microbiome. Roots are often considered as the hidden and neglected other-half of plant architecture and have not been a direct target for selection during early wheat domestication and in modern wheat breeding programmes (Waines and Ehdaie 2007).

In many environments, water availability is the main factor defining crop rotations and performance. Projections on future climate predict more variable weather events relating to the timings and intensity of precipitations which could negatively affect food security (Cattivelli et al. 2008; Rojas et al. 2019). Optimising root system architecture (RSA) for improved nutrient and water uptake under these uncertain scenarios provides a rational approach to help achieve future food and nutrition security.

The wheat root system is comprised of two main root types, seminal (embryonic) and nodal (post-embryonic) roots, that develop at different times. (Manske and Vlek 2002). As the first root type that emerges, seminal roots are entirely responsible for nutrient and water uptake in seedlings. Seminal roots are therefore important for seedling vigour and early plant establishment which also determines competitiveness against weeds. Nodal roots on the other hand are shoot-borne and develop soon after tillering to provide anchorage and support resource uptake especially during the reproductive stage of wheat growth (Manske and Vlek 2002).

Despite their early establishment, seminal roots remain functionally active through to the reproductive stage and may grow up to 2 m in length (Manschadi et al. 2013; Araki and Iijima 2001). They have also been shown to have similar nutrient uptake efficiency as nodal roots in wheat and contribute to yield potential especially under conditions of low soil moisture where nodal roots may not grow (Weaver and Zink 1945; Sebastian et al. 2016). Given their importance, seminal root traits, such as number and angle, have been linked to adaptive responses under water limiting conditions (Manschadi et al. 2008; Cane et al. 2014; Golan et al. 2018). Steep seminal root angle has been associated with increased soil water exploration at depth which is beneficial in drought condition where topsoil moisture is depleted (Richard et al. 2015; Olivares-Villegas et al. 2007; Manschadi et al. 2008).

Seminal roots develop from the root primordia in the embryo of a germinating wheat seed. There is genetic variation among wheat genotypes for the number of seminal roots that develop, this can range from three to six seminal roots per plant dependent on cultivar (Araki and Iijima 2001; Robertson et al. 1979; Golan et al. 2018). Typically, the seminal root system consists of a primary root that emerges first followed by two pairs of secondary seminal roots that emerge sequentially. A sixth seminal root may also develop in some wheat varieties. A few quantitative trait loci (QTL) have been identified to underlie variation in seminal root number in wheat germplasm (Atkinson et al. 2015; Maccaferri et al. 2016; Ren et al. 2012; Sanguineti et al. 2007; Ma et al. 2017; Iannucci et al. 2017). However, many of these QTLs are defined to broad intervals which makes their genetic dissection difficult and their use in breeding limited. Unlike other cereals (e.g. rice, maize), only one gene controlling root system architecture (RSA), *VERNALIZATION1* (*VRN1*; Voss-Fels et al. 2018), has been identified in wheat.

This delay in identifying genetic loci controlling root traits is most likely due to a series of factors which makes genetic analyses in wheat difficult. Bread wheat is a hexaploid plant with

a relatively large (16 Mb) and repeat-rich (>85%) genome comprised of three homoeologous sub-genomes (B, A and D). High sequence similarity in the coding regions of these sub-genomes results in high levels of genetic redundancy that mask the phenotypic effects of underlying natural variation for many traits, including RSA traits (Uauy et al. 2017; Borrill et al. 2015). Also, the "out-of-sight" nature and extreme phenotypic plasticity of roots under native field conditions makes root phenotyping difficult, cumbersome and time-consuming (Atkinson et al. 2019).

The use of induced variation has proven useful to uncover novel phenotypes and dissect genetic pathways underlying complex phenotypes in plants (Parry et al. 2009). Our current understanding of the genetic determinants regulating root development in many cereals have almost entirely stemmed from the isolation and characterisation of mutants defective in one or more RSA traits (Reviewed in Coudert et al. 2010; Hochholdinger et al. 2018; Marcon et al. 2013). Despite this potential, the use of mutant populations to study the genetic control of root development in wheat has not hitherto been exploited. The recent development of an *in-silico* platform for the rapid identification of mutations in 1,200 ethyl methanesulfonate (EMS) mutagenized lines in the UK hexaploid wheat cultivar 'Cadenza' now makes large-scale reverse and forward genetic investigation of traits more feasible in wheat (Krasileva et al. 2017). Progress has also been made on the root phenomics front, with the development of fast, low-cost, and flexible two-dimensional (2D) root phenotyping pipelines with sufficient throughput for phenotyping large populations (Selvara et al. 2013; Atkinson et al. 2019; Adeleke et al. 2019).

Taking advantage of these new developments, we implemented a relatively high-throughput root phenotyping pipeline to conduct a forward genetic screen for variation in seminal root number using a subset of the exome-sequenced Cadenza mutant population. From this work,

136　we describe the identification and characterisation of novel hexaploid bread wheat mutants

137　with decreased and increased numbers of seminal roots.

MATERIALS AND METHODS

Germplasm

Mutant population for primary screens: A hexaploid wheat mutant population was previously developed by EMS treatment of the UK bread wheat cultivar Cadenza (Krasileva et al. 2017; Rakszegi et al. 2010). In this study, we used 645 exome-sequenced mutants from this population (Krasileva et al. 2017). These mutant lines were selected based on the criteria that they show greater than 90% germination rate during a seed multiplication that was conducted in the field. To obtain homogenous phenotypes and reduce the variation from segregating mutations, single spikes harvested from field-grown M_4 plants were individually threshed and derived M_5 seeds (40 - 50 seeds) from each spike were divided for use in forward genetic Screen A and Screen B described below (Figure 1).

Germplasm for mutant validation and characterisation: Three additional M_4 spikes (containing M_5 seed) from the same field grown samples as those used for the primary screens (Figure 1) were used to further validate the phenotypes of the seven mutants identified in the primary screens (described in results). The spikes were individually threshed and M_5 seeds from each spike were phenotyped separately. Two validated mutants (Cadenza0900 and Cadenza0062) were further selected for genetic characterisation: M_5 plants were grown to maturity for cross-pollination with wild-type Cadenza to generate F_1 hybrids which were subsequently self-pollinated to generate F_2 progenies. M_5 plants of the selected mutants were also self-pollinated to generate M_6 seeds to characterise the stability of their phenotypes in the subsequent generation.

High-throughput Seminal Root Phenotyping

Two similar and independent screens (Screen A and B) were conducted concurrently on

subsets of the Cadenza mutant population to identify lines with altered seminal root numbers

(Figure 1). All 645 lines were phenotyped at high-throughput in Screen A, while only the first

385 of the 645 lines (in numerical order) were phenotyped in Screen B due to more limited

throughput. For mutant lines phenotyped in both screens, M_5 seeds from the same field-

grown M_4 spike were used in both screens as described in the germplasm section above.

Screen A: This screen was carried out at the John Innes Centre, UK, using a custom 2D root

phenotyping platform based on the protocol described by Atkinson et al. (2015) with some

modifications to increase the throughput from 360 to 1,800 seedlings per run. This screen

also took seed size effect on RSA traits into consideration. For each mutant line, M_5 seeds

from a M_4 single spike were first size-stratified into large, medium or small seed by passing

the seeds through two sets of calibrated graduated sieves with 2.8 mm and 3.35 mm mesh

sizes. Large-sized seeds were collected above the 3.35 mm sieve, medium-sized seeds

collected between the 2.8 mm and 3.35 mm sieves, and small-sized seeds were collected

below the 2.8 mm sieve. Each mutant line was thus designated as either being Large (524

lines), Medium (119 lines) or Small (2 lines) based on the size group where most of its seeds

were defined. Only seeds representative of each mutant line size classification were used.

The two mutants with small seed size were phenotyped for variation in seminal root number

compared to wild-type Cadenza, but were not included in the analysis of seed size effect on

RSA due to the small sample size. Seeds (15 per mutant line) were surface sterilized by

rinsing in 5% (v/v) Sodium Hypochlorite (Sigma Aldrich, UK) for 10 mins and were rinsed with

water three times before being imbibed in 1.75 mL of water for 5 days at 4°C to ensure

uniform germination. Of these, 10 seeds per mutant were placed crease facing down into

individual growth pouches made from a sheet of germination paper (21.5 cm x 28 cm; Anchor

Paper Company, St Paul, MN, USA) clipped to a black polythene sheet (22 cm x 28 cm, 75

μm thickness, Cransford Polythene LTD, Suffolk, UK) using an acrylic rod and 18 mm fold

clip (Figure 2A). The growth pouches were suspended in an upright position in plastic storage

boxes (120 cm x 27 cm x 36 cm, Really Useful Product, West Yorkshire, UK) with 60

pouches per box (Figure 2B). The sides of the box were covered in black plastic sticky back

cover film to block out light from the roots of the developing seedlings. Each box was filled

with 10 L of half-strength Hoaglands growth solution containing (Hoagland and Arnon 1950):

$NH_4H_2PO_4$, 0.6 g; $Ca(NO_3)_2$, 3.3 g; $MgSO_4$, 1.2 g; KNO_3, 1.0 g; H_3BO_3, 14.3 mg; Cu_2SO_4, 0.4

mg; $MnCl_2(H_2O)_4$, 9.1 mg; MoO_3, 0.1 mg; $ZnSO_4$, 1.1 mg; $KHCO_3$, 2.0 g, Ferric Tartrate, 2.8

g. The base of each pouch was suspended in the growth solution to supply nutrients to the

developing seedling through capillary action. A randomised complete block design was

adopted with each line replicated across 10 different boxes (blocks). The phenotyping boxes

were placed in a controlled environment room under long day conditions with 16h light (250–

400 mmol) at 20°C, 8h darkness at 15°C and at 70% relative humidity (Figure 2C). After 7

days of growth, pouches were taken out of the phenotyping box; placed on a copy stand and

the black plastic sheet covering the germination paper was gently pulled back to reveal the

roots. Images of the roots were taken with a Nikon D3400 DSLR Camera fitted to the copy

stand (Figure 2D). Phenotyping of the mutant population was done over 12 experiments with

60 lines (59 mutants and a Cadenza control) phenotyped per experiment. Mutants of the

same seed-size group were phenotyped together – large sized mutants in experiment 1 - 9

and small/medium sized mutants in experiment 10 -12. Cadenza seeds representative of the

seed-size group for each experiment were used as controls. The 12 experiments were

completed across five rounds of phenotyping with 2 -3 experiments (120 – 180 lines) set up

per round. For 96% of the mutants examined, we successfully imaged the roots of 8 -10

plants. In the remain 4% of the mutants, only 3 - 7 plants could be images due to poor seed

germination or poor seedling growth on the pouch. In total, 6,240 (6,127 mutant and 113

Cadenza) plants were phenotyped. The same phenotyping set-up was used for the M_5

validation experiments and to characterise M_6, F_1 and F_2 progenies of the two mutants

selected for genetic characterisation.

Screen B: A second screen was conducted in parallel at the University of Leeds, UK, using

commercial CYG paper pouches (Mega-International, Minnesota, USA). Due to the more

limited throughput of this screen, only the first 385 of the 645 mutant lines (in numerical

order) were examined in this screen. For each mutant line, 10 visually uniform M_5 seeds were

selected and placed onto moist filter paper in a 90 mm round petri dish. Petri dishes were

wrapped in aluminium foil to exclude light and placed at 4°C for 2 days, seeds were placed

crease side down into individual CYG seed germination pouches with the bottom removed to

allow wicking of growth solution from a reservoir of media. Pouches were wrapped in

aluminium foil in batches of five to exclude light from the roots and were placed upright in a

reservoir of full strength Hoaglands No 2 growth solution ($NH_4H_2PO_4$, 115.03 mg; $Ca(NO_3)_2$,

656.4 mg; $MgSO_4$, 240.76 mg; KNO_3, 606.6 mg; H_3BO_3, 2.86 mg; Cu_2SO_4, 0.08 mg;

$MnCl_2(H_2O)_4$, 1.81 mg; MoO_3, 0.016 mg; $ZnSO_4$, 0.22 mg; Ferric Tartrate, 5 mg. per Litre).

Pouches were placed in long day conditions (as in screen A) and were grown for 5 days

before roots were imaged. For imaging, pouches were placed onto a copy stand; the front of

the pouch carefully removed, and the root system imaged using a Sony Cybershot DSC-

RX100. Plants were screened in rounds of twenty lines with a total of 200 plants per round.

Image Analysis

High-resolution images captured from the phenotyping were pre-processed (rotated, cropped

and compressed) using ImageJ (https://imagej.nih.gov/index.html) and Caesium image

compressor before being processed in RootNav (Pound et al. 2013). Captured root

architectures were imported into a RootNav viewer database for measurement of RSA traits

using standard RootNav functions (Figure 2E).

Anatomical Characterisation of Seminal Root Primordia

Embryos of mutants validated to have reduced seminal root number phenotypes were

examined using the method described by Golan et al. (2018). In brief: embryos from mature

dry grains were fixed in FAA solution (10% formaldehyde, 5% acetic acid, 50% ethanol, and

35% distilled water by volume) overnight and dehydrated at room temperature in a graded

ethanol series (30 min each, in 50%, 70%,90%, 95%, and 100% ethanol). Then, embryos

were cleared in xylene, embedded in paraffin and sectioned (5 µM) using a microtome (Leica

Biosystems, Germany). Cross sections were de-paraffinized with histoclear, rehydrated and

stained with Harris Hematoxylin. A stereo microscope (SZX16, Olympus, Tokyo, Japan) was

used for imaging.

Statistical Analysis

All statistical analyses were performed in R 3.5.1 (R Core Team 2018) and Minitab 17

statistical software. Statistically significant seminal root number difference in the primary

screening experiments (Screen A and B) was determined by ANOVA using a Dunnett's

comparison within each phenotyping experiment with the Cadenza plants in each experiment

used as control. Adjusted probability values of $P < 0.05$ were considered statistically

significant. Statistically significant root architectural difference in the validation experiment as

well as in the M_6 and F_1 phenotyping experiments were based on Student's t-test comparison

of individual spike/line to the Cadenza control. A Chi-square test was used to examine the

goodness of fit of the segregation pattern observed in the F_2 progenies to patterns expected

for single recessive or dominant gene action.

Data Availability

Seeds of mutant lines reported in this study can be ordered through the SeedStor site at

www.seedstor.ac.uk. Supplemental materials (tables and figures) are available at *FigShare*.

Table S1 contains information on mutants with significantly different seminal root number

phenotypes to Cadenza from Screen A. Table S2 contains information on all the mutants

phenotyped in both Screen A and B. Figure S1 shows the relationship between seminal root

number and total root length in Screen A. Figure S2 and Figure S3 show representative

images and embryo size measurement of mutants with validated altered seminal root number

phenotypes, respectively. All the root images from Screen A (6,240 images) including the

original RootNav measurements for different root traits are publicly available on Zenodo

(https://doi: 10.5281/zenodo.3270726) for download and reuse.

277   **RESULTS**

Identifying induced variation for seminal root number in hexaploid wheat

We implemented a 2D root phenotyping pipeline suitable for large-scale phenotyping at a

throughput of 1,800 seedlings per run (Figure 1 and 2). Using this platform, we performed a

forward genetic screen for variation in seminal root number using 645 seed-size stratified

(small, medium and large; see Methods) M_5 mutants from the exome-sequenced Cadenza

mutant population (Krasileva et al. 2017). In our screen, Cadenza mainly displayed five

seminal roots (4.9 ± 0.05) including a primary seminal root (SR_1), as well as first ($SR_{2,3}$) and

second ($SR_{4,5}$) pairs of seminal roots (Figure 3A). We observed variation in seminal root number in the mutant population, with seminal root number ranging from 1 to 7 in individual plants and mean seminal root number per mutant (with $n \geq 4$ plants per mutant) ranging from 2.9 to 5.9.

Within the mutant lines, seed size groups (large and medium) showed significant difference in seminal root number ($P < 0.0001$) and total root length ($P < 0.0001$). Mutants with large and medium-sized grains had an average seminal root number of 4.7 and 4.4, respectively, and total root length of 481 mm and 324 mm, respectively (Figure 3B-C). In the Cadenza wild-type, significant differences between seed size groups were only observed for total root length. There were only two mutant lines with small seed size and these were not included in the analysis of seed size effect on seminal root traits. Across all mutant lines, we observed a significant positive correlation ($R = 0.56$, $P < 0.0001$) between the number of seminal roots and the total root length in the population (Figure S1).

Parallel to this, we phenotyped a subset of these Cadenza mutants for similar root traits in an independent screen (Screen B; Figure 1). Given the lower throughput of this approach (See Screen 2 in Methods), only 385 out of the 645 mutant lines were screened. We observed a significant positive correlation between seminal root number measurements in Screen A and Screen B ($R = 0.63$; $P < 0.0001$; Figure 3D). The heritability estimate of the seminal root measurement across the two screens was 0.77 suggesting a strong heritable genetic effect in the determination of seminal root number in the Cadenza mutant population.

We first assessed the statistical significance from the mutants studied in Screen A. Dunnett's multiple comparison identified 52 mutants (8 % of mutants in Screen A) with significantly different number of seminal roots relative to the Cadenza control across the three seed size groups (Table S1). Five of these mutants had significantly higher number of seminal roots

with mean seminal root number ranging from 5.7 to 5.9 and modal seminal root number of 6

per mutant (Table S1). The higher seminal root phenotype is mainly driven by the

development of an extra root, hereafter referred to as SR_6. The remaining 47 mutants

showed significantly lower number of seminal roots with mean seminal root number per

mutant of 2.9 to 4.1 and modal seminal root number between 3 and 5.

Thirty three out of the 52 significant mutants identified in Screen A were also phenotyped as

part of Screen B. This included four of the five higher root number mutants and 29 lower root

number mutants. Details of individual mutant lines phenotyped in both screens are presented

in Table S2. In Screen B we confirmed the significant phenotype of three of the four higher

root number mutants in common with Screen A; these lines displayed a mean seminal root

number of 5.1 to 5.5 in Screen B. In the case of the 29 lower root number mutants which

were investigated in both screens, we confirmed 20 mutants which displayed fewer numbers

of seminal roots (less than 4 roots) than Cadenza. Based on the results from the two

screens, we selected the three higher and 20 lower root number mutants with consistent

phenotypes in both screens for further phenotypic evaluation.

Altered root number mutants show stable homozygous seminal root number

phenotypes

To validate the 23 selected mutants, we phenotyped M_5 seeds from three additional M_4

spikes (containing M_5 seeds) from the same field-grown bulk as the spike used in the primary

screen (Figure 1). These M_4 spikes originate from successive bulking of multiple M_3 and M_4

plants. Selecting three separate spikes increases the probability of phenotyping plants with

independent background mutations thereby providing robust biological replications to

examine the stability of the mutations effects and segregation patterns (homozygous or heterozygous).

For seven of the selected mutants including one higher root number (Cadenza0900) and six lower root number mutants (Cadenza0062, Cadenza0369, Cadenza0393, Cadenza0465, Cadenza0818, and Cadenza1273), we observed the altered seminal root number phenotype in the three additional spikes phenotyped (Figure 4, Figure S2). This suggests that the phenotypes of these mutants are consistent across sibling lines and controlled by mutations that were most likely homozygous in the original single M_2 plant from which the population was derived. For the rest of the 16 mutants, we did not consistently observe the altered seminal root number phenotypes in all three additional spikes. These might represent lines with segregating phenotypes arising from heterozygous mutations in the initial M_2 plants or false-positive selection in the primary screens; these lines were not studied further.

We further characterized the seven validated mutants from 1 to 7 days post germination (dpg) to examine when the phenotype was first visible and to identify the seminal root type (primary, first pair or second pair) defective in these mutants (Figure 5). Cadenza showed fully emerged primary root (SR_1) at 1 dpg, while the first ($SR_{2,3}$) and second pairs ($SR_{4,5}$) of seminal roots emerged at 3 and 5 dpg, respectively. Similar to Cadenza, Cadenza0369, Cadenza0393, Cadenza0465, Cadenza0818, and Cadenza1273 with lower root counts also developed the primary and first pair of seminal roots at 1 dpg and 3 dpg, respectively, but are defective in the development of the second pair of seminal roots ($SR_{4,5}$ or only SR_5). Contrary to this, Cadenza0900 with higher root count showed a faster rate of seminal root development relative to Cadenza with the primary, first and second pair of seminal roots having emerged by 3 dpg and an extra sixth root emerged at 7 dpg. Cadenza0062 showed a strong dormancy phenotype and was not included in this experiment.

Embryo imaging of the lower root number mutants points to defects in primordia development and growth

Seminal roots emerge from the activation of root primordia that form in the embryo of a developing seed. The primary (SR_1) and secondary ($SR_{2,3}$ and $SR_{4,5}$) seminal roots emerge from primordia formed in the central portion and sides of the embryo, respectively. Although uncommon, the sixth seminal roots (SR_6 like in Cadenza0900) are known to emerge from a primordium located in the midpoint between the primordia of the second pair of secondary seminal root (Hoshikawa 1964). Importantly, differential activation of root primordia of $SR_{4,5}$ has been shown to account for lower numbers of seminal roots in some wild wheat species (Golan et al., 2018). We therefore examined if the seminal root phenotypes of the lower root number mutants (Cadenza0062, Cadenza0369, Cadenza0393, Cadenza0465, Cadenza0818, and Cadenza1273) lacking either SR_5 or $SR_{4,5}$ originate from defects in primordia development and/or https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4169152/the failure of developed root primordia to activate to become seminal roots. Cadenza (WT) consistently develops five fully formed root primordia (Figure 6). All the lower root number mutants examined showed altered root primordia development compared to Cadenza, with $SR_{4,5}$ (or only SR_5) primordia either absent or reduced in size (Figure 6). In addition, primordia activity was altered in the mutants, as all mutants had greater number of primordia compared to the number of roots observed in the M_5 seedlings (Figure 5, Table 1). These measurements provide an initial indication that the defects in the lower root count mutants most likely originate from a combination of both lower number and activity of root primordia in these lines. Embryo size was also significantly lower in four of the six mutants (Cadenza0393, Cadenza0465, Cadenza0818, and Cadenza1273; Figure S3), but it is not clear from these

380　results if the smaller embryo size of these mutants contributes to their lower seminal roots

381　number phenotypes.

Genetic characterisation of the altered root number phenotypes

To understand the transgenerational stability and mode of inheritance of the altered root

number mutant phenotype, we characterized M_6 plants of the higher root number mutant

(Cadenza0900) and one lower root number mutant (Cadenza0062), as well as F_1 hybrids

derived from crosses of these mutants to Cadenza. M_6 progenies of Cadenza0900 showed

significantly (P <0.0001) increased number of seminal roots compared to Cadenza and no

significant difference to the phenotypes of it M_5 parents (Figure 7) with an average root

number of 5.73 and more than 73% of the plants having six roots. F_1 hybrids of Cadenza0900

x Cadenza all had five seminal roots like Cadenza (P = 0.19), but significantly lower than their

M_5 plants (P < 0.0001; Figure 7). This suggests that the Cadenza0900 phenotype originates

from a recessive mutation or is caused by a combination of loci segregating independently.

The M_6 plants of Cadenza0062 showed a significantly lower number of seminal roots (P

<0.0001) compared to Cadenza with a mean root number of 2.6 (Figure 7). The seminal root

numbers of the M_6 progenies of Cadenza0062 were also slightly but significantly (P = 0.04)

lower than the M_5 plant, most probably due to smaller variation in the seminal root number

phenotype in the M_6 lines. With an average root number of 3.0 (Figure 7), the F_1 hybrid of

Cadenza0062 x Cadenza showed a significantly (P <0.0001) lower number of seminal roots

compared to Cadenza but a non-significant difference to the original M_5 plants (Figure 7).

This suggests that the Cadenza0062 phenotype is caused by a dominant mutation(s). Unlike

the M_5 plants, the Cadenza0062 x Cadenza F_1 did not show any reduced germination,

suggesting that the dormancy phenotype of Cadenza0062 segregates independently of its

altered root number.

To gain further insight into the genetic architecture underlying higher and lower root count

phenotypes, we further characterized F_2 progenies of the Cadenza0900 and Cadenza0062

crosses to Cadenza. We used the chi-square test-statistic to test the goodness of fit of the

inheritance pattern of Cadenza0900 and Cadenza0062 phenotypes to those consistent with

segregation of single recessive and single dominant traits, respectively. The phenotype of

Cadenza0900 F_2 progenies (238) was not consistent with the expected 3:1 wild-type:mutant

phenotype segregation ratio of a single recessive gene (χ^2 = 52.71, P <0.0001), suggesting

that multiple genes may be responsible for the Cadenza0900 phenotype. In contrast, the

segregation pattern of the Cadenza0062 F_2 population (51 plants) is consistent with the 3:1

mutant:wildtype segregation ratio expected of a single dominant gene (χ^2 = 0.53, P =

0.4669), suggesting that the Cadenza0062 phenotype is caused by a single dominant gene.

418   **DISCUSSION**

Use of sequenced mutant population to characterise RSA genes in wheat

Mutant analyses have played a key role in the identification of genes controlling key stages of

root development. For instance, most of the genes identified to control root architecture in

maize were identified via mutant analyses (Hochholdinger et al. 2018; Marcon et al. 2013).

These include *RTCS, RTCL, RUM1* and *BIGE1* which display seminal root phenotypes

(Taramino et al. 2007; Suzuki et al. 2015; von Behrens et al. 2011; Xu et al. 2015). Despite

the buffering effect of genetic redundancy that often masks single homoeolog mutations in

polyploid wheat (Borrill et al. 2015), our study highlights the usefulness of forward genetic

screens to identify heritable variation for root development traits in wheat. These results are

also consistent with recent examples of dominant mutations being identified in forward

screens of the exome-sequenced populations (Harrington et al. 2019; Mo et al. 2018).

The use of a sequenced mutant population in this study also provided the opportunity to

examine the presence of mutations in candidate genes from other species. For example, the

phenotypes of the lower root number mutants (Cadenza0062, Cadenza0369, Cadenza0393,

Cadenza0465, Cadenza0818, and Cadenza1273) are similar to the phenotypes of maize *rtcs*

and *rtcl* mutations (Taramino et al. 2007), and their orthologous rice mutants (Liu et al. 2005;

Inukai et al. 2005). *In-silico* examination of the coding regions in these mutants on the

Ensembl Plant database revealed that Cadenza1273 contains a functional mutation in

TraesCS4D01G312800, one of the three wheat homoeologs of *RTCS, RTCL* and

ARL1/CRL1. Cadenza1273 harbours a G765A mutation in *TraesCS4D01G312800* producing

a premature termination codon which results in a truncated 265 amino acid (aa) protein

instead of the 289 aa native protein. Further molecular and genetic characterisation will be

required to test if the G765A mutation in Cadenza1273 is responsible for its phenotype.

However, this exemplifies the power of combining the sequenced mutant information with

known candidate genes and a now fully annotated wheat genome (Appels et al. 2018). The

other lower root number mutants do not contain any functional EMS mutations in the three

wheat homoeologs of *RTCS, RTCL* and *ARL1/CRL1* and therefore likely represent new

variation controlling seminal root development in cereals*.*

Similarly, the higher seminal root number phenotype of Cadenza0900 is similar to the

phenotype of maize *bige1* (Suzuki et al. 2015) mutants. However, unlike *bige1* which is

monogenic, we show that the phenotype of Cadenza0900 is most likely conditioned by more

than one mutation. Also, *in-silico* examination of mutations in Cadenza0900 show that it does

not harbour any mis-sense or non-sense mutations in the coding sequences of the three

wheat orthologs of *BIGE1* (*TraesCS4A02G350200, TraesCS5B02G522900 and*

TraesCS5D02G521600) thereby reducing the likelihood that the Cadenza0900 phenotype

originates from mutations of the wheat *BIGE1* gene. It is, however, important to note that

these *in-silico* investigations are restricted to mutations in the coding region of the wheat

genome and we cannot rule out that mutations in promoter regions of these candidate genes

might be responsible for some of the altered root number mutants identified.

Relationship between grain size and seminal root traits

Size stratification of seeds in our study allowed an examination of the relationship between

grain size and root architecture. We observed a positive effect of grain size on root length

and number. Irrespective of the genetic background (wild-type Cadenza or mutant

background), large-sized grain showed longer root length compared to medium-sized grain

consistent with the rationale that bigger grains have greater carbohydrate reserves in their

endosperm to support faster root elongation. We also noticed a weak but positive effect of

grain size on the number of seminal roots developed in the mutant population. Grain size

effect on seedling traits, including seminal root traits, can be attributed to constituent

components – embryo and/or endosperm (Bremner et al. 1963; Meyer 1976). Interestingly,

four of the lower root number mutants (Cadenza0393, Cadenza0465, Cadenza0818 and

Cadenza1273) have smaller embryos compared to Cadenza. However, without further

genetic analyses, it is premature to conclude that the small embryo of these mutants directly

affects their seminal root number phenotypes. Indeed, grain size only accounts for a small

proportion (3.4%) of the total variance in root number in our study, indicating that seed size

per se is not a major determinant of root number. This is further underlined by the fact that

differently sized wild-type Cadenza seeds show similar root number averages (Figure 3B).

Although informative, the qualitative stratification of grain size (large, medium and small)

adopted in this study does not allow a quantitative modelling of grain size effect on root traits.

We propose that a finer calibration and partitioning of grain size measurement into

constituent parameters (width, length, height) and tissue (embryo and endosperm)

components (Brinton and Uauy 2019) will allow for a finer understanding of the effects of

these seed size components on root architecture.

High-throughput platforms enables population-scale screening

The high-throughput afforded by the 2D platform used in Screen A in this study enabled the

screening of a large mutant population (645 lines). Based on a previous 2D phenotyping

pipeline implemented by Atkinson et al. (2015), our platform increases the throughput from

360 to 1,800 seedlings while maintaining a similar phenotyping rate (~4.5 minutes per plant).

A similar high-throughput phenotyping platform was also recently adapted by (Adeleke et al.

2019) with a maximum phenotyping capacity of 672 plants. It is also worth noting that our

high-throughput platform is relatively inexpensive and easy to implement using widely-

available low-cost materials: storage boxes, germination paper, plastic sheets and paper

binder clips. The entire phenotyping pipeline costed £5,570 to phenotype 6,240 plants:

£3,280 to set-up the infrastructure and £2,090 recurring cost (paper and nutrient solution

reagents). Besides the initial set-up cost, this represents a recurring cost of £0.33 per plant,

making it cheaper than commercially available CYG germination papers. Also, we believe an

important feature of this set-up is the modularity of phenotyping it offers, that is, the ability to

set up experiment in small individual phenotyping units (boxes, Figure 2B and C) which can

be used as experimental blocks. This modularity also allows flexibility of phenotyping scale

as the experiments can be scaled up or down by adjusting the number of boxes used.

More than half of the processing time in Screen A was spent on image analysis. This included semi-automated analysis of RSA using RootNav as individual root tips need to be selected first before the root outline is automatically extracted. This processing time can be further improved by adopting the latest deep machine-learning algorithm recently applied to root image processing allowing fully automated extraction of root outline. This innovation will drastically reduce the time taken for image processing without compromising on accuracy (Pound et al. 2017).

Despite the high plasticity (e.g. dynamic response to varying water, nutrient and environmental conditions) associated with root traits, we obtained a high heritability estimate for seminal root number measurements across the two screens, similar to previous estimates (Maccaferri et al. 2016; Ma et al. 2017). While this high heritability may be due to the controlled hydroponic environment used in these experiments (Figure 2), it nonetheless demonstrates that seminal root number is a stable phenotype under strong genetic control and can be targeted for selection to improve RSA in wheat breeding programmes. There is also evidence to suggest that seminal root number phenotypes observed in hydroponic conditions are transferrable to soil conditions (Richard et al. 2015; Watt et al. 2013) and might therefore be useful under field conditions especially during the vegetative phase. We are currently evaluating the mutant lines under field conditions to determine this.

Developmental and genetic characterisation of seminal root formation

Natural variation in wheat seminal root number has been shown to originate from defects in root primordia development in the embryo and/or differential activation of developed primordia to form fully emerged seminal roots (Golan et al. 2018). Our anatomical characterisation of intact non-imbibed seeds of the lower root number mutants highlight a

tendency for these mutants to develop less than five root primordia, whereas wild-type

Cadenza plants consistently develop five root primordia. In addition, these mutants form even

fewer numbers of seminal roots than the root primordia they developed, suggesting that

some of the primordia developed in these mutants might be dormant/inactive or arrested

soon after activation. Together these results highlight both primordia development and

activation as two important development check points in the formation of seminal root in

wheat.

All the altered root number mutants develop SR_1 and $SR_{2,3}$ but show defects in the

development of $SR_{4,5}$ as in the lower root number mutants, or develop an extra root (SR_6) as

in Cadenza0900. This suggests that the development of the different seminal root types is

under distinct genetic control. We did not recover mutant lines defective in either SR_1 nor

$SR_{2,3}$: this could be due to the fact that defects in these root types might have severe

negative or lethal effects on seedling growth. Golan et al. (2018) show that SR_1 and $SR_{2,3}$

contribute more to water uptake in durum wheat than $SR_{4,5}$ under well-watered conditions. It

is however possible that $SR_{4,5}$ and indeed SR_6 may contribute significantly to nutrient and

water uptake under resource-limiting conditions where an increase in root surface area

maximises soil exploration. Detailed field physiological evaluation will be required to better

understand the cost-benefit relationship of the altered seminal root phenotypes.

The lower root number mutant Cadenza0062 shows a dominant mode of action with a 3:1

segregation ratio in the F_2 population, characteristic of a monogenic trait. Until this and the

other mutants are independently characterised, we cannot conclude that this dominant

monogenic behaviour is representative of all the lower root number mutants, nor can we rule

out the possibility of some of these mutations being allelic. Unlike Cadenza0062, the extra

seminal root number mutant Cadenza0900 shows a recessive, multigenic phenotype that

suggests a more complex genetic regulation of additional seminal roots in wheat. More

detailed genetic characterisation and mapping will be required to better dissect the genetic

control of these phenotypes.

Outlook

Our work provides a complementary approach to the use of natural variation in dissecting the

genetic control of seminal root development in wheat. The isolation of these mutants

represents an important step in identifying the genetic determinants controlling wheat seminal

root development and growth. These will be followed by extensive genetic characterisation to

map these mutations to defined chromosomal positions and identify the causal gene(s)

underlying their phenotypes. Given the high mutation rate in the Cadenza mutant population

(~33 mutation per Mb; Krasileva et al. 2017), it will also be necessary to reduce the mutation

load in these mutants to enable specific characterisation of each mutation. This can be

achieved through backcrossing each mutant to wild-type Cadenza and selection of F_1

progenies (for dominant mutations like in Cadenza0062) or F_2 progenies (for recessive

mutations like Cadenza0900) that retain the altered seminal root number phenotype (Uauy et

al. 2017). These backcrossed mutants will also provide the background for developing

crosses to examine interactions between the different seminal root number mutations.

ACKNOWLEDGEMENTS

We would like to thank Dr Jonathan Atkinson, Prof Malcolm Bennett, Chloe Riviere and

Guiditta Giordani for technical advice and assistance in the setting up the high-throughput 2D

root screen and Andrew Davis for help with root imaging. This research is supported by the

UK Biotechnology and Biological Sciences Research Council (BBSRC) Designing Future

Wheat programme (BB/P016855/1), a Royal Society FLAIR award (FLR\R1\1918500) to OS,

a BBSRC DTP award to RK, the Chief Scientist of the Israel Ministry of Agriculture and Rural

Development grant (#12-01-0005), and the U.S. Agency for International Development

Middle East Research and Cooperation grant (# M34-037).

LITERATURE CITED

Adeleke, E., R. Millas, W. McNeal, J. Faris, and A. Taheri, 2019 Assessing Root System Architecture of Wheat
Seedlings Using A High-Throughput Root Phenotyping System. *bioRxiv*:677955.
Appels, R., K. Eversole, C. Feuillet, B. Keller, J. Rogers *et al.*, 2018 Shifting the limits in wheat research and
breeding using a fully annotated reference genome. *Science* 361 (6403).
Araki, H., and M. Iijima, 2001 Deep Rooting in Winter Wheat: Rooting Nodes of Deep Roots in Two Cultivars
with Deep and Shallow Root Systems. *Plant Production Science* 4 (3):215-219.
Atkinson, J.A., M.P. Pound, M.J. Bennett, and D.M. Wells, 2019 Uncovering the hidden half of plants using new
advances in root phenotyping. *Current Opinion in Biotechnology* 55:1-8.
Atkinson, J.A., L.U. Wingen, M. Griffiths, M.P. Pound, O. Gaju *et al.*, 2015 Phenotyping pipeline reveals major
seedling root growth QTL in hexaploid wheat. *Journal of Experimental Botany* 66 (8):2283-2292.
Borrill, P., N. Adamski, and C. Uauy, 2015 Genomics as the key to unlocking the polyploid potential of wheat.
New Phytologist 208 (4):1008-1022.
Bremner, P., R. Eckersall, and R. Scott, 1963 The relative importance of embryo size and endosperm size in
causing the effects associated with seed size in wheat. *The Journal of Agricultural Science* 61 (1):139 -
145.
Brinton, J., and C. Uauy, 2019 A reductionist approach to dissecting grain weight and yield in wheat. *Journal of
Integrative Plant Biology* 61 (3):337-358.
Cane, M.A., M. Maccaferri, G. Nazemi, S. Salvi, R. Francia *et al.*, 2014 Association mapping for root
architectural traits in durum wheat seedlings as related to agronomic performance. *Mol Breed* 34
(4):1629-1645.
Cattivelli, L., F. Rizza, F.-W. Badeck, E. Mazzucotelli, A.M. Mastrangelo *et al.*, 2008 Drought tolerance
improvement in crop plants: An integrated view from breeding to genomics. *Field Crops Research* 105
(1):1-14.
Coudert, Y., C. Perin, B. Courtois, N.G. Khong, and P. Gantet, 2010 Genetic control of root development in rice,
the model cereal. *Trends Plant Sci* 15 (4):219-226.
Golan, G., E. Hendel, E.G.E. Méndez, N. Schwartz, and Z. Peleg, 2018 Activation of seminal root primordia
during wheat domestication reveals underlying mechanisms of plant resilience. *Plant, Cell &
Environment* 41 (4):755-766.
Harrington, S.A., N. Cobo, M. Karafiátová, J. Doležel, P. Borrill *et al.*, 2019 Identification of a dominant chlorosis
phenotype through a forward screen of the Triticum turgidum cv. Kronos TILLING
population. *bioRxiv*:622076.
Hedden, P., 2003 The genes of the Green Revolution. *Trends in Genetics* 19 (1):5-9.
Hoagland, D.R., and D.I. Arnon, 1950 The water-culture method for growing plants without soil. *Circular.
California Agricultural Experiment Station* 347 (2):32.

Hochholdinger, F., P. Yu, and C. Marcon, 2018 Genetic Control of Root System Development in Maize. *Trends*
Plant Sci 23 (1):79-88.
Hoshikawa, K., 1964 Studies on the Ripening of Wheat Grain : 5. Formation of pericarp, seed coat and hyaline
layer. *Japanese journal of crop science* 32 (4):333-337.
Iannucci, A., D. Marone, M.A. Russo, P. De Vita, V. Miullo *et al.*, 2017 Mapping QTL for Root and Shoot
Morphological Traits in a Durum Wheat x *T. dicoccum* Segregating Population at Seedling Stage.
International Journal of Genomics 2017:17.
Inukai, Y., T. Sakamoto, M. Ueguchi-Tanaka, Y. Shibata, K. Gomi *et al.*, 2005 Crown rootless1, which is essential
for crown root formation in rice, is a target of an AUXIN RESPONSE FACTOR in auxin signaling. *Plant*
Cell 17 (5):1387-1396.
Krasileva, K.V., H.A. Vasquez-Gross, T. Howell, P. Bailey, F. Paraiso *et al.*, 2017 Uncovering hidden variation in
polyploid wheat. *Proceedings of the National Academy of Sciences* 114 (6):E913-E921.
Liu, H., S. Wang, X. Yu, J. Yu, X. He *et al.*, 2005 ARL1, a LOB-domain protein required for adventitious root
formation in rice. *The Plant Journal* 43 (1):47-56.
Lynch, J.P., 2007 Roots of the Second Green Revolution. *Australian Journal of Botany* 55 (5):493-512.
Ma, J., W. Luo, H. Zhang, X.-H. Zhou, N.-N. Qin *et al.*, 2017 Identification of quantitative trait loci for seedling
root traits from Tibetan semi-wild wheat (Triticum aestivum subsp. tibetanum). *Genome* 60 (12):1068-
1075.
Maccaferri, M., W. El-Feki, G. Nazemi, S. Salvi, M.A. Canè *et al.*, 2016 Prioritizing quantitative trait loci for root
system architecture in tetraploid wheat. *Journal of Experimental Botany* 67 (4):1161-1178.
Manschadi, A.M., G.L. Hammer, J.T. Christopher, and P. deVoil, 2008 Genotypic variation in seedling root
architectural traits and implications for drought adaptation in wheat (Triticum aestivum L.). *Plant and*
Soil 303 (1):115-129.
Manschadi, A.M., G. Manske, and P.L.G. Vlek, 2013 Root Architecture and Resource Acquisition – Wheat as a
Model Plant, pp. 1 -22 in *Plant Roots: The Hidden Half*, edited by A. Eshel and T. Beeckman. CRC Press,
Taylor & Francis Group.
Manske, G.G., and P.L. Vlek, 2002 Root architecture–wheat as a model plant. *Plant roots: The hidden half*
3:249-259.
Marcon, C., A. Paschold, and F. Hochholdinger, 2013 Genetic control of root organogenesis in cereals. *Methods*
Mol Biol 959:69-81.
Meyer, W.S., 1976 Seminal roots of wheat : manipulation of their geometry to increase the availability of soil
water and to improve the efficiency of water use, pp. 217 in *Faculty of Agricultural Science* University
of Adelaide.
Mo, Y., T. Howell, H. Vasquez-Gross, L.A. de Haro, J. Dubcovsky *et al.*, 2018 Mapping causal mutations by
exome sequencing in a wheat TILLING population: a tall mutant case study. *Molecular Genetics and*
Genomics 293 (2):463-477.
Olivares-Villegas, J.J., M.P. Reynolds, and G.K. McDonald, 2007 Drought-adaptive attributes in the Seri/Babax
hexaploid wheat population. *Functional Plant Biology* 34 (3):189-203.
Parry, M.A.J., P.J. Madgwick, C. Bayon, K. Tearall, A. Hernandez-Lopez *et al.*, 2009 Mutation discovery for crop
improvement. *Journal of Experimental Botany* 60 (10):2817-2825.
Pound, M.P., J.A. Atkinson, A.J. Townsend, M.H. Wilson, M. Griffiths *et al.*, 2017 Deep machine learning
provides state-of-the-art performance in image-based plant phenotyping. *GigaScience* 6 (10).
Pound, M.P., A.P. French, J.A. Atkinson, D.M. Wells, M.J. Bennett *et al.*, 2013 RootNav: Navigating Images of
Complex Root Architectures. *Plant Physiology* 162 (4):1802-1814.
R Core Team, 2018 R: A Language and Environment for Statistical Computing. R Foundation for Statistical
Computing, Vienna, Austria.
Rakszegi, M., B.N. Kisgyörgy, K. Tearall, P.R. Shewry, L. Láng *et al.*, 2010 Diversity of agronomic and
morphological traits in a mutant population of bread wheat studied in the Healthgrain program.
Euphytica 174 (3):409-421.
Ren, Y., X. He, D. Liu, J. Li, X. Zhao *et al.*, 2012 Major quantitative trait loci for seminal root morphology of
wheat seedlings. *Molecular Breeding* 30 (1):139-148.

Richard, C.A., L.T. Hickey, S. Fletcher, R. Jennings, K. Chenu *et al.*, 2015 High-throughput phenotyping of
seminal root traits in wheat. *Plant Methods* 11 (1):13.
Robertson, B.M., J.G. Waines, and B.S. Gill, 1979 Genetic Variability for Seedling Root Numbers in Wild and
Domesticated Wheats1. *Crop Science* 19:843-847.
Rojas, M., F. Lambert, J. Ramirez-Villegas, and A.J. Challinor, 2019 Emergence of robust precipitation changes
across crop production areas in the 21st century. *Proceedings of the National Academy of Sciences* 116
(14):6673-6678.
Sanguineti, M.C., S. Li, M. Maccaferri, S. Corneti, F. Rotondo *et al.*, 2007 Genetic dissection of seminal root
architecture in elite durum wheat germplasm. *Annals of Applied Biology* 151 (3):291-305.
Sebastian, J., M.-C. Yee, W. Goudinho Viana, R. Rellán-Álvarez, M. Feldman *et al.*, 2016 Grasses suppress
shoot-borne roots to conserve water during drought. *Proceedings of the National Academy of Sciences*
113 (31):8861-8866.
Selvara, M.G., S. Ogawa, and M. Ishitani, 2013 Root Phenomics-New Windows to Understand Plant
Performance and Increase Crop Productivity. *Journal of Plant Biochemistry & Physiology* 1:116.
Suzuki, M., Y. Sato, S. Wu, B.-H. Kang, and D.R. McCarty, 2015 Conserved Functions of the MATE Transporter
BIG EMBRYO1 in Regulation of Lateral Organ Size and Initiation Rate. *The Plant Cell* 27 (8):2288-2300.
Taramino, G., M. Sauer, S.J. L., D. Multani, X. Niu *et al.*, 2007 The maize (*Zea mays L.*) *RTCS* gene encodes a LOB
domain protein that is a key regulator of embryonic seminal and post-embryonic shoot-borne root
initiation. *The Plant Journal* 50 (4):649-659.
Uauy, C., B.B.H. Wulff, and J. Dubcovsky, 2017 Combining Traditional Mutagenesis with New High-Throughput
Sequencing and Genome Editing to Reveal Hidden Variation in Polyploid Wheat. *Annual Review of
Genetics* 51 (1):435-454.
von Behrens, I., M. Komatsu, Y. Zhang, K.W. Berendzen, X. Niu *et al.*, 2011 Rootless with undetectable
meristem 1 encodes a monocot-specific AUX/IAA protein that controls embryonic seminal and post-
embryonic lateral root initiation in maize. *Plant J* 66 (2):341-353.
Voss-Fels, K.P., H. Robinson, S.R. Mudge, C. Richard, S. Newman *et al.*, 2018 *VERNALIZATION1* Modulates Root
System Architecture in Wheat and Barley. *Mol Plant* 11 (1):226-229.
Waines, J.G., and B. Ehdaie, 2007 Domestication and Crop Physiology: Roots of Green-Revolution Wheat.
Annals of Botany 100 (5):991-998.
Watt, M., S. Moosavi, S.C. Cunningham, J.A. Kirkegaard, G.J. Rebetzke *et al.*, 2013 A rapid, controlled-
environment seedling root screen for wheat correlates well with rooting depths at vegetative, but not
reproductive, stages at two field sites. *Annals of Botany* 112 (2):447-455.
Weaver, J.E., and E. Zink, 1945 Extent and Longevity of the Seminal Roots Of Certain Grasses. *Plant Physiology*
20 (3):359-379.
Xu, C., H. Tai, M. Saleem, Y. Ludwig, C. Majer *et al.*, 2015 Cooperative action of the paralogous maize lateral
organ boundaries (LOB) domain proteins RTCS and RTCL in shoot-borne root formation. *New Phytol*
207 (4):1123-1133.

Figure Legends

Figure 1: Schematic workflow of the forward genetic screens, and validation and

characterisation experiments. Single spikes for each mutant were first phenotyped in the

primary screens (light brown box) and selected mutants were subsequently phenotyped in

the validation and characterisation experiments (dark brown box). The number of lines used

in each experiment are indicated in blue text. Times indicated in Screen A are an

approximation of the time taken to set-up the high-throughput Screen A per phenotyping

batch (~1,800 plants) in 8-hours work days.

Figure 2: Root phenotyping set-up used for screen A: (A) Growth pouch showing blue blotter

germination paper and cover plastic sheet. (B) Phenotyping box containing growth pouch (60

pouches per box) and nutrient solution at the bottom. (C) Root phenotyping in controlled

environment room. (D) Nikon D3400 DSLR Camera mounted on copy stand for root imaging.

(E) Digital extraction of root architecture using RootNav.

Figure 3: Variation in seminal root number in the Cadenza mutant population. (A) Seminal

root architecture of a Cadenza seedling showing the primary root (SR_1) and the first ($SR_{2,3}$)

and second pairs of seminal roots ($SR_{4,5}$). (B-C) Distribution of the seminal root number (B)

and total root length measurements (C) phenotypes observed in Screen A across large and

medium seed-size groups. The mid-line represents the mean of the distribution. The number

of plants in each seed-size group are indicated below each distribution. *** indicates *P*

<0.0001 for statistical comparisons between the seed-size groups. (D) Regression of the

seminal root number phenotypes observed in primary Screens A and B using a linear model.

Only lines phenotyped in both screens are shown. The regression line of the two screens

(dotted diagonal line) is compared to a hypothetical perfect correlation (solid line) between

the screens.

Figure 4: Validated mutants show homozygous seminal root number phenotypes. Seminal

root number distribution in wild-type Cadenza and mutants with validated altered seminal root

number phenotypes across four spikes phenotyped in primary Screen A (spike 1) and the validation experiments (spikes 2 to 4). The number of plants phenotyped from each spike ranged from four to ten.

Figure 5: Characterisation of root development in validated mutants. Temporal characterization of seminal root development in the validated mutants at 1, 3, 5 and 7 days post germination (dpg) shows altered $SR_{4,5}$ and SR_6 seminal root type phenotypes.

Figure 6: Primordia imaging in lower root number mutants. Longitudinal cross sections of mutants with validated lower seminal root number phenotype prior to seed imbibition. The seminal root primordia of Cadenza are marked by asterisks. The inset table shows the primordia count observed in Cadenza and the lower root number mutants. The number of embryos images are indicated in parenthesis beside each line.

Figure 7: Genetic characterisation of the Cadenza0900 and Cadenza0062 seminal root phenotypes. Seminal root number distribution of M_5, M_6 and F_1 progenies of Cadenza0900 and Cadenza0062 mutants are presented alongside the Cadenza control. The number of plants phenotyped for each genotype is indicated below the distribution. * and *** indicates P ≤0.05 and P <0.0001, respectively, and the comparison groups are indicated by solid lines; black comparisons to Cadenza and grey comparisons to M_5 plants. NS indicates non-significant difference.

763 Table 1: Phenotypic summary of validated altered root number mutants with information on
764 the mutation type

TYPE	MUTANT	MEAN (SEM)
Wild-type	Cadenza	4.90 (0.05)
Higher Root Count	Cadenza0900	5.87 (0.05)
Lower Root Count Mutant	Cadenza0393	3.08 (0.04)
	Cadenza0818	3.11 (0.06)
	Cadenza0062	3.18 (0.17)
	Cadenza1273	3.18 (0.07)
	Cadenza0369	3.25 (0.09)
	Cadenza0465	3.25 (0.09)

Cadenza

Cadenza0062

Cadenza0369

Cadenza0393

Line (n)	Mean Primordia Count (StDev)
Cadenza (3)	5.00 (-)
Cadenza0062 (8)	4.50 (0.27)
Cadenza0369 (6)	4.83 (0.16)
Cadenza0393 (10)	3.90 (0.25)
Cadenza0465 (3)	4.33 (0.33)
Cadenza0818 (4)	3.75 (0.48)
Cadenza1273 (8)	4.25 (0.31)

Cadenza0465

Cadenza0818

Cadenza1273

200 μM