Self-assembling Manifolds in Single-cell RNA Sequencing Data

Alexander J. Tarashansky[1], Yuan Xue[1], Pengyang Li[1], Stephen R. Quake[1,2,3], Bo Wang[1,4,*]

[1]Department of Bioengineering, [2]Department of Applied Physics, Stanford University, Stanford,

CA, USA.

[3]Chan Zuckerberg Biohub, San Francisco, CA, USA.

[4]Department of Developmental Biology, Stanford University School of Medicine, Stanford, CA,

USA.

*Correspondence should be addressed to B.W. (wangbo@stanford.edu)

Abstract

Single-cell RNA sequencing has spurred the development of computational methods that enable

researchers to classify cell types, delineate developmental trajectories, and measure molecular

responses to external perturbations. Many of these technologies rely on their ability to detect genes

whose cell-to-cell variations arise from the biological processes of interest rather than

transcriptional or technical noise. However, for datasets in which the biologically relevant

differences between cells are subtle, identifying these genes is a challenging task. We present the

self-assembling manifold (SAM) algorithm, an iterative soft feature selection strategy to quantify

gene relevance and improve dimensionality reduction. We demonstrate its advantages over other

state-of-the-art methods with experimental validation in identifying novel stem cell populations of

Schistosoma, a prevalent parasite that infects hundreds of millions of people. Extending our

analysis to a total of 56 datasets, we show that SAM is generalizable and consistently outperforms

other methods in a variety of biological and quantitative benchmarks.

Introduction

Single-cell RNA sequencing (scRNAseq) datasets typically contain tens of thousands of genes, although many of them may not be informative for differentiating between cell types or states. Feature selection is thus commonly used to select a subset of genes prior to downstream analyses, such as manifold reconstruction and cell clustering (Crow et al., 2018; Satija et al., 2015; Vallejos et al., 2015). However, current approaches have two major limitations.

First, feature selection methods filter genes based on arbitrarily or empirically chosen thresholds, small changes in which may result in different gene sets (Vallejos et al., 2017). In addition, the selection of features typically operates under the assumption that genes with highly variable expression between individual cells capture biologically meaningful variation. Because single-cell transcriptomes are inevitably contaminated by a combination of random transcriptional and technical noise (Grün et al., 2014), the variation in biologically relevant genes may be hard to distinguish from the background noise, especially when the differences between cell populations are subtle. Resolving these differences, or "signals", is essential to study a variety of biological problems, including identifying cell subtypes (Olsson et al., 2016; Treutlein et al., 2014; Lönnberg et al., 2017; Fincher et al., 2018; Baron et al., 2016) and quantifying the effects of molecular perturbations to otherwise homogeneous populations of cells (Lane et al., 2017). In such datasets, only a small fraction of the genes, and therefore only a small fraction of the total variation, may contain the signals relevant for distinguishing cell types. Choosing these features ahead of time without *a priori* knowledge remains an unmet computational challenge.

The second limitation is that existing methods have been almost exclusively benchmarked on well-annotated, gold standard datasets with clearly distinguishable cell types (Wang et al., 2017; Kiselev et al., 2017; Duò et al., 2019; Bahlo et al., 2018). These datasets are not informative

for distinguishing the performance between methods, because the differences between cell types

are relatively straightforward to detect. However, evaluating the performance of feature selection

and/or dimensionality reduction methods on datasets with more subtle signals is difficult as their

ground truth labels are typically ambiguous or nonexistent.

To overcome the shortcomings of current feature selection approaches, here, we introduce

the Self-Assembling Manifold (SAM) method, an unsupervised, "soft feature selection" algorithm

that iteratively rescales gene expressions to refine a nearest neighbor graph of cells until the graph

converges to a stable solution. At each iteration, SAM assigns more weight to genes that are

spatially variable across the constructed graph, and this weighted gene expression is then used to

improve the next nearest neighbor assignment. SAM presents two advantages: it rescales all genes

according to their weights, solving the problem of thresholding, and it prioritizes genes that are

variable across the intrinsic manifold of the data rather than selecting genes that are variable across

individual cells.

Second, in order to better distinguish the performance between methods, we define a

network sensitivity measure to identify datasets with subtle signals. With limited annotations in

most high-sensitivity datasets, we introduce unsupervised graph-based metrics to quantify the

degree of structure within the reconstructed manifolds for comparison between methods. In

addition, we perform benchmarking using known ground truth labels on simulated datasets

spanning a wide range of sensitivities by introducing increasing levels of noise to well-annotated

datasets. These analyses reveal that SAM consistently improves feature selection and cell

clustering.

To demonstrate the utility of SAM in practice, we provide an in-depth analysis of two

datasets that are challenging to analyze using existing methods: stem cells in a human parasitic

worm, *Schistosoma*, and activated macrophages (Lane et al., 2017). We show that SAM can

capture novel biology undetectable by other methods and validate these results with experimental

evidence.

Results

The SAM algorithm

The SAM algorithm begins with a random *k*-nearest neighbor (kNN) graph and averages

the expression of each cell with its *k* nearest neighbors: $C = \frac{1}{k}NE$, where N is the directed

adjacency matrix and E is the gene expression matrix (**Figure 1a**). For each gene *i*, SAM computes

a spatial dispersion factor of the averaged expressions C_i, which measures variation across

neighborhoods of cells rather than individual cells (**Methods**). These dispersions are used to

calculate the gene weights, which then rescale the expression matrix: $\hat{E} = E\sqrt{W_D}$, where W_D is a

diagonal matrix with gene weights along the diagonal. Using the rescaled expressions $\hat{E}$, we

compute a pairwise cell distance matrix and update the assignment of each cell's *k*-nearest

neighbors accordingly. This cycle continues until the gene weights converge.

To demonstrate the implementation and utility of SAM, below we analyze a challenging

dataset comprised of a few hundred relatively homogeneous stem cells isolated from *Schistosoma*

mansoni (**Figure 1–Figure supplement 1**), a widespread human pathogen (Hoffman et al., 2014).

Using a protocol we have established previously (Wang et al., 2018), these cells were collected by

sorting dividing cells from juvenile parasites harvested from their mouse hosts at 2.5 weeks post

infection. At this stage, the parasites use an abundant stem cell population (~15-20% of the total

number of cells) for rapid organogenesis and growth (Wang et al., 2013; Wang et al., 2018).

Testing several existing methods (Wang et al., 2017; Kiselev et al., 2017; Satija et al., 2015), we

found that they were not able to identify distinct cell populations in this dataset. In contrast, SAM

finds a stable solution independent of initial conditions (**Figure 1b**). A graph structure with clearly

separated cell populations self-assembles through the iterative process (**Figure 1c**). In parallel, the

gene weights converge onto the final weight vector. Eventually, only a small fraction of genes

(~1%) are strongly weighted and useful for separating cell clusters, reflecting the inherent

difficulty of analyzing this dataset.

Figure 1d shows that SAM iteratively improves a series of graph characteristics, including

the network-average clustering coefficient (NACC), modularity, and Euclidean norm of the spatial

dispersions (**Methods**). The NACC and modularity quantify the degree of structure within the

graphs – graphs with high NACC and modularity have regions of high density separated by regions

of low density. The dispersion quantifies the spatial organization of gene expression – the higher

the spatial dispersion the less uniformly distributed the gene expressions are along the graph.

Importantly, we verified that SAM does not artificially boost these metrics in data that lack

inherent structure: when applying SAM to a randomly shuffled expression matrix, none of these

metrics increased from the random initial conditions.

SAM identifies novel subpopulations within schistosome stem cells

Visualizing the converged graph in two dimensions using Uniform Manifold

Approximation and Projection (UMAP, Becht et al., 2019), we find that cells can be separated into

three main populations, with Louvain clustering (Blondel et al., 2008) further splitting one of these

clusters into two subpopulations (**Figure 2a**). In contrast, other commonly-used dimensionality

reduction methods, such as principal component analysis (PCA), Seurat (Satija et al., 2015), and

SIMLR (Wang et al., 2017), failed to distinguish these cell populations (see **Methods** for the

selection of algorithms for comparison). **Supplementary Table 1** lists genes with high SAM

weights, which includes most markers that were previously implicated to be enriched in subsets of

schistosome stem cells (Wang et al., 2013; Wang et al., 2018).

Figure 2b shows that the three populations include previously characterized δ′-cells, which

specifically express an RNA binding protein *nanos-2* (Smp_051920), and ε-cells, which are

marked by the expression of *eledh* (*eled*, Smp_041540) (Wang et al., 2018). More importantly,

SAM reveals a novel stem cell population, μ, comprising ~30% of all sequenced cells (μ denotes

muscle progenitors as discussed below). While μ-cells express ubiquitous stem cells markers (e.g.,

ago2-1, Smp_179320; *cyclin B*, Smp_082490) and cell cycle regulators (**Figure 2–Figure**

supplement 1a) (Collins et al., 2013; Wang et al., 2013; Wang et al., 2018), they are also strongly

enriched for a large set of genes, with a calcium binding protein (*cabp*, Smp_005350), an actin

protein (Smp_161920), an annexin homolog (Smp_074140), a helix-loop-helix transcription factor

(*dhand*, Smp_062490), and a phosphatase (*dusp10*, Smp_034500) as the most specific markers of

this population in comparison to other stem cells (**Figure 2–Figure supplement 1b**).

Fluorescent *in-situ* hybridization (FISH) in conjunction with EdU labeling of dividing cells

reveals that μ-cells (*cabp*$^+$EdU$^+$) are distributed near the parasite surface right beneath a layer of

post-mitotic differentiated cells that also express *cabp* (**Figure 2c**). Close to the parasite surface,

there are two major cell types intertwined in space: the skin-like tegumental cells and the body

wall muscle cells. However, μ-cells express none of the recently identified markers in tegumental

progenitors (Wendt et al., 2018), suggesting that they may be associated with the muscle lineage.

To test this idea, we performed double FISH experiments and observed in post-mitotic *cabp*$^+$ cells

the coexpression of a set of canonical muscle markers (Witchley et al., 2013), including

tropomyosin (Smp_031770), myosin (Smp_045220), troponin (Smp_018250), and collagen

(Smp_170340) (**Figure 2d**). These results suggest that *cabp* is a specific marker for parasite body

wall muscles and μ-cells are the muscle progenitors. Why the juvenile parasites maintain such an

active pool of muscle progenitors will be an important question for future studies.

In addition, SAM identifies two subpopulations among ε-cells: ε_α-cells that are highly

enriched for an aschaete-scute transcription factor (*astf*, Smp_142120), and ε_β-cells that

abundantly express another basic helix-loop-helix protein (*bhlh*, Smp_087310) (**Figure 2b**, right

panels). FISH experiments confirm these cells to be in close spatial proximity but with no

coexpression of *astf* and *bhlh* (**Figure 2e**). Moreover, we observed with FISH that there are fewer

astf$^+$ cells in larger, more matured juveniles, suggesting ε_α-cells are a dynamic population during

development. To verify this observation, we sequenced another ~370 stem cell from juveniles at a

later developmental time point (3.5 weeks post infection). After correcting for batch effects in the

combined 2.5- and 3.5-week datasets using the mutual nearest neighbors (MNN) algorithm

(Haghverdi et al., 2018), we find that δ'-, μ-, and ε_β-cells remain relatively constant throughout

both time points, whereas ε_α-cells comprise a significantly smaller fraction of the stem cells at 3.5

weeks (7%) compared to 21% at 2.5 weeks (**Figure 2f**). Taken together, these analyses

demonstrate that SAM can identify experimentally validated stem cell populations that are

previously too subtle to separate using other methods but are closely associated with the

schistosome development.

The critical difference between SAM and other methods lies in how they select genes for

manifold reconstruction. SAM prioritizes genes with variable expressions across neighborhoods

of cells rather than individual cells as in other methods (e.g., Seurat). **Figure 2g** shows that genes

with high standardized dispersion across individual cells often have low SAM weights, indicating

that these highly variable genes (HVGs) are irrelevant to the topological relationships between

cells. Other methods (e.g. SC3, Kiselev et al., 2017) identify marker genes based on differential

gene expression between cell clusters, but this approach suffers when cell cluster assignment is

poor, especially when discrete cell groups are difficult to separate or absent. Indeed, SC3 failed in

the default mode as it incorrectly predicted there to be only one cluster in the schistosome dataset.

After we manually increased the number of clusters, SC3 could recover a few of the marker genes

associated with only one (μ-cells, blue symbols in **Figure 2h**) of the populations detected by SAM.

Furthermore, changing the number of clusters resulted in different solutions and large variability

in SC3 scores for its top ranked genes.

SAM outperforms other state-of-the-art methods in extensive quantitative benchmarking

Below, we assess the general applicability of SAM by benchmarking its performance

against state-of-the-art scRNAseq analysis methods on a large collection of datasets. We focus on

three methods, i.e., Seurat, SIMLR, and SC3, as they are mostly unsupervised, have been broadly

used, and were shown to outperform other methods through extensive benchmarking (Kiselev et

al., 2017; Wang et al., 2017; Duò et al., 2019; Bahlo et al., 2018; Tian et al., 2019, see **Methods**

for the selection of algorithms for comparison). We first benchmark against nine datasets

(**Supplementary Table 2**) that have high-confidence annotations to evaluate the accuracy of SAM

in assigning cell clusters. We find that SAM has the highest Adjusted Rand Index (ARI, a measure

of clustering accuracy) (Hubert and Arabie, 1985) on eight out of the nine datasets and does not

over cluster the data (**Figure 3a**). Furthermore, SAM converges to the same set of gene weights

for all datasets analyzed (**Figure 3b**, **Figure 3-Figure supplement 1a**) and its performance is

robust to the choice of parameters and random initial conditions (**Figure 3-Figure supplement

1b-c**). In contrast, applying SAM to randomly generated datasets (**Methods**), the resulting gene

weights are highly dissimilar across random initial conditions (**Figure 3b**), indicating that SAM

does not converge to a stable solution on datasets with no intrinsic structure. Finally, the scalability

of SAM is similar to that of Seurat, capable of analyzing hundreds of thousands of cells in minutes

(**Figure 3c**), whereas SIMLR and SC3 are orders of magnitudes slower and thus excluded from

further benchmarking which requires the analysis of many more datasets.

Because these nine datasets are all comprised of clearly distinguishable cell types, they

may not represent the performance of methods on other datasets that contain cell populations that

are only subtly different. To identify such datasets, we introduce a network sensitivity metric that

quantifies the changes in the cell-to-cell distances when randomly selecting a subset of features

from the gene expression matrices (**Methods**). High network sensitivity indicates that changes to

the selected features strongly alters the resulting topological network. Networks that are robust to

the selected features correspond to datasets that have many redundant signals or genes

corroborating that network structure. In the datasets we compiled (**Supplementary Table 2**), all

broadly-used benchmarking datasets have lower sensitivities whereas the schistosome dataset,

which we have shown to be challenging to analyze for other methods, has the highest sensitivity

(**Figure 4a**). The fraction of genes with large SAM weights (>0.5) is negatively correlated with

the network sensitivity, suggesting that the biologically relevant variation in datasets with high

sensitivity is captured by relatively fewer genes (**Figure 4b**). Analyzing all 56 datasets, we found

that SAM improves the clustering, modularity, and spatial organization of gene expression across

the graph in comparison to Seurat as the datasets become increasingly sensitive (**Figure 4c**).

Evaluating the clustering accuracy for the highly sensitive datasets, however, is

challenging, because many of them have incomplete or nonexistent cell type annotations.

Therefore, we use the nine well-annotated benchmarking datasets to simulate data across a wide

spectrum of sensitivities. For this, we corrupt the data by randomly permuting gradually increasing

fractions of the gene expressions. As illustrated by the Darmanis dataset (Darmanis et al., 2015),

Figure 5a shows that the sensitivity increases along with the corruption. SAM's ARI scores only

marginally decrease as the corruption (and thereby sensitivity) increases, whereas Seurat's

performance rapidly deteriorates. A similar contrast was observed with the NACC, modularity,

and spatial dispersion between SAM and Seurat. Importantly, passing the genes with high SAM

weights into Seurat rescued its performance across all metrics, indicating that SAM is able to

consistently capture the genes relevant to the underlying structure of the data even with increasing

levels of noise and illustrating the robustness of its feature selection strategy compared to the HVG

filtering approach of Seurat. These observations generalize to all nine benchmarking datasets,

quantified by the area under the curves (AUC) of the metrics with respect to corruption (**Figure**

5b).

SAM clusters macrophages by their activation dynamics with proper temporal ordering

We next highlight another dataset to show that SAM can recover biologically meaningful

information that other methods fail to capture. We chose this example, which contains ~600

macrophages treated with lipopolysaccharide (LPS) when individually trapped in microfluidic

channels (Lane et al., 2017), because it has high network sensitivity (**Figure 4a**) and has

accompanying single cell functional data of macrophage activation dynamics that may help

validate the results of our analysis. Applied to this dataset, SAM initially identifies two clusters

(**Figure 6a**, top). Performing gene set enrichment analysis (GSEA, **Methods**, Subramanian et al.,

2005), we find that genes with high SAM weights are dominated by cell cycle-related processes,

with one of the clusters heavily enriched for cell cycle genes (e.g., Top2a, Mki67, **Figure 6-Figure**

supplement 1a). After removing the cell cycle effects (**Methods**), SAM identifies two different

clusters in which cells are properly ordered by the time since LPS induction, with the highly

weighted genes being primarily involved in immune signaling (**Figure 6a**, bottom). These

observations demonstrate that, in conjunction with GSEA, the quantitative gene weights output by

SAM can be used to infer the biological pathways that drive the clustering of cells.

One of the two clusters is enriched for TNFα expression (**Figure 6b**). It is known that LPS

activates two independent pathways, one through the innate immune signal transduction adaptor

(Myd88) and the other through the TIR-domain-containing adapter-inducing interferon-β (TRIF)

(Lee et al., 2009). While the Myd88 pathway directly activates NF-κB, the TRIF pathway first

induces the secretion of TNFα, which subsequently binds to its receptor, TNFR, to prolong the

activation of NF-κB (**Figure 6c**). **Figure 6d** and **Figure 6-Figure supplement 1b** show examples

of genes that are highly enriched with TNFα, a number of which are inflammatory factors known

to accumulate due to prolonged NF-κB activation (Lane et al., 2017). These results suggest that

SAM grouped the cells based on their activated signaling pathways: one cluster is activated

through both Myd88 and TRIF pathways (MT) while the other is only activated through Myd88

(M).

To further verify that the separation between the MT and M clusters truly reflects the

dichotomy in cellular response to LPS induction, we noted that this dataset combines scRNAseq

with live-cell imaging of NF-κB activity in single cells. This allows us to directly test if the MT

and M clusters correspond to different signaling dynamics (**Methods**). We found that most of the

cells with prolonged NF-κB response (i.e., cells showing broad peaks of NF-κB activation in time)

are in fact in the MT cluster (**Figure 6e-f**, and **Figure 6-Figure supplement 2a**), consistent with

the expectation that TNFα signaling prolongs NF-κB activation. Although our interpretation of the

data matches that provided in the original study, we were able to analyze the dataset with almost

no *a priori* knowledge. In contrast, the original study required extensive manual curation, analyzed

only a subset of the dataset, and could not group cells by their NF-κB activation dynamics from

the gene expression data alone. Similarly, Seurat and SIMLR were unable to order the cells by the

time since LPS induction or group cells based on their activation dynamics after removing the cell

cycle effects (**Figure 6g**, and **Figure 6-Figure supplement 2b-c**).

Discussion

Here, we introduced a scRNAseq analysis method, SAM, that uses an unsupervised, robust,

and iterative strategy for feature selection and manifold reconstruction. As demonstrated by our

analysis of the schistosome stem cells and activated macrophages, SAM can capture biology that

is undetectable by other methods. While SAM has consistently higher clustering accuracy than

other state-of-the-art methods on datasets containing clearly distinct cell types, its advantages are

especially apparent on datasets in which cell states or types are only distinguishable through subtle

differences in gene expression.

The strength of SAM lies in the integration of three algorithmic components: spatial

dispersion to measure feature relevance, soft feature selection, and the iterative scheme. By

averaging the gene expression of a cell with that of its neighbors, the spatial dispersion quantifies

the variation across neighborhoods of cells rather than individual cells. Genes with high spatial

dispersion are more likely to be biologically relevant as they are capable of separating cells into

distinct topological locations. Soft feature selection includes all genes and weights their

contribution to the manifold reconstruction by their spatial dispersions. This mitigates the

shortcoming of existing approaches in which the selection of features is a binary decision: genes

are either included or not depending on arbitrarily chosen thresholds.

The conceptual challenge here is that calculating the gene weights requires the manifold,

but reconstructing the manifold requires the gene weights for feature selection. SAM thus uses an

iterative strategy to converge onto both the gene weights and the corresponding graph topology

from a random initial graph. Each successive iteration refines the gene weights and network

structure until the algorithm converges. Empirically, for all datasets analyzed we have shown that

SAM converges onto a stable solution and is robust to the random initial conditions. Practically,

we could initialize SAM using the graph output of another method such as Seurat, but using

random initial conditions avoids potential biases in the analysis and enables the evaluation of the

stability of SAM.

To demonstrate the strengths of SAM in practice, we analyzed the schistosome stem cells

and identified novel stem cell populations that were validated by FISH experiments (**Figure 2**). In

the analysis of activated macrophages, we showed that SAM can simultaneously order cells by the

time since LPS induction and group cells according to their respective activated signaling

pathways. We have validated this result using the live-cell imaging data presented in the original

study (**Figure 6**).

We expect the application of SAM is not limited to feature selection, cell clustering, and

manifold reconstruction; it can be readily integrated with existing analytical pipelines as its gene

weights and reconstructed manifolds can be used in downstream analyses. For example, we have

shown how the genes ranked by their SAM weights can be used as input to GSEA to determine

the biological processes enriched among the highly weighted genes (**Figure 6**), thus directly

testing if the weights reflect biologically relevant genes. Additionally, the manifold reconstructed

by SAM can be used as input to pseudotemporal ordering algorithms (Setty et al., 2016; Trapnell

et al., 2014).

Beyond the two example case studies, we have rigorously evaluated SAM on a total of 56

datasets. While previous studies benchmarked on datasets with clearly defined cell populations,

we defined a network sensitivity measure to rank the datasets based on the inherent difficulty of

their analysis (**Figure 4**). Using these datasets, we showed that SAM consistently outperforms

other methods in terms of both cell clustering accuracy measured by ground truth annotations, and

manifold reconstruction measured by quantitative graph characteristics. These improvements can

be attributed to the robust selection of features relevant for cell clustering and manifold

reconstruction even in the presence of significant amounts of random noise, as shown in the

corruption tests (**Figure 5**). Overall, the network sensitivity and quantitative benchmarking metrics

should help in characterizing the performance of future scRNAseq analysis methods across a wider

variety of datasets.

Materials and Methods

Code and data availability. The SAM source code and tutorials can be found at https://github.com/atarashansky/self-assembling-manifold. The schistosome stem cell scRNAseq data generated in this study is available through the Gene Expression Omnibus (GEO) under accession number GSE116920.

Data processing. Supplementary Table 2 summarizes all datasets used in this study as well as the methods used to convert raw sequence read counts to gene expression, such as TPM (transcripts per million), CPM (counts per million), RPKM (reads per kilobase per million), or FPKM (fragments per kilobase per million). Datasets with asterisks next to their accession numbers are sourced from the *conquer* database (Soneson and Robinson, 2018). The nine benchmarking datasets used with high-confidence labels are marked by crosses. Gene expression is measured in log space with a pseudocount of 1 (e.g., $\log_2(\text{TPM}+1)$). Genes expressed ($\log_2(\text{TPM}+1)>1$) in fewer than 1% or more than 99% of cells are excluded from downstream analysis as these genes lack statistical power. To reduce the influence of technical noise near the molecular detection limit, we set gene expression to zero when $\log_2(\text{TPM}+1)<1$. We denote the resulting expression matrix as E.

In the SAM algorithm (see below), we either standardize the gene expression matrix E to have zero mean and unit variance per gene (which corrects for differences in distributions between genes) or normalize the expressions such that each cell has unit Euclidean (L2) norm (which prevents cells with large variances in gene expressions from dominating downstream analyses) prior to dimensionality reduction. In the below section, we denote the standardized or normalized expression matrix as $\bar{E}$. Empirically, we have found that standardization performs well with large,

sparse datasets collected through droplet-based methods, whereas L2-normalization is more

suitable for smaller datasets with higher sequencing depth such as those prepared with the Smart-

Seq2 protocol (Picelli et al., 2013). This is likely due to the fact that standardization amplifies the

relative expression of genes specific to small populations in large datasets, thereby making them

easier to identify. In contrast, standardization decreases the relative expression of genes specific

to populations comprising larger fractions of the data, as is typically the case in smaller datasets,

thereby making distinct populations more difficult to identify. **Supplementary Table 2** documents

the preprocessing step used for each dataset.

The SAM algorithm. After first generating a random kNN adjacency matrix, the SAM algorithm

goes through three steps that are repeated until convergence.

1) Calculate the gene weights

First, the expression of each cell is averaged with its k-nearest neighbors:

$$C = \frac{1}{k}NE \tag{1}$$

349   where N is the directed adjacency matrix for the kNN graph, and E is the original n x m gene

350   expression matrix with rows as cells and columns as genes. Here, we do not use $\bar{E}$ as it may

351   contain negative values, for which dispersion is ill-defined. For each gene i, SAM computes the

352   Fano factor from the averaged expressions C_i:

$$\mu_{C_i} = \frac{1}{n}\sum_{j=1}^{n} C_{ji} \tag{2}$$

$$\sigma_{C_i}^2 = \frac{1}{n} \sum_{j=1}^{n} (C_{ji} - \mu_{C_i})^2 \qquad (3)$$

$$F_i = \frac{\sigma_{C_i}^2}{\mu_{C_i}} \qquad (4)$$

where μ_{C_i} is the mean and $\sigma_{C_i}^2$ is the variance. We use the Fano factor to measure the gene

expression variance relative to the mean in order to account for the fact that genes with high mean

expressions tend to have higher variability. Computing the Fano factors based on the kNN-

averaged expressions links gene dispersion to the cellular topological structure: Genes that have

highly variable expressions among individual cells but are homogeneously distributed across the

topological representation should have small dispersions. k, set by default to 20, determines the

topological length scale over which variations in gene expression are quantified. **Figure 3-Figure**

supplement 1b shows that the downstream analysis is robust to the specific choice of k.

Additionally, the choice of k does not significantly affect runtime complexity or scalability.

To compute the gene weights, we normalize the Fano factors to be between 0 and 1. First,

we saturate the Fano factors to ensure that outlier genes with large spatial dispersions do not skew

the distribution of weights:

$$\{F_i | F_i > z\} = z \qquad (5)$$

365 where z is by default the mean of the largest 50 dispersions. In other words, Fano factors exceeding

366 this number are saturated to be z. We then calculate the gene weights as:

$$W_i = \frac{F_i}{z} \qquad (6)$$

367 *2) Rescale the expression matrix*

368   Having calculated the gene weights, SAM multiplies them into the preprocessed expression

369 matrix:

$$\hat{E} = \bar{E}\sqrt{W_D} \tag{7}$$

where $\bar{E}$ is the standardized or normalized expression matrix and W_D is a diagonal matrix with W_i

along the diagonal. This matrix multiplication linearly rescales the gene expression variances and

gene-gene covariances by their respective weights, attenuating the influence of genes with low

dispersions across neighborhoods.

3) Updating the kNN graph

To compute pairwise cell-cell distances, we perform PCA on the rescaled expression

matrix $\hat{E}$. The variance-scaling operation in **Eq. 7** improves the robustness of PCA to variations in

genes that are uniformly distributed along the current graph (i.e., genes with low weights).

Furthermore, this weighting strategy eliminates the typical requirement of selecting a subset of

HVGs to feed into PCA, which often relies on arbitrary thresholds and heuristics. To perform PCA,

we first mean center $\hat{E}$ to form $\hat{E}_\mu$:

$$\hat{E}_\mu = \hat{E} - \frac{1}{n}ee^T\hat{E} \tag{8}$$

382 where e is a column vector of ones with dimension n. We then compute the Singular Value

383 Decomposition (SVD) of $\hat{E}_\mu$:

$$\hat{E}_\mu = USV^T \tag{9}$$

384 with the principal components defined as

$$P = US \tag{10}$$

385 The eigenvalues corresponding to the eigendecomposition of the gene-gene covariance matrix

386 are defined in terms of the singular values as

$$\Lambda = \frac{S^2}{n-1} \tag{11}$$

387   where S is a diagonal matrix with singular values along the diagonal. Using the PC matrix P, SAM

388   computes a pairwise cell-cell distance matrix. While typical dimension reduction approaches select

389   a subset of the PCs, which is often subjective or requires computationally intensive maximum-

390   likelihood approaches, we include all PCs and scale their variances by their corresponding

391   eigenvalues:

$$\hat{P} = P\sqrt{\Lambda} \tag{12}$$

As a result, PCs with small eigenvalues are weighted less in the calculation of the distance between

cells i and j, $D_{P_i P_j}$. $D_{P_i P_j}$ is the Pearson correlation or Euclidean distance between rows P_i and P_j

in the PC matrix. Pearson correlation distance is used by default, although **Figure 3-Figure**

supplement 1c shows that SAM is robust to the choice of distance metric. Using the distances to

define the k-nearest neighbors for each cell, SAM updates the kNN matrix and repeats steps 1-3.

The algorithm continues until convergence, defined as when the RMSE between gene weights in

adjacent iterations diminished as defined by:

$$\sqrt{\frac{1}{m}\sum_{j=1}^{m}(W_{i,j} - W_{i+1,j})^2} < 5 \times 10^{-3} \tag{13}$$

where m is the number of genes and $W_{i,j}$ is the weight for gene j at iteration i.

Visualization. To visualize the topological structure identified by SAM, we feed the final

weighted PCA matrix, $\hat{P}$, into UMAP (Becht et al., 2019) using Pearson correlation as the distance

metric by default. To directly visualize the final kNN adjacency matrix (**Figure 1c**), we used the

Fruchterman-Reingold force-directed layout algorithm and drawing tools implemented in the

Python package *graph-tool* (Peixoto, 2017).

Choosing the benchmarking methods. We used three main criteria for choosing the benchmark

scRNAseq analysis methods: they should be widely used, have done extensive benchmarking

against other methods, and be mostly unsupervised. We found on Web of Science that among the

highest cited scRNAseq analysis tools in 2017-2018 are Seurat, SC3, SIMLR, Reference

Component Analysis (RCA, Li et al., 2017), Monocle (Trapnell et al., 2014), zero-inflated factor

analysis (ZIFA, Pierson and Yau, 2015), and Wishbone (Setty et al., 2016), of which we chose

Seurat, SC3, and SIMLR.

SC3 is a consensus clustering algorithm that has done rigorous benchmarking against other

methods such as SINCERA (Guo et al., 2015), SNN-Cliq (Xu and Su, 2015) and pcaReduce

(Žurauskien and Yau, 2016) on 12 datasets with ground truth labels. SIMLR, a dimensionality

reduction and clustering algorithm, evaluated its clustering performance on four annotated datasets

against eight other dimensionality reduction methods, including PCA, Factor Analysis (FA), t-

SNE, multidimensional scaling (MDS), and (ZIFA). Both methods have demonstrated the highest

clustering accuracy across most of the tested datasets. Additionally, as both methods have built-in

functions to estimate the number of clusters present within the data, they are largely unsupervised.

We also selected Seurat as one of the benchmarking methods, because it is arguably the most

widely used tool for dimensionality reduction and clustering of scRNAseq data and has performed

well in rigorous benchmarking studies against various methods including SC3, SIMLR, RCA, and

pcaReduce (Duò et al., 2019; Bahlo et al., 2018).

We did not select Reference Component Analysis as it is primarily designed for cases in

which an atlas of bulk, cell-type specific, reference transcriptomes is present. Additionally, we did

not benchmark against Monocle and Wishbone, because they are pseudotime analysis methods

and are meant for datasets with continuous branching processes such as cell differentiation.

However, it is important to note that SAM can be used for dimensionality reduction upstream of

pseudotime algorithms for such datasets. Finally, we did not benchmark against ZIFA as it has

already been shown to have lower clustering accuracy than SIMLR.

In addition to measuring clustering accuracy, we also introduce the unsupervised NACC,

modularity, and spatial dispersion metrics to quantify both the degree of structure and spatial

organization of gene expression within a nearest-neighbor graph. As such, these metrics can only

be applied to dimensionality reduction methods that construct a graph representation of the dataset.

Consequently, we cannot use these metrics to evaluate SC3.

Although it does technically produce a graph representation of the data, SIMLR should be

considered as a hybrid between a clustering and dimensionality reduction method. Because its

similarity graph is assumed to have a block structure where the number of blocks is equal to the

prespecified number of clusters, the resulting nearest-neighbor graph will, by construction, tend to

have a higher degree of structure and therefore artificially inflated NACC and modularity.

Furthermore, the poor scalability of SC3 and SIMLR makes them difficult to run for many

trials across a large number of datasets. Although SIMLR, in particular, does provide an alternative

algorithm that can scale to much larger datasets, it has not been extensively benchmarked. Even

so, despite the improved speed of this large-scale implementation, estimating the number of

clusters using its built-in function remains a significant computational and memory bottleneck. For

example, when applied to the ~10,000 planarian neoblasts, neither implementations of SIMLR

could estimate the number of clusters within 2 hours. As a result, we cannot run SIMLR in an

unsupervised manner on datasets significantly larger than ~3000 cells.

As there are few practical alternatives for manifold reconstruction that have been as

extensively benchmarked and widely used, we primarily compare SAM to Seurat in tests involving

the unsupervised, graph-based metrics to highlight the key, advantageous characteristics of SAM

as a manifold reconstruction and feature selection algorithm when applied to datasets with varying

sensitivities (**Figure 4a-c**).

Benchmarking. To generate the convergence curves in **Figure 1b**, we computed the root mean

square error (RMSE) of the gene weights averaged across all pairwise comparisons of ten

replicates starting from randomly generated initial graphs. In **Figure 3b**, we extend this analysis

to all datasets analyzed and report the final error. We use randomly generated datasets of varying

sizes (ranging from 200 to 5000 cells) as a negative control to show that SAM does not converge

onto the same solution across initial conditions when the data has no intrinsic structure. These

datasets were randomly generated by sampling gene expressions from a Poisson distribution with

mean drawn from a gamma distribution. To generate the convergence curves in **Figure 3-Figure**

supplement 1a, we computed the RMSEs, which are ensemble-averaged across ten replicate runs,

between the gene weights in adjacent iterations. We compute the adjacency error between kNN

adjacency matrices N_i and N_j (**Figure 1b**) as

$$A_{i,j} = \frac{e^T |N_i - N_j| e}{2 e^T N_i e} \tag{14}$$

where e is a column vector of ones. This simply measures the fraction of total edges that are

different between the two graphs.

To compute the standardized dispersion factors in **Figure 2g**, we used Seurat's

methodology implemented in Scanpy (Wolf et al., 2018), which groups the genes into 20 bins

based on their mean expression values and computes the z-score of each gene's Fano factor with

respect to the mean and standard deviation of all Fano factors in its corresponding bin. To generate

the AUROC scores in **Figure 2h**, which quantify the likelihood of genes being cluster-specific

markers, we ran SC3 on the schistosome data with the number of clusters ranging from 2 to 12.

We used the AUROC scores corresponding to 4 clusters for the points on the scatter plot and the

standard deviations of the scores across all tested numbers of clusters for the error bars.

478   We evaluated each analysis method on nine gold standard datasets (**Figure 3a**) using the

479   Adjusted Rand Index (ARI), which measures the accuracy between two cluster assignments X and

480   Y while accounting for randomness in the clustering:

$$ARI = \frac{\Sigma \binom{n_{ij}}{2} - \left[\Sigma \binom{a_i}{2} \Sigma \binom{b_j}{2}\right]/\binom{n}{2}}{\frac{1}{2}\left[\Sigma \binom{a_i}{2} + \Sigma \binom{b_j}{2}\right] - \left[\Sigma \binom{a_i}{2} \Sigma \binom{b_j}{2}\right]/\binom{n}{2}} \tag{15}$$

481   where n is the number of cells, and n_{ij}, a_i, and b_j are elements from a contingency table that

482   summarizes the overlap between the assignments X and Y (Hubert and Arabie, 1985). n_{ij} denotes

483   the number of cells assigned to X_i that are also assigned to Y_j, while a_i and b_j are the sums of the

484   ith row jth column of the contingency table, respectively.

485   Seurat was implemented using the Scanpy package in Python (Wolf et al., 2018). For

Seurat, we selected the top 3000 variable genes according to their standardized dispersions and

chose the number of PCs (bounded between 6 and 50) which explain 30% of the variance for

dimensionality reduction. From these PCs, we calculated a cell-cell correlation distance matrix.

To keep the comparison between SAM and Seurat graphs consistent, this distance matrix was

converted into a kNN adjacency matrix with the value of k used by SAM. To assign cluster labels

for SAM and Seurat, we applied HDBSCAN (Mcinnes et al., 2017), an unsupervised, density-

based clustering algorithm to their respective PCA outputs. As HDBSCAN does not cluster any

cell it deems an outlier, we assign the remaining outlier cells to clusters using kNN classification.

For each outlier cell, we identify its 20 nearest neighbors among the clustered cells. Outliers are

assigned to the same cluster as that of the majority of its neighbors. This minor extension to

HDBSCAN is available as the built-in function *hdbknn_clustering* in SAM. SC3 was run using

default parameters. The SIMLR package was implemented in R and run with the normalization

parameter set to "True", which mean-centers gene expressions after normalizing them to be

between 0 and 1. Both SC3 and SIMLR provide their own functions to estimate the number of

clusters and cluster assignments.

To compare the quality of graphs generated by different methods, we use the NACC,

modularity, and spatial dispersion. The NACC is the average of the local clustering coefficient for

each node of a graph and quantifies the degree of structure in the graph (Watts and Strogatz, 1998).

The local clustering coefficient is defined as

$$a_i = \frac{L_i}{k_i(k_i - 1)} \tag{16}$$

505 where L_i is the number of edges between the k_i neighbors of node i and measures the degree of

506 connectedness in a particular node's local neighborhood. We calculate the NACC using the

507 implementation in *graph-tool* (Peixoto, 2017).

508   The modularity Q of a graph is defined as

$$Q = \frac{1}{4m} \sum_{i,j}^{c} \left(A_{ij} - \frac{k_i k_j}{2m} \delta_{ij} \right) \tag{17}$$

where A_{ij} is one if there is an edge from cell i to cell j, k_i is the degree of cell i, k_j is the degree of

cell j, m is the total number of edges, and δ_{ij} is one if cells i and j are in the same cluster. High

modularity indicates that clusters have on average many more edges within clusters than between

clusters. To find the optimal modularity for a particular graph, we used Louvain clustering, which

searches for a partition in which modularity is maximized.

To quantify the spatial organization of gene expression along the graph, we calculate the

Euclidean norm of the largest 500 spatial dispersions. Spatial dispersion is defined as before in the

SAM algorithm: $F_i = \frac{\sigma^2_{C_i}}{\mu_{C_i}}$, where F_i is the Fano factor of the kNN-averaged expressions, $C_i =$

$\frac{1}{k}NE_i$. N is the directed adjacency matrix output by SAM or Seurat and E_i is a column vector of

expression values for gene i.

To measure the inherent sensitivity of each dataset, we randomly perturbed the gene

expression matrices of each dataset by randomly sampling 2000 genes and applied PCA to the

subsampled data. A correlation distance matrix was calculated from the top 15 PCs and

perturbations were repeated 20 times to generate distance matrix replicates. Sensitivity is then

defined as the average error across all pairwise comparisons between replicates. The error between

two distance matrices j and k, S_{jk}, is defined as the average correlation distance between

corresponding pairs of rows in the distance matrices d_j and d_k:

$$S_{jk} = \frac{1}{n}\sum_{i=1}^{n} D\{d_{j,i}, d_{k,i}\} \tag{18}$$

where $D\{d_{j,i}, d_{k,i}\}$ is the Pearson correlation distance between the distances from cell i in distance

matrices j and k.

We simulated datasets with increasing sensitivity by introducing increasing degrees of

corruption in each of the nine annotated datasets. To corrupt a dataset, we swapped random pairs

of elements in the expression matrix. The number of swaps, p, corresponds to the degree of

corruption, with p varying from 0 to half of the total number of elements in the matrix. For each

annotated dataset, we simulated 10 replicates per value of p. SAM and Seurat were run with default

values on each corrupted dataset, clustering was performed using the *hdbknn_clustering* function

in SAM, and the ARI, NACC, modularity, and spatial dispersion metrics were recorded. The Area

Under the Curve (AUC) was calculated for each metric with respect to the fraction of elements

swapped, p, using the trapezoidal rule. Finally, to rescue the performance of Seurat, we used as

input to Seurat the top 3000 genes with the highest SAM weights.

Gene set enrichment analysis (GSEA). GSEA (Subramanian et al., 2005) is typically run on a

gene expression matrix with user-defined cluster assignments to quantify the differential

expression for each gene. By default, differential expression is quantified using a signal-to-noise

metric and the resulting scores are used to rank the genes in descending order. However, GSEA

can also run in an alternative mode in which the user provides a predefined list of gene rankings.

Therefore, we used the genes ranked by their SAM weights as input to GSEA to determine the

biological processes enriched among the highly weighted genes. As shown in **Figure 6a**, we can

directly test if SAM captures the relevant biological processes. GSEA provides a number of

statistical measures to assess the significance of enriched gene sets, of which we use the False

Discovery Rate (FDR). The FDR quantifies the likelihood that a highly enriched gene set

represents a false positive. The significance threshold typically used with FDR is 25%, which

implies that the results are likely to be valid 75% of the time.

Removal of cell cycle effects. To remove cell cycle effects from the macrophage dataset, we

adopted a simpler version of the strategy used in *ccRemover* (Barron and Li, 2016), in which we

subtract from the data PCs that are significantly associated with known cell cycle genes. Letting P

represent the PCs and L be the gene loadings, we quantify the association between the set of cell

cycle genes G and PC j as

$$A_j = \frac{1}{|G|} \sum_{i \in G} |L_{ji}| \tag{19}$$

557   PC j is selected if its association A_j is at least two standard deviations above the mean of the

558   associations for all PCs. In the particular case of the macrophage data, we identified the set of PCs

559    $S = \{P_0, P_1, P_8\}$ as being significantly associated with the cell cycle genes. We next reconstruct the

560   data using these PCs, which thus captures the cell-cycle effects, and subtract the reconstructed data

561   from the expression matrix E:

$$E_{removed} = E - \sum_{j \in S} P_j L_j \sqrt{W} \tag{20}$$

When reconstructing the data, we scale the gene loadings by the SAM weights W so that only the

highly weighted SAM genes (which are initially dominated by cell cycle genes) contribute to the

cell cycle removal, as there may be genes involved in other biological processes that could also be

correlated with the PCs in S. To run SAM on the data with cell cycle effects removed, we use E

as opposed to $E_{removed}$ for the calculation of spatial dispersions, because the latter may contain

negative values, for which dispersion is ill-defined. This method is made available as a part of the

SAM package in the functions *calculate_regression_PCs* and *regress_genes*.

Clustering the NF-κB activity time series. In the original study, Lane et al. combined imaging

and transcriptomics to link NF-κB nuclear translocation dynamics to changes in gene expression

within single cells. Macrophages stimulated with LPS were individually trapped in microfluidic

chambers and imaged for various lengths of time (75-300 min) prior to scRNAseq library

preparation. NF-κB was tagged with a fluorescent protein, and its activation was measured as the

nuclear-localized fluorescence intensity. Based on the imaging data, the authors identified three

main classes of NF-κB dynamics, the first with a transient initial response, the second with a

prolonged initial response, and the third with a recurrent response. Because the recurrent response

is found only in the 300 min time point and comprises only ~8% of these cells, we primarily

focused on clustering cells based on their initial dynamics. To do this, we used the *tslearn*

(Tavenard, 2017) python package to group cells based on their NF-κB activity time series. Because

these time series are quite noisy, we were conservative in labeling cells as having a prolonged

initial response in an effort to avoid false positives. As a result, these cells comprise only ~30% of

the dataset.

For the cells sampled at 75 and 150 min after LPS stimulation, we used the time series *k-*

means algorithm with the *softdtw* distance metric to cluster them each into three groups, which

resulted in representative time series with transient, intermediate, and prolonged responses.

Merging the cells with transient and intermediate responses into one cluster (which we simply

labeled as transient response), we obtained the 75 and 150 min (black and blue, respectively)

representative time series shown in **Figure 6e**. Because the cells sampled at 300 min displayed

much more variability in their NF-κB time series, we clustered them into 6 groups, labeling the

cluster whose representative time series had the broadest initial peak as the prolonged response

cluster (red in **Figure 6e**, right). The remaining groups were labeled as the transient response

cluster (red in **Figure 6e**, left).

Mapping the schistosome datasets. We used the Mutual Nearest Neighbors algorithm

(Haghverdi et al., 2018) with default values to generate an expression matrix $E_{corrected}$ in which

the batch effects between the 2.5-week and 3.5-week datasets were corrected for. To run SAM on

the batch-corrected data, we use E for the calculation of spatial dispersions as opposed to

$E_{corrected}$.

scRNAseq of schistosome stem cells. Schistosome stem cells were isolated from juvenile

parasites retrieved from infected mice at 2.5 and 3.5 weeks post infection. We followed the

protocol as previously described (Wang et al., 2018). Briefly, we retrieved juvenile parasites from

schistosome-infected mice (Swiss Webster NR-21963) by hepatic portal vein perfusion. Parasites

were cultured at $37°C/5\%$ CO_2 in Basch Medium 169 supplemented with 1X Antibiotic-

Antimycotic for 24-48 hr to allow complete digestions of host blood cell in parasite intestines. In

adherence to the Animal Welfare Act and the Public Health Service Policy on Humane Care and

Use of Laboratory Animals, all experiments with and care of mice were performed in accordance

with protocols approved by the Institutional Animal Care and Use Committees (IACUC) of

Stanford University (protocol approval number 30366).

Before dissociation, parasites were permeabilized in PBS containing 0.1% Triton X-100

and 0.1% NP-40 for 30 seconds and washed thoroughly to remove the surfactants. The

permeabilized parasites were dissociated in 0.25% trypsin for 20 min. Cell suspensions were

passed through a 100 μm nylon mesh (Falcon Cell Strainer) and centrifuged at 150 g for 5 min.

Cell pellets were gently resuspended, passed through a 30 μm nylon mesh, and stained with

Vybrant DyeCycle Violet (DCV; 5 μM, Invitrogen), and TOTO-3 (0.2 μM, Invitrogen) for 30–45

min. As the stem cells comprise the only proliferative population in schistosomes, we flow-sorted

cells at G_2/M phase of the cell cycle on a SONY SH800 cell sorter. Dead cells were excluded based

on TOTO-3 fluorescence. Single stem cells were gated using forward scattering (FSC), side

scattering (SSC), and DCV to isolate cells with doubled DNA content compared to the rest of the

population (Wang et al., 2018). Cells that passed these gates were sorted into 384-well lysis plates

containing Triton X-100, ERCC standards, oligo-dT, dNTP, and RNase inhibitor.

cDNA was reverse transcribed and amplified on 384-well plate following the Smart-Seq2

protocol (Picelli et al., 2013). For quality control, we quantified the histone *h2a* (Smp_086860)

levels using qPCR, as *h2a* is a ubiquitously expressed in all schistosomes stem cell (Collins et al.,

2013; Wang et al., 2013; Wang et al., 2018). We picked wells that generated C_T values within 2.5

C_T around the most probable values (~45% of total wells, **Figure 1-Figure supplement 1**). cDNA

was then diluted to 0.4 ng/µL for library preparation. Tagmentation and barcoding of wells were

prepared using Nextera XT DNA library preparation kit. Library fragments concentration and

purity were quantified by Agilent bioanalyzer and qPCR. Sequencing was performed on a NextSeq

500 using V2 150 cycles high-output kit at ~1 million reads depth per cell. Raw sequencing reads

were demultiplexed and converted to fastq files using bcl2fastq. Paired-end reads were mapped to

S. mansoni genome version WBPS9 (WormBase Parasite) using STAR. In 2.5-week dataset, 338

cells with more than 1700 transcripts expressed at >2 TPM were used for downstream analysis. In

the 3.5-weeks dataset, 338 cells with more than 1350 transcripts expressed at >2 TPM were used

for downstream analysis (**Figure 1-Figure supplement 1**).

In situ hybridization and EdU labeling. RNA FISH experiments were performed as detailed in

previous publications (Collins et al., 2013; Wang et al., 2013; Wang et al., 2018). Clones used

for riboprobe synthesis were generated as described previously, with oligonucleotide primers

listed in **Supplementary Table 3**. Juvenile parasites were cultured with 10 µM EdU overnight,

killed in 6 M $MgCl_2$ for 30s, and then fixed in 4% formaldehyde with 0.2% Triton X-100 and 1%

NP-40. Fixed parasites were sequentially dehydrated in methanol, bleached in 3% H_2O_2 for 30

min, and rehydrated. Parasites were permeabilized by 10 µg/mL proteinase K for 15 min and

post fixed with 4% formaldehyde. The hybridization was performed at 52°C with riboprobes

labeled with either digoxigenin-12-UTP (Roche) or fluorescein-12-UTP (Roche). For detection,

samples were blocked with 5% horse serum and 0.5% of Roche Western Blocking Reagent, and

then incubated with anti-digoxigenin-peroxidase (1:1000; Roche) or anti-fluorescein peroxidase

(1:1500; Roche) overnight at 4°C for tyramide signal amplification (TSA). For double FISH, the

first peroxidase was quenched for 30 min in 0.1% sodium azide solution before the detection of

the second gene. After FISH, EdU detection was performed by click reaction with 25µM Cy5-

azide conjugates (Click Chemistry Tools). Samples were mounted in *scale* solution (30%

glycerol, 0.1% Triton X-100, 4 M urea in PBS supplemented with 2 mg/mL sodium ascorbate)

and imaged on a Zeiss LSM 800 confocal microscope.

Acknowledgements. *S. mansoni* (strain: NMRI) was provided by the NIAID Schistosomiasis

Resource Center for distribution through BEI Resources, NIH-NIAID Contract

HHSN272201000005I. We thank F. Zanini, N. Neff, J. Okamoto, and D. Nanes-Sarfati for

experimental help, K. Lane, M. Covert, J. Qin, F. Horns, G. Stanley, J. Rink, and M. Chen for

conceptual input and stimulating discussions. B.W. is supported by the Burroughs Wellcome Fund

through the CASI program and a Beckman Young Investigator Award.

References

Bahlo, M., Tian, L., Lönnstedt, I., Ng, M. & Hicks, S. Comparison of clustering tools in R for
medium-sized 10x Genomics single-cell RNA-sequencing. *F1000Research* **7,** 1–26 (2018).
664
665   Baron, M. *et al*. A single-cell transcriptomic map of the human and mouse pancreas reveals inter-
666   and intra-cell population structure. *Cell syst.* **3,** 346–360 (2016).
667
668   Barron, M. & Li, J. Identifying and removing the cell- cycle effect from single-cell RNA-
669   Sequencing data. *Sci. Rep.* **6,** 33892 (2016).
670
671   Becht, E. *et al.* Dimensionality reduction for visualizing single-cell data using UMAP. *Nat.*
672   *Biotechnol.* **37,** 38–44 (2019).
673
674   Blondel, V. D., Guillaume, J., Lambiotte, R. & Lefebvre, E. Fast unfolding of communities in
675   large networks. *J. Stat. Mech. Theory Exp.* **2008,** P10008 (2008).
676
677   Collins, J. J. *et al.* Adult somatic stem cells in the human parasite Schistosoma mansoni. *Nature*
678   **494,** 476–479 (2013).
679
680   Crow, M. *et al*. Characterizing the replicability of cell types defined by single cell RNA-
681   sequencing data using MetaNeighbor. *Nat. Commun.* **9,** 884 (2018).
682
683   Darmanis, S. *et al.* A survey of human brain transcriptome diversity at the single cell level. *Proc.*
684   *Natl. Acad. Sci. U.S.A.* **112,** 7285–7290 (2015).
685
686   Duò, A., Robinson, M. D. & Soneson, C. A systematic performance evaluation of clustering
687   methods for single-cell RNA-seq data. *F1000Research* **7,** 1–21 (2019).
688
689   Fincher, C. T. *et al*. Cell type transcriptome atlas for the planarian Schmidtea mediterranea.
690   *Science* **360,** eaaq1736 (2018).
691
692   Guo, M. *et al*. SINCERA: A Pipeline for Single-Cell RNA-Seq Profiling Analysis. *PLOS*
693   *Comput. Biol.* **11,** e1004575 (2015).
694
695   Grün, D., Kester, L. & Oudenaarden, A. V. Validation of noise models for single-cell
696   transcriptomics. *Nat. Methods* **11,** 637–640 (2014).
697
698   Haghverdi, L. *et al*. Batch effects in single-cell RNA-sequencing data are corrected by matching
699   mutual nearest neighbors. *Nat. Biotechnol.* **36,** 421–427 (2018).
700
701   Hoffmann, K. F., Brindley, P. J. & Berriman, M. Halting harmful helminths. *Nature* **168,** 168–
702   169 (2014).
703
704   Hubert, L. & Arabie, P. Comparing partitions. *J. Classif.* **2,** 193–218 (1985).
705

706 Kiselev, V. Y. *et al.* SC3: Consensus clustering of single-cell RNA-seq data. *Nat. Methods* **14,**
707 483–486 (2017).
708
709 Lane, K. *et al.* Measuring signaling and RNA-seq in the same cell links gene expression to
710 dynamic patterns of NF-κB activation. *Cell Syst.* **4,** 458–469 (2017).
711
712 Lee, T. K. *et al.* A Noisy Paracrine Signal Determines the Cellular NF- k B Response to
713 Lipopolysaccharide. *Sci. Immunol.* **2,** ra65 (2009).
714
715 Li, H. *et al.* Reference component analysis of single-cell transcriptomes elucidates cellular
716 heterogeneity in human colorectal tumors. *Nat. Genet.* **49,** 708–718 (2017).
717
718 Lönnberg, T. *et al.* Single-cell RNA-seq and computational analysis using temporal mixture
719 modeling resolves TH1/TFH fate bifurcation in malaria. *Sci. Immunol.* **2,** eaal2192 (2017).
720
721 Mcinnes, L., Healy, J. & Astels, S. hdbscan: Hierarchical density based clustering. *J. Open*
722 *Source Softw.* **2,** 205 (2017).
723
724 Olsson, A. *et al.* Single-cell analysis of mixed-lineage states leading to a binary cell fate choice.
725 *Nature* **537,** 698–702 (2016).
726
727 Peixoto, T. P. The graph-tool python library. (2017).
728 doi:https://doi.org/10.6084/m9.figshare.1164194.v14
729
730 Picelli, S. *et al.* Smart-seq2 for sensitive full-length transcriptome profiling in single cells. *Nat.*
731 *Methods* **10,** 1096–1100 (2013).
732
733 Pierson, E. & Yau, C. ZIFA: Dimensionality reduction for zero-inflated single-cell gene
734 expression analysis. *Genome Biol.* **16,** 241 (2015).
735
736 Qiu, X. *et al*. Reversed graph embedding resolves complex single-cell trajectories. *Nat. Methods*
737 **14,** 979–982 (2017).
738
739 Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of single-
740 cell gene expression data. *Nat. Biotechnol.* **33,** 495–502 (2015).
741
742 Schwalie, P. C. *et al.* A stromal cell population that inhibits adipogenesis in mammalian fat
743 depots. *Nature* **559,** 103–108 (2018).
744
745 Setty, M. *et al.* Wishbone identifies bifurcating developmental trajectories from single-cell data.
746 *Nat. Biotechnol.* **34,** 637–645 (2016).
747
748 Soneson, C. & Robinson, M. D. Bias, robustness and scalability in single-cell differential
749 expression analysis. *Nat. Methods* **15,** 255–261 (2018).
750

751 Subramanian, A. *et al*. Gene set enrichment analysis: A knowledge-based approach for
752 interpreting genome-wide. *Proc. Natl. Acad. Sci. U.S.A.* **102,** 15545–15550 (2005).
753
754 Tavenard, R. tslearn: A machine learning toolkit dedicated to time-series data. (2017).
755
756 Tian, L. *et al*. Benchmarking single cell RNA-sequencing analysis pipelines using mixture
757 control experiments. *Nat. Methods* **16,** 479–487 (2019).
758
759 Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by
760 pseudotemporal ordering of single cells. *Nat. Biotechnol.* **32,** 381–386 (2014).
761
762 Treutlein, B. *et al.* Reconstructing lineage hierarchies of the distal lung epithelium using single
763 cell RNA-seq. *Nature* **509,** 371–375 (2014).
764
765 Vallejos, C. A. *et al*. BASiCS: Bayesian Analysis of Single-Cell Sequencing Data. *PLOS*
766 *Comput. Biol.* **11.6,** e1004333 (2015).
767
768 Vallejos, C. A. *et al*. Normalizing single-cell RNA sequencing data: challenges and
769 opportunities. *Nat. Methods* **14,** 565–571 (2017).
770
771 Wang, B., Collins, J. J. & Newmark, P. A. Functional genomic characterization of neoblast-like
772 stem cells in larval Schistosoma mansoni. *eLife* **2,** e00768 (2013).
773
774 Wang, B., Zhu, J., Pierson, E., Ramazzotti, D. & Batzoglou, S. Visualization and analysis of
775 single-cell rna-seq data by kernel-based similarity learning. *Nat. Methods* **14,** 414–416 (2017).
776
777 Wang, B. *et al.* Stem cell heterogeneity drives the parasitic life cycle of Schistosoma mansoni.
778 *eLife* **7,** e35449 (2018).
779
780 Watts, D. J. & Strogatz, S. H. Collective dynamics of 'small-world' networks. *Nature* **393,** 440–
781 442 (1998).
782
783 Wendt, G.R. *et al.* Flatworm-specific transcriptional regulators promote the specification of
784 tegumental progenitors in Schistosoma mansoni. *eLife* **7,** e33221 (2018).
785
786 Witchley, J. N., Mayer, M., Wagner, D. E., Owen, J. H. & Reddien, P. W. Muscle cells provide
787 instructions for planarian regeneration. *Cell Rep.* **4,** 633–641 (2013).
788
789 Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: Large-scale single-cell gene expression data
790 analysis. *Genome Biol.* **19,** 1–5 (2018).
791
792 WormBase, https://www.wormbase.org, release WS268.
793
794 Xu, C. & Su, Z. Gene expression Identification of cell types from single-cell transcriptomes
795 using a novel clustering method. *Bioinformatics* **31,** 1974–1980 (2015).
796

797   Žurauskien, J. & Yau, C. pcaReduce: hierarchical clustering of single cell transcriptional
798   profiles. *BMC Bioinform.* **17,** 140 (2016).

 Figure legends

Figure 1. The SAM algorithm. (**a**) SAM starts with a randomly initialized kNN adjacency matrix

and iterates to refine the adjacency matrix and gene weight vector until convergence. (**b**) Root

mean square error (RMSE) of the gene weights (top) and the fraction of different edges of the

nearest-neighbor adjacency matrices (bottom) between adjacent iterations (blue) and between

independent runs at the same iteration (orange) to show that SAM converges to the same solution

regardless of initial conditions. The differences between the gene weights and nearest-neighbor

graphs from independent runs are relatively small, indicating that SAM converges to the same

solution through similar paths. (**c**) Graph structures and gene weights of the schistosome stem cell

data converging to the final output over the course of 10 iterations (*i* denotes iteration number).

Top: nodes are cells and edges connect neighbors. Nodes are color-coded according to the final

clusters. Bottom: weights are sorted according to the final gene rankings. (**d**) Network properties

iteratively improve for the graphs reconstructed from the original data (red) but not on the

randomly shuffled data (blue). Dashed lines: metrics measured from the Seurat-reconstructed

graphs.

Figure 2. SAM identifies novel subpopulations within schistosome stem cells. (**a**) UMAP

projections of the manifolds reconstructed by SAM, PCA, and Seurat. SIMLR outputs its own 2D

projection based on its constructed similarity matrix using a modified version of t-SNE. The

schistosome cells are color-coded by the stem cell subpopulations μ, δ', ε_α, and ε_β determined by

Louvain clustering. (**b**) UMAP projections with gene expressions of subpopulation-specific

markers (*eledh*, *nanos-2 cabp*, *astf*, *bhlh*,) and a ubiquitous stem cell marker, *ago2-1*, overlaid.

Insets: magnified views of the expressing populations. (**c**) FISH of *cabp* and EdU labeling of

dividing stem cells in juvenile parasites at 2.5 weeks post infection show that μ-cells ($cabp^+EdU^+$,

arrowheads) are close to the parasite surface and beneath a layer of post-mitotic *cabp*$^+$ cells.

Dashed outline: parasite surface. Right: magnified views of the boxed region. (**d**) FISH of *cabp*

and a set of canonical muscle markers, *troponin*, *myosin*, *tropomyosin*, and *collagen*, shows

complete colocalization in post-mitotic *cabp*$^+$ cells. Images in (**c-d**) are single confocal slices. (**e**)

FISH of *astf* and *bhlh* shows their orthogonal expression in adjacent EdU$^+$ cells (arrowheads).

Bottom: magnified views of the boxed region. Image is a maximum intensity projection of a

confocal stack with a thickness of 12 µm. (**f**) UMAP projection of stem cells isolated from

juveniles at 2.5 and 3.5 weeks post infection. Cell subpopulation assignments based on marker

gene expressions are specified. Right: a magnified view to show the mapping of ε_α- and ε_β-cells.

(**g**) Standardized dispersions as calculated by Seurat plotted vs. the SAM gene weights. (**h**) SC3

AUROC scores plotted vs. the SAM gene weights. Error bars indicate the standard deviation of

SC3 AUROC scores between trials using different chosen numbers of clusters. In (**g**) and (**h**), the

top 20 genes specific to each subpopulation are colored according to the color scheme used in (**a**).

Figure 3. SAM improves clustering accuracy and runtime performance. (**a**) Accuracy of

cluster assignment quantified by adjusted rand index (ARI) on nine annotated datasets (left). Right:

differences between the number of clusters found by each method and the number of annotated

clusters. Smaller differences indicate more accurate clustering. (**b**) RMSE of gene weights output

by SAM averaged across ten replicate runs with random initial conditions for 56 datasets (blue)

and simulated datasets with no intrinsic structure (green, **Methods**). (**c**) Runtime of SAM, SC3,

SIMLR, and Seurat as a function of the number of cells in each dataset. SC3 and SIMLR were not

run on datasets with >3000 cells as the run time exceeds 20 minutes.

Figure 4. SAM improves the analysis of datasets with varying network sensitivities. (**a**) Network sensitivity of all 56 datasets ranked in descending order. Blue: the nine benchmarking datasets used in **Figure 3a**. Sensitivity measures the robustness of a dataset to changes in the selected features (**Methods**). (**b**) The network sensitivity plotted against the fraction of genes with SAM weight greater than 0.5 (in log scale) with Spearman correlation coefficient specified in the upper-right corner. (**c**) Fold improvement of SAM over Seurat for NACC, modularity, and spatial dispersion with respect to sensitivity for all 56 datasets. These ratios are linearly correlated with network sensitivity with Pearson correlations (r^2) specified in the upper-left corner of each plot.

Figure 5. Robust feature selection improves cell clustering and manifold reconstruction. (**a**) Network sensitivity, ARI, NACC, modularity, and spatial dispersion with respect to corruption of the Darmanis dataset, in which we swap random pairs of gene expressions with the number of swaps ranging from 0-50% of the total number of elements in the data (**Methods**). Performance is compared between SAM (blue), Seurat (red), and Seurat rescued with the top-ranked SAM genes (indigo). Error bars indicate the standard deviations across 10 replicate runs. The errors for points with no bars are too small to be seen. (**b**) Comparison of the area under curve (AUC) of the metrics in (**a**) with respect to data corruption for all nine datasets. Error bars indicate the standard deviations across 10 replicate runs. The errors for data with no error bars are too small to be seen.

Figure 6. SAM captures the cellular activation dynamics in a stimulated macrophage dataset. (**a**) GSEA analysis (left) and SAM projections (right) of the activated macrophages[7] before (top) and after (bottom) removing cell cycle effects. Teal: significantly enriched gene sets determined

by the significance threshold of 0.25 for the False Discovery Rate (FDR, dashed lines). Bottom:

the two clusters are denoted as MT and M with colors representing the time since LPS induction.

Arrows: evolution of time. (**b**) TNFα is enriched in the MT cluster. (**c**) Diagram of NF-κB

activation in response to LPS stimulation via the Myd88 and TRIF signaling pathways. (**d**) Log$_2$

fold changes of the average expressions of selected inflammatory genes in the MT cluster vs. the

M cluster. All genes are significantly differentially expressed between the two clusters according

to the Welch's two-sample t-test ($p < 5 \cdot 10^{-3}$). (**e**) Representative traces for transient (left) and

prolonged (right) NF-κB activation (**Methods**). (**f**) Cells with prolonged NF-κB response (denoted

as P) are primarily in the MT population. (**g**) Seurat and SIMLR projections show that they fail to

order the cells by time since LPS induction and do not identify cell clusters representing the

different modes of NF-κB activation.

Figure 1 – Figure supplement 1. Quality control of library preparation and sequencing of

the schistosome stem cells. (**a**) Histograms of *h2a* qPCR measurements in 2.5- (left) and 3.5-

(right) week samples. (**b**) Scatter plot of gene count (>2 TPM) vs mapped read count of individual

sequenced cells. Cells with low gene count or *h2a* expression are discarded and filtered from

analysis (red) and the remaining cells are analyzed (blue). The number of final cells kept for

analysis is annotated on the top left corner of each plot.

Figure 2 – Figure supplement 1. μ-cells express ubiquitous stem cell marker and population

specific genes. UMAP projections with gene expressions of (**a**) stem cell markers and (**b**) μ-cell-

specific genes overlaid.

Figure 3 – Figure supplement 1. SAM converges to a stable solution independent of random

initial conditions and is robust to the number of nearest neighbors and choice of distance

metric. (**a**) RMSE of gene weights between adjacent iterations within a run, averaged across ten

replicate runs. (**b-c**) Average ARI scores for the nine annotated benchmarking datasets when

varying (**b**) the number of nearest neighbors, k, from 10 to 30 or (**c**) the choice of distance metric

(Euclidean or Pearson correlation). Error bars indicate standard deviations of ARI scores across

the different values of k and distance metrics. The errors for data with no error bars are too small

to be seen.

Figure 6 – Figure supplement 1. Cluster-specific marker genes before and after removing

cell cycle effects. UMAP projections with marker genes specific to the dividing cells (**a**) and the

MT cluster (**b**) overlaid.

Figure 6 – Figure supplement 2. SAM groups cells based on NF-κB activation dynamics while

other methods cannot. (**a**) UMAP projection of the macrophage cells after the removal of cell

cycle effects. Cells with prolonged NF-κB dynamics are highlighted in red. (**b**) UMAP and t-SNE

projections for Seurat and SIMLR, respectively, after the removal of cell cycle effects. Cells with

prolonged NF-κB dynamics are highlighted in red. (**c**) UMAP projections with three MT-specific

marker gene expressions overlaid.

Figures

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

921   **Figure 6.**

923   **Figure 1 – Figure supplement 1.**

925 **Figure 2 – Figure supplement 1.**

926

927　**Figure 3 – Figure supplement 1.**

 Figure 6 – Figure supplement 1.

Figure 6 – Figure supplement 2.

Supplementary Table legends:

Supplementary Table 1: Ranked gene list with high SAM weights in the schistosome stem

cell data. Gene IDs and annotations are given in the *S. mansoni* genome version 9 (WormBase,

WS268). Genes are assigned to the cluster corresponding to the marker gene, *nanos-2*, *cabp*, *astf*,

or *bhlh*, that they have the highest correlation with. Genes found in our prior work[12] to be enriched

in subsets of stem cells are specified.

Supplementary Table 2: Datasets used in this study. Accession numbers, library size

normalization methods, data preprocessing methods, sensitivity scores, and corresponding

references are provided for each dataset. Accession numbers with asterisks indicate datasets that

are sourced from the *conquer* database (Soneson and Robinson, 2018). Accession numbers with

crosses indicate the nine well-annotated datasets that were used for benchmarking.

Supplementary Table 3: Cloning primer sequences used for generating riboprobes for the

FISH experiments. Functional annotations of the genes were given in the *S. mansoni* genome

version 9 (WormBase, WS268).