Synchronised brood transport by ants occurs without communication

Danielle P. Mersch[1,*], Jean-Pierre Eckmann[2], Alessandro Crespi[3], Laurent Keller[1]

[1] Department of Ecology and Evolution, University of Lausanne, Switzerland

[2] Département de Physique Théorique et Section des Mathématiques, Université de Genève, Switzerland

[3] Biorobotics Laboratory, Ecole polytechnique fédérale de Lausanne, Switzerland

[*] *corresponding authors*: danielle.mersch@gmail.com , laurent.keller@unil.ch

Abstract

Collective behaviours in societies such as those formed by ants are thought to be the result of distributed mechanisms of information processing and direct decision-making by well-informed individuals, but their relative importance remains unclear. Here we tracked all ants and brood movements to investigate the decision strategy underlying brood transport in nests of the ant *Camponotus fellah*. Changes in environmental conditions induced workers to quickly transport the brood to a preferred location. Only a minority of the workers, mainly nurses, participated in this task. Using a large number of statistical tests we could further show that these transporters omitted to recruit help, and relied only on private information rather than information obtained from other workers. This reveals that synchronised group behaviour, often suggestive of coordinated actions among workers, can also occur in the complete absence of communication.

Introduction

The success of group actions frequently relies on communication between individuals. Communication is manifest in animal groups as different as jellyfish that use bioluminescence to locate each other and team up[1], prairie dogs that call

to warn their family of predators[2] and honeybees that use waggle dance to signal

a food source to nest mates[3,4]. In all these cases communication serves to

enhance the efficiency and safety of the group. However, communication is

complex. It requires that the sender recognizes the appropriate circumstances

and produces a correct signal, and that the receivers are able to understand the

signal and react appropriately. These inherent difficulties constrain when and

under what conditions groups of animals might communicate.

In ant societies communication is widespread and individuals make use of an array of

olfactory, vibrational and tactile communication strategies. Therefore, communication

is often assumed to be underlying all group behaviours[5,6,7,8,9,10]. Ants optimize foraging

by creating pheromone trails[11,12], and by recruiting help to retrieve food through

tandem runs, a method whereby a knowledgeable ant induces a naive ant through

tactile and chemical signals to follow it[13]. In emergencies, ants release highly volatile

alarm pheromones[11]. If a nest is destroyed knowledgeable ants first lead tandem runs

to new nest sites before switching to brood transport[14]. In all these instances

communication is manifest and beneficial to the society. Pheromone trails and tandem

recruitment reduce the risks of random food searches and ensure that a sufficient

number of workers locate and retrieve food before it disappears, thereby enhancing

the colony's chances of survival and reproduction. Similarly in emergencies the

survival of the colony is at stake. Alarm pheromones ensure that workers are alerted

and leave the nest[15] for fight or flight. Tandem runs ensure that a sufficient number of

workers know the location of a safe alternative nest before evacuating brood[9].

However, there is a range of other group behaviours such as nest construction or

brood relocation where the advantages of communication are less apparent. For

example, many ant species regularly move brood within a nest and between nests to

raise offspring under optimal temperature and humidity[16,17,18,19,20]. Such controlled

responses to environmental variables are a central part of colony organisation in social

insects because they have direct impacts on colony growth, metabolic expenditure,

survival and reproduction[19,20,21].

In this study we conduct a detailed analysis of brood transport in the ant *Camponotus*

fellah to investigate to what extent workers communicate to displace the brood after

changes in environmental conditions. We took advantage of the fortuitous observation

that workers moved the brood in response to environmental changes in three colonies

(colony size=197, 192, and 206 workers, brood items=150, 60 and 35) to investigate

whether workers communicate about observed changes in local conditions. In *C.*

fellah, as in most other ants, workers quickly respond to environmental changes to

move the brood to the nest regions with the best conditions [22, 23, 24, 25].

Results

Colonies transport brood in synchrony

In each of the three colonies, and each of the nights, workers responded to the

environmental change, initiating brood transport 22.4±6.2 minutes (mean±SEM) after

the light was turned off in the tunnel (Fig. 1). There were neither consistent

differences across colonies, nor a change in the response delay over the three days

(ANCOVA, colony: $F=0.9$, $p=0.37$; day: $F=0.77$, $p=0.41$; interaction colony*day:

$F=0.41$, $p=0.69$). On average workers took 160.0±48.0 minutes to move all the brood

from the nest to the tunnel once transport was initiated. Workers also performed this

task in synchrony with multiple workers transporting in parallel during 66.1±28.0% of

the time. The average time taken by a worker to transport one brood item was

36.7±4.0 seconds (see Supplementary Video 1). Workers that transported more brood

items were faster to transport brood than those transporting fewer brood items

(Spearman rank correlation: $\varrho=-0.51$, $p<0.0001$; Supplementary Fig. 2). There were

again neither significant differences across colonies, nor over days, in the time

required to transport all the brood (ANCOVA on log-transformed duration: colony:

$F=1.5$, $p=0.31$; day: $F=1.3$, $p=0.24$; colony*day: $F=1.2$, $p=0.40$).

Fig 1. Brood transport dynamics on three consecutive days in three colonies. The black line indicates the cumulated number of brood transports to the tunnel of all workers. Each coloured circle represents a single brood transport event by one worker, and data are shown as cumulative transports. Different colours represent different workers.

A small minority of a colony's workforce transports brood

The number of workers involved in brood transport was consistently low, with only 8.1±1.1 workers (4.1%±0.6% of the workforce) participating in brood transport on any given day in any given colony (Fig. 2). Colonies did not differ in the distribution of the workload among workers, and there was no significant change over days in the way the workload was distributed among transporters (ANCOVA: colony: F=0.40 p=0.67; day: F=0.15 p=0.86; colony*day: F=0.14, p=0.97). However, there was variation among transporters in their relative contribution with the notable effect that more than 80% of all brood transports were performed by less than 1.8% of all workers. In addition, there was also a high worker turnover with 66.9±5.2% of the transporters working on a single night, while only 18.8±11.9% of the transporters worked on all three nights. Importantly, however, the persistent transporters were responsible for 44.3±25.3% of all transports while those that worked a single night

110   contributed together to 24.8±18.7% of the transports.

Fig 2. **The workload is distributed unevenly among the transporters.** Absolute numbers
are given in the histogram, and proportions are indicated in the pie chart. Arrows indicate
workers that transport without being privately informed (i.e. they had not visited the tunnel
before starting to transport). Transporters with ID labels in black are nurses, while those with
labels in grey belong to the cleaner or forager groups.

Transporters are nurses

To determine whether brood transport was preferentially conducted by a specific

group of workers, we used the Infomap algorithm[26] to determine the daily interaction

networks of workers and assign each of them to a specific social group[27]. Colonies

had on average 55.9%±11.3% nurses, 16.5%±4.9% cleaners and 25.1%±7.4%

foragers (Fig. 3). Nurses were 3.8 times more likely to transport than cleaners, and 7.3

more likely to transport than foragers (ANOVA, F=51.38, p<0.0002). There was also

an effect of age, with transporters being on average younger (83.5 days) than non-

transporters (119.5 days; Kruskal-Wallis: $\chi^2=12.1$, p<0.001). This effect was due to

age differences between the three groups of workers (average age nurses 93.8 days,

cleaners 124.2 days, foragers, 159.4 days; Kruskal-Wallis: $\chi^2=138.6$, p<0.00001).

When only nurses were considered, there was no significant age difference between

transporters and non-transporters (Kruskal-Wallis: $\chi^2=0.81$, p=0.37; insufficient data

was available to conduct similar tests for nest cleaners and foragers).

Fig 3. Transporters are mainly nurses. The network shown is that of colony 10 on day 4.
Each node represents a worker, and links between nodes are shown for workers who had more
than 10 interactions on that day. The network layout is a spring embedded layout. Group
membership is indicated by the node colour: nurse (blue), cleaner (green), forager (orange).
Red-shaded circles around nodes highlight transporters, with light red indicating transports on
one day, medium red indicating transports on two days, and dark red indicating transports on
three days.

Transporters gather information themselves

To determine whether workers make use of information available to others to decide

when to initiate brood transport, we tracked the information available to each worker

after the light was turned off. Because the nest entrance was constructed with two 90°

bends and painted in matt black on the inside thereby preventing light from entering

the nest, the only means for workers to know whether there was light in the tunnel

was to access it. Workers were therefore considered as having private information

once they had left the nest for at least three seconds, which is the minimum amount of

time an ant needs to reach the tunnel and return to the nest. Ants were considered as

socially informed once they interacted with a privately informed worker.

At the start of brood transport, only 31.6%±2.9% of all workers and 37.8%±8.7% of

the nurses had private information. However, almost all transports (99.8%) were

performed by privately informed ants. Of the seven workers, which had not visited the

tunnel before initiating brood transport, four had transported brood on previous days

(Fig. 2). The three remaining workers had visited the tunnel the nights before when it

contained brood. Thus, these transporters may have used this information together

with circadian timing to initiate transport[24,25]. Therefore, these observations suggest

that private information is the primary or only source of information workers use to

decide when and where to transport the brood.

Transporters neither communicate nor recruit help

Five lines of evidence further support the view that workers do not use information

obtained from other workers to initiate brood transport. First, transporters did not

increase their interaction frequency with other workers once it was dark in the tunnel.

The rate of interactions in the hour preceding light-off was not significantly different

from the rate during the interval between light-off and the first brood transport

(Kruskal-Wallis: χ^2=0.05, p=0.82; Supplementary Fig. 3). Second, transporters did not

change their activity after interacting with a privately informed ant. Their increase in

speed —a signature of information transfer in ants[28]— was similar after interacting

with a privately informed or an uninformed ant (Kruskal-Wallis: χ^2=2.8, p=0.09, see

Supplementary Table 1). Third, brood accumulating in the tunnel did not speed up the

recruitment of additional transporters. The average time elapsed before one additional

worker contributed to brood transport was 16.6±3.4 min. The number of workers

already participating in brood transport did not alter the time needed to rally an

additional worker (Spearman rank correlation: ϱ=0.06, p=0.60; Supplementary Fig.

4). Fourth, the first interaction with a privately informed ant did not trigger a change

in behaviour. After interacting with a privately informed ant, transporters and non-

transporters were neither more likely to approach the nest entrance (Wilcoxon signed

rank test: transporters: V=1232, p=0.79; non-transporters: V=495789, p=0.97) nor to

orient towards it (Rao's spacing test for uniformity: transporters: Test

Statistic=139.98, p>0.05 with a critical value=148.34; for non-transporters: Test

Statistic=134.13, p>0.05 with a critical value=136.94; Fig. 4A, 4B). Simulations were

conducted to determine the expected effect if 90%, 50%, 10% or 0% of the

transporters were able to understand a message that they should go to the tunnel after

interacting with a privately informed ant (Fig 4C). These simulations revealed that the

observed pattern was consistent with a complete lack of communication between

privately informed ants and non-informed transporters. Finally, and most importantly

we did not observe any successful recruitment through tandem running although these

ants are capable of tandem running (see Supplementary Videos 2, 3).

Fig 4. No evidence for communication between workers. (A) Changes of trajectory
following the first interaction with a privately informed ant. Arrows indicate the trajectories
after the first interaction with a privately informed ant and the dotted lines the trajectories just
before this interaction. Transporter trajectories are in red and those of other ants in black. The
blue circle indicates the nest entrance. Data shown are those of colony 1 on day 5.
(B) Distribution of directions after the first interaction with a privately informed ant. Each dot
represents the direction relative to the nest entrance of a single worker on a given day. Red
dots indicate transporters and blue dots (forming a ring) indicate other ants. The arrow
indicates the direction of the nest entrance. (C) Expected change in direction from simulated
data in which 0% (blue), 10% (cyan), 50% (red) or 90% (black) of the ants understood a
message. Each dot is the average movement towards the nest entrance of 66 simulated
transporters. The cross and ellipse show the average and the standard deviation across 40
simulations with the same set of parameters. The star shows the average of the observed data.

Colonies do not use quorum sensing to initiate brood transport

At the colony level there was also no indication of a system of quorum sensing

leading to the onset of brood transport. At the time of first transport, the percentage of

privately and socially informed workers and the percentage of workers in the tunnel

varied greatly (privately informed: 0.6% to 12.0%; socially informed: 1.9% to 47.5%,

ants in tunnel: 6.0% to 19.4%; Fig. 5). Furthermore, the use of a quorum would imply

that colonies deferred the onset of brood transport on some days for almost one hour

after reaching the quorum, while starting to transport just minutes after reaching the

quorum on other days (delays for privately informed: 4.3–59.8 minutes; socially

informed: 2.8–58.8 minutes; ants in tunnel: 5.4–59.1 minutes). Given that the

variability was large for both the quorum threshold and the delay until transport onset,

it seems unlikely that a minimum colony level information threshold or a minimum

ant proportion in the tunnel needs to be reached for brood transport to be initiated.

Fig 5. No evidence for a quorum threshold triggering brood transport. Each line shows
the percentage of ants: privately informed ants in red, socially informed ants in blue, ants in
the tunnel in grey. The vertical line indicates when the first transport occurred, and the dashed
lines highlight the percentages of ants at the time of first transport.

Finally, our analyses also revealed high consistency in the direction of brood transport

(Supplementary Fig. 5). Overall, there were only 20 return-transports (2.3%) among

the 859 transports recorded. Interestingly, the majority of the workers (69.2%)

performing return transports did not transport brood to the tunnel while the vast

majority (91.7%) of the workers transporting brood to the tunnel did not perform

return-transports.

Discussion

The use of an automated system allowed us to obtain detailed and individual-level

information on the processes regulating brood transport in response to environmental

changes, a process central to the organization of social insect colonies. Overall,

workers quickly transported the brood to the preferable location after the light was

turned off, and workers almost never transported brood in the wrong direction.

However, this seemingly coordinated transport occurred without any detectable sign

of communication among workers. While workers frequently interacted, these

interactions resulted in no visible change in the behaviour of the transporters, even if

the interaction partner had knowledge about the tunnel being dark. Instead,

transporters appeared to rely exclusively on self-gathered information, because they

initiated brood transport only after having noticed the change of state of the tunnel

themselves. Together, these data indicate that synchronised behaviour at the colony

level can occur without communication.

Visual inspections of our videos also revealed no evidence that workers relied on

chemical signals to initiate and communicate brood transport. Transporters never

dragged their gaster over the ground, as ants typically do when depositing trails. There

were also no instances of worker tandem running, thereby excluding targeted

recruitment that could have been mediated by secretions from a gland[13]. The only

targeted recruitment that we observed was that of the queen and in one instance that of

non-transporting workers (see Supplementary Videos 2, 3). In these cases a worker

approached the head of the queen or worker and pulled on her mandibles, with the

effect that the pulled ant became active and followed the worker in a tandem-run to

the tunnel.

The observed lack of communication is likely due to the inherent difficulty of reliably

communicating a message in a noisy environment. Communication requires that an

informed individual intentionally encodes a message, transmits it successfully, and

that an uninformed individual is able to receive it, decode it, and act upon it[29]. Ants

have a limited ability to convey a message through tactile communication alone[28, 30,31].

In addition, the density of workers is extremely high in the nest, resulting in numerous

interactions not only with informed individuals but also with uninformed ones. Such a

situation leads to a very noisy system where conflicting feedbacks may readily

compromise any attempts of communication. Moreover, investing time in recruiting a

helper would only beneficial if the time needed for successful recruitment is short,

and if recruitment occurs early on (see Figure 6).

Figure 6. Cost and benefit of successful recruitment. The time invested in recruiting help is
indicated in red. The time gained from recruiting a helper is shown in blue, and the time lost
due to recruiting help in orange. (A) Recruiting a helper early on after the task is initiated and
with little time investment enables faster completion of the brood transport than without a
helper. (B) Recruiting a helper early on but with high time investment delays the completion
of the brood transport compared to a situation without a helper. (C) Recruiting a helper later
while the task is performed also delays the completion of the brood transport compared to a
situation without a helper.

Our observation that transporters check the state of the tunnel themselves, before

starting to transport brood, suggests that individual workers gather cues from the

environment before deciding to transport brood. The most likely cues used by the

transporters in our experiments are the confinement, absence of light and presence of

workers in the tunnel[32,33]. The use of cues for decision-making also occurs in other

ants, and for processes unrelated to brood transport. For instance, in harvester ants,

potential foragers decide whether or not to initiate a foraging trip based on the

frequency with which they meet returning foragers[34,35]. Workers of the black garden

ant *Lasius niger* use the chemical profile of the nest wall and their own body size

compared to the height of nest pillars as cues to decide whether to switch from wall

building to building a roof[36]. These data, together with our results, suggest that the use

of cues as a mean to obtain private information might be more widespread and easier

to implement in ant colonies than information exchange through tactile

communication.

The use of cues combined with the lack of communication and the absence of a
quorum means that transporters most likely decide independently of each other
whether, when and where to transport the brood. Such individual-led decisions are
further supported by rare instances in which a worker mistakenly returned brood from
the tunnel to the nest, while transporters were already moving brood to the tunnel.
Interestingly, the vast majority of transporters arrived at the same decision and
transported brood from the nest to the tunnel. This strong uniformity in behaviour
suggests that there is high homogeneity in preferences among group members.

Our results indicate that colonies can display synchronized behaviour without
communicating thus emphasizing that not all group-level behaviours in social insects
are driven by communication. We suspect that communication is context-dependant
and only used when cue-based options are insufficient. For instance, the
communication that precedes brood transport in house-hunting ants occurs in the
context of an emergency after their nest has been destroyed[9,14]. In contrast,
synchronization without communication is optimal when reliable communication is
expensive, hard to achieve, or when perfect synchrony is not needed[29,37]. It can be
achieved if workers share similar preferences and react to the same cues, which are
limited in time. In our experiments light in the tunnel acted as this strong time-limited
cue. Synchronized group behaviour exists also in solitary bees, who congregate at
nesting sites for reproduction[38], bats and starlings that converge at seasonal feeding
and sleeping spots[39,40] and Mormon crickets, who migrate in masses in search for salt
and proteins[41]. In ants simulations further suggest that food choice during foraging
could be achieved without communication through individual learning and
preference[42].
320

Our results also revealed that only a tiny fraction of the individuals, 1.5%–6.6% of the
colonies' workforce —as few as three workers in some cases— contributed to brood
transport. Moreover, within colonies there was strong variation in the relative
contribution of workers with more than 80% of all transports being carried out by less
than 1.8% of the workers. Similar fractions of transporters and workload disparities
were observed in colony emigrations of *Formica sanguinea* and *Camponotus*

sericeus[43]. The large variability in behaviour is puzzling and we offer two possible

explanations. There could be specialist nurses that focus on brood transport. Indeed

nine out of 48 transporters moved brood every single night and did slightly less than

half of the work, thus acting as key individuals[44] during the brood displacement.

Similar specialization has been reported for foraging, brood care, stone

collection[45,46,47] and could result from inherent and consistent differences between

workers, for example in motivation, physiology, or sensory threshold[48,49]. Another

explanation is that transporters represent a varying subset of the nurses, whose

likelihood to transport depends on the individual's state in the early night. This idea is

supported by the observation that two thirds of the transporters only worked a single

night.

Importantly, a small minority of transporters imposed their transport decision on the

colony. Such an outcome was only possible because the other workers did not oppose

the brood transports or if they did so initially, never persisted in their opposition.

Minority-driven behaviour occurs also in *Paratrechina longicornis* ants, where a

single worker can temporarily decide the pull direction during collective transport[50].

Our results therefore highlight that a small minority of the workforce can determine

the colony fate through persistent activity in a largely indifferent society. Similar

observations exist for fish schools and human crowds where few knowledgeable

individuals can lead large groups of uninformed individuals to a new location[51,52].

Ultimately, the social unresponsiveness of the majority might be the optimal strategy

because social unresponsiveness can ensure that the colonies react to environmental

change while also being robust to noise and avoiding losses in information accuracy

resulting from an over-reliance on social information[53].

Acknowledgements. We thank M. Chapuisat, O. Feinerman, N. Stroeymeyt, T.

Richardson, T. Kay and S. McGregor for comments on an earlier version, and B.

Sutcliffe for proofreading. DPM was supported by a grant from HFSP, JPE was

supported by an advanced ERC grant, LK was supported by grants from the Swiss

NSF and an advanced ERC grant.

Contribution. DPM and LK planned the experiment. DPM and AC designed the experimental system and performed the experiment. DPM and JPE analysed and interpreted the data. DPM wrote the paper with input from JPE and LK. All authors revised the paper.

Material and Methods

The three colonies were each established from a single queen collected after a mating flight in Tel Aviv on March 23rd 2007. The experiment started when queens were 3 years old, out of a maximum life span of 26 years[54]. At the start of the experiment, colonies each comprised a queen, brood and 197, 192 and 206 workers, for colonies 1, 2 and 3 respectively. The colony sizes were those naturally reached by queens of that age, and reflect normal growth rates in the laboratory; no data are available for field colonies. All workers were the offspring of a single queen, which in *Camponotus fellah* is usually singly-mated[55].

To determine workers' age, new-born workers were paint-marked on a weekly basis during the 12 months preceding the experiment. Because 38 out of the 45 transporters were nurses, we limited the analysis on the effect of age to nurses only.

During experiments colonies were kept in a dark nest chamber connected by a 60 cm long and 1cm wide tunnel to a foraging chamber. The tunnel and the foraging box had 12h light-12h dark cycles, and the ants had access to food (gelatinous sugary water) and water in the foraging box. The temperature (30 °C), humidity (60%), light (~500 Lux), and food supply were computer-controlled, and both chambers were filmed from above with high-resolution monochrome cameras operating under infrared light, as previously described[27] (Supplementary Fig. 1). All colony members were video-tracked using fiducial identification labels over 14 consecutive days. We recorded the position and orientation of all individuals twice per second.

During the night, workers transported the brood to the tunnel and brought it back to the nest at dawn, presumably because they prefer to keep the brood in a confined environment rather than an open environment when both are dark. We tracked the transport of brood items manually during three consecutive nights. A brood transport

was defined as the time interval from when an ant collected one (or several) brood
items from the nest box, to when the ant disappeared with it into the tunnel. We also
recorded cases where brood was transported from the tunnel to the nest. In these
return-transports, the transport was defined as the time interval from when the ant
entered the nest with brood until the ant dropped the brood. For each transporter and
each night we defined its workload as the number of transports during that night and
its work time as the time from the start of its first transport until the end of its last
transport. Using the work times of all workers, we estimated synchrony as the
percentage of time during which at least two workers worked in parallel. We also
visually inspected the videos for instances of tandem running, that is events where one
ant guides another ant to the tunnel. A tandem-run results in successful recruitment if
the follower ant subsequently starts transporting brood. We did not track brood
transports in the mornings when the lights turned on in the tunnel, because in these
conditions all ants in the tunnel were immediately informed of the environmental
change, thus making the question of communication inane.

To determine group membership of each worker, *i.e.* nurse, cleaner or forager, we
used the same approach as in Mersch *et al*. (2013)[27]. In brief, we inferred all social
interactions between workers based on their distance and orientation, and analysed the
social networks with the Infomap algorithm[26] to assign each worker to a group.
Because the majority of workers were in the tunnel at night and thus undetectable
with our tracking setup, we built daily interaction networks using only data collected
between 8am and 7pm, when the majority of workers were detectable.

To measure the speed change following interactions, we calculated the speed during
the 10 seconds prior to the interaction and during the 10 seconds after the interaction.
We included only those interactions for which we had data on the speed before the
interaction for both partners and on the speed after the interaction for the focal ant. As
a consequence, 50 interactions (10.2%) were excluded from the analysis. Excluding
these interactions had neither an impact on the average duration of an interaction
(10.5±29.9 s *vs*. 10.3±30.5 s) nor on the proportion of interactions with privately
informed partners (7.72% *vs*. 7.69%). To further ensure that our results are not
influenced by the chosen interval (10 s), we repeated the same analyses for shorter

(5 s) and longer (20 s) time intervals. Because the results were the same for all time
intervals (see Supplementary Table 1), we only report data for the 10-second interval.

To investigate whether a privately informed ant can communicate information about
the change of state in the tunnel to its interaction partner we estimated the change in
trajectory of each worker following its first interaction with a privately informed ant.
We calculated the heading of the ant's trajectory after it had moved away from the
interaction point, transforming data of all colonies so that an orientation of 0°
corresponds to an orientation towards the nest entrance. We also calculated the
distance to the entrance at the time of the interaction and after the ant had moved at
least 2 cm ($\approx$ queen body length) away from the interaction point. Workers who did
not interact with a privately informed ant before the end of the brood transport were
not included in the analysis (351 out of 1785 ant-days excluded).

To estimate how communication about the change of state in the tunnel could modify
the trajectory of workers, we generated simulated datasets in which 0%, 10%, 50% or
90% of the transporters moved toward the nest entrance after interacting with a
privately informed ant. Understanding the message meant that one bit —that is, one
unit of information— was transferred from the privately informed ant to the
transporter. Such one-bit information could convey two options —towards and away
from nest entrance— and signal to the transporter to move towards the nest entrance.
Each dataset was the average of 66 simulated direction vectors v_j defined as

$v_j = (\cos(\alpha_j), \sin(\alpha_j))$

with α_j being the angle of the direction relative to the line connecting the interaction
point with the nest entrance. For each informed transporter, we randomly chose a
direction from a uniform distribution limited to angles between $-\pi/2$ and $\pi/2$, for all
other transporters we randomly chose an angle from a uniform distribution between $-$
π and π. We repeated this process 40 times for each information level. We also
calculated the average direction of the 66 transporters from the observed data.

To test whether a quorum triggered the observed brood transport, we determined the
number of ants, the number of informed ants, and the number of ants in the tunnel at

the time of the first brood transport. To estimate whether the quorum induced brood

transport, we also calculated the duration between the time the quorum was reached

for the first time and the first brood transport. Because the estimated quorum varied

between colonies and days, we calculated the delays for all colonies and days using

the smallest estimated quorum threshold.

We performed all statistical analysis in R (Version 3.4.0)[56]. When the test assumptions

were met, we used two-tailed parametric tests and included the colony ID as a random

factor in our analysis; otherwise we used non-parametric tests. For statistical tests on

colonies, each colony was one replicate. For statistical tests on individual workers,

each transporter on each day was a replicate. The data analysis code will be available

as a zip file.

The data used to prepare all figures and perform statistical tests will be available on

Dryad DOI after publication in a journal.

References

1. Kaartvedt S, Ugland KI, Klevjer TA, Røstad A, Titelman J, Solberg I (2015)
Social behaviour in mesopelagic jellyfish. Scientific Reports 5:11310. DOI:
10.1038/srep11310

2. Kiriazis J, Slobodchikoff CN (2006) Perceptual specificity in the alarm calls
of Gunnison's prairie dogs. Behavioral Processes. 73:29–35. DOI:
10.1016/j.beproc2006.01.015

3. Rohrseitz K, Tautz J (1999) Honey bee dance communication: Waggle run
direction coded in antennal contacts? J Comp Physiol A 184:463–470.

4. von Frisch K. Tanzsprache und Orientierung der Bienen. (Springer Verlag,
Heidelberg 1965) 566

5. Conrad L, List C (2009) Group decisions in humans and animals: a survey.
Philosophical Transactions of the Royal Society of London B: Biological
Sciences 364:719–742. DOI: 10.1098/rstb.2008.0276

6. Pratt SC (2005) Quorum sensing by encounter rates in the ant *Temnothorax*
albipennis. Behav Ecol 16:488–496. DOI: 10.1093/beheco/ari020

7.   Conrad L, Roper TJ (2005) Group decision-making in animals. Nature
421:155-8. DOI: 10.1038/nature01294

8.   Seeley TD (2003) Consensus building during nest-site selection in honey bee
swarms: the expiration of dissent. Behav Ecol Sociobiol 53:417. DOI:
10.1007/s00265-003-0598-z

9.   Franks NR, Pratt SC, Mallon EB, Britton NF, Sumpter DJT (2002)
Information flow, opinion polling and collective intelligence in house-hunting
social insects. Philosophical Transactions of the Royal Society B: Biological
Sciences 357:1567–1583.

10. Britton NF, Franks NR, Pratt SC, Seeley TD (2002) Deciding on a new home:
how do honeybees agree? Proceedings of the Royal Society of London B:
Biological Sciences 269:1383–1388. DOI: 10.1098/rspb.2002.2001

11. Hölldobler B, Wilson EO, *The Ants* (Springer, Berlin, 1990).

12. Attygalle AB, Morgan ED, Berridge MJ, Treherne JE, Wigglesworth VB. Ant
Trail Pheromones, in Advances in Insect Physiology (Academic Press 1985)

13. Möglich M, Maschwitz U, Hölldobler B (1974) Tandem calling: A new kind
of signal in ant communication. Science 186:1046–1047. DOI:
10.1126/science.186.4168.1046

14. Stroeymeyt N, Franks NR, Giurfa M (2011) Knowledgeable individuals lead
collective decisions in ants. J Exp Biol 214:3046–3054. DOI:
10.1242/jeb.059188

15. Amoore JE, Palmieri G, Wanke E, Blum MS (1969) Ant alarm pheromone
activity: Correlation with molecular shape by scanning computer. Science 165:
1266–1269. DOI: 10.1126/science.165.3899.1266

16. Kadochová S, Frouz J (2013) Thermoregulation strategies in ants in
comparison to other social insects, with a focus on *Formica rufa*.
F1000Research 2:280. DOI: 10.12688/f1000research.2-280.v2

17. Abril S, Oliveras J, Gómez C (2010) Effect of Temperature on the
Development and Survival of the Argentine Ant, *Linepithema humile*. J Insect
Science 10:97. DOI: 10.1673/031.010.9701

18. Fielde AM (1905) Temperature as a factor in the development of ants. With
further observations on ants deprived of food. Biological Bulletin 9:361–367.
DOI: 10.2307/1535632

19. Porter SD (1988) Impact of temperature on colony growth and developmental
rates of the ant, *Solenopsis invicta*. J Insect Physiol 34:1127–1133

20. Porter SD, Tschinkel WR (1993) Fire ant thermal preferences: behavioral
control of growth and metabolism. Behav Ecol Sociobiol 32:321–329.

21. Penick CA, Diamond SE, Sanders NJ, Dunn RR (2016) Beyond thermal
limits: comprehensive metrics of performance identify key axes of thermal
adaptation in ants. Functional Ecology 31:1091–1100. DOI: 10.1111/1365-
2435.12818

22. Steiner A (1929) Temperaturuntersuchungen in Ameisennestern mit
Erdkuppeln, im Nest von *Formica exsecta* Nyl. und in Nestern unter Steinen.
Z vergl Physiol 9:l–66.

23. Kay CAR (1978) Preferred temperatures of desert honey ants (Hymenoptera:
Formicidae). J Thermal Biol 3:213–217

24. Roces F, Núñez JA (1989) Brood translocation and circadian variation of
temperature preference in the ant *Camponotus mus*. Oecologia 81:33–37. DOI:
10.1007/BF00377006

25. Roces F, Núñez JA (1995) Thermal sensitivity during brood care in workers of
two *Camponotus* ant species: Circadian variation and its ecological correlates.
J Insect Physiol 41:659–669. DOI: 10.1016/0022-1910(95)00019-Q

26. Rosvall M, Bergstrom CT (2008) Maps of random walks on complex
networks reveal community structure. Proc Natl Acad Sci USA 105:1118–
1123

27. Mersch DP, Crespi A, Keller L (2013) Tracking individuals shows spatial
fidelity is a key regulator of ant social organization. Science 340:1090–1093.
DOI: 10.1126/science.1234316s

28. Razin N, Eckmann JP, Feinerman O (2013) Desert ants achieve reliable
recruitment across noisy interactions. J R Soc Interface 10:20130079. DOI:
10.1098/rsif.2013.0079

29. Maynard-Smith J, Harper D. Animal Signals. 1st ed. Oxford: Oxford
University Press; 2003.

30. Hölldobler B (1971) Recruitment Behavior in *Camponotus socius* (Hym.
Formicidae). Z vergl Physiol 75:123–142.

31. Traniello JFA (1977) Recruitment behavior, orientation, and the organization
of foraging in the carpenter ant *Camponotus pennsylvanicus* degeer
(Hymenoptera: Formicidae). Beh Ecol Soc 2:61–79. DOI:
10.1007/BF00299289.

32. Franks NR, Mallon EB, Bray HE, Hamilton MJ, Mischler TC (2003)
Strategies for choosing between alternatives with different attributes:
exemplified by house-hunting ants. Anim Behav 65:215–223. DOI:
10.1006/anbe.2002.2032.

33. Symonowicz B, Kieruzel M, Szczuka A, Korczyńska J, Wnuk A,
Mazurkiewicz PJ, Chiliński M, Godzińska EJ (2005) Behavioral reversion and
dark–light choice behavior in workers of the red wood ant *Formica polyctena*.
J Insect Behav 28:245–256. DOI: 10.1007/s10905-015-9496-2.

34. Greene MJ, Gordon DM (2007) Interaction rate informs harvester ant task
decisions. Behav Ecol 18:451–455. DOI: 10.1093/beheco/arl105.

35. Pinter-Wollman N, Bala A, Merrell A, Queirolo J, Stumpe MC, Stumpe,
Holmes S, Gordon DM (2013) Harvester ants use interactions to regulate
forager activation and availability. Anim Behav 86:197–207. DOI:
10.1016/j.anbehav.2013.05.012.

36. Khuong A, Gautrais J, Perna A, Sbaï C, Combe M, Kuntz P, Jost C, Theraulaz
G (2016) Stigmergic construction and topochemical information shape ant nest
architecture. Proc Natl Acad Sci USA 113:1303–1308. DOI:
10.1073/pnas.1509829113

37. Dostálková I, Špinka M (2007) Synchronization of behaviour in pairs: the role
of communication and consequences in timing. Animal Behav 74:1735e1742.
DOI:10.1016/j.anbehav.2007.04.014

38. Kierat J, Filipiak M, Szentgyörgyi H, Woyciechowski M (2017) Predation
Cues in Solitary bee Nests. Journal of Insect Behavior 30:385–393. DOI:
10.1007/s10905-017-9626-0

39. Lewis SE (1994) Night Roosting Ecology of Pallid Bats (*Antrozous pallidus*)
in Oregon. The American Midland Naturalist 132: 219–226

40. Caccamise DF, Morrison DW (1986) Avian Communal Roosting:
Implications of Diurnal Activity Centers. The American Naturalist 128:191–
198. DOI: 10.1086/284553

41. Simpson SJ, Sword GA, Lorch PD, Couzin ID (2006) Cannibal crickets on a
forced march for protein and salt. Proc Natl Acad Sci USA 103:4152–4156.
DOI: 10.1073/pnas.0508915103

42. Oberhauser FB, Koch A, Czaczkes TJ (2018) Small differences in learning
speed for different food qualities can drive efficient collective foraging in ant
colonies. bioRxiv preprint. DOI: 10.1101/274209

43. Möglich M, Hölldobler B (1974) Social carrying behavior and division of
labor during nest moving in ants. Psyche 81:219–236. DOI:
10.1155/1974/25763

44. Robson S, Traniello J. Key individuals and the organisation of labor in ants, in
Information Processing in Social Insects (Birkhäuser ed. 1999) pp. 239–259

45. Hammel B, Vollet-Neto A, Menezes C, Nascimento FS, Engels W, Grüter C
(2016) Soldiers in a Stingless Bee. The American Naturalist 187:120–129.
DOI: 10.1086/684192

46. Dornhaus A (2008) Specialization Does Not Predict Individual Efficiency in
an Ant. PLoS Biology 6:e285. DOI: 10.1371/journal.pbio.0060285

47. Robson SKA, Traniello JFA (2002) Transient division of labor and behavioral
specialization in the ant *Formica schaufussi*. Naturwissenschaften 89:128–131

48. Pruitt JN, Keiser CN (2014) The personality types of key catalytic individuals
shape colonies' collective behaviour and success. Animal Behav 93:87–
95.DOI: 10.1016/j.anbehav.2014.04.017

49. Robinson GE (1992) Regulation of Division of Labor in Insect Societies.
Annual Review of Entomology 37:637–665. DOI:
10.1146/annurev.en.37.010192.003225

50. Gelblum A, Pinkoviezky I, Fonio E, Ghosh A, Gov N, Feinerman O (2015)
Ant groups optimally amplify the effect of transiently informed individuals.
Nature Communications 6: 7729. DOI: 10.1038/ncomms8729

51. Reebs SG (2000) Can a minority of informed leaders determine the foraging
movements of a fish shoal? Anim. Behav 59:403–409.
DOI:10.1006/anbe.1999.1314.

52. Dyer JRG, Johansson A, Helbing D, Couzin ID, Krause J (2009) Leadership,
consensus decision making and collective behaviour in humans. Phil Trans R
Soc B 364:781–789. DOI: 10.1098/rstb.2008.0233.

53. Torney CJ, Lorenzi T, Couzin ID, Levin SA (2015) Social information use
and the evolution of unresponsiveness in collective systems. J R Soc Interface
12:20140893. DOI: 10.1098/rsif.2014.0893.

54. Vonshak M, Shlagman A (2009) A *Camponotus fellah* queen sets a record for
Israeli ant longevity. Isr J Entomology 39: 165–168

55. Mersch DP, La Mendola C, Keller L (2017) *Camponotus fellah* queens are
singly mated. Insect Soc 64:269–276. DOI: 10.1007/s00040-017-0543-1

56. R Core Team (2017) R: A language and environment for statistical computing.
R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-
project.org/.

Supplementary material

Supplementary Figure 1: Tracking setup (A) Lateral view (B) Top view; reproduced with

permission from Mersch *et al.* (2013)[27]

Supplementary Figure 2. Individual workers transport brood rapidly. Each black dot

shows the average transport time needed by a single transporter.

Supplementary Figure 3. No change in interaction frequencies after light-off. Grey

dots show data in the hour preceding light-off. Yellow dots show data between light-off

and the first transport. The black line shows the average relationship between the number

of ants in the nest and the number of interactions before light-off, and the red line shows

the same relationship in the interval between light-off and the first brood transport.

Supplementary Figure 4. Brood accumulation in the tunnel does not speed up

transporter recruitment. A. Each black dot shows the recruitment delay. For all but the

first transporter, recruitment delays are with regard to the transport start of the previous

transporter. For the first transporter, recruitment delays are with regard to light-off. Red

circles indicate the median recruitment delay for each transporter rank. B. The recruitment

delays are the same as in A. Blues dots show data for night 4, magenta dots data for night

5, and cyan dots data for night 6. Data are shown separately for each colony.

Supplementary Figure 5. Workers transport almost exclusively from the nest to the

tunnel. Grey dots show transports from the nest to the tunnel. Red dots show transports

from the tunnel to the nest.

Interval length for speed estimate (s)	#interactions excluded due to missing speed data(%)	Speed increase after an interaction with a privately informed ant (mm/s);mean±std	Speed increase after an interaction with a non privately informed ant(mm/s); mean±std	Kruskal-Wallis Chi2	p-value
5	79 (12.9)	0.89±2.3	0.26±3.1	3.25	0.071
10	50 (10.2)	0.25±2.8	0.15±2.9	2.77	0.096
20	20 (6.3)	0.23±1.8	0.07±2.3	3.09	0.079

Supplementary Table 1. Speed change after an interaction with a privately informed

ant.

Supplementary Video 1. Worker transporting brood. Worker 62 transports brood to the

tunnel. At 16s in the video, ant 62 takes brood directly from another worker without this

worker changing its behaviour. Data is from colony 2 and the frame rate is accelerated 5

times. The green line shows the worker's trajectory in the previous minute.

Supplementary Video 2. Targeted queen recruitment to the tunnel. Worker 632 (in

pink) approaches the queen, pulls on her mandibles, and then returns to the tunnel with

the queen (in blue) following her. The data are from colony 1.

Supplementary Video 3. Recruitment of two non-transporters to the tunnel. Worker

458 (in green) interacts with workers 607 (in blue) and 278 (in cream), and both then follow

worker 458 to the tunnel. The trajectories are shown for all three workers after the

interactions finished. The data are from colony 1.