Resource limitation modulates the fate of dissimilated nitrogen in a dual-pathway

Actinobacterium

David C. Vuono[a], Robert W. Read[a], James Hemp[b], Benjamin W. Sullivan[c], John A. Arnone III[a],

Iva Neveux[a], Bob Blank[d], Carl Staub[e], Evan Loney[a], David Miceli[a], Mari Winkler[f], Romy

Chakraborty[g], David A. Stahl[f], Joseph J. Grzymski[a,*]

[a]Division of Earth and Ecosystem Sciences, Desert Research Institute, Reno, NV 89512, USA

[b]Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena,

CA, USA

[c]Department of Natural Resources and Environmental Science, University of Nevada, Reno, NV

89557, USA

[d]Agricultural Research Service, U.S. Department of Agriculture, Reno, NV, 89512, USA

[e]Agtron, Inc. 9395 Double R Blvd, Reno, NV 89521, USA

[f]Department of Civil and Environmental Engineering, University of Washington, Seattle, WA,

98195, USA

[g]Lawrence Berkeley National Laboratory, Berkeley, CA, 94729, USA

[*]Correspondence: Joseph J. Grzymski, Division of Earth and Ecosystem Sciences, Desert

Research Institute, Reno, 2215 Raggio Parkway, NV 89512, USA

Email: Joe.Grzymski@dri.edu

Abstract

Respiratory ammonification and denitrification are two evolutionarily unrelated dissimilatory

nitrogen (N) processes central to the global N cycle, the activity of which is thought to be

controlled by carbon (C) to nitrate (NO_3^-) ratio. Here, we find that *Intrasporangium calvum* C5, a

novel menaquinone-based dual-pathway denitrifier/respiratory ammonifier, disproportionately

utilizes ammonification rather than denitrification when grown under carbon or nitrate limitation,

not $C:NO_3^-$ ratio. Higher growth rates are promoted by ammonification and metabolite and

transcriptional profiles during growth show that the bacterium produces its own formate from a

fermentable carbon source (lactate) to further generate a proton motive force for the

ammonification pathway. Transcript abundances encoding for nitrite reducing enzymes, NrfAH

and NirK, also significantly increase in response to nitrite production. Mechanistically, our

results suggest that pathway selection is driven by intracellular redox potential (redox poise),

which is lowered during resource limitation, thereby decreasing catalytic activity of upstream

electron transport steps needed for denitrification enzymes. Our work advances our

understanding of the biogeochemical tapestry of N-cycle dynamics and pathway evolution.

Globally, respiratory ammonification and denitrification are vital nitrogen (N) dissimilation

pathways that either retain reactive N to support net primary productivity or close the N-cycle

through the release of gaseous N, respectively[1]. The environmental controls on these two

pathways, particularly the ratio of electron-donor to electron-acceptor (e.g., $C:NO_3^-$)[2], have

gained attention[3–7] due to increased anthropogenic N inputs into the environment[8]. However, the

effects of resource limitation on growth and pathway selection (i.e., allocation of C and N to

dissimilatory and assimilatory processes), which are often confounded by $C:NO_3^-$ ratio, have not

been tested. Strong selective pressures from Earth's shifting biogeochemistry and oxidation-state

have driven evolutionary adaptions to microbial electron transport chains (ETC)[9,10], respiratory

chain redox potentials[11–13], and protein atomic composition[14,15], may shed light on how these

pathways are regulated in contemporary organisms. Here, by identifying the biochemical and

evolutionary differences between respiratory ammonification and denitrification, we disentangle

the functional significance and molecular mechanisms of electron transfer through either

pathway in a dual pathway organism.

From a biochemical standpoint, the primary difference between respiratory ammonification and

denitrification is their respective source of reducing equivalents in the ETC: 1) heme-based

cytochrome c nitrite reductase used in respiratory ammonification receive electrons directly from

the quinone (Q) pool[16] while 2) copper and cd_1 nitrite reductases used in denitrification receive

electrons from a soluble electron carrier (e.g., cytochrome c) via the bc_1 complex[17]. From an

evolutionary standpoint, we can place each N-module's origin to a putative time in Earth history

based on the metal co-factors that would have been bioavailable: heme-based cytochromes in an

ancient, more reduced, environment compared to the copper-containing nitrite reductases in an

oxidizing environment[18]. The bioenergetic chains of microorganisms also underwent selective

pressure to shift from low-potential (LP) to high-potential (HP) quinones in response to Earth's

oxygenation[11,12]. Menaquinone (MK) is thought to be the ancestral type of LP quinone[19].

Organisms that use ubiquinone (UQ) are thought to have evolved under high O_2 tensions with α-,

β-, γ-proteobacteria as the only bacterial clades to use UQ[12]. Surprisingly, our understanding for

the biochemistry of denitrification is based predominantly on HP UQ-based systems[20], leaving a

significant knowledge gap in the physiology and biochemistry of LP MK-based denitrifiers and

how they link electron transfer with energy capture under resource limitation[21–23].

As a predicted environmental control for respiratory ammonification versus denitrification,

$C:NO_3^-$ ratio, the predicted end-products should remain identical at different substrate

concentrations of the same ratio. In order to resolve the mechanisms of $C:NO_3^-$ control on

pathway selection and better understand branched respiratory chains in LP-based nitrate-

reducing organisms, we undertook the characterization of the novel Gram-positive

Actinobacterium strain *Intrasporangium calvum* C5: a dual-pathway nitrite reducer that uses MK

as sole pool quinone. Here we show that over a range of $C:NO_3^-$ ratios, duplicated at two

substrate concentrations, *I. calvum* disproportionately utilizes its ammonia-forming pathway

during C limitation (≤ 0.4mM lactate), when $C:NO_3^-$ ratios are < 1 (an observation contrary to the

current paradigm). Using a genome-guided approach coupled to time-series transcriptomics and

metabolite profiles, we identified differentially expressed genes in the bacterium's ETC and

central metabolic pathways. Using this information to inform a metabolic reconstruction of the

ETC we infer mechanisms by which these two pathways are regulated at the biochemical level.

Results

Genomic analysis of *I. calvum* C5. We sequenced and analyzed the genome of *I. calvum* C5 to

first compare its similarity to the type species *I. calvum* 7KIP. We identified a high degree of

sequence similarity to 7KIP based on three homologous sequence regions as locally collinear

blocks (SI Results). Genome size of C5 was 4,025,044 base pairs (bp), only 662 bp longer than

7KIP. Genomic analysis of the ETC revealed the typical suite of complexes common to

facultative aerobes, including primary dehydrogenases (*nuo* complex, succinate dehydrogenase),

alternative *NDH-2* NADH dehydrogenase, cytochrome bc_1 complex, high-oxygen adapted

cytochrome c oxidase (A-family), and low-oxygen adapted cytochrome *bd* oxidase. The bc_1

complex subunits are also located immediately upstream of cytochrome c oxidase, suggesting

that these enzymes are encoded in a single operon creating a supercomplex. Despite *I. calvum*'s

seeming propensity for aerobic growth on a number of growth media[24], its bioenergetic system

uses MK as its sole pool quinone. *I. calvum* also possesses multiple pathways for supplying

electrons into the MK-pool, such as formate, malate, hydroxybutyrate, and glycerophosphate

dehydrogenases. Once in the MK-pool, there are alternative pathways for MKH_2 oxidation that

can circumvent the bc_1 complex, such as a membrane-bound respiratory nitrate reductase module

(NarG). In addition, to NarG, its dissimilatory N module composition consists of a truncated

denitrification pathway (N_2O is a terminal product) using a copper nitrite reductase NirK and

quinol-dependent nitric oxide reductase qNor. *I. calvum* also possesses both catalytic and

membrane anchor subunits (NrfA and NrfH, respectively) for a pentaheme cytochrome c module

involved in respiratory nitrite ammonification.

***I. calvum* encodes for a functional NrfAH complex and assimilates NH_4^+ via respiratory**

nitrite ammonification. To gain insight into possible function of the NrfAH complex, we

aligned the NrfA protein sequences from C5 and 7KIP to a collection of 33 recognized

cytochrome c nitrite reductases from published annotated genomes (Table *S1*). This confirmed

that NrfA from *I. calvum* is a member of the CxxCH 1st heme motif group (Figure *1*A), which

forms one of four clades on the NrfA phylogenetic tree. We then queried the genomes of the taxa

in our phylogeny for other annotated N-reducing modules used in nitrate reduction, nitrite

reduction, NO-forming nitrite reduction, and primary pool quinone. Among the three major

clades of NrfA, at least 5 additional taxa are noted having dissimilatory N-module inventories

containing dual respiratory pathways: *S. thermophilum, B. azotoformans, B. bataviensis*, *B.*

bacteriovorus, and *Candidatus* N. inopinata, (Figure *1*A). None of the taxa in

Figure 1. (A) Maximum likelihood phylogenetic tree of NrfA amino acid sequences from known
respiratory ammonifiers and accompanying N-module composition for each organism. Pool
quinone is also noted for dual-pathway nitrite reducers and model species. Colors of the main
branches denote the 1st heme motif type: CxxCK and CxxCH. (B) Protein atomic composition
for N and C normalized to protein length for NirK and NrfA nitrite reductases. (C) State-
transition from ammonium-replete to ammonium-deplete for I. calvum C5 grown under 8mM
lactate 12mM nitrate minimal media at 30 ºC. Metabolite profiles for ammonium-deplete are
shown.

our NrfA phylogeny harbored the cd_1 nitrite reductases (NirS). Due to the exclusive NirK

representation in dual-pathway membership, we asked whether there might be differences in

protein atomic composition between NirK and NrfA, given the disparate evolutionary origins of

these modules[25]. We collected 20 additional publicly available NirK protein sequences from

nondual-pathway denitrifiers (Table *S1*) and calculated the protein C and N composition for our

NirK/NrfA collection as atoms per residue side-chain (Figure *1*B). These results showed a

significant depletion in C and N per residue side-chain for NirK compared to NrfA (C and N:

$p<0.001$; t-test), indicating that resource constraints are imprinted on the evolution of these

proteins.

We next tested the functionality of *I. calvum*'s Nrf complex by growing the bacterium under

reducing conditions (8 mM lactate, 12 mM nitrate, ammonium-replete). We then performed a

state-transition where biomass from late-exponential growth phase was collected and

anaerobically inoculated into ammonia-deplete media (Figure 1C; SI Results). Despite no

detectable amounts of ammonium produced in the media over time, cell counts increased

$5.4x10^5 \pm 8.9x10^4$ cells/mL (0.126 ± 0.02 optical absorbance units, OD_{600}) over a 48-hour

incubation, indicating consumption of ammonium produced by NrfA. Net ammonium production

was 13 ± 2.7 µmoles with the remainder of dissimilated N being used by the denitrification

pathway (24 ± 4.2 µmoles N_2O-N), resulting in a recovery of 97.4% dissimilated N. These results

confirmed that *I. calvum* C5 has a functional Nrf complex and also consumes the product

(ammonium) of respiratory nitrite ammonification.

Respiratory nitrite ammonification exceeds denitrification under C-limitation. We

investigated $C:NO_3^-$ control on respiratory ammonification versus denitrification on cultures of *I.*

calvum C5 over a high resource $C:NO_3^-$ range (16-0.4 mM lactate, 12 mM nitrate; ratio 4-0.1)

and low resource $C:NO_3^-$ range (1.6-0.04 mM lactate, 1.2 mM nitrate; ratio 4-0.1). This

experimental design enabled us to evaluate $C:NO_3^-$ control over a broader range than previous

studies that only considered ratios ≥ 1.5[3,4,26], while also testing the effects of resource

concentration on pathway selection. Under all the treatments tested, gas and ion chromatography

measurements showed products of both respiratory pathways, differing only in the relative

fraction of N_2O versus ammonium production across treatments (Figure 2). At high resource

concentrations, respiratory ammonification did not prevail at high $C:NO_3^-$ ratios (Figure 2A,

Figure 2B, left panels; Table $S2$). Instead, significantly greater amounts of N_2O were produced

over ammonium, though nitrite was still the major extracellular end-product of nitrate

respiration. Despite the predominance of N_2O production under the high resource concentrations,

ammonium production exceeded N_2O production only at the lowest $C:NO_3^-$ ratio (0.4 mM

lactate, ratio=0.1) (Figure 2) and accounted for $76.2 \pm 0.1\%$ of dissimilated N.

Results from the low resource dataset provided weak support for the strict stoichiometry

hypothesis that $C:NO_3^-$ controls pathway selection. Ammonia exceeded N_2O production only

under one high $C:NO_3^-$ ratio treatment (ratio=4; 1.6 mM lactate; Figure 2A, Figure 2B, right

panels). However, at ratios ≤ 1 (≤ 0.4 mM lactate), significantly more ammonium than N_2O was

produced. On average, respiratory ammonification accounted for $78.1 \pm 8.9\%$ of dissimilated N

for lactate concentrations ≤ 0.4 mM. When these results are taken in context with cell physiology,

we observed a significant and positive relationship between specific growth rate (μ) and the

Figure 2. The effects of high resource (left; range of lactate concentrations with 12 mM NO_3^-) and low resource (right; range of lactate concentrations with 1.2 mM NO_3^-) concentrations with the same C:NO_3^- ratio on pathway selection in I. calvum C5. (A) Production of N_2O-N and net change of NH_4^+ over a 100-hour incubation period at 30 ºC. Each bar represents the average of 8-10 replicates per treatment (Table S5). (B) Fraction of dissimilated N by pathway. (C) Growth rates for each corresponding treatment. The x-axis label defines lactate concentration and C:NO_3^- ratio in parentheses.

fraction of N dissimilated by respiratory ammonification (R^2=0.5; p<0.001) (Figure *2*C, S1; Table S3).

Resource concentration influences the metabolite profiles of ammonium and N_2O production. Given the co-occurrence of end products from both pathways during the end-point experiments (Figure *2*), we next investigated the timing of ammonium and N_2O production relative to metabolite profiles for lactate, nitrate/nitrite, and growth phase at two resource concentrations with the same ratio (8 mM and 0.8 mM lactate, ratio=2;Figure 3A, Figure S2, Figure *S3*). Despite ample e-donor and e-acceptor available for growth, the high resource cultures entered a quasi-stationary phase at ~50 hours, after which there was continued slow growth (Figure 3A). Metabolite profiles showed that ammonium and N_2O production began simultaneously, as soon as nitrite was produced from nitrate reduction. The low resource cultures entered stationary phase at ~40 hours (Figure S2) after nitrate had been fully utilized. No further cell growth was observed after stationary phase was reached. These results show that cell growth occurred primarily on the reduction of nitrate, while nitrite reduction to ammonium and N_2O

Figure 3. (A) Time-series metabolite profiles for lactate, nitrate, and nitrite (top pane), production of dissimilated end-products as N_2O-N and net change in NH_4^+ ammonium production (middle pane), and corresponding growth curve of *I. calvum* cells grown under 8mM lactate 12mM nitrate (C:NO_3^- ratio = 2) (bottom pane). Sampling points during growth phases are marked for transcriptomic analysis. (B) Metabolic reconstruction of the ETC from *I. calvum* with transcriptional changes for genes participating in dual-pathway dissimilatory nitrite reduction. Log_2 fold changes in transcript abundance are shown for late exponential relative to early exponential growth phase (EE vs. LE), stationary phase relative to early exponential growth phase (EE vs. ST), and stationary phase relative to late exponential growth phase (LE vs. ST). Locus IDs for each gene product correspond to heat map subplots in the order shown (left-to-right for each growth phase and top-to-bottom for each locus ID specified). Higher transcript abundance is represented in red, lower transcript abundance in blue, and no change in transcript abundance in white. Significant changes in transcript abundance ($p < 0.01$) are marked as a red box. Value of log_2 fold change is specified within each subplot. The log_2 fold changes of 14 NADH dehydrogenase subunits (Intca_03465-03530) were averaged as transcriptional changes were all shifted in the same direction.

occurred during a stationary growth phase, demonstrating that microbial activity is not always correlated with growth. The metabolite profiles for ammonium and N_2O at low resources (Figure S2) did not mirror those observed at high resources (Figure 3A). The rate of N_2O production significantly decreased and ammonium production oscillated rather than steadily increase through time. These differences in metabolite profiles, further demonstrate that concentration influences the activities of pathway bifurcation. Repeated time series experiments that were extended up to 300 hours show that nitrite is slowly depleted, but does not get fully consumed

(Figure *S3*). When cultures were given nitrite, instead of nitrate as a terminal electron acceptor (8

mM lactate, 12 mM nitrite; ratio=2), we observed no immediate growth (as was observed with

nitrate) but measured more N_2O than ammonium production (33.4±4.8 µmoles N_2O-N and

8.0±2.5 µmoles NH_4^+, respectively) (Figure S4), demonstrating respiratory ammonification does

not exceed denitrification when nitrite is supplied as the sole acceptor in *I. calvum*.

Nitrite-reducing modules are up-regulated during late exponential- and stationary-phase

growth. In order to gain insight into mechanisms of gene regulation and transcriptional

organization of *I. calvum*, we conducted RNA-Seq in parallel with the high resource time-series

metabolite profile (Figure 3A). This approach enabled us to compare genome-wide differential

expression based on $\log_2$ fold change (lfc) of RNA extracted from three growth phases: early

exponential (EE), late exponential (LE), and stationary (ST) (Figure 3B, Figure *S5*). Within the

central metabolic pathway beginning with the conversion of lactate to pyruvate, we observed a

moderate decrease in transcript abundance of L-lactate dehydrogenase (LDH) (Intca_16740)

between EE-LE and -ST (lfc = -1.6±0.7; -1.9±0.7), respectively. Lactate utilization protein C

(LUP) (Intca_04080), an enzyme involved in lactate degradation, also showed a moderate and

significant decrease in transcript abundance between EE-LE and -ST (lfc = -1.6±0.6; -2.4±0.6,

p=0.002), respectively. *I. calvum* encodes for two parallel metabolic pathways for pyruvate

conversion to acetyl-CoA: pyruvate dehydrogenase (PDH) (Intca_01255) and pyruvate

ferredoxin oxidoreductase (PFOR) (Intca_15510). For PDH, there was a significant and

moderate increase in transcript abundance between EE-LE and -ST (lfc = 2.1±0.6, *p*=0.002;

1.5±0.6), respectively. For PFOR, there was a minor decrease in transcript abundance between

EE-LE (lfc = -0.43±0.5), and then a moderate increase in transcript abundance between EE-ST

(1.1±0.5). Citrate synthase (Intca_04135), the enzyme catalyzing the conversion of acetyl-CoA

to citrate and the first step of the tricarboxylic acid (TCA) cycle, showed a highly significant

increase in transcript abundance between EE-LE and -ST (lfc = 4.3±0.5, *p*<0.001; 6.9±0.5,

p<0.001).

Within the ETC, there was moderate and significant decrease in transcript abundance for all

subunits from the primary dehydrogenase (*nuo* complex; Intca_03465-03539) between EE-LE

and -ST (lfc = -1.2±0.3; -2.4±0.6, *p*<0.001), respectively. Nitrate reductase subunits showed no

change in transcript abundance between EE-LE (lfc = 0.01±0.07) and moderately decreased in

abundance by ST (lfc = -1.2±0.1), which was corroborated by the depletion of nitrate during

stationary phase. There was a significant increase in transcript abundance of *nirK* (Intca_17170)

(lfc = 2.2±0.6, *p*=0.003; 2.4±0.6, *p*<0.001) and quinol dehydrogenase/membrane anchor subunit

nrfH (Intca_09465) (lfc = 2.5±0.6, *p*=0.001; 2.1±0.6, *p*=0.003) by EE-LT and EE-ST,

respectively, which coincided with nitrite production (Figure 3A). The catalytic subunit of the

cytochrome c nitrite reductase complex (*nrfA*) (Intca_09460) also increased moderately in

transcript abundance by EE-LT and EE-ST (lfc = 1.6±0.6; 1.0±0.6), respectively (Figure 3B).

Contrary to the transcript abundance patterns of *nirK* and *nrfAH*, nitric oxide reductase (qNor;

Intca_01525) transcripts moderately increased between EE-LT (lfc = 1.6±0.6) but decreased in

the successive time periods (lfc = 0.43±0.6 between EE-ST; lfc = -1.2±0.6 between LE-ST)

(Figure 3B).

There was a significant increase in transcript abundance of formate transporter *focA*

(Intca_17150) between EE-ST, as well as LE-ST (lfc = 4.9±0.7, *p*=0.002; 4.8±0.7, *p*=0.002;

respectively). We verified the production of formate in our ion chromatography measurements in

the range of 100-200μM following late exponential growth. We also observed a moderate

increase in transcript abundance of formate dehydrogenase (FDH) subunits (Intca_11150-

11160). These results implicate the activity of formate oxidation, which would contribute to a Δp

in the periplasm via a Q-loop mechanism and the reduction of MK for electron transfer to nitrite

via cytochrome c nitrite reductase. Considering that formate was not provided in our media

recipe, an alternative pathway for formate production must exist in *I. calvum*. We also observed

acetate production in similar concentrations as formate (100-200μM). In *E. coli*, formate is

produced anaerobically from the action of pyruvate formate lyase (PFL). We identified a putative

PFL based on genome annotation (Intca_12230), where transcript abundance also significantly

increased by ST. PFL is also highly sensitive to oxygen[27], which was also in agreement with a

significant increase in transcript abundance between EE-ST and LE-ST (Figure 3B) of

cytochrome *bd* oxidase (Intca_01110 and Intca_01115), which is thought to protect anaerobic

enzymes against oxidative stress[28].

Discussion

We challenge the paradigm that $C:NO_3^-$ ratio controls pathway selection in a dual-pathway

organism based on a simple principle: ratios do not account for the abundance of growth-limiting

resources. We hypothesized that limitation in C or NO_3^- should better predict pathway selection

in a dual-pathway denitrifier/respiratory ammonifier. To test this hypothesis, we systematically

measured the response of the Gram-positive Actinobacterium *Intrasporangium calvum* C5 to the

same range of $C:NO_3^-$ ratios at both high and low resource loadings to better resolve mechanisms

of pathway selection. We demonstrated that resource concentration, not $C:NO_3^-$ ratio, influences

pathway selection. We found stronger support for respiratory ammonification preference under

C-limitation (at low $C:NO_3^-$ ratios), which also grew at significantly higher growth rates (Figure

2). These results suggest that the NrfA complex, which receives electrons directly from the MK-

pool, is optimized to maximize power when one or more resources are limiting. These data,

together with metabolic reconstructions from metabolite and transcriptional profiles (Figure 3),

suggest that $C:NO_3^-$ ratio alone is insufficient to explain pathway selection.

The theoretical basis for pathway selection is explained by the law of the minimum (LM) and the

maximum power principle (MPP), which state that growth is limited by the least abundant

resource and that biological systems are designed to maximize power in order to effectively

allocate energy to reproduction and survival[29,30], respectively. Here, it appears these two natural

theories are working together: when resources are limited, the cell utilizes the respiratory

pathway for growth that is optimized to maximize power. Power, in this case, is realized as

higher growth rates from the cultures exhibiting disproportionately higher ammonium production

than N_2O production (Figure *2*: high resources: $C:NO_3^-$ ratio = 0.1; low resources: $C:NO_3^-$ ratios

= 4, 1, 0.5, 0.1). More specifically, the bacterium must generate a greater Δp in order to

maximize power when starved for a growth limiting resource. This may help to further explain

how respiratory ammonification, which is overall energetically less favorable than denitrification

(lactate with nitrite: ΔG^o = -763.98 versus ΔG^o = -1196.93, respectively), can have higher growth

yields[31] and growth rates (Figure *2*, Figure S1) under C- and N-limitation due to the higher

energy yield on a per-nitrite basis (denitrification: -217 KJ per mole nitrite; respiratory

ammonification: -399 KJ per mole nitrite). For comparison, a total of 8 H^+ are translocated

during denitrification by *I. calvum* (not including nitrate reduction since both pathways share this

step) (Figure 3): NADH dehydrogenase translocates 4 H^+ per MKH_2 oxidized and the bc_1

complex translocates an additional 4 H^+ per MKH_2 oxidized. However, 2 H^+ must be consumed

in the periplasm to reduce nitrite to $NO^{[32]}$. qNor has a net zero H^+ release (consumes 2 H^+ to

make N_2O but releases 2 H^+) without MKH_2 regeneration[33]. Thus, a net total of 6 H^+ are

translocated per nitrite reduced in denitrification with added biosynthetic costs of making the bc_1

complex and qNor. In respiratory ammonification, MK/MKH$_2$ redox pair is cycled between

NADH dehydrogenase and formate dehydrogenase. 6 electrons and 8 H$^+$ are needed to reduce

nitrite to ammonium, thus 3 MKH$_2$ are needed[16]. If MKH$_2$ is received from NADH

dehydrogenase, 12 H$^+$ are translocated plus 2 H$^+$ from FDH. As each MKH$_2$ is oxidized at the

binding site of NrfH, 2 H$^+$ are liberated[16], resulting in a net total of 12 H$^+$ translocated per nitrite

reduced for respiratory ammonification. This implies that the cell might deplete its NADH pool

more rapidly on a per nitrite basis. However, if more protons are pumped in the early stages of

growth, the cell would be allocating the ATP generated for anabolism, as evidenced by higher

growth rates in the cultures exhibiting higher amounts of respiratory ammonification (Figure *2*),

which is supported by the MPP.

Under our high resource conditions (Figure *2*; left panels), at C:NO$_3^-$ ratios $\geq$ 1, we observed that

denitrification prevailed and these cultures had lower growth rates than the predominantly

ammonium producing cultures. These high resource circumstances resulted in the production of

toxic intermediates (i.e., NO$_2^-$ and possibly NO, albeit at undetectable levels), which may explain

why these cultures had lower growth rates (Figure *2*; left panels) and quasi-steady state growth

curves in our high resource metabolite profile (Figure 3A). Rowley and colleagues[34] reported

that at least 20% of the N$_2$O released during high C conditions were produced by competition

between nitrite and nitrate in the active-site of NarG. Under excess C concentrations, NarG

produces intracellular NO from NO$_2^-$ and these intermediates are likely inhibitory to cell growth,

which may explain why our growth curves (Figure 3A) reached a quasi-steady state before

nitrate had been fully utilized (as compared to the low resource metabolite profile, Figure S2).

Furthermore, resources were not limiting growth under these conditions. Rather, the cells were

likely experiencing toxicity from NO and NO$_2^-$ and thus the metabolic outcomes would be

beyond the scope of the LM and MPP. Nonetheless, these results clearly demonstrate that end-

product formation from the two resource concentrations tested, with the same C:NO$_3^-$ ratios, are

not identical thereby refuting the C:NO$_3^-$ control hypothesis.

We selected a single treatment (8 mM lactate, 12 mM nitrate; C:NO$_3^-$ ratio = 2), in which we

observed both denitrification and respiratory ammonification occurring simultaneously, for

RNA-Seq in order to gain insight into the transcriptional organization of actively growing *I.*

calvum cells (Figure 3). Strangely, we saw a decrease in transcript abundance encoding for two

enzymes known to convert lactate to pyruvate, LDH and LUP. While normalized read counts

(Figure *S5*) were generally consistent across growth phases, indicative of constitutive expression,

further research investigating the mode of anaerobic lactate oxidation in *I. calvum* would

illuminate how reducing equivalents are fed into its central metabolic pathway. For example, *S.*

loihica PV-4 is known to use lactate for both denitrification and respiratory ammonification, but

only uses acetate for denitrification[35]. Nonetheless, our transcriptomic data suggests that

pyruvate plays a central role in providing reducing equivalents to the TCA cycle as Acetyl-CoA,

as evidenced by significant upregulation in the genes encoding for pyruvate dehydrogenase and

citrate synthase, as well as apparent "leaking" via incomplete lactate oxidation through the

release of acetate and formate. Such leaking may be produced by a putative PFL, adding to the

diversity of C utilization pathways feeding the ETC, and thereby driving pathway selection for

nitrite reduction. Our transcriptomic results, coupled with a parallel metabolite profile (Figure 3),

also suggest that the dual-pathway is induced by the presence of nitrite, and is not constitutively

expressed like nitrate reductase, *narG*. Furthermore, it appears that the significant increase in

transcript abundance for the gene encoding the *bd* oxidase helps to protect the anaerobic-

dependent biochemical machinery against oxidative stress, thereby scavenging any residual

oxygen during anaerobic growth.

Our metabolite profiles for N oxyanion respiration and N_2O versus ammonium production show

conflicting patterns relative to previous studies (Figure 3A, Figure S2). Yoon and colleagues[36]

reported complete reduction of nitrate, production of nitrite, and then rapid consumption of

nitrite, with N_2O as the main end-product, by *S. loihica* PV-4 (5 mM lactate, 1 mM nitrate;

ratio=0.6). When Yoon and colleagues[36] replaced nitrate with nitrite as the dominant electron

acceptor (5 mM lactate, 1 mM nitrite, ratio=0.6), ammonification prevailed. Other research has

shown the same response to nitrite replacement and ammonification dominance using non-

fermentable C-sources (i.e., acetate) in chemostat enrichments of *Geobacter lovleyi*[37]. In our

work, nitrite was never fully depleted (Figure 3A, Figure S2, Figure *S3*) and when nitrite was

given as the only electron acceptor, the bacterium predominantly used denitrification but without

concurrent growth (Figure S4). Similar to our work, Kraft and colleagues[26] also reported

denitrification dominance when nitrite was supplied as the terminal acceptor. These differences

highlight an incomplete understanding for the molecular mechanisms underlying the framework

put forth by the LM and MPP.

A detailed look into the biochemistry of ETC complexes helps to shed light on the molecular

mechanisms modulating pathway bifurcation. For example, Yoon and colleagues[3] demonstrated

that elevated pH selects for ammonification in *S. loihica* PV-4. This phenotypic response is due

to a decrease in the midpoint potential of the Rieske protein at higher pH[38–40]. Thus, any

hindrance of electron flow through the bc$_1$ complex would ultimately reduce the activity of

downstream processes and promote alternative respiratory pathways. Nitrogen and C limitation

have also been shown to influence flux distributions in redox sensitive proteins, including those

found in electron transport[41]. A drop in the intracellular redox potential (redox poise) of the cell

due to resource limitation may decrease the midpoint potential of the Rieske protein and reduce

the activity of any downstream electron exit modules, such as NirK[42–44]. Thus, based on

fundamental principles of protein redox chemistry and thermodynamics, it becomes clear that

denitrification versus ammonification are likely not modulated by an arbitrary ratio of C:NO$_3^-$,

but rather by thermodynamic constraints of the Q-cycle[11,12]. The phenotypic response of higher

rates of denitrification over ammonification at high C:NO$_3^-$ ratios in other published studies[3,4]

may also be due to enrichment bias for organisms that utilize quinones with higher midpoint

potentials in their bioenergetic chains (Figure *1*). Bergdoll and colleagues[11] suggested that

comparisons of Rieske/cyt*b* complexes from organisms with high- and low-potential quinones

may help to reconcile the thermodynamic properties of Q-cycle function. However, most of our

understanding of denitrification bioenergetics is based on evolutionarily recent UQ-based HP

bioenergetic chains from Gram-negative *α*-, *β*-, *γ*-proteobacteria. Because *I. calvum* uses a MK-

based LP bioenergetic chain it may be possible that the differences in pathway selection across

treatments are unique to LP chains.

Piecing together the evolutionary history of the N-cycle using isotopic signatures for

geochemically available N module cofactors (i.e., Ni, Fe, and Mo) coupled to molecular

evolutionary analysis has revealed respiratory ammonification was likely a major component of

the Archean N-cycle[25]. Abiotic nitrite formation and depletion of ammonia through

photodissociation[45] would have created selective pressures for a dissimilatory N pathway that

also produced assimilatory N. We demonstrate that NrfA proteins are significantly enriched in N

compared to NirK (i.e., no evolutionary constraints to cost minimize N in the *nrfA* gene

product[15]) (Figure *1*B) and that ammonium production (without accumulation in the medium)

supports growth in *I. calvum* (Figure *1*C). The Nrf module is also relatively simplistic in that it

receives electrons directly from the quinol pool and not the bc_1 complex used in denitrification.

The early exit of electrons from the ETC (i.e., before reaching the bc_1 complex) suggests that Nrf

may have originated prior to the bc_1 complex. Furthermore, the quinol oxidation site (Q_o) of

cytochrome *b* contains a PDWY motif, indicative of an ancestral LP respiratory chain found in

many Gram-positive organisms[46]. However, there is still debate regarding the presence of a

cytochrome *bc* complex in the last universal common ancestor[46,47]. Lastly, the Nrf module is

wired to operate via a q-loop with formate dehydrogenase whose Mo-cofactors would have also

been bioavailable during the Archean, further supporting an early evolution.

In summary, we employ a new predictive framework that accounts for the biochemistry and

evolutionary history of N modules, ETC complexes, and pool quinones to suggest the

mechanisms by which these two pathways are regulated at the molecular level. With this

understanding, it may be possible to extend our framework to environmental microbial

populations and accelerate model development across different ecosystem scales (i.e., cross-scale

systems biology).

Materials and Methods

Culture Conditions

<u>Sample collection</u>: *I. calvum* was isolated from a groundwater well (GW247) collected on

2/18/2013 at Oak Ridge National Laboratory (Lat: 35.97990, Long: 84.27059) with a

groundwater temperature of 15.57 ºC, conductivity of 521.9 µS/cm, and pH of 7.71. *I. calvum*

was isolated and obtained from Dr. R. Chakraborty (Lawrence Berkeley National Laboratory).

We began our experimental process by first picking single colonies from LB agar plates and

transferring the clonal isolates to LB broth.

<u>Media preparation</u>: All subsequent cultures were grown at 30 ºC and shaken at 250 rpm. Nitrate

reducing minimal media was prepared with the following final concentrations: NaCl (0.6mM),

NH_4Cl (1.75mM) (for ammonium replete conditions but not used in NH4-deplete conditions),

MgCl2 (0.2mM), CaCl2 (0.04mM), KCl (0.1mM), K2HPO4 (0.01mM), NaHCO3- (0.3mM),

cysteine (1mM) as reducing agent, resazurin as redox indicator, and trace elements and trace

vitamin solutions as reported[35,48]. 1M sterile filtered (0.2µm) Concentrated stocks of 60% w/w

sodium DL-lactate solution (Sigma-Aldrich, St. Louis, MO, USA), sodium-nitrate and sodium-

nitrite (≥99%, Fisher Scientific, Pittsburg, PA, USA) were diluted into media prior to

autoclaving to achieve the desired $C:NO_3^-$ ratio. $C:NO_3^-$ ratio was calculated based on[3] where the

number of C atoms (n) in the e-donor is multiplied by the concentration of the e-donor, divided

by the number of N atoms in the e-acceptor multiplied by the concentration of the e-acceptor

(Table *S4*). Media preparation was conducted in a 2L Widdel Flask. After autoclaving, the media

was immediately put under an anoxic headspace (N2/CO2 80:20 mix) and sterile filtered

(0.2µm) trace elements, trace vitamins, and reducing agent were added. The media was cooled

under an anoxic headspace and buffered with bicarbonate to maintain a pH of 7.2. Hungate

technique was used to dispense media into culture tubes (20 mL) and serum vials (100 mL) pre-

flushed with a sterile stream of ultra-high purity (UHP) N_2 and sealed with blue 1" butyl rubber

stoppers. End-point cultures were grown in Balch tubes (18x150-mm glass tube) sealed with

butyl rubber stoppers. Cultures for time-course sampling were grown in 160ml serum vials. All

end-point experiments were terminated after 100 hours unless otherwise noted. Mean pH for all

culture vessels (time series and end-point; Table *S5*), measured at the end of each experiment,

was 7.3±0.05 (n=144)

Analytical procedures

<u>Growth Curve/Cell counts/Yield Measurements</u>: Growth curves were measured from scratch-

free Balch-tubes grown cultures using an automated optical density reader at OD_{600} nm

(Lumenautix LLC, Reno, NV). End-point cultures were monitored until all replicates reached

stationary phase (65-100 hours depending on $C:NO_3^-$ treatment) (Figure *S6*).

Cell counts were performed by fixing cells in 4% paraformaldehyde (final concentration) for 20

minutes, filtered onto 0.2µm pore-sized black polycarbonate filters, and washed three times with

phosphate buffered saline (PBS, pH 7.2). Filtered cells captured on the black polycarbonate

filters were stained with SYBR® Gold nucleic acid stain (10-minute incubation) (ThermoFisher

Scientific) and counted manually with a fluorescence microscope (Olympus BX60, Tokyo,

Japan). We collected cells from during lag phase, exponential phase, and stationary phase in

order to create a standard curve of cell counts versus optical density (OD_{600}). We fit a linear

model to cell count versus OD_{600} (R^2=0.99) and used the resulting linear equation for cell count

enumeration for growth curves during our various treatment conditions.

Biomass concentrations were measured by filtration and drying as per standard protocol[49] for

8mM lactate/12mM nitrate and 0.8mM lactate/1.2mM nitrate treatments and conducted in

parallel with growth curve/cell counts as described above. Analysis from triplicate cultures

yielded (0.064 ± 0.003) and (0.016 ± 0.001) mg of biomass (dry weight) ml^{-1} for 8mM and 0.8

mM lactate cultures, respectively. Cell counts from stationary phase cultures were (1.5 ± 0.05) x

10^7 and (1.16 ± 0.09) x 10^6 for 8mM and 0.8 mM lactate cultures, respectively. From these

values the dry weight of a single *I. calvum* cell was estimated to be 1.09 x 10^{-10} g. Growth yield

(*Y*) (Table S3) was calculated by dividing biomass (g) by lactate mass (g) and moles consumed,

as described by[50]. Lactate measurements are described below.

<u>Ion and Gas Chromatography Measurements</u>: A dual channel Dionex ICS-5000+ (Thermo

Scientific) was used to measure all anions (AS11-HC) and cations (CS-16) from the bacterial

growth media. New glass IC vials were used for every sample in order to ensure no cross

contamination of analytes. Headspace gas from Balch tubes and serum vials was sampled with

volume appropriate gastight syringes (Hamilton Company, Reno, NV) pre-flushed with UHP N_2.

For high and low nutrient treatments, 10µl and 100µl of headspace were sampled and diluted

into 12ml exetainers (Labco, Lampter, Wales, UK) over-pressurized with 15ml UHP N_2,

respectively. Similar dilutions were performed for nitrite as e-acceptor experiments, ammonium-

deplete experiments, and time-series experiments. For time-series experiments, an equal volume

of headspace gas that was removed at each time-point was replaced with sterile UHP N_2. N_2O

and NO were measured by gas chromatography (Shimadzu Greenhouse Gas Analyzer GC-2014)

using a 500µl injection volume. The rubber septa on the injection port of the GC was replaced

after 100 injections in order to prevent leakage of the sample after the injection needle was lifted

out from the injection port. Aqueous concentrations of N_2O were calculated using a Henry's

constant of 1.751 (mM (g)/mM (aq)) corrected for the medium's ionic strength and temperature.

A total of 8-11 replicates per treatment were analyzed for all experiments discussed in this work

(Table *S5*).

<u>Ammonium production and thermodynamic calculations</u>: All nitrogen needed for assimilation

must be first converted into ammonium[50]. This fact makes ammonium production measurements

via respiratory nitrite ammonification difficult since the cells simultaneous consume ammonium

for growth. Therefore, direct measurements of ammonium from IC or spectrophotometric

analyses would not yield accurate production values. Because *I. calvum* simultaneously

consumes ammonium as its sole N source for growth and produces ammonium via respiratory

nitrite ammonification, we first measured ammonium consumption with O_2 and lactate by

calculating the difference between starting and ending ammonium concentrations. These

ammonium consumption values were then normalized to lactate consumed (0.31μmols

NH_4^+/lactate) and to ammonium consumed per cell, which was calculated from average cell

number of stationary phase biomass (Figure *S6*) (7.07×10^{-7} μmols NH_4^+/cell). These two

ammonium consumption constants were used as two separate methods to compare ammonium

production based on actual starting and ending ammonium/lactate measurements and cell

number during nitrate reducing minimal media experiments.

For ammonium production normalized to lactate consumption, calculations are based on a mass

balance approach from[51] for Total Belowground Carbon Allocation (TBCA) but adapted for

nitrogen flux instead of carbon flux:

$$\text{Net NH}_4^+{}_{(APL)} = (\Delta\text{lactate}_{\text{start-end}} \times 0.31\mu\text{mols NH}_4^+/\text{lactate}) + \Delta\text{NH}_4^+{}_{\text{end-start}} \qquad (1)$$

Here, the $\Delta\text{lactate}_{\text{start-end}}$ variable (μmols) is multiplied by the ammonium consumed per lactate

consumed constant (APL). This value is added to the $\Delta\text{NH}_4^+{}_{\text{end-start}}$ variable (μmols), denoted as

ending minus starting concentration, which defines whether the change in ammonium is positive

(more ammonium produced than consumed) or negative (more ammonium consumed than

produced). The ammonium production method using cell count (APC) is calculated as reported

by[35], where :

$$\text{Net NH}_4^+{}_{(APC)} = (\text{Cell Yield} \times 7.07 \times 10^{-7} \mu\text{mols NH}_4^+/\text{cell}) + \Delta\text{NH}_4^+{}_{\text{end-start}} \qquad (2)$$

Where Cell Yield is the number of cells at stationary phase (Figure *S6*). Comparison of these two

ammonia production methods revealed that they are in agreement with each other for anaerobic

nitrate reducing media experiments (Figure *S7*). However, because the cell counts under low

nutrient treatments were at the lower limit of our standard curve (OD versus cell count), we

observed slightly more variability using the APC method. Lactate measurements by IC gave us

higher sensitivity over the large range of treatments tested, thus we report the APL method in the

main text as the most sensitive method for reporting ammonium production by respiratory

ammonification.

Thermodynamic calculations for anaerobic lactate oxidation with nitrate and nitrite were carried

out using standard Gibbs free-energy values defined by Thauer *et al.*,[52].

Nucleic Acid Extraction and Sequencing

DNA for genome sequencing was extracted with phenol-chloroform. The DNA quality and

integrity was verified on Agilent 2200 TapeStation and DNA was sequenced on the PacBio

platform (Duke Center for Genomic and Computational Biology).

For gene expression analysis, *I. calvum* cells were grown in 300 mL batch cultures under 8mM

and 0.8mM lactate (12mM and 1.2mM nitrate, respectively) (n=3). Growth curves were

monitored in real-time to collect cells during early-exponential, late-exponential, and stationary

phase. Upon sampling, culturing vessels were immediately incubated on ice, aseptically

transferred to 50 mL falcon tubes and immediately centrifuged at 15000 G for 5 minutes at 4 ºC.

Supernatant was immediately poured off and the cell pellets were harvested and stored in TRIzol

Reagent (Ambion RNA by Life Technologies) at -80 ºC until further processing. RNA was

extracted with TRIzol Reagent (Ambion RNA by Life Technologies), following the

manufacturer's protocol. The extraction was scaled up to 20 ml of TRIzol and performed in 50

ml conical tubes. RNA was resuspended in water, treated with Baseline-ZERO DNase

(Epicentre) and cleaned up with RNeasy MinElute Cleanup Kit (QIAGEN). Total extracted RNA

from each growth phase was quantified (RiboGreen RNA quantification), analyzed for RNA

integrity number (RIN) (RIN≥0.8 were used for sequencing), treated with RiboZero rRNA

removal kit (Illumina, San Diego, CA). Library preparation was performed with the TruSeq

Stranded mRNA Library Prep kit (Illumina, San Diego, CA) and sequenced using Illumina

NextSeq 500 2x150 Mid Output v2 platform, resulting in 10-15 million reads per sample.

Phylogenetic, Genomic, and Transcriptomic Analysis

Genomic DNA was assembled using Canu (version 1.7.1) with an estimated genome size of 5

million base pairs[53]. The resulting single contiguous fragment was aligned to the *I. calvum* 7KIP

genome (Acc: NC_014830.1) to compare sequence similarity in Mauve[54,55]. Genome annotation

for C5 was performed through the NCBI Prokaryotic Genome pipeline

(www.ncbi.nlm.nih.gov/genome/annotation_prok/). Additional gene prediction analysis and

functional annotation was performed by the DOE Joint Genome Institute (JGI) using the Isolate

Genome Gene Calling method (Prodigal V2.6.3 February, 2016) under the submission ID

172966. The complete genome sequence and annotation is available in the NCBI database under

the BioProject number PRJNA475609.

A set of 34 NrfA amino acid sequences, representing 33 complete genome sequences and 1

octaheme nitrite reductase (ONR) from known respiratory ammonification organisms were

downloaded from GenBank (Table *S1*). A multiple sequence alignment (MSA) was generated

from the sequences annotated as cytochrome c nitrite reductase and ONR using MUSCLE[56]. The

resulting alignment was visualized within MEGA5[57] where the alignment was manually screened

for the presence of conserved amino acid residues consistent with those found in NrfA (i.e.,

heme motifs). A maximum likelihood tree was created from the alignment using RAxML[58] with

500 bootstrap iterations. The presence of NapA, NarG, NirK, and Nor modules were manually

queried from each NCBI genome in our set and confirmed by MSA, as described above.

Metabolic pathway for pool quinone type was queried on BioCyc Pathway/Genome Database

(biocyc.org) for each organism in our set. The structure of *I. calvum*'s NirK protein was

predicted using the protein structure predicting algorithm Phyre2[59]. Protein atomic composition

for C and N was calculated from amino acid sequences as input files, as described by[14,15], using

custom python scripts for each element separately (github.com/dvuono/Cost_minimization).

For transcriptomic analysis, the resulting raw reads were inspected using FastQC[60] to determine

quality, read length, and ambiguous read percentage. Reads were trimmed based on quality score

with a sliding window of 5 base pairs, quality cutoff of 28, trailing cutoff quality score of 10, as

well as adapter contamination removal in Trimmomatic[61]. Due to the high similarity of C5 to

7KIP, reads were aligned to the *Intrasporangium calvum* genomic reference sequence and gtf file

(Acc: NC_014830.1) using the STAR RNA-seq aligner[62], with the --limitBAMsortRam

parameter set to the recommended value by STAR. Sequence reads were mapped to genomic

features to obtain count data using featureCounts[63]. Systematic changes across experimental

conditions were performed on normalized read counts in DESeq2[64]. The RNA-seq data reported

in this study are available within the NCBI BioProject number PRJNA475609.

Statistical analyses were conducted in the R environment for statistical computing (r-

project.org). Data that was tested using parametric statistical analysis were first validated for

normality by visualizing the data as a histogram and testing via Shapiro-Wilks test for normality.

ACKNOWLEDGMENTS. We thank F. von Netzer, K. Hunt, S Morales, N. Stopnisek, K. Meinhardt, W. Qin, N

Elliot, T. Hazen, H. Carlson, B. Ramsey, A. Murray, Z. Harrold, T. Morgan, P. Longley, and R. Hallnan. This

research was supported by a grant from the Nevada Governor's office of Economic Development (JG), by the

Desert Research Institute (DRI) postdoctoral research fellowship program, and in part by Ecosystems and Networks

Integrated with Genes and Molecular Assemblies (ENIGMA) (http://enigma.lbl.gov)—a Scientific Focus Area

Program at Lawrence Berkeley National Laboratory under contract number DE-AC02-05CH11231 and funded in

part by Oak Ridge National Laboratory under contract DE-AC05-00OR22725, and is based upon work supported by

the U.S. Department of Energy, Office of Science, Office of Biological & Environmental Research.

References

1.   Knowles, R. Denitrification. *Microbiol. Rev.* **46,** 43–70 (1982).

2.   Tiedje, J. M., Sexstone, A. J., Myrold, D. D. & Robinson, J. A. Denitrification: ecological niches ,
competition and survival. *Antonie Van Leeuwenhoek* **48,** 569–583 (1982).

3.   Yoon, S., Cruz-García, C., Sanford, R., Ritalahti, K. M. & Löffler, F. E. Denitrification versus respiratory
ammonification: Environmental controls of two competing dissimilatory NO 3-/NO 2-reduction pathways in
Shewanella loihica strain PV-4. *ISME J.* **9,** 1093–1104 (2015).

4.   van den Berg, E. M., van Dongen, U., Abbas, B. & van Loosdrecht, M. C. Enrichment of DNRA bacteria in
a continuous culture. *ISME J.* **9,** 2153–2161 (2015).

5.   Schmidt, C. S., Richardson, D. J. & Baggs, E. M. Constraining the conditions conducive to dissimilatory
nitrate reduction to ammonium in temperate arable soils. *Soil Biol. Biochem.* **43,** 1607–1611 (2011).

6.   Hardison, A. K., Algar, C. K., Giblin, A. E. & Rich, J. J. Influence of organic carbon and nitrate loading on
partitioning between dissimilatory nitrate reduction to ammonium (DNRA) and N2 production. *Geochim.*
Cosmochim. Acta **164,** 146–160 (2015).

7.   Fazzolari, É., Nicolardot, B. & Germon, J. C. Simultaneous effects of increasing levels of glucose and
oxygen partial pressures on denitrification and dissimilatory nitrate reduction to ammonium in repacked soil
cores. *Eur. J. Soil Biol.* **34,** 47–52 (1998).

8.   Galloway, J. N. *et al.* The Nitrogen Cascade. *Bioscience* **53,** 341 (2003).

9.   Moparthi, V. K. & Hägerhäll, C. The evolution of respiratory chain complex i from a smaller last common
ancestor consisting of 11 protein subunits. *J. Mol. Evol.* **72,** 484–497 (2011).

10.   Dibrova, D. V., Cherepanov, D. A., Galperin, M. Y., Skulachev, V. P. & Mulkidjanian, A. Y. Evolution of

cytochrome bc complexes: From membrane-anchored dehydrogenases of ancient bacteria to triggers of
apoptosis in vertebrates. *Biochim. Biophys. Acta - Bioenerg.* **1827,** 1407–1427 (2013).

11. Bergdoll, L., Ten Brink, F., Nitschke, W., Picot, D. & Baymann, F. From low- to high-potential bioenergetic
chains: Thermodynamic constraints of Q-cycle function. *Biochim. Biophys. Acta* **1857,** 1569–1579 (2016).

12. Schoepp-Cothenet, B. *et al.* Menaquinone as pool quinone in a purple bacterium. *Proc. Natl. Acad. Sci.* **106,**
8549–54 (2009).

13. Soo, R. M., Hemp, J., Parks, D. H., Fischer, W. W. & Hugenholtz, P. On the origins of oxygenic
photosynthesis and aerobic respiration in Cyanobacteria. *Science (80-.).* **355,** 1436–1440 (2017).

14. Baudouin-Cornu, P., Schuerer, K., Marlière, P. & Thomas, D. Intimate evolution of proteins: Proteome
atomic content correlates with genome base composition. *J. Biol. Chem.* **279,** 5421–5428 (2004).

15. Grzymski, J. J. & Dussaq, A. M. The significance of nitrogen cost minimization in proteomes of marine
microorganisms. *ISME J.* **6,** 71–80 (2012).

16. Einsle, O., Messerschmidt, A., Huber, R., Kroneck, P. M. H. & Neese, F. Mechanism of the six-electron
reduction of nitrite to ammonia by cytochrome c nitrite reductase. *J. Am. Chem. Soc.* **124,** 11737–11745
(2002).

17. Kraft, B., Strous, M. & Tegetmeyer, H. E. Microbial nitrate respiration - Genes, enzymes and environmental
distribution. *J. Biotechnol.* **155,** 104–117 (2011).

18. Godfrey, L. V. & Falkowski, P. G. The cycling and redox state of nitrogen in the Archaean ocean. *Nat.*
Geosci. **2,** 725–729 (2009).

19. Nitschke, W. & Dracheva, S. Reaction center associated cytochromes in Anoxygenic Photosynthetic
Bactera. in (eds. Blankenship, R., Madigan, M. & Bauer, C.) 775–805 (Kluwer Academic, 1995).

20. Lalucat, J., Bennasar, A., Bosch, R., Garcia-Valdes, E. & Palleroni, N. J. Biology of Pseudomonas stutzeri.
Microbiol. Mol. Biol. Rev. **70,** 510–547 (2006).

21. Suharti & de Vries, S. Membrane-bound denitrification in the Gram-positive bacterium Bacillus
azotoformans. *Biochem. Soc. Trans.* **33,** 130–133 (2005).

22. Heylen, K., Keltjens, J. & Stein, L. Y. Redundancy and modularity in membrane-associated dissimilatory
nitrate reduction in Bacillus. *Front. Microbiol.* **3,** 1–27 (2012).

23. Decleyre, H., Heylen, K., Bjorn, T. & Willems, A. Highly diverse nirK genes comprise two major clades
that harbour ammonium-producing denitrifiers. *BMC Genomics* **17,** 1–13 (2016).

24. Kalakoutskii, L. V., Kirillova, I. P. & Krassilnikov, N. A. a New Genus of the. *J. Gen. Microbiol.* **48,** 373-
79–85 (1967).

25. Klotz, M. G. & Stein, L. Y. Nitrifier genomics and evolution of the nitrogen cycle. *FEMS Microbiol. Lett.*
278, 146–156 (2008).

26. Kraft, B. *et al.* The environmental controls that govern the end product of bacterial nitrate respiration.
Science (80-.). **345,** 676–679 (2014).

27. Abbe, K. S., Takahashi, S. & Yamada, T. Invovlement of oxygen-sensitive pyruvate-formate lysase in
mixed-acid fermetnation by Streptococcus mutans under strictly anaerobic conditions. *J. Bacteriol.* **152,** 175
(1982).

28. Das, A., Silaghi-Dumitrescu, R., Ljungdahl, L. G. & Kurtz, D. M. Cytochrome *bd* oxidase, oxidative stress,
and dioxygen tolerance of the strictly anaerobic bacterium *Moorella thermoacetica*. *J Bacteriol* **187,** 2020–

2029 (2005).

29.   Lotka, A. J. Contribution to the Energetics of Evolution. *Proc. Natl. Acad. Sci.* **8,** 147–151 (1922).

30.   DeLong, J. P. The maximum power principle predicts the outcomes of two-species competition experiments.
638   *Oikos* **117,** 1329–1336 (2008).

639   31.   Strohm, T. O., Griffin, B., Zumft, W. G. & Schink, B. Growth yields in bacterial denitrification and nitrate
640   ammonification. *Appl. Environ. Microbiol.* **73,** 1420–1424 (2007).

641   32.   Zumft, W. G. Cell biology and molecular basis of denitrification. *Microbiol. Mol. Biol. Rev.* **61,** 533–616
642   (1997).

643   33.   Hemp, J. & Gennis, R. B. Diversity of the Heme-Copper Superfamily in Archaea: Insights from Genomics
644   and Structural Modeling. in *Results and Problems in Cell Differentiation* (eds. Schäfer, G. & Penefsky, H.
645   S.) 1–28 (Springer Science+Business Media, 2008).

646   34.   Rowley, G. *et al.* Resolving the contributions of the membrane-bound and periplasmic nitrate reductase
647   systems to nitric oxide and nitrous oxide production in Salmonella enterica serovar Typhimurium. *Biochem.*
648   *J.* **762,** 755–762 (2012).

649   35.   Yoon, S., Sanford, R. A. & Löffler, F. E. Shewanella spp. Use acetate as an electron donor for denitrification
650   but not ferric iron or fumarate reduction. *Appl. Environ. Microbiol.* **79,** 2818–2822 (2013).

651   36.   Yoon, S., Sanford, R. A. & Löffler, F. E. Nitrite control over dissimilatory nitrate/nitrite reduction pathways
652   in Shewanella loihica strain PV-4. *Appl. Environ. Microbiol.* **81,** 3510–3517 (2015).

653   37.   van den Berg, E. M., Rombouts, J. L., Kuenen, J. G., Kleerebezem, R. & van Loosdrecht, M. C. M. Role of
654   nitrite in the competition between denitrification and DNRA in a chemostat enrichment culture. *AMB*
655   *Express* **7,** 1–7 (2017).

656   38.   Link, T. A., Hagen, W. R., Pierik, A. J., Assmann, C. & von Jagow, G. Determination of the redox
657   properties of the Rieske cluster of bovine heart bc1 complex by direct electrochemistry of a water- soluble
658   fragment. *Eur.J.Biochem.* **208,** 685–691 (1992).

659   39.   Ugulava, N. B. & Crofts, A. R. CD-monitored redox titration of the Rieske Fe-S protein of Rhodobacter
660   sphaeroides: pH dependence of the midpoint potential in isolated bc1 complex and in membranes. *FEBS*
661   *Lett.* **440,** 409–413 (1998).

662   40.   Zu, Y. *et al.* Reduction Potentials of Rieske Clusters: Importance of the Coupling between Oxidation State
663   and Histidine Protonation State. *Biochemistry* **42,** 12400–12408 (2003).

664   41.   Ansong, C. *et al.* Protein redox dynamics during light-to-dark transitions in cyanobacteria and impacts due
665   to nutrient limitation. *Front. Microbiol.* **5,** 1–10 (2014).

666   42.   Snyder, C. H., Merbitz-Zahradnik, T., Link, T. A. & Trumpower, B. L. Role of the Rieske Iron – Sulfur
667   Protein Midpoint Potential in the Protonmotive Q-Cycle Mechanism of the Cytochrome bc 1 Complex. *J.*
668   *Bioenerg. Biomembr.* **31,** 235–242 (1999).

669   43.   Denke, E. *et al.* Alteration of the midpoint potential and catalytic activity of the Rieske iron-sulfur protein
670   by changes of amino acids forming hydrogen bonds to the iron-sulfur cluster. *J. Biol. Chem.* **273,** 9085–
671   9093 (1998).

672   44.   Hunte, C., Solmaz, S., Palsdóttir, H. & Wenz, T. A structural perspective on mechanism and function of the
673   cytochrome bc1 complex. in *Results and Problems in Cell Differentiation* (eds. Schäfer, G. & Penefsky, H.
674   S.) 253–270 (Springer Science+Business Media, 2008).

45.   Falkowski, P. G. Evolution of the nitrogen cycle and its influence on the biological sequestration of CO2 in

the ocean. *Nature* **387,** 272–275 (1997).

46. Kao, W. C. & Hunte, C. The molecular evolution of the Qo Motif. *Genome Biol. Evol.* **6,** 1894–1910 (2014).

47. Dibrova, D. V., Cherepanov, D. A., Galperin, M. Y., Skulachev, V. P. & Mulkidjanian, A. Y. Evolution of
cytochrome bc complexes: From membrane-anchored dehydrogenases of ancient bacteria to triggers of
apoptosis in vertebrates. *Biochim. Biophys. Acta - Bioenerg.* **1827,** 1407–1427 (2013).

48. Hillesland, K. L. *et al.* Erosion of functional independence early in the evolution of a microbial mutualism.
Proc. Natl. Acad. Sci. **111,** 14822–14827 (2014).

49. APHA. *Standard Methods for the Examination of Water and Wastewater*. (American Public Health
Association: Washington, DC, 2012). doi:ISBN 9780875532356

50. White, D. *The physiology and biochemistry of prokaryotes*. (Oxford University Press, 2000).

51. Giardina, C. P. & Ryan, M. G. Total belowground carbon allocation in a fast-growing Eucalyptus plantation
estimated using a carbon balance approach. *Ecosystems* **5,** 487–499 (2002).

52. Thauer, R. K., Jungermann, K. & Decker, K. Energy conservation in chemotrophic anaerobic bacteria.
Bacteriol. Rev. **41,** 100–180 (1977).

53. Koren, S. *et al.* Canu : scalable and accurate long- - - read assembly via adaptive k - - - mer weighting and
repeat separation. *Genome Res.* 1–35 (2017). doi:10.1101/gr.215087.116.Freely

54. Darling, A. C. E., Mau, B., Blattner, F. R. & Perna, N. T. Mauve : Multiple Alignment of Conserved
Genomic Sequence With Rearrangements. *Genome Res.* **14,** 1394–1403 (2004).

55. Darling, A. E., Mau, B. & Perna, N. T. Progressivemauve: Multiple genome alignment with gene gain, loss
and rearrangement. *PLoS One* **5,** e11147 (2010).

56. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids*
Res. **32,** 1792–7 (2004).

57. Tamura, K. *et al.* MEGA5: Molecular evolutionary genetics analysis using maximum likelihood,
evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* **28,** 2731–2739 (2011).

58. Stamatakis, A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies.
Bioinformatics **30,** 1312–1313 (2014).

59. Kelley, L. A., Mezulis, S., Yates, C., Wass, M. & Sternberg, M. The Phyre2 web portal for protein
modelling, prediction, and analysis. *Nat. Protoc.* **10,** 845–858 (2015).

60. Andrews, S. FastQC: A quality control tool for high throughput sequence data. *Available online at: http://*
www.bioinformatics.babraham.ac.uk/projects/fastqc. (2010).

61. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: A flexible trimmer for Illumina sequence data.
Bioinformatics **30,** 2114–2120 (2014).

62. Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29,** 15–21 (2013).

63. Liao, Y., Smyth, G. K. & Shi, W. FeatureCounts: An efficient general purpose program for assigning
sequence reads to genomic features. *Bioinformatics* **30,** 923–930 (2014).

64. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data
with DESeq2. *Genome Biol.* **15,** 1–21 (2014).