Resource limitation modulates the fate of dissimilated nitrogen in a dual-pathway

Actinobacterium

David C. Vuono[a], Robert W. Read[a], James Hemp[b], Benjamin W. Sullivan[c], John A. Arnone III[a],

Iva Neveux[a], Bob Blank[d], Carl Staub[e], Evan Loney[a], David Miceli[a], Mari Winkler[f], Romy

Chakraborty[g], David A. Stahl[f], Joseph J. Grzymski[a,1]

[a]Division of Earth and Ecosystem Sciences, Desert Research Institute, Reno, NV 89512, USA

[b]Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena,

CA, USA

[c]Department of Natural Resources and Environmental Science, University of Nevada, Reno, NV

89557, USA

[d]Agricultural Research Service, U.S. Department of Agriculture, Reno, NV, 89512, USA

[e]Agtron, Inc. 9395 Double R Blvd, Reno, NV 89521, USA

[f]Department of Civil and Environmental Engineering, University of Washington, Seattle, WA,

98195, USA

[g]Lawrence Berkeley National Laboratory, Berkeley, CA, 94729, USA

[1]Correspondence: Joseph J. Grzymski, Division of Earth and Ecosystem Sciences, Desert

Research Institute, Reno, 2215 Raggio Parkway, NV 89512, USA

Email: Joe.Grzymski@dri.edu

Abstract

Carbon to nitrate ($C:NO_3^-$) ratios are thought to modulate pathway selection between respiratory ammonification and denitrification, two nitrogen (N) dissimilatory processes vital to the global N budget. However, the molecular mechanisms that enable the selection of these pathways remains unclear. Here we evaluate $C:NO_3^-$ control on pathway selection in *Intrasporangium calvum* C5, a Gram-positive menaquinone-based dual-pathway denitrifier/respiratory ammonifier and show that $C:NO_3^-$ control theory is insufficient to explain pathways selection. We demonstrate that the bacterium disproportionately utilizes ammonification rather than denitrification (with nitrous oxide as its terminal end-product) when grown under carbon or nitrate limitation, not $C:NO_3^-$ ratio. The ammonification pathway also promoted higher bacterial growth rates. Time series analysis of metabolite and transcriptional profiles during growth showed that transcript abundances for nitrite reducing complexes, NrfAH and NirK, significantly increased in response to nitrite production. Although lactate was the only carbon source provided to the organism, we detected formate production during growth ($\sim$200 μM), a five-fold upregulation in formate transporters, and a simultaneous up-regulation of formate dehydrogenase used to translocate protons via a quinol-loop. These results suggest that additional reducing equivalents can be obtained from a single carbon source and used for ammonification during resource limitation. Mechanistically, pathway selection may be driven by intracellular redox potential (redox poise), which is lowered during resource limitation, thereby decreasing catalytic activity of the bc_1 complex (an upstream electron transport step needed for denitrification). Our work advances our understanding of the conditions and underlying mechanisms that select for denitrification and respiratory ammonification in environmental systems.

Significance

The mechanisms regulating electron transfer through branched anaerobic respiratory chains, such as dual-pathway nitrite reduction, have remained elusive. In this work, we tested the response of resource-limitation and -excess versus $C:NO_3^-$ ratio on denitrification versus respiratory ammonification in a dual-pathway denitrifying/respiratory ammonifier from the phylum *Actinobacteria*. We discovered that carbon and nitrate scarcity promoted a higher nitrogen flux through the respiratory ammonification pathway. Thus, control of respiratory pathway, denitrification versus respiratory ammonification, is not modulated by $C:NO_3^-$ ratio, as

previously thought, but rather the cell's need to maximize energy recovery when either resource

is in short supply. This discovery has significant relevance to a developing understanding of

factors controlling retention or release of reactive nitrogen from environmental systems.

Introduction

Globally, respiratory ammonification and denitrification are vital nitrogen (N) dissimilation

pathways that either retain reactive N to support net primary productivity or close the N-cycle

through the release of N_2, respectively. The environmental controls on these two pathways have

gained attention due to increased anthropogenic N inputs into the environment (1). Several

environmental controls have been posited, particularly the ratio of electron-donor to electron-

acceptor (e.g., $C:NO_3^-$) (2), but have received limited experimental and systematic validation (3,

4). Furthermore, the underlying molecular mechanisms for pathway selection are still unknown.

Over geologic time, strong selective pressures from Earth's shifting biogeochemistry and

oxidation-state have driven evolutionary adaptions to microbial electron transport chains (ETC)

(5, 6), respiratory chain redox potentials (7–9), and protein atomic composition (10, 11), which

also shed light on how these pathways are regulated in contemporary organisms. Here, by

identifying the biochemical and evolutionary differences between respiratory ammonification

and denitrification, we disentangle the functional significance and molecular mechanisms of

electron transfer through either pathway in a dual pathway organism.

From a biochemical standpoint, the primary difference between respiratory ammonification and

denitrification is their respective source of reducing equivalents in the ETC: 1) heme-based

cytochrome c nitrite reductase used in respiratory ammonification receive electrons directly from

the quinone (Q) pool (12) while 2) copper-containing and cytochrome cd_1 nitrite reductases used

in denitrification receive electrons from a soluble electron carrier (e.g., cytochrome c) via the bc_1

complex (13). From an evolutionary standpoint, we can place each N-module's origin to a

putative time in Earth history based on the metal co-factors that would have been bioavailable:

heme-based cytochromes in an ancient, more reduced, environment compared to the copper-

containing nitrite reductases in an oxidizing environment (14). The bioenergetic chains of

microorganisms also underwent selective pressure to shift from low-potential (LP) to high-

potential (HP) quinones in response to Earth's oxygenation (7, 8). Menaquinone (MK) is thought

to be the ancestral type of LP quinone (15) typically having a redox midpoint potential ~150 mV

lower than the evolutionarily more recent ubiquinone (UQ) (8). Organisms that use UQ are

thought to have evolved under high O_2 tensions and the only clades within the bacterial domain

to use UQ are the α-, β-, γ-proteobacteria (8). Surprisingly, our understanding for the

biochemistry of denitrification is based predominantly on HP UQ-based systems (16), leaving a

significant knowledge gap in the physiology and biochemistry of LP MK-based denitrifiers and

how they link electron transfer with energy capture under resource limitation.

Dual-pathway respiratory nitrite reducers, where respiratory ammonification and denitrification

modules are encoded in the same genome, provide a unique system to investigate the molecular

mechanisms of $C:NO_3^-$ control on pathway selection as they blend the presumptively ancient

(NrfA) and modern (NirK) N-reducing modules. While many dual-pathway denitrifiers and

respiratory ammonifiers have been identified (many of which use LP ETCs), most still lack

phenotypic characterization (17–19). Only *Shewanella loihica* PV-4, a Gram-negative, γ-

proteobacterium that uses UQ and dimethylmenaquinone (DMK)-based bioenergetics has been

thoroughly characterized (3, 20, 21). Nonetheless, these dual-pathway nitrite reducers are

capable of differential energy conservation through pathway bifurcation of nitrite. Yet no

mechanistic or metabolic models exist to explain how these dual-pathway organisms partition

electron flow and proton translocation between pathways to maximize resource-use efficiency. It

is not known if these organisms differentially distribute electron flow through alternative

respiratory chains in response to shifting resource availability or control each pathway

independently in response to resource thresholds. Is there an evolutionary and biochemical basis

for pathway selection?

It has long been postulated that $C:NO_3^-$ ratio modulates the activity of respiratory

ammonification versus denitrification (2), with numerous studies in recent years differing in the

methods used to calculate $C:NO_3^-$ ratio, carbon sources used, range of C:N ratios tested,

magnitudes of C and N concentrations, sample types (enrichment cultures vs isolates), and

culturing mode (chemostat/batch culture) (Table S1) (3, 4, 22–24). The hypothesis for $C:NO_3^-$

control states that high $C:NO_3^-$ ratios (stoichiometric limitation of nitrate relative to C) drive

respiratory ammonification while low $C:NO_3^-$ ratios (stoichiometric limitation of C relative to

nitrate) drive denitrification. This hypothesis is based on the rationale that denitrification

theoretically yields more free energy per electron, but respiratory ammonification yields more

free energy per nitrate due to the stoichiometry of each reaction: one nitrate is needed to make

ammonium while two are needed to make N_2O/N_2 (2, 25). Thus, when $C:NO_3^-$ ratio is low, cells

select denitrification to maximize the free energy advantage per electron. When $C:NO_3^-$ is high,

cells select respiratory ammonification to maximize the free energy advantage per nitrate.

However, the theoretical basis for this prediction is inconsistent with the observation that growth

yields of pure denitrifiers are significantly lower than expected and lower than growth yields of

respiratory ammonification (25). This observation suggests that despite a lower free energy yield

(ΔG) compared to denitrification, respiratory ammonification conserves more energy during

catabolism, through the generation of a proton motive force (Δp) in the ETC, to build more

biomass during anabolism. Thus, there is a need to better understand the effects of different

resource limitations on growth and pathway selection (i.e., allocation of C and N to dissimilatory

and assimilatory processes), which are often confounded by $C:NO_3^-$. For example, reportedly

"low" $C:NO_3^-$ ratios rarely fall below 1.5 have rarely been tested (Table S1), conditions in which

C would be a growth-limiting resource. Instead, pathway selection should be better predicted by

1) Liebig's law of the minimum (LM) and 2) the maximum power principle (MPP) (26). Under

the MPP, which states that biological systems are designed to maximize power intake and energy

transformation, the cell's aim is therefore to maximize power (i.e., realized in the form of growth

rate and yield) given the constraints of a growth-limiting resource (i.e., LM). Therefore,

limitation of a growth-limiting resource, whether it be C or NO_3^-, should dictate the selection of

the most efficient respiratory module to maximize power (i.e., MPP).

If indeed there is regulatory control by $C:NO_3^-$ ratio, the predicted end-products should also

remain identical at different substrate concentrations of the same ratio. In order to resolve the

inconsistencies and molecular mechanisms of $C:NO_3^-$ control on pathway selection and better

understand branched respiratory chains in LP-based nitrate-reducing organisms, we undertook

the characterization of the novel Gram-positive actinobacterium strain *Intrasporangium calvum*

C5: a dual-pathway nitrite reducer that uses MK as sole pool quinone. Physiological and

genomic analysis revealed that the bacterium is a close relative to type species *I. calvum* 7KIP,

reduces nitrate and nitrite using a range of carbohydrates, and uses ammonium as its sole

assimilatory N source. *I. calvum*'s dissimilatory N modules are composed of a membrane-bound

nitrate reductase (NarG), a copper nitrite reductase (NirK), a nitric oxide (NO) reductase (qNor),

and a cytochrome c nitrite reductase (CxxCH 1[st] heme motif variety) with a *nrfAH* operon

organization. Here we show that over a range of $C:NO_3^-$ ratios, duplicated at two substrate

concentrations, *I. calvum* disproportionately utilizes its ammonia-forming pathway during C

limitation (≤ 0.4mM lactate), when $C:NO_3^-$ ratios are < 1 (an observation contrary to the current

paradigm). Using a genome-guided approach coupled to time-series transcriptomics and

metabolite profiles, we identified differentially expressed genes in the bacterium's ETC and

central metabolic pathways. Using this information to inform a metabolic reconstruction of the

ETC we infer mechanisms by which these two pathways are regulated at the biochemical level.

Results

Genomic analysis of *I. calvum* C5. *I. calvum* C5 was isolated from a nitrate contaminated well

(>200mM nitrate) at the Oak Ridge National Laboratory Field Research Station in Oak Ridge,

TN. The strain was selected for further analysis due to its nitrate reducing phenotype in minimal

media. We sequenced and analyzed the genome of *I. calvum* C5 to first compare its similarity to

the type species *I. calvum* 7KIP. We identified a high degree of sequence similarity to 7KIP

based on three homologous sequence regions as locally collinear blocks. Genome size of C5 was

4,025,044 base pairs (bp), only 662 bp longer than 7KIP. Genomic analysis of the ETC revealed

the typical suite of complexes common to facultative aerobes, including primary dehydrogenases

(*nuo* complex, succinate dehydrogenase), alternative *NDH-2* NADH dehydrogenase, cytochrome

bc_1 complex, high-oxygen adapted cytochrome c oxidase (A-family), and low-oxygen adapted

cytochrome *bd* oxidase. The bc_1 complex subunits are also located immediately upstream of

cytochrome c oxidase, suggesting that these enzymes are encoded in a single operon creating a

supercomplex. Despite *I. calvum*'s seeming propensity for aerobic growth on a number of

growth media (27), its bioenergetic system uses MK as its sole pool quinone. *I. calvum* also

possesses multiple pathways for supplying electrons into the MK-pool, such as formate, malate,

hydroxybutyrate, and glycerophosphate dehydrogenases. Once in the MK-pool, there are

alternative pathways for MKH_2 oxidation that can circumvent the bc_1 complex, such as a

membrane-bound respiratory nitrate reductase module (NarG). In addition, to NarG, its

dissimilatory N module composition consists of a truncated denitrification pathway (N_2O is a

terminal product) using a copper nitrite reductase NirK and quinol-dependent nitric oxide

reductase qNor. The NirK peptide sequence length in *I. calvum* is significantly longer than that

of canonical NirKs: 909 versus 300-500 amino acids long and has 13 predicted transmembrane

regions on its N-terminus. *I. calvum* also possesses both catalytic and membrane anchor subunits

(NrfA and NrfH, respectively) for a pentaheme cytochrome c module involved in respiratory

nitrite ammonification.

I. calvum encodes for a functional NrfAH complex and assimilates NH$_4^+$ via respiratory

nitrite ammonification. To gain insight into possible function of the NrfAH complex, we

aligned the NrfA protein sequences from C5 and 7KIP to a collection of 33 recognized

cytochrome c nitrite reductases from published annotated genomes (Table S2). This phylogenetic

analysis confirmed that NrfA from *I. calvum* is a member of the CxxCH 1st heme motif group

(Fig.1A), which forms one of four clades on the NrfA phylogenetic tree. The other three clades

consisted of taxa possessing the common CxxCK 1st heme motif. We then queried the genomes

of the taxa in our phylogeny for other annotated N-reducing modules used in nitrate reduction,

nitrite reduction, NO-forming nitrite reduction, and primary pool quinone. Among the three

major clades of NrfA, at least 5 additional taxa are noted having dissimilatory N-module

inventories containing dual respiratory pathways: *Symbiobacterium thermophilum, Bacillus*

azotoformans, Bacillus bataviensis, *Bdellovibrio bacteriovorus*, and *Candidatus* Nitrospira

inopinata, (Fig.1A). Also, none of the taxa in our NrfA phylogeny harbored the cd_1 nitrite

reductases (NirS). Due to the exclusive presence of NirK representation in dual-pathway

membership, we asked whether there might be differences in protein atomic composition

between NirK and NrfA, given the disparate evolutionary origins of these modules (28). We

collected 20 additional publicly available NirK protein sequences from nondual-pathway

denitrifiers (Table S2) and calculated the protein C and N composition for our NirK/NrfA

collection as atoms per residue side-chain (Fig. 1B). These results showed a significant depletion

in C and N per residue side-chain for NirK compared to NrfA (C and N: $p<0.001$; t-test),

indicating that resource constraints are imprinted on the evolution of these proteins.

We next tested the functionality of *I. calvum*'s Nrf complex by growing the bacterium under

reducing conditions using Hungate-technique prepared minimal media with reducing agent (1

mM cysteine) to ensure no C was consumed with residual O$_2$ (8 mM lactate, 12 mM nitrate,

ammonium-replete). We then performed a state-transition where biomass from late-exponential

growth phase was collected and inoculated into ammonia-deplete media (Fig. 1C). Despite no

detectable amounts of ammonium produced in the media over time, cell counts increased

$5.4 \times 10^5 \pm 8.9 \times 10^4$ cells/mL (0.126 ± 0.02 optical absorbance units, OD_{600}) over a 48-hour

incubation, indicating consumption of ammonium produced by NrfA. Net ammonium production

was 13 ± 2.7 μmoles with the remainder of dissimilated N being used by the denitrification

pathway (24 ± 4.2 μmoles N_2O-N), resulting in a recovery of 97.4% dissimilated N. These results

confirmed that *I. calvum* C5 has a functional Nrf complex and also consumes the product

(ammonium) of respiratory nitrite ammonification.

Respiratory nitrite ammonification exceeds denitrification under C-limitation. We

investigated $C:NO_3^-$ control on respiratory ammonification versus denitrification on cultures of *I.*

calvum C5 over a high resource $C:NO_3^-$ range (16-0.4 mM lactate, 12 mM nitrate; ratio 4-0.1)

and low resource $C:NO_3^-$ range (1.6-0.04 mM lactate, 1.2 mM nitrate; ratio 4-0.1). This

experimental design enabled us to evaluate $C:NO_3^-$ control over a broader range than previous

studies that only considered ratios ≥ 1.5 (3, 4, 22), while also testing the effects of resource

concentration on pathway selection. Under all the treatments tested, gas and ion chromatography

measurements showed products of both respiratory pathways, differing only in the relative

fraction of N_2O versus ammonium production across treatments (Fig. 2). At high resource

concentrations, respiratory ammonification did not prevail at high $C:NO_3^-$ ratios (Fig. 2A & 2B,

left panels; Table S3). Instead, significantly greater amounts of N_2O were produced over

ammonium, though nitrite was still the major extracellular end-product of nitrate respiration.

Despite the predominance of N_2O production under the high resource concentrations, ammonium

production exceeded N_2O production only at the lowest $C:NO_3^-$ ratio (0.4 mM lactate, ratio=0.1)

(Table S3) and accounted for $76.2 \pm 0.1\%$ of dissimilated N.

Results from the low resource dataset provided weak support for the strict stoichiometry

hypothesis that $C:NO_3^-$ controls pathway selection. Ammonia exceeded N_2O production only

under one high $C:NO_3^-$ ratio treatment (ratio=4; 1.6 mM lactate; Fig. 2AB, right panels).

However, at ratios ≤ 1 (≤ 0.4 mM lactate), significantly more ammonium than N_2O was produced.

On average, respiratory ammonification accounted for $78.1 \pm 8.9\%$ of dissimilated N for lactate

concentrations ≤ 0.4 mM. When these results are taken in context with cell physiology, we

observed a significant and positive relationship between specific growth rate (μ) and the fraction

of N dissimilated by respiratory ammonification ($R^2=0.5$; $p<0.001$) (Fig. 2C, Fig. 3; Table S4).

Resource concentration influences the metabolite profiles of ammonium and N$_2$O

production. Given the co-occurrence of end products from both pathways during the end-point experiments (Fig. 2), we next investigated the timing of ammonium and N$_2$O production relative to metabolite profiles for lactate, nitrate/nitrite, and growth phase at two resource concentrations with the same ratio (8 mM and 0.8 mM lactate, ratio=2; Fig. 4A, S2, respectively). Despite ample e-donor and e-acceptor available for growth, the high resource cultures entered a quasi-stationary phase at ~50 hours, after which there was continued slow growth (Fig. 4A). Metabolite profiles showed that ammonium and N$_2$O production began simultaneously, as soon as nitrite was produced from nitrate reduction. The low resource cultures entered stationary phase at ~40 hours (Fig. S2) after nitrate had been fully utilized. No further cell growth was observed after stationary phase was reached. These results show that cell growth occurs primarily on the reduction of nitrate, while nitrite reduction to ammonium and N$_2$O takes place during a stationary growth phase, demonstrating that microbial activity is not always correlated with growth. The metabolite profiles for ammonium and N$_2$O at low resources did not mirror those observed at high resources. The rate of N$_2$O production significantly decreased and ammonium production oscillated rather than steadily increased through time. These differences in metabolite profiles clearly demonstrate that concentration influences the activities of pathway bifurcation. Furthermore, repeated time series experiments that were extended up to 300 hours show that nitrite is slowly depleted, but does not get fully consumed (Fig. S3). When cultures were given nitrite, instead of nitrate as a terminal electron acceptor (8 mM lactate, 12 mM nitrite; ratio=2), we observed no immediate growth (as was observed with nitrate) but measured more N$_2$O than ammonium production (33.4±4.8 µmoles N$_2$O-N and 8.0±2.5 µmoles NH$_4^+$, respectively) (Fig. S4), demonstrating respiratory ammonification does not exceed denitrification when nitrite is supplied as the sole acceptor in *I. calvum*.

Nitrite-reducing modules are up-regulated during late exponential- and stationary-phase

growth. In order to gain insight into mechanisms of gene regulation and transcriptional organization of *I. calvum*, we conducted RNA-Seq in parallel with the high resource time-series metabolite profile. This approach enabled us to compare genome-wide differential expression based on log$_2$ fold change (lfc) of RNA extracted from three growth phases: early exponential (EE), late exponential (LE), and stationary (ST) (Figs. 4B, S5). Within the central metabolic pathway beginning with the conversion of lactate to pyruvate, we observed a moderate decrease

in transcript abundance of L-lactate dehydrogenase (LDH) (Intca_16740) between EE-LE and -

ST (lfc = -1.6±0.7; -1.9±0.7), respectively. Lactate utilization protein C (LUP) (Intca_04080), an

enzyme involved in lactate degradation, also showed a moderate and significant decrease in

transcript abundance between EE-LE and -ST (lfc = -1.6±0.6; -2.4±0.6, *p*=0.002), respectively. *I.*

calvum encodes for two parallel metabolic pathways for pyruvate conversion to acetyl-CoA:

pyruvate dehydrogenase (PDH) (Intca_01255) and pyruvate ferredoxin oxidoreductase (PFOR)

(Intca_15510). For PDH, there was a significant and moderate increase in transcript abundance

between EE-LE and -ST (lfc = 2.1±0.6, *p*=0.002; 1.5±0.6), respectively. For PFOR, there was a

minor decrease in transcript abundance between EE-LE (lfc = -0.43±0.5), and then a moderate

increase in transcript abundance between EE-ST (1.1±0.5). Citrate synthase (Intca_04135), the

enzyme catalyzing the conversion of acetyl-CoA to citrate and the first step of the tricarboxylic

acid (TCA) cycle, showed a highly significant increase in transcript abundance between EE-LE

and -ST (lfc = 4.3±0.5, *p*<0.001; 6.9±0.5, *p*<0.001). Within the ETC, there was moderate and

significant decrease in transcript abundance for all subunits from the primary dehydrogenase

(*nuo* complex; Intca_03465-03539) between EE-LE and -ST (lfc = -1.2±0.3; -2.4±0.6, *p*<0.001),

respectively. Nitrate reductase subunits showed no change in transcript abundance between EE-

LE (lfc = 0.01±0.07) and moderately decreased in abundance by ST (lfc = -1.2±0.1), which was

corroborated by the depletion of nitrate during stationary phase. There was a significant increase

in transcript abundance of *nirK* (Intca_17170) (lfc = 2.2±0.6, *p*=0.003; 2.4±0.6, *p*<0.001) and

quinol dehydrogenase/membrane anchor subunit *nrfH* (Intca_09465) (lfc = 2.5±0.6, *p*=0.001;

2.1±0.6, *p*=0.003) by EE-LT and EE-ST, respectively, which coincided with nitrite production

(Fig. 4A). The catalytic subunit of the cytochrome c nitrite reductase complex (*nrfA*)

(Intca_09460) also increased moderately in transcript abundance by EE-LT and EE-ST (lfc =

1.6±0.6; 1.0±0.6), respectively (Fig. 4B). Contrary to the transcript abundance patterns of *nirK*

and *nrfAH*, nitric oxide reductase (qNor; Intca_01525) transcripts moderately increased between

EE-LT (lfc = 1.6±0.6) but decreased in the successive time periods (lfc = 0.43±0.6 between EE-

ST; lfc = -1.2±0.6 between LE-ST) (Fig. 4B).

There was a significant increase in transcript abundance of formate transporter *focA*

(Intca_17150) between EE-ST, as well as LE-ST (lfc = 4.9±0.7, *p*=0.002; 4.8±0.7, *p*=0.002;

respectively). We verified the production of formate in our ion chromatography measurements in

the range of 100-200μM following late exponential growth. We also observed a moderate

increase in transcript abundance of formate dehydrogenase (FDH) subunits (Intca_11150-

11160). These results implicate the activity of formate oxidation, which would contribute to a Δp

in the periplasm via a Q-loop mechanism and the reduction of MK for electron transfer to nitrite

via cytochrome c nitrite reductase. Considering that formate was not provided in our media

recipe, an alternative pathway for formate production must exist in *I. calvum*. We also observed

acetate production in similar concentrations as formate (100-200μM). In *E. coli*, formate is

produced anaerobically from the action of pyruvate formate lyase (PFL). We identified a putative

PFL based on genome annotation (Intca_12230), where transcript abundance also significantly

increased by ST. PFL is also highly sensitive to oxygen (29), which was also in agreement with a

significant increase in transcript abundance between EE-ST and LE-ST (Fig. 4B) of cytochrome

bd oxidase (Intca_01110 and Intca_01115), which is thought to protect anaerobic enzymes

against oxidative stress (30).

Discussion

We challenge the paradigm that $C:NO_3^-$ ratio strictly controls pathway selection in a dual-

pathway organism based on a simple principle: ratios do not account for the abundance of a

growth-limiting resource. We hypothesized that limitation in C or NO_3^- should better predict

pathway selection in a dual-pathway denitrifier/respiratory ammonifier. To test this hypothesis,

we systematically measured the response of the Gram-positive Actinobacterium

Intrasporangium calvum C5 to the same range of $C:NO_3^-$ ratios at both high and low resource

loadings to better resolve mechanisms of pathway selection. We demonstrated that resource

concentration, not $C:NO_3^-$ ratio, influences pathway selection. We found stronger support for

respiratory ammonification preference under C-limitation (at low $C:NO_3^-$ ratios), which also

grew at significantly higher growth rates (Fig. 2). These results suggest that the NrfA complex,

which receives electrons directly from the MK-pool, is optimized to maximize power when one

or more resources are limiting. These data, together with metabolic reconstructions from

metabolite and transcriptional profiles (Fig. 4), suggest that $C:NO_3^-$ ratio alone is insufficient to

explain pathway selection.

The theoretical basis for pathway selection is explained by the law of the minimum (LM) and the

maximum power principle (MPP), which state that growth is limited by the least abundant

resource and that biological systems are designed to maximize power in order to effectively

allocate energy to reproduction and survival (26), respectively. Here, it appears these two natural

theories are working together: when resources are limited, the cell utilizes the respiratory

pathway for growth that is optimized to maximize power. Power, in this case, is realized as

higher growth rates from the cultures exhibiting disproportionately higher ammonium production

than N_2O production (Fig. 2: high resources: $C:NO_3^-$ ratio = 0.1; low resources: $C:NO_3^-$ ratios =

4, 1, 0.5, 0.1; Fig. 3). More specifically, the bacterium must generate a greater Δp in order to

maximize power when starved for a growth limiting resource. This may help to further explain

how respiratory ammonification, which is overall energetically less favorable than denitrification

(lactate with nitrite: ΔG^o = -763.98 versus ΔG^o = -1196.93, respectively), can have higher growth

yields (25) and growth rates (Fig. 2, Fig. 3) under C- and N-limitation due to the higher energy

yield on a per-nitrite basis (denitrification: -217 KJ per mole nitrite; respiratory ammonification:

-399 KJ per mole nitrite). For comparison, a total of 8 H^+ are translocated during denitrification

by *I. calvum* (not including nitrate reduction since both pathways share this step) (Fig. 4): NADH

dehydrogenase translocates 4 H^+ per MKH_2 oxidized and the bc_1 complex translocates an

additional 4 H^+ per MKH_2 oxidized. However, 2 H^+ must be consumed in the periplasm to reduce

nitrite to NO (31). qNor has a net zero H^+ release (consumes 2 H^+ to make N_2O but releases 2

H^+) without MKH_2 regeneration (32). Thus, a net total of 6 H^+ are translocated per nitrite reduced

in denitrification with added biosynthetic costs of making the bc_1 complex and qNor. In

respiratory ammonification, MK/MKH_2 redox pair is cycled between NADH dehydrogenase and

formate dehydrogenase. 6 electrons and 8 H^+ are needed to reduce nitrite to ammonium, thus 3

MKH_2 are needed (12). If MKH_2 is received from NADH dehydrogenase, 12 H^+ are translocated

plus 2 H^+ from FDH. As each MKH_2 is oxidized at the binding site of NrfH, 2 H^+ are liberated

(12), resulting in a net total of 12 H^+ translocated per nitrite reduced for respiratory

ammonification. This implies that the cell might deplete its NADH pool more rapidly on a per

nitrite basis. However, if more protons are pumped in the early stages of growth, the cell would

be allocating the ATP generated for anabolism, as evidenced by higher growth rates in the

cultures exhibiting higher amounts of respiratory ammonification (Fig. 2), which is supported by

the MPP.

Following in the same line of reasoning for proton accounting in the ETC, *S. loihica* PV-4 (a

Gram-negative dual-pathway γ-proteobacterium) and *I. calvum* use different types of nitrate

reducing modules. *S. loihica* PV-4 utilizes NapA whereas *I. calvum* utilizes NarG (Fig. 1). The

latter translocates two H^+ per nitrate reduced while NapA consumes two H^+ in the periplasm

(13). Both reductases would generate a Δp via NADH dehydrogenase H^+ translocation, but the

NapA module would result in a net loss of two H^+, which may impact the selection of

downstream respiratory modules. For example, given the lower-than-expected observed growth

yields of denitrifiers compared to respiratory ammonifiers (25), if a NapA module is used, it

would make sense for respiratory ammonification to be selected under high C:N ratios because

the cell would need to compensate for less energy conservation during nitrate reduction. Nitric

oxide reductase composition may also impact pathway selection. For example, qNor does not

translocate H^+, while sNor, eNor, and gNor are predicted to conserve energy through H^+

translocation (32). Thus, the modularity of dissimilatory N reduction processes, and whether

those modules conserve energy or not, may impose certain constraints on pathway selection in

different organisms and should be further investigated as these considerations add significantly

to the biogeochemical tapestry of N-cycle dynamics and pathway evolution.

Under our high resource conditions (Fig. 2; left panels), at C:NO_3^- ratios ≥ 1, we observed that

denitrification prevailed and these cultures had lower growth rates than the predominantly

ammonium producing cultures. These high resource circumstances resulted in the production of

toxic intermediates (i.e., NO_2^- and possibly NO, albeit at undetectable levels), which may explain

why these cultures had lower growth rates (Fig. 2; left panels) and quasi-steady state growth

curves in our high resource metabolite profile (Fig. 4A). Rowley and colleagues (33) reported

that at least 20% of the N_2O released during high resource conditions were produced by

competition between nitrite and nitrate in the active-site of NarG. Under excess C

concentrations, NarG produces intracellular NO from NO_2^- and these intermediates are likely

inhibitory to cell growth, which may explain why our growth curves (Fig. 4A) reached a quasi-

steady state before nitrate had been fully utilized (as compared to the low resource metabolite

profile, Fig. S2). Furthermore, resources were not limiting growth under these conditions.

Rather, the cells were likely experiencing toxicity from NO and NO_2^- and thus the metabolic

outcomes would be beyond the scope of the LM and MPP. Nonetheless, these results clearly

demonstrate that end-product formation from the two resource concentrations tested, with the

same C:NO_3^- ratios, are not identical thereby refuting the C:NO_3^- control hypothesis.

We selected a single treatment (8 mM lactate, 12 mM nitrate; C:NO$_3^-$ ratio = 2), in which we

observed both denitrification and respiratory ammonification occurring simultaneously, for

RNA-Seq in order to gain insight into the transcriptional organization of actively growing *I.*

calvum cells. Strangely, we saw a decrease in transcript abundance encoding for two enzymes

known to convert lactate to pyruvate, LDH and LUP. While normalized read counts (Fig. S5)

were generally consistent across growth phases, indicative of constitutive expression, further

research investigating the mode of anaerobic lactate oxidation in *I. calvum* would illuminate how

reducing equivalents are fed into its central metabolic pathway. For example, *S. loihica* PV-4 is

known to use lactate for both denitrification and respiratory ammonification, but only uses

acetate for denitrification (20). Nonetheless, our transcriptomic data suggests that pyruvate plays

a central role in providing reducing equivalents to the TCA cycle as Acetyl-CoA, as evidenced

by significant upregulation in the genes encoding for PDH and citrate synthase, as well as

apparent "leaking" via incomplete lactate oxidation through the release of acetate and formate.

Such leaking may be produced by a putative PFL, adding to the diversity of C utilization

pathways feeding the ETC, and thereby driving pathway selection for nitrite reduction. Our

transcriptomic results, coupled with a parallel metabolite profile (Fig. 4A & 4B), also suggest

that the dual-pathway is induced by the presence of nitrite, and is not constitutively expressed

like nitrate reductase, *narG*. Furthermore, it appears that the significant increase in transcript

abundance for the gene encoding the *bd* oxidase helps to protect the anaerobic-dependent

biochemical machinery against oxidative stress, thereby scavenging any residual oxygen during

anaerobic growth. Undoubtedly, more conditions must be tested to better understand the

transcriptional organization of this unique bacterium.

Our metabolite profiles for N oxyanion respiration and N$_2$O versus ammonium production show

conflicting patterns relative to previous studies. Yoon and colleagues (21) reported complete

reduction of nitrate, production of nitrite, and then rapid consumption of nitrite, with N$_2$O as the

main end-product, by *S. loihica* PV-4 (5 mM lactate, 1 mM nitrate; ratio=0.6). When Yoon and

colleagues (21) replaced nitrate with nitrite as the dominant electron acceptor (5 mM lactate, 1

mM nitrite, ratio=0.6), ammonification prevailed. Other research has shown the same response to

nitrite replacement and ammonification dominance using non-fermentable C-sources (i.e.,

acetate) in chemostat enrichments of *Geobacter lovleyi* (34). In our work, nitrite was never fully

depleted (Figs. 3A, S1, S2) and when nitrite was given as the only electron acceptor, the

bacterium predominantly used denitrification but without concurrent growth (Fig. S3). Similar to

our work, Kraft and colleagues (22) also reported denitrification dominance when nitrite was

supplied as the terminal acceptor in incubations of coastal, sandy tidal flat sediments. These

differences highlight an incomplete understanding for the molecular mechanisms underlying the

framework put forth by the LM and MPP.

A detailed look into the biochemistry of ETC complexes helps to shed light on the molecular

mechanisms modulating pathway bifurcation. For example, Yoon and colleagues (3)

demonstrated that elevated pH selects for ammonification in *S. loihica* PV-4. This phenotypic

response is due to a decrease in the midpoint potential of the Rieske protein at higher pH (35–

37). Thus, any hindrance of electron flow through the bc_1 complex would ultimately reduce the

activity of downstream processes and promote alternative respiratory pathways. Nitrogen and C

limitation have also been shown to influence flux distributions in redox sensitive proteins,

including those found in electron transport (38). A drop in the intracellular redox potential (redox

poise) of the cell due to resource limitation may decrease the midpoint potential of the Rieske

protein and reduce the activity of any downstream electron exit modules, such as NirK (39–41).

Thus, based on fundamental principles of protein redox chemistry and thermodynamics, it

becomes clear that denitrification versus ammonification are likely not modulated by an arbitrary

ratio of $C:NO_3^-$, but rather by thermodynamic constraints of the Q-cycle (7, 8). The phenotypic

response of higher rates of denitrification over ammonification at high C:N ratios in other

published studies (3, 4) may also be due to enrichment bias for organisms that utilize quinones

with higher midpoint potentials in their bioenergetic chains (Fig. 1). Bergdoll and colleagues (7)

suggested that comparisons of Rieske/cyt*b* complexes from organisms with high- and low-

potential quinones may help to reconcile the thermodynamic properties of Q-cycle function.

However, most of our understanding of denitrification bioenergetics is based on evolutionarily

recent UQ-based HP bioenergetic chains from Gram-negative *α-, β-, γ*-proteobacteria. Because *I.*

calvum uses a MK-based LP bioenergetic chain it may be possible that the differences in

pathway selection across treatments are unique to LP chains.

Piecing together the evolutionary history of the N-cycle using isotopic signatures for

geochemically available N module cofactors (i.e., Ni, Fe, and Mo) coupled to molecular

evolutionary analysis has revealed respiratory ammonification was likely a major component of

the Archean N-cycle (28). Abiotic nitrite formation and depletion of ammonia through

photodissociation (42) would have created selective pressures for a dissimilatory N pathway that

also produced assimilatory N. We demonstrate that NrfA proteins are significantly enriched in N

compared to NirK (i.e., no evolutionary constraints to cost minimize N in the *nrfA* gene product

(11)) (Fig. 1B) and that ammonium production (without accumulation in the medium) supports

growth in *I. calvum* (Fig. 1C). The Nrf module is also relatively simplistic in that it receives

electrons directly from the quinol pool and not the bc_1 complex used in denitrification. The early

exit of electrons from the ETC (i.e., before reaching the bc_1 complex) suggests that Nrf may have

originated prior to the bc_1 complex. Furthermore, the quinol oxidation site (Q_o) of cytochrome *b*

contains a PDWY motif, indicative of an ancestral LP respiratory chain found in many Gram-

positive organisms (43). However, there is still debate regarding the presence of a cytochrome *bc*

complex in the last universal common ancestor (LUCA) (43, 44). Lastly, the Nrf module is wired

to operate via a q-loop with formate dehydrogenase whose Mo-cofactors would have also been

bioavailable during the Archean, further supporting an early evolution.

In summary, we employ a new predictive framework that accounts for the biochemistry and

evolutionary history of N modules, ETC complexes, and pool quinones to suggest the

mechanisms by which these two pathways are regulated at the molecular level. With this

understanding, it may be possible to extend our framework to environmental microbial

populations and accelerate model development across different ecosystem scales (i.e., cross-scale

systems biology).

Materials and Methods

Culture Conditions

Sample collection: *I. calvum* was isolated from a groundwater well (GW247) collected on

2/18/2013 at Oak Ridge National Laboratory (Lat: 35.97990, Long: 84.27059) with a

groundwater temperature of 15.57 ºC, conductivity of 521.9 µS/cm, and pH of 7.71. *I. calvum*

was isolated and obtained from Dr. R. Chakraborty (Lawrence Berkeley National Laboratory).

We began our experimental process by first picking single colonies from LB agar plates and

transferring the clonal isolates to LB broth.

<u>Media preparation</u>: All subsequent cultures were grown at 30 ºC and shaken at 250 rpm. Nitrate

reducing minimal media was prepared with the following final concentrations: NaCl (0.6mM),

NH_4Cl (1.75mM) (for ammonium replete conditions but not used in NH4-deplete conditions),

MgCl2 (0.2mM), CaCl2 (0.04mM), KCl (0.1mM), K2HPO4 (0.01mM), NaHCO3- (0.3mM),

cysteine (1mM) as reducing agent, resazurin as redox indicator, and trace elements and trace

vitamin solutions as reported (20, 45). 1M sterile filtered (0.2µm) Concentrated stocks of 60%

w/w sodium DL-lactate solution (Sigma-Aldrich, St. Louis, MO, USA), sodium-nitrate and

sodium-nitrite (≥99%, Fisher Scientific, Pittsburg, PA, USA) were diluted into media prior to

autoclaving to achieve the desired $C:NO_3^-$ ratio. $C:NO_3^-$ ratio was calculated based on (3) where

the number of C atoms (n) in the e-donor is multiplied by the concentration of the e-donor,

divided by the number of N atoms in the e-acceptor multiplied by the concentration of the e-

acceptor (Table S1). Media preparation was conducted in a 2L Widdel Flask. After autoclaving,

the media was immediately put under an anoxic headspace (N2/CO2 80:20 mix) and sterile

filtered (0.2µm) trace elements, trace vitamins, and reducing agent were added. The media was

cooled under an anoxic headspace and buffered with bicarbonate to maintain a pH of 7.2.

Hungate technique was used to dispense media into culture tubes (20 mL) and serum vials (100

mL) pre-flushed with a sterile stream of ultra-high purity (UHP) N_2 and sealed with blue 1" butyl

rubber stoppers. End-point cultures were grown in Balch tubes (18x150-mm glass tube) sealed

with butyl rubber stoppers. Cultures for time-course sampling were grown in 160ml serum vials.

All end-point experiments were terminated after 100 hours unless otherwise noted. Mean pH for

all culture vessels (time series and end-point; Table S5), measured at the end of each experiment,

was 7.3±0.05 (n=144)

Analytical procedures

<u>Growth Curve/Cell counts/Yield Measurements</u>: Growth curves were measured from scratch-

free Balch-tubes grown cultures using an automated optical density reader at OD_{600} nm

(Lumenautix LLC, Reno, NV). End-point cultures were monitored until all replicates reached

stationary phase (65-100 hours depending on $C:NO_3^-$ treatment) (Fig. S7).

Cell counts were performed by fixing cells in 4% paraformaldehyde (final concentration) for 20

minutes, filtered onto 0.2µm pore-sized black polycarbonate filters, and washed three times with

phosphate buffered saline (PBS, pH 7.2). Filtered cells captured on the black polycarbonate

filters were stained with SYBR® Gold nucleic acid stain (10-minute incubation) (ThermoFisher

Scientific) and counted manually with a fluorescence microscope (Olympus BX60, Tokyo,

Japan). We collected cells from during lag phase, exponential phase, and stationary phase in

order to create a standard curve of cell counts versus optical density (OD_{600}). We fit a linear

model to cell count versus OD_{600} (R^2=0.99) and used the resulting linear equation for cell count

enumeration for growth curves during our various treatment conditions.

Biomass concentrations were measured by filtration and drying as per standard protocol (46) for

8mM lactate/12mM nitrate and 0.8mM lactate/1.2mM nitrate treatments and conducted in

parallel with growth curve/cell counts as described above. Analysis from triplicate cultures

yielded (0.064 ± 0.003) and (0.016 ± 0.001) mg of biomass (dry weight) ml^{-1} for 8mM and 0.8

mM lactate cultures, respectively. Cell counts from stationary phase cultures were (1.5 ± 0.05) x

10^7 and (1.16 ± 0.09) x 10^6 for 8mM and 0.8 mM lactate cultures, respectively. From these

values the dry weight of a single *I. calvum* cell was estimated to be 1.09 x 10^{-10} g. Growth yield

(*Y*) (Table S4) was calculated by dividing biomass (g) by lactate mass (g) and moles consumed,

as described by (47). Lactate measurements are described below.

<u>Ion and Gas Chromatography Measurements</u>: A dual channel Dionex ICS-5000+ (Thermo

Scientific) was used to measure all anions (AS11-HC) and cations (CS-16) from the bacterial

growth media. New glass IC vials were used for every sample in order to ensure no cross

contamination of analytes. Headspace gas from Balch tubes and serum vials was sampled with

volume appropriate gastight syringes (Hamilton Company, Reno, NV) pre-flushed with UHP N_2.

For high and low resource treatments, 10μl and 100μl of headspace were sampled and diluted

into 12ml exetainers (Labco, Lampter, Wales, UK) over-pressurized with 15ml UHP N_2,

respectively. Similar dilutions were performed for nitrite as e-acceptor experiments, ammonium-

deplete experiments, and time-series experiments. For time-series experiments, an equal volume

of headspace gas that was removed at each time-point was replaced with sterile UHP N_2. N_2O

was measured by gas chromatography (Shimadzu Greenhouse Gas Analyzer GC-2014) using a

500μl injection volume. The rubber septa on the injection port of the GC was replaced after 100

injections in order to prevent leakage of the sample after the injection needle was lifted out from

the injection port. Aqueous concentrations of N_2O were calculated using a Henry's constant of

529 1.751 (mM (g)/mM (aq)) corrected for the medium's ionic strength and temperature. A total of

530 8-11 replicates per treatment were analyzed for all experiments discussed in this work.

531 <u>Ammonium production calculations</u>:

532 All nitrogen needed for assimilation must be first converted into ammonium (47). This fact

533 makes ammonium production measurements via respiratory nitrite ammonification difficult since

534 the cells simultaneous consume ammonium for growth. Therefore, direct measurements of

535 ammonium from IC or spectrophotometric analyses would not yield accurate production values.

536 Because *I. calvum* simultaneously consumes ammonium as its sole N source for growth and

537 produces ammonium via respiratory nitrite ammonification, we first measured ammonium

538 consumption with O_2 and lactate by calculating the difference between starting and ending

539 ammonium concentrations. These ammonium consumption values were then normalized to

540 lactate consumed (0.31μmols NH_4^+/lactate) and to ammonium consumed per cell, which was

541 calculated from average cell number of stationary phase biomass (Fig. S5) (7.07×10^{-7} μmols

542 NH_4^+/cell). These two ammonium consumption constants were used as two separate methods to

543 compare ammonium production based on actual starting and ending ammonium/lactate

544 measurements and cell number during nitrate reducing minimal media experiments.

545 For ammonium production normalized to lactate consumption, calculations are based on a mass

546 balance approach from (48) for Total Belowground Carbon Allocation (TBCA) but adapted for

547 nitrogen flux instead of carbon flux:

548   Net $NH_4^+{}_{(APL)}$ = (Δlactate$_{start-end}$ x 0.31μmols NH_4^+/lactate) + $\Delta NH_4^+{}_{end-start}$

549 Here, the Δlactate$_{start-end}$ variable (μmols) is multiplied by the ammonium consumed per lactate

550 consumed constant (APL). This value is added to the $\Delta NH_4^+{}_{end-start}$ variable (μmols), denoted as

551 ending minus starting concentration, which defines whether the change in ammonium is positive

552 (more ammonium produced than consumed) or negative (more ammonium consumed than

553 produced). The ammonium production method using cell count (APC) is calculated as reported

554 by (20), where :

555   Net $NH_4^+{}_{(APC)}$ = (Cell Yield x 7.07×10^{-7} μmols NH_4^+/cell) + $\Delta NH_4^+{}_{end-start}$

Where Cell Yield is the number of cells at stationary phase (Fig. S7). Comparison of these two

ammonia production methods revealed that they are in agreement with each other for anaerobic

nitrate reducing media experiments (Fig. S8). However, because the cell counts under low

resource treatments were at the lower limit of our standard curve (OD versus cell count), we

observed slightly more variability using the APC method. Lactate measurements by IC gave us

higher sensitivity over the large range of treatments tested, thus we report the APL method in the

main text as the preferred method for reporting ammonium production by respiratory

ammonification.

Nucleic Acid Extraction and Sequencing

DNA for genome sequencing was extracted with phenol-chloroform. The DNA quality and

integrity was verified on Agilent 2200 TapeStation and DNA was sequenced on the PacBio

platform (Duke Center for Genomic and Computational Biology).

For gene expression analysis, *I. calvum* cells were grown in 300 mL batch cultures under 8mM

and 0.8mM lactate (12mM and 1.2mM nitrate, respectively) (n=3). Growth curves were

monitored in real-time to collect cells during early-exponential, late-exponential, and stationary

phase. Upon sampling, culturing vessels were immediately incubated on ice, aseptically

transferred to 50 mL falcon tubes and immediately centrifuged at 15000 G for 5 minutes at 4 ºC.

Supernatant was immediately poured off and the cell pellets were harvested and stored in TRIzol

Reagent (Ambion RNA by Life Technologies) at -80 ºC until further processing. RNA was

extracted with TRIzol Reagent (Ambion RNA by Life Technologies), following the

manufacturer's protocol. The extraction was scaled up to 20 ml of TRIzol and performed in 50

ml conical tubes. RNA was resuspended in water, treated with Baseline-ZERO DNase

(Epicentre) and cleaned up with RNeasy MinElute Cleanup Kit (QIAGEN). Total extracted RNA

from each growth phase was quantified (RiboGreen RNA quantification), analyzed for RNA

integrity number (RIN) (RIN≥0.8 were used for sequencing), treated with RiboZero rRNA

removal kit (Illumina, San Diego, CA). Library preparation was performed with the TruSeq

Stranded mRNA Library Prep kit (Illumina, San Diego, CA) and sequenced using Illumina

NextSeq 500 2x150 Mid Output v2 platform, resulting in 10-15 million reads per sample.

Phylogenetic, Genomic, and Transcriptomic Analysis

Genomic DNA was assembled using Canu with an estimated genome size of 5 million base pairs

(49). The resulting single contiguous fragment was aligned to the *I. calvum* 7KIP genome (Acc:

NC_014830.1) to compare sequence similarity in Mauve (50, 51). Genome annotation for C5

was performed by the DOE Joint Genome Institute (JGI) using the Isolate Genome Gene Calling

method (Prodigal V2.6.3 February, 2016) under the submission ID 172966. The complete

genome sequence is deposited into NCBI BioProject SUB4141755.

A set of 34 NrfA amino acid sequences, representing 33 complete genome sequences and 1

octaheme nitrite reductase (ONR) from known respiratory ammonification organisms were

downloaded from GenBank (Table S1). A multiple sequence alignment (MSA) was generated

from the sequences annotated as cytochrome c nitrite reductase and ONR using MUSCLE (52).

The resulting alignment was visualized within MEGA5 (53) where the alignment was manually

screened for the presence of conserved amino acid residues consistent with those found in NrfA

(i.e., heme motifs). A maximum likelihood tree was created from the alignment using RAxML

(54) with 500 bootstrap iterations. The presence of NapA, NarG, NirK, and Nor modules were

manually queried from each NCBI genome in our set and confirmed by MSA, as described

above. Metabolic pathway for pool quinone type was queried on BioCyc Pathway/Genome

Database (biocyc.org) for each organism in our set. The structure of *I. calvum*'s NirK protein

was predicted using the protein structure predicting algorithm Phyre2 (55). Protein atomic

composition for C and N was calculated from amino acid sequences as input files, as described

by (10, 11), using custom python scripts for each element separately

(github.com/dvuono/Cost_minimization).

For transcriptomic analysis, the resulting raw reads were inspected using FastQC (56) to

determine quality, read length, and ambiguous read percentage. Reads were trimmed based on

quality score with a sliding window of 5 base pairs, quality cutoff of 28, trailing cutoff quality

score of 10, as well as adapter contamination removal in Trimmomatic (57). Due to the high

similarity of C5 to 7KIP, reads were aligned to the *Intrasporangium calvum* genomic reference

sequence and gtf file (Acc: NC_014830.1) using the STAR RNA-seq aligner (58), with the --

limitBAMsortRam parameter set to the recommended value by STAR. Sequence reads were

mapped to genomic features to obtain count data using featureCounts (59). Systematic changes

across experimental conditions were performed on normalized read counts in DESeq2 (60). The

RNA-seq data reported in this study have been deposited in NCBI BioProject SUB4141755.

Statistical analyses were conducted in the R environment for statistical computing (r-

project.org). Data that was tested using parametric statistical analysis were first validated for

normality by visualizing the data as a histogram and testing via Shapiro-Wilks test for normality.

ACKNOWLEDGMENTS. We thank F. von Netzer, K. Hunt, S Morales, N. Stopnisek, K. Meinhardt, W. Qin, N

Elliot, T. Hazen, H. Carlson, B. Ramsey, A. Murray, Z. Harrold, T. Morgan, P. Longley, and R. Hallnan. This

research was supported by a grant from the Nevada Governor's office of Economic Development (JG), by the

Desert Research Institute (DRI) postdoctoral research fellowship program, and in part by Ecosystems and Networks

Integrated with Genes and Molecular Assemblies (ENIGMA) (http://enigma.lbl.gov)—a Scientific Focus Area

Program at Lawrence Berkeley National Laboratory under contract number DE-AC02-05CH11231 and funded in

part by Oak Ridge National Laboratory under contract DE-AC05-00OR22725, and is based upon work supported by

the U.S. Department of Energy, Office of Science, Office of Biological & Environmental Research.

1. Galloway JN, et al. (2003) The Nitrogen Cascade. *Bioscience* 53(4):341.

2. Tiedje JM, Sexstone AJ, Myrold DD, Robinson JA (1982) Denitrification: ecological niches , competition
and survival. *Antonie Van Leeuwenhoek* 48(10598):569–583.

3. Yoon S, Cruz-García C, Sanford R, Ritalahti KM, Löffler FE (2015) Denitrification versus respiratory
ammonification: Environmental controls of two competing dissimilatory NO 3-/NO 2-reduction pathways in
Shewanella loihica strain PV-4. *ISME J* 9(5):1093–1104.

4. van den Berg EM, van Dongen U, Abbas B, van Loosdrecht MC (2015) Enrichment of DNRA bacteria in a
continuous culture. *ISME J* 9(10):2153–2161.

5. Moparthi VK, Hägerhäll C (2011) The evolution of respiratory chain complex i from a smaller last common
ancestor consisting of 11 protein subunits. *J Mol Evol* 72(5–6):484–497.

6. Dibrova D V., Cherepanov DA, Galperin MY, Skulachev VP, Mulkidjanian AY (2013) Evolution of
cytochrome bc complexes: From membrane-anchored dehydrogenases of ancient bacteria to triggers of
apoptosis in vertebrates. *Biochim Biophys Acta - Bioenerg* 1827(11–12):1407–1427.

7. Bergdoll L, Ten Brink F, Nitschke W, Picot D, Baymann F (2016) From low- to high-potential bioenergetic
chains: Thermodynamic constraints of Q-cycle function. *Biochim Biophys Acta* 1857(9):1569–1579.

8. Schoepp-Cothenet B, et al. (2009) Menaquinone as pool quinone in a purple bacterium. *Proc Natl Acad Sci*
106(21):8549–54.

9. Soo RM, Hemp J, Parks DH, Fischer WW, Hugenholtz P (2017) On the origins of oxygenic photosynthesis
and aerobic respiration in Cyanobacteria. *Science (80-)* 355(6332):1436–1440.

10. Baudouin-Cornu P, Schuerer K, Marlière P, Thomas D (2004) Intimate evolution of proteins: Proteome
atomic content correlates with genome base composition. *J Biol Chem* 279(7):5421–5428.

11. Grzymski JJ, Dussaq AM (2012) The significance of nitrogen cost minimization in proteomes of marine
microorganisms. *ISME J* 6(1):71–80.

12. Einsle O, Messerschmidt A, Huber R, Kroneck PMH, Neese F (2002) Mechanism of the six-electron
reduction of nitrite to ammonia by cytochrome c nitrite reductase. *J Am Chem Soc* 124(39):11737–11745.

13. Kraft B, Strous M, Tegetmeyer HE (2011) Microbial nitrate respiration - Genes, enzymes and environmental
distribution. *J Biotechnol* 155(1):104–117.

14. Godfrey L V., Falkowski PG (2009) The cycling and redox state of nitrogen in the Archaean ocean. *Nat
Geosci* 2(10):725–729.

15. Nitschke W, Dracheva S (1995) Reaction center associated cytochromes in Anoxygenic Photosynthetic
Bactera. eds Blankenship R, Madigan M, Bauer C (Kluwer Academic, Dordrecht, The Netherlands), pp
775–805.

16. Lalucat J, Bennasar A, Bosch R, Garcia-Valdes E, Palleroni NJ (2006) Biology of Pseudomonas stutzeri.
Microbiol Mol Biol Rev 70(2):510–547.

17. Suharti, de Vries S (2005) Membrane-bound denitrification in the Gram-positive bacterium Bacillus
azotoformans. *Biochem Soc Trans* 33(Pt 1):130–133.

18. Heylen K, Keltjens J, Stein LY (2012) Redundancy and modularity in membrane-associated dissimilatory
nitrate reduction in Bacillus. *Front Microbiol* 3(371):1–27.

19. Decleyre H, Heylen K, Bjorn T, Willems A (2016) Highly diverse nirK genes comprise two major clades
that harbour ammonium-producing denitrifiers. *BMC Genomics* 17(1):1–13.

20. Yoon S, Sanford RA, Löffler FE (2013) Shewanella spp. Use acetate as an electron donor for denitrification
but not ferric iron or fumarate reduction. *Appl Environ Microbiol* 79(8):2818–2822.

21. Yoon S, Sanford RA, Löffler FE (2015) Nitrite control over dissimilatory nitrate/nitrite reduction pathways
in Shewanella loihica strain PV-4. *Appl Environ Microbiol* 81(10):3510–3517.

22. Kraft B, et al. (2014) The environmental controls that govern the end product of bacterial nitrate respiration.
Science (80-) 345(6197):676–679.

23. Schmidt CS, Richardson DJ, Baggs EM (2011) Constraining the conditions conducive to dissimilatory
nitrate reduction to ammonium in temperate arable soils. *Soil Biol Biochem* 43(7):1607–1611.

24. Hardison AK, Algar CK, Giblin AE, Rich JJ (2015) Influence of organic carbon and nitrate loading on
partitioning between dissimilatory nitrate reduction to ammonium (DNRA) and N_2 production.
Geochim Cosmochim Acta 164:146–160.

25. Strohm TO, Griffin B, Zumft WG, Schink B (2007) Growth yields in bacterial denitrification and nitrate
ammonification. *Appl Environ Microbiol* 73(5):1420–1424.

26. Lotka AJ (1922) Contribution to the Energetics of Evolution. *Proc Natl Acad Sci* 8(6):147–151.

27. Kalakoutskii LV, Kirillova IP, Krassilnikov NA (1967) a New Genus of the. *J Gen Microbiol* 48:373-79–85.

28. Klotz MG, Stein LY (2008) Nitrifier genomics and evolution of the nitrogen cycle. *FEMS Microbiol Lett*
278:146–156.

29. Abbe KS, Takahashi S, Yamada T (1982) Invovlement of oxygen-sensitive pyruvate-formate lysase in
mixed-acid fermetnation by Streptococcus mutans under strictly anaerobic conditions. *J Bacteriol*
152(1):175.

30. Das A, Silaghi-Dumitrescu R, Ljungdahl LG, Kurtz DM (2005) Cytochrome *bd* oxidase, oxidative stress,
and dioxygen tolerance of the strictly anaerobic bacterium *Moorella thermoacetica*. *J Bacteriol*
187(6):2020–2029.

31. Zumft WG (1997) Cell biology and molecular basis of denitrification. *Microbiol Mol Biol Rev* 61(4):533–
616.

32. Hemp J, Gennis RB (2008) Diversity of the Heme-Copper Superfamily in Archaea: Insights from Genomics
and Structural Modeling. *Results and Problems in Cell Differentiation*, eds Schäfer G, Penefsky HS
(Springer Science+Business Media), pp 1–28.

33. Rowley G, et al. (2012) Resolving the contributions of the membrane-bound and periplasmic nitrate
reductase systems to nitric oxide and nitrous oxide production in Salmonella enterica serovar Typhimurium.
Biochem J 762:755–762.

34. van den Berg EM, Rombouts JL, Kuenen JG, Kleerebezem R, van Loosdrecht MCM (2017) Role of nitrite
in the competition between denitrification and DNRA in a chemostat enrichment culture. *AMB Express*
7(91):1–7.

35. Link TA, Hagen WR, Pierik AJ, Assmann C, von Jagow G (1992) Determination of the redox properties of
the Rieske cluster of bovine heart bc1 complex by direct electrochemistry of a water- soluble fragment.
EurJBiochem 208:685–691.

36. Ugulava NB, Crofts AR (1998) CD-monitored redox titration of the Rieske Fe-S protein of Rhodobacter
sphaeroides: pH dependence of the midpoint potential in isolated bc1 complex and in membranes. *FEBS Lett*
440(3):409–413.

37. Zu Y, et al. (2003) Reduction Potentials of Rieske Clusters: Importance of the Coupling between Oxidation
State and Histidine Protonation State. *Biochemistry* 42:12400–12408.

38. Ansong C, et al. (2014) Protein redox dynamics during light-to-dark transitions in cyanobacteria and
impacts due to resource limitation. *Front Microbiol* 5(July):1–10.

39. Snyder CH, Merbitz-Zahradnik T, Link TA, Trumpower BL (1999) Role of the Rieske Iron – Sulfur Protein
Midpoint Potential in the Protonmotive Q-Cycle Mechanism of the Cytochrome bc 1 Complex. *J Bioenerg*
Biomembr 31(3):235–242.

40. Denke E, et al. (1998) Alteration of the midpoint potential and catalytic activity of the Rieske iron-sulfur
protein by changes of amino acids forming hydrogen bonds to the iron-sulfur cluster. *J Biol Chem*
273(15):9085–9093.

41. Hunte C, Solmaz S, Palsdóttir H, Wenz T (2008) A structural perspective on mechanism and function of the
cytochrome bc1 complex. *Results and Problems in Cell Differentiation*, eds Schäfer G, Penefsky HS
(Springer Science+Business Media), pp 253–270.

42. Falkowski PG (1997) Evolution of the nitrogen cycle and its influence on the biological sequestration of
CO2 in the ocean. *Nature* 387(6630):272–275.

43. Kao WC, Hunte C (2014) The molecular evolution of the Qo Motif. *Genome Biol Evol* 6(7):1894–1910.

44. Dibrova D V., Cherepanov DA, Galperin MY, Skulachev VP, Mulkidjanian AY (2013) Evolution of
cytochrome bc complexes: From membrane-anchored dehydrogenases of ancient bacteria to triggers of
apoptosis in vertebrates. *Biochim Biophys Acta - Bioenerg* 1827(11–12):1407–1427.

45. Hillesland KL, et al. (2014) Erosion of functional independence early in the evolution of a microbial
mutualism. *Proc Natl Acad Sci* 111(41):14822–14827.

46. APHA (2012) *Standard Methods for the Examination of Water and Wastewater* (American Public Health
Association: Washington, DC). 23rd Ed. doi:ISBN 9780875532356.

47. White D (2000) *The physiology and biochemistry of prokaryotes* (Oxford University Press). 3rd ed.

48.   Giardina CP, Ryan MG (2002) Total belowground carbon allocation in a fast-growing Eucalyptus plantation
733   estimated using a carbon balance approach. *Ecosystems* 5(5):487–499.

734   49.   Koren S, et al. (2017) Canu : scalable and accurate long- - - read assembly via adaptive k - - - mer weighting
735   and repeat separation. *Genome Res*:1–35.

736   50.   Darling ACE, Mau B, Blattner FR, Perna NT (2004) Mauve : Multiple Alignment of Conserved Genomic
737   Sequence With Rearrangements. *Genome Res* 14:1394–1403.

738   51.   Darling AE, Mau B, Perna NT (2010) Progressivemauve: Multiple genome alignment with gene gain, loss
739   and rearrangement. *PLoS One* 5(6):e11147.

740   52.   Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic
741   Acids Res* 32(5):1792–7.

742   53.   Tamura K, et al. (2011) MEGA5: Molecular evolutionary genetics analysis using maximum likelihood,
743   evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 28(10):2731–2739.

744   54.   Stamatakis A (2014) RAxML version 8: A tool for phylogenetic analysis and post-analysis of large
745   phylogenies. *Bioinformatics* 30(9):1312–1313.

746   55.   Kelley LA, Mezulis S, Yates C, Wass M, Sternberg M (2015) The Phyre2 web portal for protein modelling,
747   prediction, and analysis. *Nat Protoc* 10(6):845–858.

748   56.   Andrews S (2010) FastQC: A quality control tool for high throughput sequence data. *Available online at:
749   http:// www.bioinformatics.babraham.ac.uk/projects/fastqc*.

750   57.   Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: A flexible trimmer for Illumina sequence data.
751   *Bioinformatics* 30(15):2114–2120.

752   58.   Dobin A, et al. (2013) STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* 29(1):15–21.

753   59.   Liao Y, Smyth GK, Shi W (2014) FeatureCounts: An efficient general purpose program for assigning
754   sequence reads to genomic features. *Bioinformatics* 30(7):923–930.

755   60.   Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-seq data
756   with DESeq2. *Genome Biol* 15(12):1–21.

Figure 1. (A) Maximum likelihood phylogenetic tree of NrfA amino acid sequences from known respiratory ammonifiers and accompanying N-module composition for each organism. Pool quinone is also noted for dual-pathway nitrite reducers and model species. Colors of the main branches denote the 1st heme motif type: CxxCK and CxxCH. (B) Protein atomic composition for N and C normalized to protein length for NirK and NrfA nitrite reductases. (C) State-transition from ammonium-replete to ammonium-deplete for *I. calvum* C5 grown under 8mM lactate 12mM nitrate minimal media at 30 ºC. Metabolite profiles for ammonium-deplete are shown.

Figure 2. The effects of high resource (left; range of lactate concentrations with 12 mM NO_3^-) and low resource (right; range of lactate concentrations with 1.2 mM NO_3^-) concentrations with the same C:NO_3^- ratio on pathway selection in *I. calvum* C5. (A) Production of N_2O-N and net change of NH_4^+ over a 100-hour incubation period at 30 ºC. Each bar represents the average of 8-10 replicates per treatment (Table S5). (B) Fraction of dissimilated N by pathway. (C) Growth rates for each corresponding treatment. The x-axis label defines lactate concentration and C:NO_3^- ratio in parentheses.

Figure 3. (A) Time-series metabolite profiles for lactate, nitrate, and nitrite (top pane), production of dissimilated end-products as N_2O-N and net change in NH_4^+ ammonium production (middle pane), and corresponding growth curve of *I. calvum* cells grown under 8mM lactate 12mM nitrate (C:NO_3^- ratio = 2) (bottom pane). Sampling points during growth phases are marked for transcriptomic analysis. (B) Metabolic reconstruction of the ETC from *I. calvum* with transcriptional changes for genes participating in dual-pathway dissimilatory nitrite reduction. Log_2 fold changes in transcript abundance are shown for late exponential relative to early exponential growth phase (EE vs. LE), stationary phase relative to early exponential growth phase (EE vs. ST), and stationary phase relative to late exponential growth phase (LE vs. ST). Locus IDs for each gene product correspond to heat map subplots in the order shown (left-to-right for each growth phase and top-to-bottom for each locus ID specified). Higher transcript abundance is represented in red, lower transcript abundance in blue, and no change in transcript abundance in white. Significant changes in transcript abundance ($p < 0.01$) are marked as a red box. Value of log_2 fold change is specified within each subplot. The log_2 fold changes of 14 NADH dehydrogenase subunits (Intca_03465-03530) were averaged as transcriptional changes were all shifted in the same direction.