Resource limitation modulates the fate of dissimilated nitrogen in a dual-pathway

Actinobacterium

David C. Vuono[1,6], Robert W. Read[1], James Hemp[2], Benjamin W. Sullivan[3], John A. Arnone

III[1], Iva Neveux[1], Bob Blank[4], Carl Staub[5], Evan Loney[1], David Miceli[1], Mari Winkler[6], Romy

Chakraborty[7], David A. Stahl[6], Joseph J. Grzymski[1,*]

[1]Division of Earth and Ecosystem Sciences, Desert Research Institute, Reno, NV 89512, USA

[2]Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena,

CA, USA

[3]Department of Natural Resources and Environmental Science, University of Nevada, Reno, NV

89557, USA

[4]Agricultural Research Service, U.S. Department of Agriculture, Reno, NV, 89512, USA

[5]Agtron, Inc. 9395 Double R Blvd, Reno, NV 89521, USA

[6]Department of Civil and Environmental Engineering, University of Washington, Seattle, WA,

98195, USA

[7]Lawrence Berkeley National Laboratory, Berkeley, CA, 94729, USA

[*]Correspondence: Joseph J. Grzymski, Division of Earth and Ecosystem Sciences, Desert

Research Institute, Reno, 2215 Raggio Parkway, NV 89512, USA

Email: Joe.Grzymski@dri.edu

Abstract

Respiratory ammonification and denitrification are two evolutionarily unrelated dissimilatory

nitrogen (N) processes central to the global N cycle, the activity of which is thought to be

controlled by carbon (C) to nitrate (NO_3^-) ratio. Here we find that *Intrasporangium calvum* C5, a

novel menaquinone-based dual-pathway denitrifier/respiratory ammonifier, disproportionately

utilizes ammonification rather than denitrification when grown under carbon or nitrate limitation,

not $C:NO_3^-$ ratio. Instead, $C:NO_3^-$ ratio is a confounding variable for resource limitation. We find

that the protein atomic composition for denitrification modules (NirK) are significantly cost

minimized for C and N compared to ammonification modules (NrfA), indicating that resource

limitation is a major selective pressure imprinted in the architecture of these proteins. The

evolutionary precedent for these findings suggests ecological and biogeochemical importance as

evidenced by higher growth rates when *I. calvum* grows predominantly using its ammonification

pathway and by assimilating its end-product (ammonium) for growth under ammonium-deplete

conditions. Genomic analysis of *I. calvum* further reveals a versatile ecophysiology to cope with

nutrient stress and redox conditions. Metabolite and transcriptional profiles during growth

indicate that transcript abundances encoding for its nitrite reducing enzyme modules, NrfAH and

NirK, significantly increase in response to nitrite production. Mechanistically, our results suggest

that pathway selection is driven by intracellular redox potential (redox poise), which may be

lowered during resource limitation, thereby decreasing catalytic activity of upstream electron

transport steps needed for denitrification enzymes. Our work advances our understanding of the

biogeochemical flexibility of N-cycling organisms, pathway evolution, and ecological food-

webs.

Introduction

Globally, respiratory ammonification and denitrification are vital nitrogen (N) dissimilation

pathways that either retain reactive N to support net primary productivity or close the N-cycle

through the release of gaseous N, respectively [1]. The environmental controls on these two

pathways, particularly the ratio of electron-donor to electron-acceptor (e.g., $C:NO_3^-$) [2], have

gained attention [3–7] due to increased anthropogenic N inputs into the environment [8].

However, the effects of resource limitation on growth and pathway selection (i.e., allocation of C

and N to dissimilatory and assimilatory processes), which are often confounded by C:NO_3^- ratio,

have not been tested. Strong selective pressures from Earth's shifting biogeochemistry and

oxidation-state have driven evolutionary adaptions to microbial electron transport chains (ETC)

[9, 10], respiratory chain redox potentials [11–13], and protein atomic composition [14, 15], may

shed light on how these pathways are regulated in contemporary organisms. Here, by identifying

the biochemical and evolutionary differences between respiratory ammonification and

denitrification, we disentangle the ecological significance and molecular mechanisms of electron

transfer through either pathway in a dual pathway organism.

From a biochemical standpoint, the primary difference between respiratory ammonification and

denitrification is their respective source of reducing equivalents in the ETC: 1) heme-based

cytochrome c nitrite reductase used in respiratory ammonification receive electrons directly from

the quinone (Q) pool [16] while 2) copper and cd_1 nitrite reductases used in denitrification

receive electrons from a soluble electron carrier (e.g., cytochrome c) via the bc_1 complex [17].

From an evolutionary standpoint, we can place each N-module's origin to a putative time in

Earth history based on the metal co-factors that would have been bioavailable: heme-based

cytochromes in an ancient, more reduced, environment compared to the copper-containing nitrite

reductases in an oxidizing environment [18]. The bioenergetic chains of microorganisms also

underwent selective pressure to shift from low-potential (LP) to high-potential (HP) quinones in

response to Earth's oxygenation [11, 12]. Menaquinone (MK) is thought to be the ancestral type

of LP quinone [19]. Organisms that use ubiquinone (UQ) are thought to have evolved under high

O_2 tensions with $α$-, $β$-, $γ$-proteobacteria as the only bacterial clades to use UQ [12]. Surprisingly,

our understanding for the biochemistry of denitrification is based predominantly on HP UQ-

based systems [20], leaving a significant knowledge gap in the physiology and biochemistry of

LP MK-based denitrifiers and how they link electron transfer with energy capture under resource

limitation [21–23].

In order to resolve the mechanisms of C:NO_3^- control on pathway selection and better understand

branched respiratory chains in LP-based nitrate-reducing organisms, we undertook the

characterization of the novel Gram-positive Actinobacterium strain *Intrasporangium calvum* C5:

a dual-pathway nitrite reducer that uses MK as sole pool quinone. Here we show that over a

range of C:NO_3^- ratios, duplicated at two substrate concentrations, *I. calvum* disproportionately

utilizes its ammonia-forming pathway during C limitation (≤ 0.4mM lactate), when $C:NO_3^-$ ratios

are < 1 (an observation contrary to the current paradigm). Using a genome-guided approach

coupled to time-series transcriptomics and metabolite profiles, we identified differentially

expressed genes in the bacterium's ETC and central metabolic pathways. Using this information

to inform a metabolic reconstruction of the ETC and extensive literature on the biochemistry of

the bc_1 complex, we propose a new mechanism by which these two pathways are regulated at the

biochemical level.

Materials and Methods

Culture Conditions

<u>Media preparation</u>: All cultures were grown at 30 ºC and shaken at 250 rpm. Nitrate reducing

minimal media was prepared with the following final concentrations: NaCl (0.6mM), NH_4Cl

(1.75mM) (for ammonium replete conditions but not used in NH4-deplete conditions), MgCl2

(0.2mM), CaCl2 (0.04mM), KCl (0.1mM), K2HPO4 (0.01mM), NaHCO3- (0.3mM), cysteine

(1mM) as reducing agent, resazurin as redox indicator, and trace elements and trace vitamin

solutions as reported [24, 25]. 1M sterile filtered (0.2μm) Concentrated stocks of 60% w/w

sodium DL-lactate solution (Sigma-Aldrich, St. Louis, MO, USA), sodium-nitrate and sodium-

nitrite ($\geq$99%, Fisher Scientific, Pittsburg, PA, USA) were diluted into media prior to

autoclaving to achieve the desired $C:NO_3^-$ ratio. $C:NO_3^-$ ratio was calculated based on [3] where

the number of C atoms (n) in the e-donor is multiplied by the concentration of the e-donor,

divided by the number of N atoms in the e-acceptor multiplied by the concentration of the e-

acceptor (Table S4). See SI Materials and Methods for complete description of Hungate

technique prepared media. Mean pH for all culture vessels (time series and end-point; Table S5),

measured at the end of each experiment, was 7.3±0.05 (n=144).

Analytical procedures

<u>Growth Curve/Cell counts/Yield Measurements</u>: Growth curves were measured from scratch-

free Balch-tubes grown cultures using an automated optical density reader at OD_{600} nm

(Lumenautix LLC, Reno, NV). End-point cultures were monitored until all replicates reached

stationary phase (65-100 hours depending on $C:NO_3^-$ treatment) (Figure S6). Cell counts were

performed by fixing cells in 4% paraformaldehyde (final concentration) for 20 minutes, filtered

onto 0.2μm pore-sized black polycarbonate filters. A complete description is provided in SI

Materials and Methods. Biomass concentrations were measured by filtration and drying as per

standard protocol [26]. A complete description is provided in SI Materials and Methods.

<u>Ion and Gas Chromatography Measurements</u>: A dual channel Dionex ICS-5000+ (Thermo

Scientific) ion chromatograph (IC) was used to measure organic (lactate, acetate, and formate)

and inorganic (nitrite and nitrate) anions on an AS11-HC column and cations (ammonium) on a

CS-16 column from the bacterial growth media. A complete description is provided in SI

Materials and Methods.

Phylogenetic, Genomic, and Transcriptomic Analysis

Genomic DNA was assembled using Canu (version 1.7.1) with an estimated genome size of 5

million base pairs [27]. The resulting single contiguous fragment was aligned to the *I.* calvum

7KIP genome (Acc: NC_014830.1) to compare sequence similarity in Mauve[28, 29]. Genome

annotation for C5 was performed through the NCBI Prokaryotic Genome pipeline

(www.ncbi.nlm.nih.gov/genome/annotation_prok/). Additional gene prediction analysis and

functional annotation was performed by the DOE Joint Genome Institute (JGI) using the Isolate

Genome Gene Calling method (Prodigal V2.6.3 February, 2016) under the submission ID

172966. The complete genome sequence and annotation is available in the NCBI database under

the BioProject number PRJNA475609. A complete description of the phylogenetic, pathway

analysis, and cost-minimization calculations is provided in SI Material and Methods. For

transcriptomic analysis, the resulting raw reads were inspected using FastQC [30] to determine

quality, read length, and ambiguous read percentage. Reads were trimmed based on quality score

with a sliding window of 5 base pairs, quality cutoff of 28, trailing cutoff quality score of 10, as

well as adapter contamination removal in Trimmomatic [31]. A complete description is provided

in SI Materials and Methods. Statistical analyses were conducted in the R environment for

statistical computing (r-project.org). Data that was tested using parametric statistical analysis

were first validated for normality by visualizing the data as a histogram and testing via Shapiro-

Wilks test for normality.

Results

Genomic analysis of *I. calvum* C5. We sequenced and analyzed the genome of *I. calvum* C5 to

first compare its similarity to the type species *I. calvum* 7KIP. We identified a high degree of

sequence similarity to 7KIP based on three homologous sequence regions as locally collinear

blocks (SI Results). Genome size of C5 was 4,025,044 base pairs (bp), only 662 bp longer than

7KIP. Genomic analysis of the ETC revealed the typical suite of complexes common to

facultative aerobes, including primary dehydrogenases (*nuo* complex, succinate dehydrogenase),

alternative *NDH-2* NADH dehydrogenase, cytochrome bc_1 complex, high-oxygen adapted

cytochrome c oxidase (A-family), and low-oxygen adapted cytochrome *bd* oxidase. The bc_1

complex subunits are also located immediately upstream of cytochrome c oxidase, suggesting

that these enzymes are encoded in a single operon creating a supercomplex. Despite *I. calvum*'s

seeming propensity for aerobic growth on a number of growth media [32], its bioenergetic

system uses MK as its sole pool quinone. *I. calvum* also possesses multiple pathways for

supplying electrons into the MK-pool, such as formate, malate, hydroxybutyrate, and

glycerophosphate dehydrogenases. Once in the MK-pool, there are alternative pathways for

MKH_2 oxidation that can circumvent the bc_1 complex, such as a membrane-bound respiratory

nitrate reductase module (NarG). In addition, to NarG, its dissimilatory N module composition

consists of a truncated denitrification pathway (N_2O is a terminal product) using a copper nitrite

reductase NirK and quinol-dependent nitric oxide reductase qNor. *I. calvum* also possesses both

catalytic and membrane anchor subunits (NrfA and NrfH, respectively) for a pentaheme

cytochrome c module involved in respiratory nitrite ammonification.

***I. calvum* encodes for a functional NrfAH complex and assimilates NH_4^+ via respiratory

nitrite ammonification.** To gain insight into possible function of the NrfAH complex, we

aligned the NrfA protein sequences from C5 and 7KIP to a collection of 33 recognized

cytochrome c nitrite reductases from published annotated genomes (Table S1). This confirmed

that NrfA from *I. calvum* is a member of the CxxCH 1^{st} heme motif group (Figure 1A), which

forms one of four clades on the NrfA phylogenetic tree. We then queried the genomes of the taxa

in our phylogeny for other annotated N-reducing modules used in nitrate reduction, nitrite

reduction, NO-forming nitrite reduction, and primary pool quinone. Among the three major

clades of NrfA, at least 5 additional taxa are noted having dissimilatory N-module inventories

containing dual respiratory pathways: *S. thermophilum, B. azotoformans, B. bataviensis*, *B.

bacteriovorus*, and *Candidatus* N. inopinata, (Figure 1A). None of the taxa in our NrfA

phylogeny harbored the cd_1 nitrite reductases (NirS). Due to the exclusive NirK representation in

dual-pathway membership, we asked whether there might be differences in protein atomic

composition between NirK and NrfA, given the disparate evolutionary origins of these modules

[33]. We collected 20 additional publicly available NirK protein sequences from nondual-

pathway denitrifiers (Table S1) and calculated the protein C and N composition for our

NirK/NrfA collection as atoms per residue side-chain (Figure 1B). These results showed a

significant depletion in C and N atoms per residue side-chain (ARSC) for NirK compared to

NrfA (C and N: $p<0.001$; t-test), indicating that resource constraints are imprinted on the

evolution of these proteins.

We next tested the functionality of *I. calvum*'s Nrf complex by growing the bacterium under

reducing conditions (8 mM lactate, 12 mM nitrate, ammonium-replete). We then performed a

state-transition where biomass from late-exponential growth phase was collected and

anaerobically inoculated into ammonia-deplete media (Figure 1C; SI Results). Despite no

detectable amounts of ammonium produced in the media over time, cell counts increased

$5.4 \times 10^5 \pm 8.9 \times 10^4$ cells/mL (0.126 ± 0.02 optical absorbance at OD_{600}) over a 48-hour incubation,

indicating consumption of ammonium produced by NrfA. Net ammonium production was

13 ± 2.7 µmoles with the remainder of dissimilated N being used by the denitrification pathway

(24 ± 4.2 µmoles N_2O-N), resulting in a recovery of 97.4% dissimilated N. These results

confirmed that *I. calvum* C5 has a functional Nrf complex and also consumes the product

(ammonium) of respiratory nitrite ammonification.

Respiratory nitrite ammonification exceeds denitrification under C-limitation. We

investigated $C:NO_3^-$ control on respiratory ammonification versus denitrification on cultures of *I.*

calvum C5 over a high resource $C:NO_3^-$ range (16-0.4 mM lactate, 12 mM nitrate; ratio 4-0.1)

and low resource $C:NO_3^-$ range (1.6-0.04 mM lactate, 1.2 mM nitrate; ratio 4-0.1). This

experimental design enabled us to evaluate $C:NO_3^-$ control over a broader range than previous

studies that only considered ratios ≥ 1.5 [3, 4, 34], while also testing the effects of resource

concentration on pathway selection. Under all the treatments tested, gas and ion chromatography

measurements showed products of both respiratory pathways, differing only in the relative

fraction of N_2O versus ammonium production across treatments (Figure 2). At high resource

concentrations, respiratory ammonification did not prevail at high $C:NO_3^-$ ratios (Figure 2A,

Figure 2B, left panels; Table S2). Instead, significantly greater amounts of N_2O were produced

over ammonium, though nitrite was still the major extracellular end-product of nitrate

respiration. Despite the predominance of N_2O production under the high resource concentrations,

ammonium production exceeded N_2O production only at the lowest $C:NO_3^-$ ratio (0.4 mM

lactate, ratio=0.1) (Figure 2) and accounted for 76.2±0.1% of dissimilated N.

Results from the low resource dataset provided weak support for the strict stoichiometry

hypothesis that $C:NO_3^-$ controls pathway selection. Ammonia exceeded N_2O production only

under one high $C:NO_3^-$ ratio treatment (ratio=4; 1.6 mM lactate; Figure 2A, Figure 2B, right

panels). However, at ratios ≤1 (≤0.4 mM lactate), significantly more ammonium than N_2O was

produced. On average, respiratory ammonification accounted for 78.1±8.9% of dissimilated N

for lactate concentrations ≤0.4 mM. When these results are taken in context with cell physiology,

we observed a significant and positive relationship between specific growth rate (μ) and the

fraction of N dissimilated by respiratory ammonification (R^2=0.5; p<0.001) (Figure 2C, S1;

Table S3).

Resource concentration influences the metabolite profiles of ammonium and N_2O

production. Given the co-occurrence of end products from both pathways during the end-point

experiments (Figure 2), we next investigated the timing of ammonium and N_2O production

relative to metabolite profiles for lactate, nitrate/nitrite, and growth phase at two resource

concentrations with the same ratio (8 mM and 0.8 mM lactate, ratio=2; Figure 1A, Figure S2,

Figure S3). Despite ample e-donor and e-acceptor available for growth, the high resource

cultures entered a quasi-stationary phase at ~50 hours, after which there was continued slow

growth (Figure 1A). Metabolite profiles showed that ammonium and N_2O production began

simultaneously, as soon as nitrite was produced from nitrate reduction. The low resource cultures

entered stationary phase at ~40 hours (Figure S2) after nitrate had been fully utilized. No further

cell growth was observed after stationary phase was reached. These results show that cell growth

occurred primarily on the reduction of nitrate, while nitrite reduction to ammonium and N_2O

occurred during a stationary growth phase, demonstrating that microbial activity is not always

correlated with growth. The metabolite profiles for ammonium and N_2O at low resources (Figure

S2) did not mirror those observed at high resources (Figure 2A). The rate of N_2O production

significantly decreased and ammonium production oscillated rather than steadily increase

through time. These differences in metabolite profiles, further demonstrate that concentration

influences the activities of pathway bifurcation. Repeated time series experiments that were

extended up to 300 hours show that nitrite is slowly depleted, but does not get fully consumed

(Figure S3). When cultures were given nitrite, instead of nitrate as a terminal electron acceptor (8

mM lactate, 12 mM nitrite; ratio=2), we observed no immediate growth (as was observed with

nitrate) but measured more N_2O than ammonium production (33.4±4.8 µmoles N_2O-N and

8.0±2.5 µmoles NH_4^+, respectively) (Figure S4), demonstrating respiratory ammonification does

not exceed denitrification when nitrite is supplied as the sole acceptor in *I. calvum*.

Nitrite-reducing modules are up-regulated during late exponential- and stationary-phase

growth. In order to gain insight into mechanisms of gene regulation and transcriptional

organization of *I. calvum*, we conducted RNA-Seq in parallel with the high resource time-series

metabolite profile (Figure 3A). This approach enabled us to compare genome-wide differential

expression based on $\log_2$ fold change (lfc) of RNA extracted from three growth phases: early

exponential (EE), late exponential (LE), and stationary (ST) (Figure 3B, Figure S5). Within the

central metabolic pathway beginning with the conversion of lactate to pyruvate, we observed a

moderate decrease in transcript abundance of L-lactate dehydrogenase (LDH) (Intca_16740)

between EE-LE and -ST (lfc = -1.6±0.7; -1.9±0.7), respectively. Lactate utilization protein C

(LUP) (Intca_04080), an enzyme involved in lactate degradation, also showed a moderate and

significant decrease in transcript abundance between EE-LE and -ST (lfc = -1.6±0.6; -2.4±0.6,

p=0.002), respectively. *I. calvum* encodes for two parallel metabolic pathways for pyruvate

conversion to acetyl-CoA: pyruvate dehydrogenase (PDH) (Intca_01255) and pyruvate

ferredoxin oxidoreductase (PFOR) (Intca_15510). For PDH, there was a significant and

moderate increase in transcript abundance between EE-LE and -ST (lfc = 2.1±0.6, *p*=0.002;

1.5±0.6), respectively. For PFOR, there was a minor decrease in transcript abundance between

EE-LE (lfc = -0.43±0.5), and then a moderate increase in transcript abundance between EE-ST

(1.1±0.5). Citrate synthase (Intca_04135), the enzyme catalyzing the conversion of acetyl-CoA

to citrate and the first step of the tricarboxylic acid (TCA) cycle, showed a highly significant

increase in transcript abundance between EE-LE and -ST (lfc = 4.3±0.5, *p*<0.001; 6.9±0.5,

p<0.001).

Within the ETC, there was moderate and significant decrease in transcript abundance for all
subunits from the primary dehydrogenase (*nuo* complex; Intca_03465-03539) between EE-LE
and -ST (lfc = -1.2±0.3; -2.4±0.6, *p*<0.001), respectively. Nitrate reductase subunits showed no
change in transcript abundance between EE-LE (lfc = 0.01±0.07) and moderately decreased in
abundance by ST (lfc = -1.2±0.1), which was corroborated by the depletion of nitrate during
stationary phase. There was a significant increase in transcript abundance of *nirK* (Intca_17170)
(lfc = 2.2±0.6, *p*=0.003; 2.4±0.6, *p*<0.001) and quinol dehydrogenase/membrane anchor subunit
nrfH (Intca_09465) (lfc = 2.5±0.6, *p*=0.001; 2.1±0.6, *p*=0.003) by EE-LT and EE-ST,
respectively, which coincided with nitrite production (Figure 3A). The catalytic subunit of the
cytochrome c nitrite reductase complex (*nrfA*) (Intca_09460) also increased moderately in
transcript abundance by EE-LT and EE-ST (lfc = 1.6±0.6; 1.0±0.6), respectively (Figure 3B).
Contrary to the transcript abundance patterns of *nirK* and *nrfAH*, nitric oxide reductase (qNor;
Intca_01525) transcripts moderately increased between EE-LT (lfc = 1.6±0.6) but decreased in
the successive time periods (lfc = 0.43±0.6 between EE-ST; lfc = -1.2±0.6 between LE-ST)
(Figure 3B).

There was a significant increase in transcript abundance of formate transporter *focA*
(Intca_17150) between EE-ST, as well as LE-ST (lfc = 4.9±0.7, *p*=0.002; 4.8±0.7, *p*=0.002;
respectively). We verified the production of formate in our ion chromatography measurements in
the range of 100-200μM following late exponential growth. We also observed a moderate
increase in transcript abundance of formate dehydrogenase (FDH) subunits (Intca_11150-
11160). These results implicate the activity of formate oxidation, which would contribute to a Δ*p*
in the periplasm via a Q-loop mechanism and the reduction of MK for electron transfer to nitrite
via cytochrome c nitrite reductase. Considering that formate was not provided in our media
recipe, an alternative pathway for formate production must exist in *I. calvum*. We also observed
acetate production in similar concentrations as formate (100-200μM). In *E. coli*, formate is
produced anaerobically from the action of pyruvate formate lyase (PFL). We identified a putative
PFL based on genome annotation (Intca_12230), where transcript abundance also significantly
increased by ST. PFL is also highly sensitive to oxygen [35], which was also in agreement with a
significant increase in transcript abundance between EE-ST and LE-ST (Figure 3B) of
cytochrome *bd* oxidase (Intca_01110 and Intca_01115), which is thought to protect anaerobic
enzymes against oxidative stress [36].

Discussion

We challenge the paradigm that $C:NO_3^-$ ratio controls pathway selection in a dual-pathway organism based on a simple principle: ratios do not account for the abundance of growth-limiting resources. We hypothesized that limitation in C or NO_3^- should better predict pathway selection in a dual-pathway denitrifier/respiratory ammonifier. To test this hypothesis, we systematically measured the response of the Gram-positive Actinobacterium *Intrasporangium calvum* C5 to the same range of $C:NO_3^-$ ratios at both high and low resource loadings to better resolve mechanisms of pathway selection. We demonstrated that resource concentration, not $C:NO_3^-$ ratio, influences pathway selection. We found stronger support for respiratory ammonification preference under C-limitation (at low $C:NO_3^-$ ratios), which also grew at significantly higher growth rates (Figure 2). These results suggest that the NrfA complex, which receives electrons directly from the MK-pool, is optimized to maximize power when one or more resources are limiting. The enrichment of C and N ARSC in publically available NrfA over NirK protein seuqences (Figure 1B) provides further support and evolutionary precedence to ammonification preference over denitrification under resource limitation. This is because the end-product of ammonification can be used as an assimilatory N-source (Figure 1C), indicating no evolutionary constraint to cost minimize N. These data, together with metabolic reconstructions from metabolite and transcriptional profiles (Figure 3), suggest that $C:NO_3^-$ ratio alone is insufficient to explain pathway selection.

The theoretical basis for pathway selection is explained by the law of the minimum (LM) and the maximum power principle (MPP), which state that growth is limited by the least abundant resource and that biological systems are designed to maximize power in order to effectively allocate energy to reproduction and survival [37, 38], respectively. Here, it appears these two theories are working together: when resources are limited, the cell utilizes the respiratory pathway for growth that is optimized to maximize power. Power, in this case, is realized as higher growth rates from the cultures exhibiting disproportionately higher ammonium production than N_2O production (Figure 2: high resources: $C:NO_3^-$ ratio = 0.1; low resources: $C:NO_3^-$ ratios = 4, 1, 0.5, 0.1). More specifically, the bacterium must generate a greater Δp in order to maximize power when starved for a growth limiting resource. This may help to further explain how respiratory ammonification, which is overall energetically less favorable than denitrification

(lactate with nitrite: $\Delta G° = -763.98$ versus $\Delta G° = -1196.93$, respectively), can have higher growth

yields [39] and growth rates (Figure 2, Figure S1) under C- and N-limitation due to the higher

energy yield on a per-nitrite basis (denitrification: -217 KJ per mole nitrite; respiratory

ammonification: -399 KJ per mole nitrite). For comparison, a total of 8 H^+ are translocated

during denitrification by *I. calvum* (not including nitrate reduction since both pathways share this

step) (Figure 3): NADH dehydrogenase translocates 4 H^+ per MKH_2 oxidized and the bc_1

complex translocates an additional 4 H^+ per MKH_2 oxidized. However, 2 H^+ must be consumed

in the periplasm to reduce nitrite to NO [40]. qNor has a net zero H^+ release (consumes 2 H^+ to

make N_2O but releases 2 H^+) without MKH_2 regeneration [41]. Thus, a net total of 6 H^+ are

translocated per nitrite reduced in denitrification with added biosynthetic costs of making the bc_1

complex and qNor. In respiratory ammonification, MK/MKH_2 redox pair is cycled between

NADH dehydrogenase and formate dehydrogenase. 6 electrons and 8 H^+ are needed to reduce

nitrite to ammonium, thus 3 MKH_2 are needed [16]. If MKH_2 is received from NADH

dehydrogenase, 12 H^+ are translocated plus 2 H^+ from FDH. As each MKH_2 is oxidized at the

binding site of NrfH, 2 H^+ are liberated [16], resulting in a net total of 12 H^+ translocated per

nitrite reduced for respiratory ammonification. This implies that the cell might deplete its NADH

pool more rapidly on a per nitrite basis. However, if more protons are pumped in the early stages

of growth, the cell would be allocating the ATP generated for anabolism, as evidenced by higher

growth rates in the cultures exhibiting higher amounts of respiratory ammonification (Figure 2),

which is supported by the MPP.

Under our high resource conditions (Figure 2; left panels), at $C:NO_3^-$ ratios ≥ 1, we observed that

denitrification prevailed and these cultures had lower growth rates than the predominantly

ammonium producing cultures. These high resource circumstances resulted in the production of

toxic intermediates (i.e., NO_2^- and possibly NO, albeit at undetectable levels), which may explain

why these cultures had lower growth rates (Figure 2; left panels) and quasi-steady state growth

curves in our high resource metabolite profile (Figure 3A). Rowley and colleagues [42] reported

that at least 20% of the N_2O released during high C conditions were produced by competition

between nitrite and nitrate in the active-site of NarG. Under excess C concentrations, NarG

produces intracellular NO from NO_2^- and these intermediates are likely inhibitory to cell growth,

which may explain why our growth curves (Figure 3A) reached a quasi-steady state before

nitrate had been fully utilized (as compared to the low resource metabolite profile, Figure S2).

Furthermore, resources were not limiting growth under these conditions. Rather, the cells were

likely experiencing toxicity from NO and NO_2^- and thus the metabolic outcomes would be

beyond the scope of the LM and MPP. Nonetheless, these results clearly demonstrate that end-

product formation from the two resource concentrations tested, with the same $C:NO_3^-$ ratios, are

not identical thereby refuting the $C:NO_3^-$ control hypothesis.

We selected a single treatment (8 mM lactate, 12 mM nitrate; $C:NO_3^-$ ratio = 2), in which we

observed both denitrification and respiratory ammonification occurring simultaneously, for

RNA-Seq in order to gain insight into the transcriptional organization of actively growing *I.*

calvum cells (Figure 3). Strangely, we saw a decrease in transcript abundance encoding for two

enzymes known to convert lactate to pyruvate, LDH and LUP. While normalized read counts

(Figure S5) were generally consistent across growth phases, indicative of constitutive expression,

further research investigating the mode of anaerobic lactate oxidation in *I. calvum* would

illuminate how reducing equivalents are fed into its central metabolic pathway. For example, *S.*

loihica PV-4 is known to use lactate for both denitrification and respiratory ammonification, but

only uses acetate for denitrification [24]. Nonetheless, our transcriptomic data suggests that

pyruvate plays a central role in providing reducing equivalents to the TCA cycle as Acetyl-CoA,

as evidenced by significant upregulation in the genes encoding for pyruvate dehydrogenase and

citrate synthase, as well as apparent "leaking" via incomplete lactate oxidation through the

release of acetate and formate. Such leaking may be produced by a putative PFL, adding to the

diversity of C utilization pathways feeding the ETC, and thereby driving pathway selection for

nitrite reduction. Our transcriptomic results, coupled with a parallel metabolite profile (Figure 3),

also suggest that the dual-pathway is induced by the presence of nitrite, and is not constitutively

expressed like nitrate reductase, *narG*. Furthermore, it appears that the significant increase in

transcript abundance for the gene encoding the *bd* oxidase helps to protect the anaerobic-

dependent biochemical machinery against oxidative stress, thereby scavenging any residual

oxygen during anaerobic growth.

Our metabolite profiles for N oxyanion respiration and N_2O versus ammonium production show

conflicting patterns relative to previous studies (Figure 3A, Figure S2). Yoon and colleagues [43]

reported complete reduction of nitrate, production of nitrite, and then rapid consumption of

nitrite, with N_2O as the main end-product, by *S. loihica* PV-4 (5 mM lactate, 1 mM nitrate;

ratio=0.6). When Yoon and colleagues [43] replaced nitrate with nitrite as the dominant electron

acceptor (5 mM lactate, 1 mM nitrite, ratio=0.6), ammonification prevailed. Other research has

shown the same response to nitrite replacement and ammonification dominance using non-

fermentable C-sources (i.e., acetate) in chemostat enrichments of *Geobacter lovleyi*[44]. In our

work, nitrite was never fully depleted (Figure 3A, Figure S2, Figure S3) and when nitrite was

given as the only electron acceptor, the bacterium predominantly used denitrification but without

concurrent growth (Figure S4). Similar to our work, Kraft and colleagues[34] also reported

denitrification dominance when nitrite was supplied as the terminal acceptor. These differences

highlight an incomplete understanding for the molecular mechanisms underlying the framework

put forth by the LM and MPP.

A detailed look into the biochemistry of ETC complexes helps to shed light on the molecular

mechanisms modulating pathway bifurcation. For example, Yoon and colleagues [3]

demonstrated that elevated pH selects for ammonification in *S. loihica* PV-4. This phenotypic

response is due to a decrease in the midpoint potential of the Rieske protein at higher pH [45–

48]. Thus, any hindrance of electron flow through the bc_1 complex would ultimately reduce the

activity of downstream processes and promote alternative respiratory pathways. Nitrogen and C

limitation have also been shown to influence flux distributions in redox sensitive proteins,

including those found in electron transport [49]. A drop in the intracellular redox potential (redox

poise) of the cell due to resource limitation may decrease the midpoint potential of the Rieske

protein and reduce the activity of any downstream electron exit modules, such as NirK [50–52].

Thus, based on fundamental principles of protein redox chemistry and thermodynamics, it

becomes clear that denitrification versus ammonification are likely not modulated by an arbitrary

ratio of $C:NO_3^-$, but rather by thermodynamic constraints of the Q-cycle [11, 12]. The phenotypic

response of higher rates of denitrification over ammonification at high $C:NO_3^-$ ratios in other

published studies [3, 4] may also be due to enrichment bias for organisms that utilize quinones

with higher midpoint potentials in their bioenergetic chains (Figure 1). Bergdoll and colleagues

[11] suggested that comparisons of Rieske/cyt*b* complexes from organisms with high- and low-

potential quinones may help to reconcile the thermodynamic properties of Q-cycle function.

However, most of our understanding of denitrification bioenergetics is based on evolutionarily

recent UQ-based HP bioenergetic chains from Gram-negative *α-, β-, γ*-proteobacteria. Because *I.*

calvum uses a MK-based LP bioenergetic chain it may be possible that the differences in

pathway selection across treatments are unique to LP chains.

Piecing together the evolutionary history of the N-cycle using isotopic signatures for

geochemically available N module cofactors (i.e., Ni, Fe, and Mo) coupled to molecular

evolutionary analysis has revealed respiratory ammonification was likely a major component of

the Archean N-cycle [33]. Abiotic nitrite formation and depletion of ammonia through

photodissociation [53] would have created selective pressures for a dissimilatory N pathway that

also produced assimilatory N. We demonstrate that NrfA proteins are significantly enriched in N

compared to NirK (i.e., no evolutionary constraints to cost minimize N in the *nrfA* gene product

[15]) (Figure 1B) and that ammonium production (without accumulation in the medium)

supports growth in *I. calvum* (Figure 1C). The Nrf module is also relatively simplistic in that it

receives electrons directly from the quinol pool and not the bc_1 complex used in denitrification.

The early exit of electrons from the ETC (i.e., before reaching the bc_1 complex) suggests that Nrf

may have originated prior to the bc_1 complex. Furthermore, the quinol oxidation site (Q_o) of

cytochrome *b* contains a PDWY motif, indicative of an ancestral LP respiratory chain found in

many Gram-positive organisms [54]. However, there is still debate regarding the presence of a

cytochrome *bc* complex in the last universal common ancestor [54, 55]. Lastly, the Nrf module is

wired to operate via a q-loop with formate dehydrogenase whose Mo-cofactors would have also

been bioavailable during the Archean, further supporting an early evolution.

In summary, we employ a new predictive framework that accounts for the biochemistry and

evolutionary history of N modules, ETC complexes, and pool quinones to suggest the

mechanisms by which these two pathways are regulated at the molecular level. With this

understanding, it may be possible to extend our framework to environmental microbial

populations and accelerate model development across different ecosystem scales (i.e., cross-scale

systems biology).

ACKNOWLEDGMENTS. We thank F. von Netzer, K. Hunt, S Morales, N. Stopnisek, K. Meinhardt, W. Qin, N

Elliot, T. Hazen, H. Carlson, B. Ramsey, A. Murray, Z. Harrold, T. Morgan, and P. Longley for thoughtful feedback

and discussions. This research was supported by a grant from the Nevada Governor's Office of Economic

Development (JG), by the Desert Research Institute (DRI) postdoctoral research fellowship program, and in part by

Ecosystems and Networks Integrated with Genes and Molecular Assemblies (ENIGMA) (http://enigma.lbl.gov)—a

Scientific Focus Area Program at Lawrence Berkeley National Laboratory under contract number DE-AC02-

05CH11231 and funded in part by Oak Ridge National Laboratory under contract DE-AC05-00OR22725, and is

based upon work supported by the U.S. Department of Energy, Office of Science, Office of Biological &

Environmental Research.

Competing Interests

443 The authors declare no conflicts of interest

References

1. Knowles R. Denitrification. *Microbiol Rev* 1982; **46**: 43–70.

2. Tiedje JM, Sexstone AJ, Myrold DD, Robinson JA. Denitrification: ecological niches , competition and
survival. *Antonie Van Leeuwenhoek* 1982; **48**: 569–583.

3. Yoon S, Cruz-García C, Sanford R, Ritalahti KM, Löffler FE. Denitrification versus respiratory
ammonification: Environmental controls of two competing dissimilatory NO 3 -/NO 2 -reduction pathways in
Shewanella loihica strain PV-4. *ISME J* 2015; **9**: 1093–1104.

4. van den Berg EM, van Dongen U, Abbas B, van Loosdrecht MC. Enrichment of DNRA bacteria in a
continuous culture. *ISME J* 2015; **9**: 2153–2161.

5. Schmidt CS, Richardson DJ, Baggs EM. Constraining the conditions conducive to dissimilatory nitrate
reduction to ammonium in temperate arable soils. *Soil Biol Biochem* 2011; **43**: 1607–1611.

6. Hardison AK, Algar CK, Giblin AE, Rich JJ. Influence of organic carbon and nitrate loading on partitioning
between dissimilatory nitrate reduction to ammonium (DNRA) and N2 production. *Geochim Cosmochim
Acta* 2015; **164**: 146–160.

7. Fazzolari É, Nicolardot B, Germon JC. Simultaneous effects of increasing levels of glucose and oxygen
partial pressures on denitrification and dissimilatory nitrate reduction to ammonium in repacked soil cores.
Eur J Soil Biol 1998; **34**: 47–52.

8. Galloway JN, Aber JD, Erisman JW, Seitzinger SP, Howarth RW, Cowling EB, et al. The Nitrogen
Cascade. *Bioscience* 2003; **53**: 341.

9. Moparthi VK, Hägerhäll C. The evolution of respiratory chain complex i from a smaller last common
ancestor consisting of 11 protein subunits. *J Mol Evol* 2011; **72**: 484–497.

10. Dibrova D V., Cherepanov DA, Galperin MY, Skulachev VP, Mulkidjanian AY. Evolution of cytochrome
bc complexes: From membrane-anchored dehydrogenases of ancient bacteria to triggers of apoptosis in
vertebrates. *Biochim Biophys Acta - Bioenerg* 2013; **1827**: 1407–1427.

11. Bergdoll L, Ten Brink F, Nitschke W, Picot D, Baymann F. From low- to high-potential bioenergetic chains:
Thermodynamic constraints of Q-cycle function. *Biochim Biophys Acta* 2016; **1857**: 1569–1579.

12. Schoepp-Cothenet B, Lieutaud C, Baymann F, Verméglio A, Friedrich T, Kramer DM, et al. Menaquinone
as pool quinone in a purple bacterium. *Proc Natl Acad Sci* 2009; **106**: 8549–54.

13. Soo RM, Hemp J, Parks DH, Fischer WW, Hugenholtz P. On the origins of oxygenic photosynthesis and
aerobic respiration in Cyanobacteria. *Science (80-)* 2017; **355**: 1436–1440.

14. Baudouin-Cornu P, Schuerer K, Marlière P, Thomas D. Intimate evolution of proteins: Proteome atomic
content correlates with genome base composition. *J Biol Chem* 2004; **279**: 5421–5428.

15. Grzymski JJ, Dussaq AM. The significance of nitrogen cost minimization in proteomes of marine
microorganisms. *ISME J* 2012; **6**: 71–80.

16. Einsle O, Messerschmidt A, Huber R, Kroneck PMH, Neese F. Mechanism of the six-electron reduction of
nitrite to ammonia by cytochrome c nitrite reductase. *J Am Chem Soc* 2002; **124**: 11737–11745.

17. Kraft B, Strous M, Tegetmeyer HE. Microbial nitrate respiration - Genes, enzymes and environmental
distribution. *J Biotechnol* 2011; **155**: 104–117.

18. Godfrey L V., Falkowski PG. The cycling and redox state of nitrogen in the Archaean ocean. *Nat Geosci*
2009; **2**: 725–729.

19. Nitschke W, Dracheva S. Reaction center associated cytochromes in Anoxygenic Photosynthetic Bactera.
In: Blankenship R, Madigan M, Bauer C (eds).1995. Kluwer Academic, Dordrecht, The Netherlands, pp
775–805.

20. Lalucat J, Bennasar A, Bosch R, Garcia-Valdes E, Palleroni NJ. Biology of Pseudomonas stutzeri.
Microbiol Mol Biol Rev 2006; **70**: 510–547.

21. Suharti, de Vries S. Membrane-bound denitrification in the Gram-positive bacterium Bacillus azotoformans.
Biochem Soc Trans 2005; **33**: 130–133.

22. Heylen K, Keltjens J, Stein LY. Redundancy and modularity in membrane-associated dissimilatory nitrate
reduction in Bacillus. *Front Microbiol* 2012; **3**: 1–27.

23. Decleyre H, Heylen K, Bjorn T, Willems A. Highly diverse nirK genes comprise two major clades that
harbour ammonium-producing denitrifiers. *BMC Genomics* 2016; **17**: 1–13.

24. Yoon S, Sanford RA, Löffler FE. Shewanella spp. Use acetate as an electron donor for denitrification but
not ferric iron or fumarate reduction. *Appl Environ Microbiol* 2013; **79**: 2818–2822.

25. Hillesland KL, Lim S, Flowers JJ, Turkarslan S, Pinel N, Zane GM, et al. Erosion of functional
independence early in the evolution of a microbial mutualism. *Proc Natl Acad Sci* 2014; **111**: 14822–14827.

26. APHA. Standard Methods for the Examination of Water and Wastewater, 23rd ed. 2012. American Public
Health Association: Washington, DC.

27. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu : scalable and accurate long- -
- read assembly via adaptive k - - - mer weighting and repeat separation. *Genome Res* 2017; 1–35.

28. Darling ACE, Mau B, Blattner FR, Perna NT. Mauve : Multiple Alignment of Conserved Genomic
Sequence With Rearrangements. *Genome Res* 2004; **14**: 1394–1403.

29. Darling AE, Mau B, Perna NT. Progressivemauve: Multiple genome alignment with gene gain, loss and
rearrangement. *PLoS One* 2010; **5**: e11147.

30. Andrews S. FastQC: A quality control tool for high throughput sequence data. *Available online at: http://
www.bioinformatics.babraham.ac.uk/projects/fastqc* 2010.

31. Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina sequence data.
Bioinformatics 2014; **30**: 2114–2120.

32. Kalakoutskii LV, Kirillova IP, Krassilnikov NA. a New Genus of the. *J Gen Microbiol* 1967; **48**: 373-79–
85.

33. Klotz MG, Stein LY. Nitrifier genomics and evolution of the nitrogen cycle. *FEMS Microbiol Lett* 2008;
278: 146–156.

34. Kraft B, Tegetmeyer HE, Sharma R, Klotz MG, Ferdelman TG, Hettich RL, et al. The environmental
controls that govern the end product of bacterial nitrate respiration. *Science (80-)* 2014; **345**: 676–679.

35. Abbe KS, Takahashi S, Yamada T. Invovlement of oxygen-sensitive pyruvate-formate lysase in mixed-acid
fermetnation by Streptococcus mutans under strictly anaerobic conditions. *J Bacteriol* 1982; **152**: 175.

36. Das A, Silaghi-Dumitrescu R, Ljungdahl LG, Kurtz DM. Cytochrome *bd* oxidase, oxidative stress, and
dioxygen tolerance of the strictly anaerobic bacterium *Moorella thermoacetica*. *J Bacteriol* 2005; **187**:
2020–2029.

37. Lotka AJ. Contribution to the Energetics of Evolution. *Proc Natl Acad Sci* 1922; **8**: 147–151.

38. DeLong JP. The maximum power principle predicts the outcomes of two-species competition experiments.
Oikos 2008; **117**: 1329–1336.

39. Strohm TO, Griffin B, Zumft WG, Schink B. Growth yields in bacterial denitrification and nitrate
ammonification. *Appl Environ Microbiol* 2007; **73**: 1420–1424.

40. Zumft WG. Cell biology and molecular basis of denitrification. *Microbiol Mol Biol Rev* 1997; **61**: 533–616.

41. Hemp J, Gennis RB. Diversity of the Heme-Copper Superfamily in Archaea: Insights from Genomics and
Structural Modeling. In: Schäfer G, Penefsky HS (eds). *Results and Problems in Cell Differentiation*. 2008.
Springer Science+Business Media, pp 1–28.

42. Rowley G, Hensen D, Felgate H, Arkenberg A, Appia-ayme C, Prior K, et al. Resolving the contributions of
the membrane-bound and periplasmic nitrate reductase systems to nitric oxide and nitrous oxide production
in Salmonella enterica serovar Typhimurium. *Biochem J* 2012; **762**: 755–762.

43. Yoon S, Sanford RA, Löffler FE. Nitrite control over dissimilatory nitrate/nitrite reduction pathways in
Shewanella loihica strain PV-4. *Appl Environ Microbiol* 2015; **81**: 3510–3517.

44. van den Berg EM, Rombouts JL, Kuenen JG, Kleerebezem R, van Loosdrecht MCM. Role of nitrite in the
competition between denitrification and DNRA in a chemostat enrichment culture. *AMB Express* 2017; **7**:
1–7.

45. Link TA, Hagen WR, Pierik AJ, Assmann C, von Jagow G. Determination of the redox properties of the
Rieske cluster of bovine heart bc1 complex by direct electrochemistry of a water- soluble fragment.
EurJBiochem 1992; **208**: 685–691.

46. Ugulava NB, Crofts AR. CD-monitored redox titration of the Rieske Fe-S protein of Rhodobacter
sphaeroides: pH dependence of the midpoint potential in isolated bc1 complex and in membranes. *FEBS Lett*
1998; **440**: 409–413.

47. Zu Y, Couture MMJ, Kolling DRJ, Crofts AR, Eltis LD, Fee JA, et al. Reduction Potentials of Rieske
Clusters: Importance of the Coupling between Oxidation State and Histidine Protonation State. *Biochemistry*
2003; **42**: 12400–12408.

48. Trumpower BL. Cytochrome bc1 complexes of microorganisms. *Microbiol Rev* 1990; **54**: 101–129.

49. Ansong C, Sadler NC, Hill EA, Lewis MP, Zink EM, Smith RD, et al. Protein redox dynamics during light-
to-dark transitions in cyanobacteria and impacts due to nutrient limitation. *Front Microbiol* 2014; **5**: 1–10.

50. Snyder CH, Merbitz-Zahradnik T, Link TA, Trumpower BL. Role of the Rieske Iron – Sulfur Protein
Midpoint Potential in the Protonmotive Q-Cycle Mechanism of the Cytochrome bc 1 Complex. *J Bioenerg*
Biomembr 1999; **31**: 235–242.

51. Denke E, Merbitz-Zahradnik T, Hatzfeld OM, Snyder CH, Link TA, Trumpower BL. Alteration of the
midpoint potential and catalytic activity of the Rieske iron-sulfur protein by changes of amino acids forming

hydrogen bonds to the iron-sulfur cluster. *J Biol Chem* 1998; **273**: 9085–9093.

52.   Hunte C, Solmaz S, Palsdóttir H, Wenz T. A structural perspective on mechanism and function of the
cytochrome bc1 complex. In: Schäfer G, Penefsky HS (eds). *Results and Problems in Cell Differentiation*.
2008. Springer Science+Business Media, pp 253–270.

53.   Falkowski PG. Evolution of the nitrogen cycle and its influence on the biological sequestration of CO2 in
the ocean. *Nature* 1997; **387**: 272–275.

54.   Kao WC, Hunte C. The molecular evolution of the Qo Motif. *Genome Biol Evol* 2014; **6**: 1894–1910.

55.   Dibrova D V., Cherepanov DA, Galperin MY, Skulachev VP, Mulkidjanian AY. Evolution of cytochrome
bc complexes: From membrane-anchored dehydrogenases of ancient bacteria to triggers of apoptosis in
vertebrates. *Biochim Biophys Acta - Bioenerg* 2013; **1827**: 1407–1427.

56.   White D. The physiology and biochemistry of prokaryotes, 3rd ed. 2000. Oxford University Press.

57.   Thauer RK, Jungermann K, Decker K. Energy conservation in chemotrophic anaerobic bacteria. *Bacteriol
Rev* 1977; **41**: 100–180.

58.   Giardina CP, Ryan MG. Total belowground carbon allocation in a fast-growing Eucalyptus plantation
estimated using a carbon balance approach. *Ecosystems* 2002; **5**: 487–499.

59.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids
Res* 2004; **32**: 1792–7.

60.   Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA5: Molecular evolutionary genetics
analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol
Evol* 2011; **28**: 2731–2739.

61.   Stamatakis A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies.
Bioinformatics 2014; **30**: 1312–1313.

62.   Kelley LA, Mezulis S, Yates C, Wass M, Sternberg M. The Phyre2 web portal for protein modelling,
prediction, and analysis. *Nat Protoc* 2015; **10**: 845–858.

63.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: Ultrafast universal RNA-seq
aligner. *Bioinformatics* 2013; **29**: 15–21.

64.   Liao Y, Smyth GK, Shi W. FeatureCounts: An efficient general purpose program for assigning sequence
reads to genomic features. *Bioinformatics* 2014; **30**: 923–930.

65.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with
DESeq2. *Genome Biol* 2014; **15**: 1–21.

 Figure legends

Figure 1. (A) Maximum likelihood phylogenetic tree of NrfA amino acid sequences from known
respiratory ammonifiers and accompanying N-module composition for each organism. Pool
quinone is also noted for dual-pathway nitrite reducers and model species. Colors of the main
branches denote the 1st heme motif type: CxxCK and CxxCH. (B) Protein atomic composition
for N and C normalized to protein length for NirK and NrfA nitrite reductases. (C) State-
transition from ammonium-replete to ammonium-deplete for *I. calvum* C5 grown under 8mM
lactate 12mM nitrate minimal media at 30 ºC. Metabolite profiles for ammonium-deplete are
shown.

Figure 2. The effects of high resource (left; range of lactate concentrations with 12 mM NO_3^-)
and low resource (right; range of lactate concentrations with 1.2 mM NO_3^-) concentrations with
the same C:NO_3^- ratio on pathway selection in *I. calvum* C5. (A) Production of N_2O-N and net
change of NH_4^+ over a 100-hour incubation period at 30 ºC. Each bar represents the average of
8-10 replicates per treatment (Table S5). (B) Fraction of dissimilated N by pathway. (C) Growth
rates for each corresponding treatment. The x-axis label defines lactate concentration and C:NO_3^-
ratio in parentheses.

Figure 3. (A) Time-series metabolite profiles for lactate, nitrate, and nitrite (top pane),
production of dissimilated end-products as N_2O-N and net change in NH_4^+ ammonium
production (middle pane), and corresponding growth curve of *I. calvum* cells grown under 8 mM
lactate 12 mM nitrate (C:NO_3^- ratio = 2) (bottom pane). Sampling points during growth phases
are marked where transcriptomic profiling was performed (red arrows). (B) Metabolic
reconstruction of the ETC from *I. calvum* with transcriptional changes for genes participating in
dual-pathway dissimilatory nitrite reduction. Log$_2$ fold changes in transcript abundance are
shown for late exponential relative to early exponential growth phase (EE vs. LE), stationary
phase relative to early exponential growth phase (EE vs. ST), and stationary phase relative to late
exponential growth phase (LE vs. ST). Locus IDs for each gene product correspond to heat map
subplots in the order shown (left-to-right for each growth phase and top-to-bottom for each locus
ID specified). Higher transcript abundance is represented in red, lower transcript abundance in
blue, and no change in transcript abundance in white. Significant changes in transcript
abundance ($p < 0.01$) are marked as a red box. Value of log$_2$ fold change is specified within each
subplot. The log$_2$ fold changes of 14 NADH dehydrogenase subunits (Intca_03465-03530) were
averaged as transcriptional changes were all shifted in the same direction.

Figure 1. (A) Maximum likelihood phylogenetic tree of NrfA amino acid sequences from known respiratory ammonifiers and accompanying N-module composition for each organism. Pool quinone is also noted for dual-pathway nitrite reducers and model species. Colors of the main branches denote the 1st heme motif type: CxxCK and CxxCH. (B) Protein atomic composition for N and C normalized to protein length for NirK and NrfA nitrite reductases. (C) State-transition from ammonium-replete to ammonium-deplete for *I. calvum* C5 grown under 8mM lactate 12mM nitrate minimal media at 30 ºC. Metabolite profiles for ammonium-deplete are shown.

Figure 2. The effects of high resource (left; range of lactate concentrations with 12 mM NO$_3^-$) and low resource (right; range of lactate concentrations with 1.2 mM NO$_3^-$) concentrations with the same C:NO$_3^-$ ratio on pathway selection in *I. calvum* C5. (A) Production of N$_2$O-N and net change of NH$_4^+$ over a 100-hour incubation period at 30 ºC. Each bar represents the average of 8-10 replicates per treatment (Table S5). (B) Fraction of dissimilated N by pathway. (C) Growth rates for each corresponding treatment. The x-axis label defines lactate concentration and C:NO$_3^-$ ratio in parentheses.

Figure 3. (A) Time-series metabolite profiles for lactate, nitrate, and nitrite (top pane), production of dissimilated end-products as N_2O-N and net change in NH_4^+ ammonium production (middle pane), and corresponding growth curve of *I. calvum* cells grown under 8mM lactate 12mM nitrate (C:NO_3^- ratio = 2) (bottom pane). Sampling points during growth phases are marked for transcriptomic analysis. (B) Metabolic reconstruction of the ETC from *I. calvum* with transcriptional changes for genes participating in dual-pathway dissimilatory nitrite reduction. Log_2 fold changes in transcript abundance are shown for late exponential relative to early exponential growth phase (EE vs. LE), stationary phase relative to early exponential growth phase (EE vs. ST), and stationary phase relative to late exponential growth phase (LE vs. ST). Locus IDs for each gene product correspond to heat map subplots in the order shown (left-to-right for each growth phase and top-to-bottom for each locus ID specified). Higher transcript abundance is represented in red, lower transcript abundance in blue, and no change in transcript abundance in white. Significant changes in transcript abundance ($p < 0.01$) are marked as a red box. Value of log_2 fold change is specified within each subplot. The log_2 fold changes of 14 NADH dehydrogenase subunits (Intca_03465-03530) were averaged as transcriptional changes were all shifted in the same direction.