Internal noise in contrast discrimination propagates forwards from early visual cortex

Greta Vilidaite, Emma Marsh & Daniel H. Baker

Department of Psychology, University of York

Correspondence (corresponding author)
Daniel H Baker, email: daniel.baker@york.ac.uk

Abstract

Human contrast discrimination performance is limited by transduction nonlinearities and variability of the neural representation (noise). Whereas the nonlinearities have been well characterised, there is less agreement about the specifics of internal noise. Psychophysical models assume that it impacts late in sensory processing, whereas neuroimaging and intracranial electrophysiology studies suggest that the noise is much earlier. We investigated whether perceptually-relevant internal noise arises in early visual areas or later decision making areas. We recorded EEG and MEG during a two-interval-forced choice contrast discrimination task and used multivariate pattern analysis to decode target/non-target and selected/non-selected intervals from evoked responses. We found that perceptual decisions could be decoded from both EEG and MEG signals, even when the stimuli in both intervals were physically identical. Above-chance decision classification started <100ms after stimulus onset, suggesting that neural noise affects sensory signals early in the visual pathway. Classification accuracy increased over time, peaking at ~700ms. Applying multivariate analysis to separate anatomically-defined brain regions in MEG source space, we found that occipital regions were informative early on but then information spreads forwards across temporal and frontal regions. This is consistent with neural noise affecting sensory processing at multiple stages of perceptual decision making. We suggest how early sensory noise might be resolved with Birdsall's linearisation, in which a dominant noise source obscures subsequent nonlinearities, to allow the visual system to preserve the wide dynamic range of early areas whilst still benefitting from contrast-invariance at later stages.

Keywords: contrast discrimination; MEG; source space; pattern classification; internal noise

1 Introduction

The ability to make comparisons between sensory stimuli of different intensities has profound survival value for most organisms. Animals might benefit from choosing the ripest fruit based on colour, swimming towards the warmest patch of ocean, or selecting the mate with the loudest roar. Understanding the features of the central nervous system that limit such sensory discriminations has been a focus of research in all areas of psychology and neuroscience, from early work in humans (c.f. Weber's law, Fechner, 1912) and model organisms (Busse et al., 2011; Hecht & Wald, 1934) to studies using contemporary neuroimaging techniques (Boynton, Demb, Glover, & Heeger, 1999).

A widely studied perceptual task is the ability to discriminate between visual stimuli of different contrasts. Human contrast discrimination performance is constrained by the nonlinearity that maps physical contrast to neural response, and the intrinsic variability of the neural representation ('internal noise'). Psychophysical, neurophysiological and neuroimaging work have converged on a nonlinearity that is expansive at low contrasts and compressive at higher contrasts (Boynton et al., 1999; Busse, Wade, & Carandini, 2009; Legge & Foley, 1980). However, there is substantially less agreement regarding the details of performance-limiting internal noise.

Most psychophysical models make the assumption that the dominant source of noise for contrast discrimination is additive (i.e. independent of signal strength) and impacts at a late stage of processing. The primary justification for this arrangement is the observation that a dominant source of noise occurring before a nonlinearity will neutralise the effects of that nonlinearity, rendering it invisible to inspection (Klein & Levi, 2009; Smith & Swift, 1985). Since contrast transduction is observably nonlinear (Boynton et al., 1999; Busse et al., 2009; Legge & Foley, 1980), any early sources of noise must be negligible in comparison to the magnitude of late additive noise.

On the other hand, most electrophysiological and neuroimaging studies have suggested that perceptually relevant noise is located in early sensory areas (Campbell & Kulikowski, 1972; Carandini, 2004; Roelfsema & Spekreijse, 2001). Ress and Heeger (2003) demonstrated the influence of early sensory noise by measuring fMRI blood-oxygen-level dependent (BOLD) responses in areas V1-V4 during contrast detection. They found that *false alarms* evoked higher responses than *misses*, suggesting that these areas encoded conscious percepts of the stimuli rather than the presence of the stimulus itself. The origin of the spurious activity in the case of *false alarms* must be neural noise in these early areas. Similarly, several intracranial primate electrophysiology studies have been able to predict the perceptual decisions of monkeys from neural activity recorded in early visual areas (Britten, Newsome, Shadlen, Celebrini, & Movshon, 1996; Britten, Shadlen, Newsome, & Movshon, 1992; Michelson, Pillow, & Seidemann, 2017). This suggests that sensory decisions are influenced by neural noise at an early stage of processing.

In this study, we attempt to understand how neural activity governs observer responses in a two-interval-forced-choice (or 2IFC) contrast discrimination paradigm, using methods typical of such studies. This paradigm involves several complicating factors that must be considered including: (i) the observer must retain a neural representation of the first stimulus for

comparison with the second stimulus, (ii) individuals might have idiosyncratic biases to prefer one or other interval, and (iii) fast acting adaptation (often termed repetition suppression) effects will reduce the neural response to the second stimulus (and perhaps also its appearance). We recorded evoked responses using both EEG (Experiment 1) and MEG (Experiment 2), and perform both traditional univariate analyses, and also employ multivariate pattern analysis to decode participants' percepts. The high temporal resolution (~1ms) of both techniques enabled us to closely examine the timecourse of perceptual decision making, and the spatial resolution of MEG source space allowed us to investigate the involvement of discrete anatomical brain areas. Our primary motivation was to determine whether the dominant source of neural noise is located in early sensory brain areas, or later (more frontal) areas involved in making decisions.

2 Methods

2.1 Participants

Twenty-two adults with normal or corrected-to-normal vision took part in Experiment 1 and ten took part in Experiment 2. All participants gave written informed consent. Experiment 1 was approved by the Ethics Committee of the Department of Psychology at the University of York, and Experiment 2 was approved by the York Neuroimaging Centre Ethics Committee.

2.2 Stimuli and psychophysical task

Stimuli were horizontally oriented sine wave gratings with a spatial frequency of 1c/deg and a diameter of 10 degrees. The edges of the gratings were blurred by a cosine function. On each trial, two stimuli were presented: a pedestal stimulus of 50% contrast (where percent contrast is defined as $100*(L_{max}-L_{min})/(L_{max}+L_{min})$, where L is luminance), and a pedestal+target stimulus consisting of the 50% contrast pedestal plus a target contrast increment. Five target contrast conditions were used in Experiment 1: 0% (no target), 2%, 4%, 8% and 16%. In Experiment 2 only the 0% (no target) and 16% target contrast conditions were used.

The two stimuli on each trial were presented sequentially for 100ms each, with a random inter-stimulus interval between 400ms and 600ms. The inter-trial interval followed the participant's response, and was of variable length between 1000ms and 1200ms to avoid distortion of ERP averages (Woldorff, 1993). The order of target and non-target intervals within trials was counterbalanced. Trials of different target contrasts were intermixed and the order was randomized. Stimulus onsets and participant responses were recorded on the M/EEG trace using low-latency digital triggers.

2.3 EEG data collection

Event-related potentials were recorded using an ANT Neuroscan EEG system and a 64-channel Waveguard cap with electrodes arranged according to the 10/20 system. Data were digitised at 1kHz using the ASALab software. Stimuli were presented on a gamma corrected ViewPixx display (VPixx Technologies Inc., Quebec, Canada) with a mean luminance of

51cd/m^2 and a refresh rate of 120Hz, using Matlab and elements of the Psychophysics Toolbox (Brainard, 1997; Kleiner, Brainard, & Pelli, 2007; Pelli, 1997).

Participants were seated in a darkened room 57cm away from the display. Instructions for the task were to 'indicate the grating that appeared higher in contrast'. They were asked to fixate on a central fixation cross throughout the task and used a mouse to indicate their responses. There were 200 trials per target contrast (1000 trials total, yielding 2000 stimulus-locked ERPs). The task was run in 5 blocks of approximately 8 minutes, with short breaks in between.

2.4 MEG data collection

MEG data were recorded using a 4D Neuroimaging Magnes 3600 Whole Head 248 Channel MEG scanner housed in a purpose-built Faraday cage. The data were recorded at 1017.25Hz, with 400Hz Bandwith using a High Pass DC filter. Nine channels were identified as having failed and were removed from all analyses. The location of the head inside the dewar was continuously monitored throughout the experiment using 5 position indicator head coils. Stimuli were presented on an Epson EB-G5900 3LCD projector (refresh rate 60Hz; mean luminance 160cd/m^2) with a 2-stop ND filter, using Psychopy v1.84 (Peirce, 2007).

Participants were seated in a hydraulic chair in front of the projector screen in a dark room. Prior to the task the three dimensional shape of the participant's head was registered using a Polhemus fast-track headshape digitization system. Five fiducial points were used for this over two registration rounds. If the distance in location between the first and second round was >2mm, the registration was repeated. When successful, the headshape was then traced and recorded using a digital wand. This was later coregistered with T1-weighted anatomical MRI scans of each participant acquired in separate sessions using a 3T GE Signa Excite HDx scanner (GE Healthcare).

Participants fixated on a small central cross throughout the task. The experiment was completed in a single block consisting of 240 trials per contrast condition (480 trials in total, yielding 960 stimulus-locked ERPs), with a total acquisition duration of around 20 minutes. A single hand response pad was used to make responses in the experiment.

2.5 EEG data analysis

EEG recordings were bandpass filtered (from 0.1Hz to 30Hz) and then epoched into 1 second-long windows (200ms before stimulus onset to 800ms after) for each interval of every trial. Each epoch was then baselined at each electrode independently by subtracting the mean response over the 200ms preceding stimulus onset. ERPs were then sorted by target/non-target intervals for stimulus classification analysis and then again by selected/non-selected intervals for decision classification.

To perform univariate analyses, ERPs were averaged across a cluster of 10 posterior electrodes (Oz, O1, O2, POz, PO3-8), and significance was determined using cluster corrected paired-samples t-tests across participants (Maris & Oostenveld, 2007). The significance of each cluster was determined by comparing to a null distribution of summed t-values derived

by randomly permuting the labels of the largest cluster 1000 times. To perform multivariate analyses, a support vector machine (SVM) was used to classify the data independently at each sample point (i.e. in 1ms steps). A second stage of normalization was applied at each time-point and each electrode by subtracting the mean response across all intervals and conditions for that time/sensor combination. The data were then randomly averaged in five subsets of 40 trials for each category (target/non-target or selected/non-selected), of which four subsets were used to train the model and one was used to test it. The classifier algorithm creates a parameter space of all data points and then fits a hyperplane boundary that maximizes the distances between the support vectors of each category. Classifier accuracy for categorising the test data was averaged across 1000 repetitions of the analysis (with different random allocations of trials on each repetition), and was repeated for each target contrast condition. Timecourses of classifier accuracy were then averaged across participants, and periods of above-chance performance were determined using the same non-parametric cluster correction procedure as used in the univariate analyses (Maris & Oostenveld, 2007).

2.6 MEG data analysis

Cortical reconstruction and volumetric segmentation was performed with the Freesurfer image analysis suite (http://surfer.nmr.mgh.harvard.edu/) using each individual participant's anatomical MRI scan. Initial MEG analyses were then performed in *Brainstorm* (Tadel, Baillet, Mosher, Pantazis, & Leahy, 2011). First the MEG sensor array was aligned with the anatomical model of the participant's head using an automated error minimisation procedure. Covariance matrices were estimated from the data, and a head model comprising overlapping spheres was generated. A minimum norm solution was used to calculate a source model, with dipole orientations constrained to be orthogonal to the cortical surface. The model consisted of a set of linear weights at each location on the cortical surface that transformed the sensor space representation into source space.

MEG data were then imported into Matlab using *Fieldtrip* (Oostenveld, Fries, Maris, & Schoffelen, 2011), bandpass filtered (from 0.1Hz to 30Hz) and epoched. Univariate and multivariate analyses performed in the same way as described for the EEG data in section 2.5. This was done using the sensor space representation (with 239 working sensors), the source space representation at approximately 500 vertices evenly spaced across the cortical mesh, and also within discrete regions of cortex defined by the Mindboggle atlas (A. Klein et al., 2017). For this latter analysis, the mean number of vertices in each cortical region is given in Table A1 in the Appendix.

3 Results

3.1 Experiment 1: EEG reveals above-chance classification of percepts

Mean event-related potentials (ERPs) averaged over the ten occipital electrodes where the changes in response from baseline were greatest (Figure 1a) showed a typical response to brief visual stimulation (black curve, Figure 1a). Clear ERPs were evident for all individual participants (thin traces, Figure 1a). In the grand average (black curve), two successive positive responses were evident over occipital electrodes at early time-points (126ms and 225ms after stimulus onset). A later time-point (594ms after stimulus onset) showed negative

voltages in occipital areas and positive voltages in frontal electrodes (see upper headplots for voltage distributions).

Task performance in the five target contrast conditions ranged from chance in the 0% target contrast condition (where there was no correct answer as the 'target' interval was determined arbitrarily) to close to ceiling in the 16% target contrast condition (94% correct). Average data (black line) and results for individual participants (thin traces) are shown in Figure 1b, where it is evident that increasing target contrast improved performance for all participants. We fitted cumulative Gaussians to each participant's data to estimate threshold contrast at 75% correct, which was 4.25% on average, with the distribution shown at the foot of Figure 1b.

Figure 1: Grand mean ERPs (a) and summary of psychophysical performance (b). The black trace in panel (a) shows the grand mean across all conditions and participants (N=22), with the grey shaded region giving 95% confidence intervals derived from 10000 bootstrap resamples. Thinner coloured traces show results for individual participants. In all cases the evoked responses were averaged across the 10 posterior electrodes shown in the lower left insert. The grey rectangle along the lower axis indicates the period during which the stimulus was presented. Distributions of voltages at three time points (126ms, 251ms and 594ms, marked by dashed vertical lines) are shown at the top of the plot. The black line in panel (b) shows the mean psychophysical performance in each condition, averaged across participants (N=22), with the grey shaded region giving 95% confidence intervals derived by bootstrapping. Thinner coloured traces show results for individual participants, and symbol colour corresponds to those used to indicate target contrast conditions in subsequent figures. The grey curve at the foot shows the distribution of individual thresholds at the 75% correct point, with the black circle giving the mean, and error bars giving 95% confidence intervals.

We first divided ERP data by contrast, and compared evoked responses in the null (pedestal only) and target (pedestal + target) intervals. The upper row of Figure 2a shows the ERPs averaged across occipital electrodes, with the null interval responses shown in black, and the target interval responses in colour. The middle row of Figure 2a shows the differences between these two ERPs, with horizontal lines at y=-1.5 indicating time points showing cluster-corrected significant differences. For a target contrast of 0%, the two stimuli are identical, and there are no meaningful differences between the waveforms (the two brief periods of significance are type I errors by definition). As target contrast increases, significant differences emerge between 100ms and 700ms post stimulus onset. These likely reflect both differences in early evoked responses, and also later decision-related components. Multivariate analyses across all 64 electrodes showed significant decoding only at the highest two target contrast levels (lower row of Figure 2a) within the same time window.

Figure 2: Univariate and multivariate analyses of EEG data. Panel (a) shows results for data partitioned according to the stimulus contrast (pedestal vs pedestal+target), and panel (b) shows results for data partitioned according to the participants' perceptual decisions (selected vs non-selected). The upper section of each sub-plot contains grand averages of the ERPs being compared, in which the coloured curve indicates the target (or selected) waveform, and the black curve indicates the pedestal (or non-selected waveform). The middle section of each sub-plot is the difference waveform. The lower section of each sub-plot shows classifier performance at each timepoint, where the baseline is 50% correct. In each panel, shaded regions show 95% confidence intervals across participants (N=22), calculated by bootstrapping. The coloured horizontal lines in the lower two sections indicate periods of time when the difference waveforms were significantly different from 0 (middle plots) or exceeded 50% correct (lower plots), calculated using a nonparametric cluster correction procedure (Maris & Oostenveld, 2007).

Next, we repeated the analyses on data organised according to the participant's decisions, rather than the physical stimulus contrast. In other words, we took ERPs from the intervals selected by the participants as appearing higher in contrast, and compared these with ERPs from the non-selected intervals. This analysis revealed additional time periods where the ERPs were significantly different, particularly in the 0% target condition, where differences were observed at around 100ms post stimulus onset. This finding was echoed in the multivariate analyses, which showed above chance decoding at early time points (around 100ms), as well as a sustained period of above chance decoding at all target contrasts from around 400-600ms post stimulus onset. The 0% target condition is of particular interest for this analysis, as any differences between evoked responses are not determined by the stimulus (which is identical in both intervals), and must be a consequence of differences in neural activity. The early significant clusters in both univariate and multivariate analyses indicate differences in the amplitude of the evoked response that influence subsequent perceptual decisions.

We tested the generality of the multivariate results in two ways. First, we took the classifier trained to discriminate between perceptual decisions at the highest target contrast (16%), and used this model to predict performance at lower target contrasts. This analysis (shown in Figure 3a) replicates the early periods of above chance decoding for 0% target contrast trials, suggesting that observers use a similar decision strategy for very challenging discriminations as for easier ones. Next, we took the classifier trained at each time point, and used it to predict selected and non-selected trials at all other time points (King & Dehaene, 2014). The results of this temporal generalization analysis (shown in Figure 3b,c) reveal isolated early structures

around 100ms and 200ms, and a sustained pattern from 400-600ms (in the 0% condition) and from 200-800ms (in the 16% condition). We propose (see Discussion) that the early periods of above chance decoding may represent neural noise at the initial stages of processing, and the later periods could reflect perceptual decisions or memory traces from the first temporal interval.

Figure 3: Multivariate generalization analyses across condition (a) and time (b,c), for data partitioned according to the participants' perceptual decisions. Panel (a) shows classifier accuracy at the four lower target contrasts after training the algorithm at the highest target contrast. Plotting conventions are as described for the multivariate analyses shown in Figure 2. Panels (b,c) show classifier accuracy when trained at each time point independently, and then tested using data at all time points. Regions enclosed by green lines indicate clusters when classifier accuracy differed significantly from chance (50% correct).

3.2 Interval biases and decoding within the first or second interval

The temporal structure of a 2IFC trial is necessarily asymmetric, as the observer has knowledge of the first interval by the time they experience the second interval. In addition, repetition suppression effects can affect the amplitudes of the second presentation (Grill-Spector, Henson, & Martin, 2006). We first compared the average ERPs for all pedestal-only presentations (where the stimulus contrast was 50%) across the two intervals. We find both subtle and gross differences between these waveforms (see Figure 4a). Before stimulus onset, the waveforms differ as the second interval (green trace) has a decreasing voltage in the 200ms before the stimulus is presented. This likely originates from the tail end of the evoked response from the first interval (see Woldorff, 1993), which is decreasing from 400-600ms (the time window in which the second interval occurred). The second interval then has a more generally negative voltage throughout the 800ms following stimulus onset. The magnitude of this difference is much greater than that at stimulus onset, and so would persist even with a different baseline normalization regime (e.g. if the voltages were normalized to those at t=0). Furthermore, the differences become much more substantial at later time points, from 400-800ms. This may relate to the perceptual decision and motor response that the participant must make following the second interval.

Do these substantial differences in the evoked response to two physically identical stimuli affect the observer's perception of the stimulus, or their decision over which interval to choose? We estimated interval bias for all participants by calculating the proportion of trials on which the second interval was selected, for the 200 trials in the 0% target contrast condition (where the two stimuli are identical). If this index is significantly below 0.5, it indicates a bias towards the first interval, and if it is significantly above 0.5 it indicates a bias towards the second interval. Despite individuals showing idiosyncratic biases (indices ranged from 0.23 to 0.92), the mean bias index was precisely 0.5 (SD: 0.14, t_{21}=0.11, p=0.91). The substantial voltage differences therefore do not appear to reflect group level differences in the appearance of the stimuli across intervals, and any idiosyncratic biases would presumably only reduce the power of our decision decoding analyses (Figure 2b), which are nevertheless significant.

The size of the voltage differences across intervals prompted us to investigate the extent to which decisions can be decoded within one or other interval, making comparisons across trials (within an interval) instead of across intervals (within a trial). The presence of interval biases for some observers (see above) meant that there were often different numbers of trials available in the two intervals, so it was necessary to train and test the classifier on averages of fewer than 40 trials in some cases. The results of this multivariate analysis are shown in Figure 4b/c for decoding perceptual decisions at 0% contrast (Fig 4b) and at 16% contrast (Fig 4c), and for all conditions in Figure A1. In each sub-plot, the upper trace shows the classifier performance for data from interval 1 (with ERPs aligned to t=0), and the lower trace shows the classifier performance for data from interval 2 (with ERPs aligned to t=500). The yellow shaded regions indicate the time window when the stimulus in the other interval was displayed (the jittered inter-stimulus interval means that this time window is probabilistic rather than exact). Overall, we find increased decoding accuracy in the second interval compared with the first. This presumably reflects the increased information available for making a decision following the second stimulus.

Figure 4: Comparisons across trial intervals. Panel (a) shows evoked responses in the first (red curve) and second (green curve) intervals of a 2IFC trial (upper plot) and their difference (lower plot). The evoked response is generally more negative in the second interval, particularly at time points >250ms after stimulus onset, despite the contrasts being physically identical (both 50%). Panels (b,c) show multivariate pattern classifier accuracy when comparing evoked potentials time-locked to either the first interval (upper plots) or the second interval (lower plots), for target contrasts of 0% (panel b) or 16% (panel c). In each plot, the shaded grey region shows the presentation of the stimulus of that interval and the yellow shaded region shows the range of time points when the stimulus of the other interval was presented (the precise inter-stimulus interval was jittered on each trial to reduce distortion of ERP averages). In all panels shaded regions around each curve show 95% confidence intervals across participants, and horizontal coloured lines indicate significant clusters, consistent with conventions in previous figures.

3.3 Experiment 2: source space decoding is more sensitive than sensor space decoding

We confirmed that our MEG data replicated the key effects from Experiment 1 in several ways. First, we performed univariate and multivariate analyses in sensor space, using a cluster of posterior sensors for the univariate analysis, and all working sensors (N=237) for the multivariate analysis. The results of this analysis are shown in Figure 5a for the data split by participants' perceptual decisions. Consistent with the EEG results, we find above chance pattern classification at early time points (~100ms) as well as later >200ms. Second, we performed complementary analyses in MEG source space, using ERPs from pericalcarine cortex (corresponding to early visual cortex) for the univariate analyses, and a subset of 500 vertices across the entire cortical surface for the multivariate analyses. The results of this analysis are shown in Figure 5b. The general pattern of results is consistent with the sensor space analysis, though the shape of the ERP waveforms from pericalcarine cortex is somewhat different from those recorded in sensor space, with the peaks of the onset and offset response appearing more prominent, and the response returning to baseline by around 500ms. Interestingly, we found that the multivariate analysis produced greater classification accuracy in source space (maximum of 80% correct) versus sensor space (maximum of 72% correct). We discuss possible reasons for this in the Discussion. Having confirmed that the multivariate source space analysis can discriminate between conditions, we next asked which different brain regions contained information relevant to the task.

Figure 5: Sensor space and source space MEG analysis. Panel (a) shows the grand average ERP for all conditions and participants, pooled across a subset of MEG sensors indicated in black in the leftmost insert. Magnetic field distributions across the sensor array are shown at three time points at the top of the plot. Panel (b) shows a similar analysis in source space, for a region of cortex around the calcarine sulcus (highlighted black in the leftmost insert). The evoked response at each vertex on the cortical mesh was normalised such that the 110ms was always positive, to avoid signal cancellation due to polarity inversions. In both panels, thin coloured curves represent individual participants (N=10). Panels (c,d) show univariate and multivariate comparisons between selected and non-selected ERPs in both contrast conditions, in the same format as described for Figure 2. Panel (c) shows this analysis in sensor space, and panel (d) show the same analysis in sensor space. The source space multivariate analyses used a matrix of around 500 points distributed across the surface of the cortex.

3.4 Classification in anatomically-defined brain regions

We divided the cortex into 31 discrete non-overlapping anatomical regions using the Mindboggle atlas (Klein et al., 2017). Maximal evoked potentials in these regions showed clear differentiation (see Figure A2). Because regions differed in size, each area contributed a different number of vertices on the cortical mesh for pattern classification (see Table A1).

At early time points, around 100ms, information in three adjacent regions around the occipital pole (the peri-calcarine region, the cuneus and the lateral occipital cortex) could be used to decode the observer's percept in the 0% target contrast condition (final three traces in Figure 6a). Over time, this information spreads forward to frontal and temporal cortex (see Figure 6c). By 300ms following stimulus onset, almost the entire brain contains information relevant to the task. This includes regions that do not appear to respond directly to presentation of visual stimuli (i.e. there is no measurable evoked response, see Figure A2). A similar pattern of results is evident in the 16% target contrast condition (see Figure A3), confirming our earlier finding that differences in physical and perceived contrast are processed in a similar fashion.

Figure 6: Atlas-based classification of decisions in the 0% target condition. Timecourses in panel (a) indicate classifier performance for each brain region, organised from anterior (top) to posterior (bottom) (see legend in panel b). Shaded regions in panel (a) indicate clusters in which classification performance was significantly above chance. In panel (c), regions containing significant clusters within a given time window are shown in blue.

4 Discussion

The present study investigated the timecourse and location of perceptually relevant neural noise in sensory processing, using univariate and multivariate analysis of EEG and MEG data. Our results show that perceptual decisions are encoded in early visual cortex even when the two stimuli in a discrimination task are physically identical. This indicates that perceptually relevant neural noise impacts at the initial stages of processing and affects stimulus encoding in the visual system. However the best classifier performance occurred at later time points (>400ms), suggesting that later sources of noise might also be involved. Analysis of differences across trial intervals revealed that neural activity in the second interval was more closely associated with subsequent decisions. We will now discuss the implications of these finding for our understanding of how neural activity (both evoked and spontaneous) influences the perceptual decisions involved in sensory discrimination.

4.1 Superior classification in MEG source space

Classifier performance overall was much higher for MEG data than for EEG data in identical conditions, despite the larger sample size of the EEG study (N=22 for EEG vs. N=10 for MEG). This is presumably due to the greater intrinsic sensitivity of MEG sensors, and the greater sampling density across the scalp (N=64 for EEG vs. N=239 for MEG). Classifier accuracy was also consistently higher in source space than in the sensor space representation primarily used in previous MEG studies (Cichy, Pantazis, & Oliva, 2014; Clarke, Devereux, Randall, & Tyler, 2015; Mostert, Kok, & de Lange, 2016). Since the source space representation is a weighted linear combination of activity at the sensors, this might be somewhat surprising. However, the source reconstruction presumably weights out signals from outside the brain (e.g. heart rate, breathing and blinking artefacts, and noise from outside of the scanner), resulting in a cleaner signal. Some form of source localisation may therefore be a useful processing step in future studies attempting multivariate classification of MEG signals. Additionally, combining the source space representation with atlas-based multivariate analysis permits questions to be asked about the information contained in specific brain regions at different points in time.

4.2 Single interval versus 2IFC

One distinction between this and most previous studies on the neural correlates of perceptual decision making is that previous work has used single interval (yes/no) paradigms (Hesselmann, Kell, Eger, & Kleinschmidt, 2008; Hillyard, Squires, Bauer, & Lindsay, 1971; Jolij, Meurs, & Haitel, 2011; Mostert et al., 2016; Ress & Heeger, 2003; Schölvinck, Friston, & Rees, 2012; Squires, Squires, & Hillyard, 1975), whereas here we used a 2IFC design. Since most psychophysical studies of contrast discrimination have used 2IFC, this choice has more direct relevance to previous work. Additional benefits are that the number of evoked potentials in the selected and non-selected categories were necessarily balanced; and it was possible to analyse perceptual decisions based on two physically identical stimuli. In addition, 2IFC

designs avoid problems with differences in bias (or response criteria) between participants, as pairs of stimuli are compared directly on a given trial (rather than against an internal standard). However, 2IFC cannot distinguish between hits and correct rejections (as these comprise 'correct' trials) or between misses and false alarms (incorrect trials), so direct comparisons of these trial categories is not possible in our design.

Another feature of 2IFC paradigms is that participants must hold information about the stimulus from the first interval in memory until after the second stimulus has been presented. This process may account for the sustained patterns of activity that permit classification long after stimulus presentation (see Figures 2-6). In particular, our analysis of interval-specific effects (see Figure 4b,c) shows greater multivariate decoding accuracy in the second interval, presumably because at this point in the trial the observer has obtained all information necessary to make a decision.

4.3 Resolving early noise and Birdsall's theorem

Early noise has typically been considered at very early stages, including photoreceptor noise in the retina (Barlow, 1962). Late additive noise is often assumed (either implicitly or explicitly) to be added at the decision stage, long after the nonlinearities of early visual processing (Cabrera, Lu, & Dosher, 2015; Mueller & Weidemann, 2008). The results here point to a perceptually-relevant source of noise that is present in the early evoked response, at around 100ms or earlier. However we note that classification performance improves after this point in processing, reaching a maximum around 700ms after target onset (see Figure 3e). This is consistent with a cascade of multiple noise sources at different stages of processing. Since mathematical treatment of complex systems involving multiple nonlinearities and noise sources is currently lacking, it is unclear what implications this would have for the visibility of early nonlinearities.

One possibility is that a strong source of noise occurs immediately after the initial contrast transduction nonlinearity in V1, leaving that nonlinearity visible but obscuring later ones. This would explain why psychophysical contrast perception maps closely onto the neural response from early visual areas (Baker & Wade, 2017; Barlow, Hawken, Parker, & Kaushal, 1987; Boynton et al., 1999), but not the highly compressive contrast-invariant response in later regions (Avidan et al., 2002; Rolls & Baylis, 1986). Indeed, this might enable the visual system to harness the properties of Birdsall linearisation to preserve the dynamic range of early representations through later processing (that is more compressive) when making comparisons across stimuli (as in a discrimination paradigm). Object recognition, and other operations that benefit from invariance to features such as contrast, position and size, but do not require comparisons across multiple stimuli, would be immune to the Birdsall effect and benefit from the later nonlinearities. Furthermore, a strong early source of noise would make the study of later 'mid-level' visual processes much more challenging, perhaps explaining why vision research has typically focussed on earlier mechanisms and can be caricatured as being 'stuck' in V1 (Graham, 2011; Peirce, 2007).

In order to link the internal fluctuations measured in our experiments with a psychophysical measure of internal noise, we correlated classifier accuracy with the contrast discrimination threshold estimated from the psychophysical responses in Experiment 1. Since high internal

noise should result in both higher discrimination thresholds and increased classifier accuracy (because, counterintuitively, it should be easier to predict the perceptual responses of participants whose neural activity is more influenced by noise), we predicted that the two measures would be correlated at time points where the neural fluctuations were most relevant to perception. This analysis is shown in Figure 7, and reveals a time window with significant correlations around 450-650ms. We speculate that neural noise within this time window most closely corresponds to the 'late' additive noise that is a feature of contemporary models of contrast discrimination. It is worth noting that the interval biases observed for some participants should act to reduce the magnitude of this correlation, since a strong interval bias should increase psychophysical thresholds, but decrease classifier accuracy. Therefore interval bias cannot explain the positive correlation we report here.

Figure 7: Correlation between individual contrast discrimination thresholds (see distribution in Figure 1b) and classifier accuracy in the 0% target contrast condition of Experiment 1 (N=22). The horizontal black lines at r=-0.5 denote clusters of significant effects (two-tailed), according to a nonparametric cluster correction procedure (Maris & Oostenveld, 2007). Grey shaded regions represent bootstrapped 95% confidence intervals calculated across participants, and the period when the stimulus was displayed (lower rectangle)

4.4 Multiplicative noise

An alternative account of contrast discrimination performance at high pedestal contrasts is that transduction is linear but internal noise is signal-dependent (Pelli, 1985). If the dominant source of noise were early and multiplicative, this would avoid any issues relating to Birdsall's theorem, as the transducer could be linear. It has proven difficult to distinguish between the multiplicative and additive noise accounts purely from contrast discrimination experiments (Georgeson & Meese, 2006; Kontsevich, Chen, & Tyler, 2002). At a single neuron level there is well-established evidence of multiplicative noise (Tolhurst, Movshon, & Dean, 1983), yet it appears that across populations of neurons with different sensitivities the overall noise is effectively additive (Chen, Geisler, & Seidemann, 2006). Since evidence from fMRI (Boynton et al., 1999), EEG (Busse et al., 2009) and psychophysics (Kingdom, 2016) all argue strongly against a linear transducer, we think this explanation is unlikely to account for the body of available data.

4.5 Conclusion

To summarise, in this study we investigated the timecourse of the neural operations involved in contrast discrimination. We demonstrated that internal noise impacting early in time and in the visual pathway can affect sensory processing and perceptual decisions. Our novel application of multivariate analysis methods to spatially-parcellated MEG source space

representations offers the capability of studying how the brain represents information in both space and time.

5 References

Avidan, G., Harel, M., Hendler, T., Ben-Bashat, D., Zohary, E., & Malach, R. (2002). Contrast sensitivity in human visual areas and its relationship to object recognition. *Journal of Neurophysiology*, *87*(6), 3102–3116.

Baker, D. H., & Wade, A. R. (2017). Evidence for an Optimal Algorithm Underlying Signal Combination in Human Visual Cortex. *Cerebral Cortex*, *27*, 254–264. http://doi.org/10.1093/cercor/bhw395

Barlow, H. B. (1962). A method of determining the over-all quantum efficiency of visual discriminations. *The Journal of Physiology*, *160*, 155–168.

Barlow, H. B., Hawken, M., Parker, A. J., & Kaushal, T. P. (1987). Human contrast discrimination and the threshold of cortical neurons. *Journal of the Optical Society of America A*, *4*(12), 2366. http://doi.org/10.1364/JOSAA.4.002366

Boynton, G. M., Demb, J. B., Glover, G. H., & Heeger, D. J. (1999). Neuronal basis of contrast discrimination. *Vision Research*, *39*(2), 257–269. http://doi.org/10.1016/S0042-6989(98)00113-8

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*, 433–436.

Britten, K. H., Newsome, W. T., Shadlen, M. N., Celebrini, S., & Movshon, J. A. (1996). A relationship between behavioral choice and the visual responses of neurons in macaque MT. *Visual Neuroscience*, *13*(01), 87–100. http://doi.org/10.1017/S095252380000715X

Britten, K. H., Shadlen, M. N., Newsome, W. T., & Movshon, J. A. (1992). The analysis of visual motion: a comparison of neuronal and psychophysical performance. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *12*(12), 4745–4765.

Busse, L., Ayaz, A., Dhruv, N. T., Katzner, S., Saleem, A. B., Scho, M. L., … Carandini, M. (2011). The Detection of Visual Contrast in the Behaving Mouse, *31*(31), 11351–11361. http://doi.org/10.1523/JNEUROSCI.6689-10.2011

Busse, L., Wade, A. R., & Carandini, M. (2009). Representation of Concurrent Stimuli by Population Activity in Visual Cortex. *Neuron*, *64*(6), 931–942. http://doi.org/10.1016/j.neuron.2009.11.004

Cabrera, C. A., Lu, Z.-L., & Dosher, B. A. (2015). Separating decision and encoding noise in signal detection tasks. *Psychological Review*, *122*(3), 429–460. http://doi.org/10.1037/a0039348

Campbell, F. W., & Kulikowski, J. J. (1972). The visual evoked potential as a function of contrast of a grating pattern. *The Journal of Physiology*, *222*(2), 345–356. http://doi.org/10.1113/jphysiol.1972.sp009801

Carandini, M. (2004). Amplification of trial-to-trial response variability by neurons in visual cortex. *PLoS Biology*, *2*(9). http://doi.org/10.1371/journal.pbio.0020264

Chen, Y., Geisler, W. S., & Seidemann, E. (2006). Optimal decoding of correlated neural population responses in the primate visual cortex. *Nature Neuroscience*, *9*(11), 1412–1420. http://doi.org/10.1038/nn1792

Cichy, R. M., Pantazis, D., & Oliva, A. (2014). Resolving human object recognition in space and time. *Nature Neuroscience*, *17*(3), 455–462. http://doi.org/10.1038/nn.3635

Clarke, A., Devereux, B. J., Randall, B., & Tyler, L. K. (2015). Predicting the Time Course of Individual Objects with MEG. *Cerebral Cortex*, *25*(10), 3602–3612.

http://doi.org/10.1093/cercor/bhu203

Fechner, G. T. (1912). The classical psychologists: Selections illustrating psychology from Anaxagoras to Wundt. In B. Rand (Ed.), *Elements of psychophysics* (pp. 562–572). Boston: Houghton Miffin.

Georgeson, M. A., & Meese, T. S. (2006). Fixed or variable noise in contrast discrimination? The jury's still out.. *Vision Research*, *46*(25), 4294–4303. http://doi.org/10.1016/j.visres.2005.08.024

Graham, N. V. (2011). Beyond multiple pattern analyzers modeled as linear filters (as classical V1 simple cells): useful additions of the last 25 years. *Vision Research*, *51*(13), 1397–1430. http://doi.org/10.1016/j.visres.2011.02.007

Grill-spector, K., Henson, R., & Martin, A. (2006). Repetition and the brain : neural models of stimulus-specific effects, *10*(1), 17–19. http://doi.org/10.1016/j.tics.2005.11.006

Hecht, S., & Wald, G. (1934). The visual acuity and intensity discrimi- nation of drosophila. *The Journal of General Physiology*, *17*(4), 517–547.

Hesselmann, G., Kell, C. A., Eger, E., & Kleinschmidt, A. (2008). Spontaneous local variations in ongoing neural activity bias perceptual decisions. *Proceedings of the National Academy of Sciences of the United States of America*, *105*(31), 10984–10989. http://doi.org/10.1073/pnas.0712043105

Hillyard, S. A., Squires, K. C., Bauer, J. W., & Lindsay, P. H. (1971). Evoked potential correlates of auditory signal detection. *Science (New York, N.Y.)*, *172*(3990), 1357–1360.

Jolij, J., Meurs, M., & Haitel, E. (2011). Why do we see what's not there? *Communicative & Integrative Biology*, *4*(6), 764–767.

King, J., & Dehaene, S. (2014). Characterizing the dynamics of mental representations : the temporal generalization method, *18*(4), 203–210. http://doi.org/10.1016/j.tics.2014.01.002.Characterizing

Kingdom, F. A. A. (2016). Fixed versus variable internal noise in contrast transduction: The significance of Whittle's data. *Vision Research*, *128*, 1–5. http://doi.org/10.1016/j.visres.2016.09.004

Klein, A., Ghosh, S. S., Bao, F. S., Giard, J., Häme, Y., Stavsky, E., … Keshavan, A. (2017). Mindboggling morphometry of human brains. *PLOS Computational Biology*, *13*(2), e1005350. http://doi.org/10.1371/journal.pcbi.1005350

Klein, S. A., & Levi, D. M. (2009). Stochastic model for detection of signals in noise. *Journal of Optical Society of America*, *26*(11), 110–126.

Kleiner, M., Brainard, D. H., & Pelli, D. G. (2007). What's new in Psychtoolbox-3? *Perception*, *36*(S), 14.

Kontsevich, L. L., Chen, C.-C., & Tyler, C. W. (2002). Separating the effects of response nonlinearity and internal noise psychophysically. *Vision Research*, *42*(14), 1771–1784.

Legge, G. E., & Foley, J. M. (1980). Contrast masking in human vision. *Journal of the Optical Society of America*, *70*(12), 1458–1471. http://doi.org/10.1364/JOSA.70.001458

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data. *Journal of Neuroscience Methods*, *164*(1), 177–190. http://doi.org/10.1016/j.jneumeth.2007.03.024

Michelson, C. A., Pillow, J. W., & Seidemann, E. (2017). Majority of choice-related variability in perceptual decisions is present in early sensory cortex. *Preprint*.

Mostert, P., Kok, P., & de Lange, F. P. (2016). Dissociating sensory from decision processes in human perceptual decision making. *Scientific Reports*, *5*(1). http://doi.org/10.1038/srep18253

Mueller, S. T., & Weidemann, C. T. (2008). Decision noise: An explanation for observed violations of signal detection theory. *Psychonomic Bulletin & Review*, *15*(3), 465–494. http://doi.org/10.3758/PBR.15.3.465

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open Source Software for Advanced Analysis of MEG, EEG, and Invasive Electrophysiological Data. *Computational Intelligence and Neuroscience*, *2011*, 1–9. http://doi.org/10.1155/2011/156869

Peirce, J. W. (2007). PsychoPy—Psychophysics software in Python. *Journal of Neuroscience Methods*, *162*(1–2), 8–13. http://doi.org/10.1016/j.jneumeth.2006.11.017

Pelli, D. G. (1985). Uncertainty explains many aspects of visual contrast detection and discrimination. *Journal of the Optical Society of America. A, Optics and Image Science*, *2*(9), 1508–1532. http://doi.org/10.1364/JOSAA.2.001508

Pelli, D. G. (1997). The VideoToolbox software for visual psychophysics: Transforming numbers into movies. *Spatial Vision*, *10*(437–442).

Ress, D., & Heeger, D. J. (2003). Neuronal correlates of perception in early visual cortex. *Nature Neuroscience*, *6*(4), 414–420. http://doi.org/10.1038/nn1024

Roelfsema, P. R., & Spekreijse, H. (2001). The representation of erroneously perceived stimuli in the primary visual cortex. *Neuron*, *31*(5), 853–863. http://doi.org/10.1016/S0896-6273(01)00408-1

Rolls, E. T., & Baylis, G. C. (1986). Size and contrast have only small effects on the responses to faces of neurons in the cortex of the superior temporal sulcus of the monkey. *Experimental Brain Research*, *65*(1), 38–48.

Schölvinck, M. L., Friston, K. J., & Rees, G. (2012). The influence of spontaneous activity on stimulus processing in primary visual cortex. *NeuroImage*, *59*, 2700–2708. http://doi.org/10.1016/j.neuroimage.2011.10.066

Smith, R. A., & Swift, D. J. (1985). Spatial-frequency masking and Birdsall's theorem. *Journal of the Optical Society of America. A, Optics and Image Science*, *2*(9), 1593–1599.

Squires, N. K., Squires, K. C., & Hillyard, S. A. (1975). Two varieties of long-latency positive waves evoked by unpredictable auditory stimuli in man. *Electroencephalography and Clinical Neurophysiology*, *38*(4), 387–401.

Tadel, F., Baillet, S., Mosher, J. C., Pantazis, D., & Leahy, R. M. (2011). Brainstorm: A User-Friendly Application for MEG/EEG Analysis. *Computational Intelligence and Neuroscience*, *2011*, 1–13. http://doi.org/10.1155/2011/879716

Tolhurst, D. J., Movshon, J. A., & Dean, A. F. (1983). The statistical reliability of signals in single neurons in cat and monkey visual cortex. *Vision Research*, *23*(8), 775–785.

Woldorff, M. (1993). Distortion of ERP averages due to overlap from temporally adjacent ERPs: Analysis and correction. *Psychophysiology*, *30*, 98–119.

6 Appendices

Table A1: Numbers of vertices on the cortical mesh. Individual regions were taken from a mesh consisting of around 3000 vertices, and pooled across hemispheres. The 'whole brain' mesh (final row) was subsampled to around 500 vertices. Precise numbers of vertices varied across individual participants owing to individual differences in brain size and morphology. Entries in the 'Colour' column correspond to the colours used in Figures 6, A2 & A3.

Region	Colour	Mean size	Minimum size	Maximum size
Rostral Middle Frontal		157	145	173
Superior Frontal		342	317	384
Rostral Anterior Cingulate		31	27	37
Lateral Orbitofrontal		100	83	113
Medial Orbitofrontal		53	45	62
Pars Triangularis		63	56	69
Pars Orbitalis		31	27	35
Caudal Anterior Cingulate		29	25	34
Pars Opercularis		53	43	61
Caudal Middle Frontal		75	59	91
Insula		60	53	69
Entorhinal		16	9	25
Pre-central		140	124	155
Superior Temporal		166	150	183
Posterior Cingulate		38	32	45
Transverse Temporal		9	6	11
Post-central		142	130	156
Para-central		48	41	61
Middle Temporal		134	123	152
Parahippocampal		21	17	25
Inferior Temporal		118	96	149
Supramarginal		123	102	155
Isthmus Cingulate		28	23	34
Fusiform		81	73	87
Precuneous		119	93	140
Superior Parietal		148	136	168
Inferior Parietal		149	131	157
Lingual		101	69	125
Cuneus		67	58	74
Peri-calcarine		38	24	45
Lateral Occipital		162	139	181
Whole brain		503	503	504

Figure A1: Interval-based MVPA analysis for all target contrast conditions, and for both contrast-based decoding (a) and decision-based decoding (b). Plotting conventions are consistent with Figure 4b,c.

Figure A2: Maximal evoked responses in different anatomical regions. Each trace in panel (a) plots the timecourse of the vertex in the named region (see legend in panel (b)) with the largest absolute deflection from baseline. Panel (c) shows absolute activity averaged across four time windows, demonstrating that the majority of activity occurs in occipito-temporal regions.

.

Figure A3: Atlas-based classification of decisions in the 16% target condition. Plotting conventions mirror those of Figure 6.