A novel interplay between GEFs orchestrates Cdc42 activation in cell polarity and cytokinesis

Brian Hercyk, Julie Rich, and Maitreyi Das[§]

Department of Biochemistry & Cellular and Molecular Biology, University of Tennessee, Knoxville, TN, USA.

§ Corresponding Author: mdas@utk.edu

ABSTRACT

The small GTPase Cdc42, a conserved regulator of cell polarity in eukaryotes, is activated by two GEFs, Gef1 and Scd1, in fission yeast. Gef1 and Scd1 localize sequentially to the division site to activate Cdc42 for efficient cytokinesis. The significance of multiple Cdc42 GEFS is not well understood. Here we report a novel interplay between Gef1 and Scd1 that fine-tunes Cdc42 activation during two cellular programs: cytokinesis and polarized growth. Gef1 promotes Scd1 localization to the division site during cytokinesis. During polarized growth, Gef1 is required for bipolar Scd1 localization. We find that Gef1 recruits Scd1 through the recruitment of the scaffold Scd2; we propose this facilitates polarized cell growth at a second site. In turn, Scd1 restricts Gef1 localization to the division site and to the cell cortex, thus maintaining polarity. Our results suggest that crosstalk between GEFs is a conserved mechanism that orchestrates Cdc42 activation during complex processes.

INTRODUCTION

Growth and division are fundamental processes of all cells, and are essential for proper function and proliferation. In most multicellular organisms, these two processes are precisely tuned to control cell shape and function, to specify cell fate and differentiation, and to enable cell adhesion and migration(Feigin and Muthuswamy, 2009; Godde et al., 2010; Halaoui and McCaffrey, 2014; Lauffenburger and Horwitz, 1996). These properties are dependent on proper cell polarization. Cell polarization relies on the ability of the cytoskeleton to establish unique domains at the cell cortex that govern the local function and activity of specific proteins (Drubin and Nelson, 1996; Nance and Zallen, 2011). The Rho family of small GTPases serves as the primary regulator of the actin cytoskeleton, and thereby cell polarity and movement (Ridley, 2006). Active Rho GTPases bind and activate downstream targets which regulate actin cytoskeleton organization. GTPases are active when GTP-bound and inactive once they hydrolyze GTP to GDP. GTPases are activated by Guanine nucleotide Exchange Factors, or GEFs, which promote GTP binding, while GTPase Activating Proteins, or GAPs, inactivate GTPases by promoting GTP hydrolysis (Bos et al., 2007). Unraveling the regulation of these GEFs and GAPs is at the crux of understanding how cell polarity is established, maintained, and altered. One conserved member of the Rho family of small GTPases, Cdc42, is a master regulator of cell polarity in eukaryotes(Etienne-Manneville, 2004; Johnson, 1999). In most eukaryotes, Cdc42 is regulated by numerous GEFs and GAPs, which complicates our understanding of GTPase regulation (Bos et al., 2007). In the fission yeast *Schizosaccharomyces pombe,* Cdc42 is activated by two GEFs, Gef1 and Scd1 (Chang et al., 1994; Coll et al., 2003). The presence of only two Cdc42 GEFs, and the well-documented process of cell polarization makes fission yeast an excellent model system to understand the mechanistic details of cell shape establishment. Here we report that the two Cdc42 GEFs regulate each other in fission yeast. This finding provides new insights into the spatiotemporal regulation of Cdc42 during critical cellular events.

Cdc42, like other small GTPases, serves as a binary molecular switch and can respond to and initiate multiple signaling pathways. In the budding yeast *Saccharomyces cerevisiae*, cells develop only a single polarized site through a winner-take-all mechanism during bud emergence (Irazoqui et al., 2003; Kozubowski et al., 2008; Slaughter et al., 2009a; Wedlich-Soldner et al., 2004). However, a winner-take-all mechanism cannot explain how cells develop multiple polarized sites often observed in higher eukaryotes. In contrast, fission yeast grows in a bipolar manner, offering a model to understand how a cell regulates polarized growth from multiple sites. In fission yeast, active Cdc42 displays anti-correlated oscillations between the two ends (Das et al., 2012). These oscillations arise from both positive and time-delayed negative feedback as well as competition between the two ends (Das et al., 2012). This oscillatory pattern regulates cell dimensions and promotes bipolar growth in fission yeast. Similar Cdc42 oscillations have been observed in natural killer cells during immunological synapse formation (Carlin et al., 2011) and in budding yeast during bud emergence (Howell et al., 2012). In plant cells, the ROP GTPases show oscillatory behaviour during pollen tube growth (Hwang et al., 2005). Furthermore, during migration in animal cells, the GTPases Rho, Rac, and Cdc42 fluctuate and are sequentially activated to enable protrusion (Machacek et al., 2009). These

observations together suggest that oscillatory behavior may be an intrinsic property of GTPases that drives polarity, and which is likely conserved in most organisms (Das and Verde, 2013).

Most models of polarized growth propose the existence of Cdc42 positive feedback loops that facilitate symmetry breaking through a winner-take-all mechanism (Slaughter et al., 2009b). Our understanding of the molecular nature of these positive feedbacks is primarily based on studies performed in budding yeast (Kozubowski et al., 2008; Slaughter et al., 2009a; Slaughter et al., 2009b). In one model, Cdc42 activation via actin organization and membrane trafficking amplifies its own localization to the polarized tip (Wedlich-Soldner et al., 2004). In a second model, a ternary complex consisting of the GEF Cdc24, the scaffold Bem1, and the Pak kinase Cla4, amplifies Cdc42 activation at the cell's growth sites (Kozubowski et al., 2008). Both models propose that active Cdc42 participates in the generation of the positive feedback. In fission yeast, a similar ternary complex with the GEF Scd1, scaffold Scd2, and kinase Pak1 has been reported (Endo et al., 2003). However, it is not clear how this complex mediates a positive feedback loop during polarized growth.

To explain Cdc42 activation during polarized growth, it is important to first understand how Cdc42 regulators function. Gef1 and Scd1 are partially redundant but exhibit unique phenotypes when deleted (Chang et al., 1994; Coll et al., 2003), indicating that they may regulate Cdc42 in different manners. Cells lacking *scd1* appear depolarized (Chang et al., 1994), and Scd1 oscillates between the two cell ends, much like Cdc42 (Das et al., 2012). In contrast, *gef1* mutants become narrower and grow in a monopolar, rather than a bipolar, manner (Coll et al., 2003). Furthermore, Cdc42 activity is reduced at the new end in *gef1* mutants (Das et al., 2012). Gef1 shows sparse localization at the cortex, making it difficult to determine whether it oscillates between cell ends (Das et al., 2015). Understanding how two different Cdc42 GEFs yield distinct phenotypes will provide valuable insights into Cdc42 regulation.

Investigations into the behaviors of Gef1 and Scd1 are complicated since these GEFs overlap at sites of polarized growth. In addition to the ends, these GEFs also localize to the site of cell division during cytokinesis (Wei et al., 2016). Cytokinesis, the final step in cell division, involves the formation of an actomyosin ring that constricts concurrent with cell wall (septum) deposition to enable membrane ingression and furrow formation (Pollard, 2010). The temporal localization and function of the two GEFs are discernible during cytokinesis since they are recruited to the division site in succession to activate Cdc42. During cytokinesis, Gef1 localizes first to the actomyosin ring to activate Cdc42 and promote ring constriction (Wei et al., 2016). Scd1 then localizes to the ingressing membrane and regulates septum formation (Wei et al., 2016). The temporal difference between Gef1 and Scd1 localization at the division site allows us to investigate the significance of multiple GEFs in Cdc42 regulation, which is unclear from studies solely at the growing ends.

Using cytokinesis as a paradigm, here we identify a novel crosstalk between the GEFs, Gef1 and Scd1, that regulates Cdc42 activity during multiple cellular programs. We find that Gef1 and Scd1 regulate each other during both cytokinesis and cell polarization. Our data indicates that Gef1 promotes the localization of Scd1 to the division site. Contrary to previously proposed models, constitutively active Cdc42 is not sufficient to rescue Scd1 localization in *gef1* mutants. Instead, we find that Gef1 promotes the localization of the scaffold Scd2 to the division site

during cytokinesis, which then recruits Scd1. Next, we show that Scd1 promotes the removal of Gef1 from the division site after completion of ring constriction. Furthermore, we show that actin cables are involved in Gef1 removal from the division site, suggesting that Scd1 promotes Gef1 removal via an actin-mediated process. We extend these observations to the sites of polarized growth, where we show that Gef1 promotes bipolar Scd1 and Scd2 localization. We report that Gef1 is necessary to recruit Scd1 to the non-dominant pole to initiate bipolar growth. In turn, Scd1 and actin are necessary to prevent isotropic localization of Gef1 at the cell cortex during interphase, thus maintaining polarity. By this manner of regulation, Cdc42 activation is promoted at nascent sites but restricted from random regions of the cell. Gef1 allows the establishment of a nascent site through the recruitment of Scd1, while Scd1 prevents ectopic Gef1 localization. To the best of our knowledge, such a crosstalk has not been reported to function between GEFs of the same GTPase. The interplay between the Cdc42 GEFs operates in the same manner during both cytokinesis and polarized growth, suggesting that this may be a conserved feature of Cdc42 regulation.

RESULTS

Gef1 promotes Scd1 recruitment to the division site

We have reported that Gef1 localizes to the assembled actomyosin ring before Scd1 (Wei et al., 2016). It is not clear why the cell recruits two distinct GEFs to the same site in a sequential manner, given that they both activate Cdc42. Previous reports have demonstrated crosstalk between GTPases via the modulation of their regulators (Guilluy et al., 2011). While there is no report of GEFs of the same GTPase regulating each other, such an interaction may explain the temporal relationship detected between Gef1 and Scd1 localization at the division site. Since Scd1 arrives at the division site soon after Gef1, we posited that Gef1 may promote Scd1 localization. To test this, we examined whether Scd1 localization to actomyosin rings is Gef1-dependent. Both Gef1 and Scd1 are low-abundance proteins and are not suitable for live cell imaging over time. This complicates the investigation of these proteins in a temporal manner. To overcome this limitation, we used the actomyosin ring as a temporal marker. The actomyosin ring undergoes visibly distinct phases during cytokinesis: assembly, maturation, constriction, and disassembly. We determined the timing of protein localization to the division site by comparing it to the corresponding phase of the actomyosin ring. We have previously reported that ring constriction is delayed in *gef1Δ* mutants (Wei et al., 2016). To eliminate any bias in protein localization due to this delay, we only analyzed cells in which the rings had initiated constriction. In *gef1Δ* mutants, the number of constricting rings that recruited Scd1-3xGFP decreased to 15% (Figure 1A-B) from 96% in *gef1+* (p<0.0001). Furthermore, the *gef1Δ* cells that managed to recruit Scd1-3xGFP did not do so as efficiently as *gef1+* cells, given the 15% decrease in Scd1-3xGFP fluorescence intensity at the division site (Figure 1A,C, p=0.0098). This indicates that Gef1 promotes Scd1 localization to the division site.

To better understand how Gef1 recruits Scd1 to the division site, we explored previous reports of GEF recruitment. GEF recruitment to sites of Cdc42 activity occurs via positive feedback, as reported in budding yeast (Butty et al., 2002; Kozubowski et al., 2008). In this model, activation of Cdc42 leads to further recruitment of the scaffold Bem1, which then recruits the GEF Cdc24 to the site of activity, thus helping to break symmetry and promote polarized growth. A similar positive feedback may also exist in fission yeast (Das et al., 2012; Das and Verde, 2013). We hypothesized that once Cdc42 is activated by Gef1, it acts as a seed for Scd1 recruitment to the division site. To test this, we asked whether constitutive activation of Cdc42 could rescue the Scd1 recruitment defect exhibited by *gef1Δ*. In order for this approach to work, the constitutively active Cdc42 must localize to the division site. Localization of active Cdc42 is visualized via the bio-probe CRIB-3xGFP that specifically binds GTP-Cdc42. Since our previous work reported that Cdc42 activity is reduced at the division site in *gef1Δ* cells (Wei et al., 2016) we validated this approach by first testing whether constitutively active Cdc42 restores CRIB-3xGFP localization at the division site in *gef1Δ* cells. The empty control vector or the vector expressing the constitutively active allele *cdc42G12V* under the control of the thiamine-repressible *nmt41* promoter was integrated into the genome of *gef1+* and *gef1Δ* cells expressing CRIB-3xGFP. Mild induction of *cdc42G12V* expression was sufficient to restore CRIB-3xGFP intensity at the division site to physiological levels in *gef1Δ*, but not in *gef1Δ* with the control vector (Figure 1D,F, p<0.0001). Surprisingly, although expression of *cdc42G12V* was able to restore Cdc42 activity at the division site in *gef1Δ* cells, it was unable to rescue Scd1-3xGFP localization to the

division site in *cdc42G12V gef1Δ* (Figure 1E,G). This indicates that active Cdc42 alone is not sufficient to recruit Scd1, and that Gef1 is required for this process.

Gef1 promotes localization of Scd2 to the division site, which in turn recruits Scd1

Next we asked if other members of the Cdc42 complex are involved in the recruitment of Scd1. The Cdc42 ternary complex consists of the GEF Scd1, the scaffold protein Scd2, and the downstream effector Pak1 kinase (Endo et al., 2003). Observations in budding yeast suggest that the PAK kinase may mediate GEF recruitment (Kozubowski et al., 2008). Contrary to this hypothesis, we find that Scd1-3xGFP intensity increases in the *nmt1 switch-off* mutant allele of *pak1*, relative to *pak1+* cells (Figure S1). These findings support similar observations reported in the hypomorphic temperature-sensitive *pak1* allele, *orb2-34* (Das et al., 2012).

Previous reports have shown that the scaffold Scd2 is required for Scd1 localization to the sites of polarized growth (Kelly and Nurse, 2011). We hypothesized that Gef1 recruits Scd1 to the division site through the scaffold Scd2. To test this, we examined whether Scd2-GFP localization to the division site is Gef1-dependent. *gef1Δ* cells displayed a significant decrease in the number of assembled rings that recruited Scd2-GFP relative to *gef1+*. In *gef1Δ* mutants, the number of rings that recruited Scd2-GFP prior to ring constriction decreased to 8% relative to 88% in *gef1+*, indicating a delay in Scd2 recruitment (Figure 2A-B, p>0.0001). Although *gef1Δ* cells managed to recruit Scd2 to the division site once ring constriction began, the fluorescence intensity of Scd2-GFP at the division site was reduced by 61% relative to *gef1+* cells (Figure 2A,C, p>0.0001, Fig. S2). This indicates that Gef1 promotes Scd2 localization to the division site.

Since a previous report suggested that Scd1 and Scd2 require each other for their localization (Kelly and Nurse, 2011), we asked if a decrease in Scd1 at the division site in *gef1* mutants results in a reciprocal decrease in Scd2 localization. Contrary to this previous report, we find that Scd2-GFP localization at the division site is not impaired in *scd1Δ* cells (Figure 2E). In contrast, Scd1-3xGFP localization is completely abolished at the division site in *scd2Δ* cells (Figure 2E). We find that while Scd1 requires Scd2 for its localization to the division site, Scd2 localization is independent of Scd1. Altogether, this indicates that Gef1 promotes Scd2 localization to the division site, which is required for Scd1 localization. To further validate these findings, we examined the temporal localization of Gef1, Scd1, and Scd2 to the division site. A well-established temporal marker for cells in cytokinesis is the distance between the spindle pole bodies. The spindle pole body distance increases as mitosis progresses until the cell reaches anaphase B (Nabeshima et al., 1998), at which time the actomyosin ring starts to constrict (Wu et al., 2003). The distance between the two spindle pole bodies can thus act as an internal clock that helps to time the recruitment of other proteins. We acquired numerous still images and calculated the distance between the spindle pole bodies, marked by Sad1-mCherry, during anaphase A or anaphase B. We report the spindle pole body distance at which Gef1-mNG (monomeric NeonGreen), Scd1-3xGFP, and Scd2-GFP signals are visible at the non-constricting actomyosin ring (Figure 2F). Next, we calculated the mean spindle pole body distance of the first 50th percentile of our data. The protein that localizes earliest to the

actomyosin ring during mitosis will display the smallest mean spindle pole body distance. We find that Gef1-mNG localized to the actomyosin ring with a mean spindle pole body distance of 3.2μm, Scd2-GFP with a mean distance of 4.1μm and Scd1-3xGFP with a mean distance of 5.1μm (Figure 2G). This suggests that Gef1-mNG is recruited to the actomyosin ring first, followed by Scd2-GFP, and finally Scd1-3xGFP. The sequence in which these proteins localize to the division site agrees with our experiments, which indicate that Gef1 recruits Scd1 indirectly through Scd2.

Scd1 promotes Gef1 removal from the division site at the end of ring constriction

Once the actomyosin ring constricts, Gef1 constricts with it and is lost from the division site when the ring disassembles (Wei et al., 2016). At this stage Scd1 is still localized to the membrane barrier. Since our data indicate that Gef1 promotes Scd1 localization, we asked if Scd1 mediates Gef1 localization to the division site. We did not detect any aberrant Gef1 behavior during early cytokinetic events in cells lacking *scd1*. However, at the end of ring constriction, we observed prolonged Gef1 localization in *scd1* mutants. In *scd1+* cells, Gef1 localizes to the membrane adjacent to the ring throughout constriction. In cells that have completed constriction, Gef1 is lost as the ring disassembles (Figure 3A). In *scd1Δ* mutants, after completion of ring constriction and disassembly, Gef1 remains at the membrane that was adjacent to the ring. In 70% of *scd1Δ* cells, post-ring-disassembly, Gef1-mNG persists at the newly formed membrane barrier, as confirmed by the absence of Rlc1-tdTomato (Figure 3A-B). Similar Gef1-mNG localization was observed in only 20% of *scd1+* cells (Figure 3B, p<0.0001).

To understand how Scd1 mediates Gef1 removal from the membrane barrier after constriction, we analyzed the phenotype of *scd1Δ* mutants. We find that the actin cytoskeleton is disrupted in *scd1Δ* cells, as observed by Alexa Fluor Phalloidin staining. *scd1Δ* cells accumulate actin patches and have fewer and more disorganized actin cables (Figure S3). Therefore, we next looked at the role of actin in Gef1 removal after constriction. We treated the cells with Latrunculin A (LatA) to disrupt the actin cytoskeleton. In LatA treated cells that were fully septated following completion of constriction, we observed persistent Gef1 localization at the division site. Gef1-mNG persists on both sides of the septum barrier in 40% of cells treated with LatA, but not in mock DMSO-treated cells (Figure 3C). Cells undergoing ring constriction and septum formation display actin cables as well as Arp2/3-complex-dependent patches at the division site (Coffman et al., 2013; Gachet and Hyams, 2005; Huang et al., 2012; Wang et al., 2016). LatA treatment removes all types of filamentous actin structures (Spector et al., 1983). To determine which actin-mediated process regulates Gef1 removal, we treated cells with CK666 to block only Arp2/3-mediated branched actin filaments (Sun et al., 2011). In these cells, Gef1-mNG removal was unhindered, as in DMSO-treated control cells, or localized to random sites along the cortex, but did not persist at the division site (Figure 3C). This suggests that Gef1 removal at the end of ring constriction is independent of branched actin. We next examined the role of filamentous actin cables in the removal of Gef1 from the membrane barrier. Since these cables are made by the formin For3 (Feierbach and Chang, 2001), we investigated whether Gef1 localization at the membrane barrier is *for3*-dependent. We find that in *for3Δ*, Gef1-3xYFP lingers at the membrane adjacent to the ring after completion of

constriction, just as in *scd1Δ.* To determine whether Gef1 removal by Scd1 and by actin operates in the same or parallel pathways, we treated *scd1+* and *scd1Δ* cells expressing Gef1-mNG with LatA. We find that in cells treated with DMSO, Gef1-mNG persists in 20% of septated *scd1+* cells and in 63% of septated *scd1Δ* cells. In cells treated with LatA, Gef1-mNG persists in 40% of septated *scd1+* cells and in 61% of septated *scd1Δ* cells (Figure 3D). The extent of Gef1 persistence in *scd1Δ* cells therefore does not increase with the addition of LatA, indicating that Scd1 is epistatic to actin-mediated removal (Figure 3D). Together, these data indicate that Scd1 removes Gef1 from the division site after ring disassembly through an actin-mediated process involving the formin For3.

Gef1 is required for bipolar Scd1 localization

Our data indicate an interesting interplay between the two Cdc42 GEFs in which they regulate each other's localization during cytokinesis. We inquired whether this novel interaction is intrinsic to the regulation of Cdc42 in other cellular processes. Cdc42 and its GEFs play a supporting role in cytokinesis, but are central players in the regulation of polarized growth. Thus, we asked whether a similar interaction occurs at sites of polarized growth. Gef1 promotes bipolar growth in fission yeast (Coll et al., 2003; Das et al., 2012). Cells lacking *gef1* show increased monopolarity, with polarized growth only occurring at the old end. In fission yeast cells, Scd1 localizes to sites of polarized growth (Das et al., 2009; Kelly and Nurse, 2011) accordingly, cells in early G2 phase display Scd1 localization at the old end. Cells in late G2, which have undergone new-end-take-off (NETO), grow in a bipolar manner and display Scd1 localization at both the old and new end of the cell. We inquired whether Gef1 promotes Scd1 localization at sites of polarized growth. Scd1, like active Cdc42, undergoes oscillations between the two competing ends (Das et al., 2012); thus, a cell undergoing bipolar growth does not always display bipolar Scd1 localization. We found that Scd1-3xGFP levels at the old end were comparable in *gef1+* and *gef1Δ* cells (Figure S4). However, Scd1-3xGFP did not localize to both ends during interphase in *gef1Δ* cells. We observed bipolar Scd1-3xGFP in 30% of interphase *gef1+* cells, but only in 14% of *gef1Δ* cells (Figure 4A,B, p=0.0004). Similarly, we also observed a decrease in bipolar Scd2 in cells lacking *gef1*; 70% of *gef1+* cells displayed bipolar Scd2-GFP localization, but this was reduced to 30% in *gef1Δ* cells (Figure 4A,B, p<0.0001). Similar to what was reported for the site of cell division, Scd2 is required for Scd1 localization to sites of polarized growth, but Scd2 localization is independent of Scd1 (Figure 4F). In *scd1Δ* mutants, Scd2-GFP signal was observed either at cell ends or ectopically at the cell cortex. In contrast, *scd2Δ* mutants failed to localize Scd1-3xGFP to the cell cortex, rather it was observed within the cytoplasm or in the nucleus. This indicates that Gef1 promotes Scd2 localization, which in turn recruits Scd1 to sites of polarized growth.

Next, we asked whether active Cdc42 can restore bipolar Scd1 localization in *gef1Δ* cells. To test this, we first checked to see whether expression of constitutively active Cdc42 results in bipolar localization of active Cdc42, as indicated by CRIB-3xGFP localization. Low-level expression of *cdc24G12V* was sufficient to restore bipolar CRIB-3xGFP localization in *gef1Δ*, relative to the empty-vector-containing *gef1Δ* mutants (Figure 4B,E, p<0.0001). We observed bipolar CRIB-3xGFP in 75% of *gef1+* cells transformed with the empty vector and in 93% of

cells expressing *cdc14G12V*. In *gef1Δ* mutants transformed with the empty vector, we observed CRIB-3xGFP in only 50% of cells. In contrast, in *gef1Δ* mutants, low levels of *cdc42G12V* expression restored bipolar CRIB-3xGFP in 92% of cells. Next we investigated whether *cdc42G12V* restored bipolar Scd1-3xGFP localization in *gef1Δ* cells. Expression of *cdc42G12V* was unable to restore bipolar Scd1-3xGFP localization to the cell ends in *gef1Δ* mutants, just as it did not rescue Scd1 localization to the division site. We observed bipolar Scd1-3xGFP in 28% of *gef1+* cells transformed with the empty vector, and in 31% of cells expressing *cdc42G12V*. In *gef1Δ* mutants transformed with the empty vector, we observed bipolar Scd1-3xGFP in only 12.5% of cells. Further, in *gef1Δ* mutants expressing low levels of *cdc42G12V*, bipolar Scd1-3xGFP remained in only 12.6% of cells (Figure 4C,E). This indicates that *cdc42G12V*, while sufficient to restore bipolar growth, cannot promote bipolar Scd1 localization in the absence of *gef1*. This further demonstrates that Gef1 is required for bipolar Scd1 localization.

Gef1 establishes polarized growth at a nascent site

Our data suggest that Gef1 promotes bipolar growth in fission yeast by enabling bipolar Scd1 localization. However, previous reports have shown that while *gef1Δ* mutants are mainly monopolar, about 40**%** of interphase cells show bipolar growth (Figure 5B) (Coll et al., 2003; Das et al., 2012; Das et al., 2015). If Gef1 is required for Scd1 localization to the new end, how does bipolar growth occur in some *gef1Δ* cells? To address this, we investigated the nature of bipolar growth in *gef1Δ* mutants. Fission yeast cells have an old end that existed in the previous generation and a new end that was formed as a result of cell division. The old end initiates growth immediately after completion of division and cell separation. As the cell grows, it eventually initiates growth at the new end, resulting in bipolar growth (Figure 5A) (Mitchison and Nurse, 1985). The two ends in fission yeast compete for active Cdc42, and initially the old end wins this competition (Das et al., 2012). The old end can thus be said to be dominant over the new end in a newborn cell, and always initiates growth first. The new end must overcome the old end's dominance in order to initiate its own growth.

We find that 68% of monopolar *gef1Δ* mutant cells exhibit a growth pattern in which one daughter cell is monopolar and the other daughter cell is prematurely bipolar (Figures 5B and S5). In monopolar *gef1Δ* cells, growth predominantly occurs at the old end, which grew in the previous generation (Figures 5B and S5). In these monopolar cells, the new end frequently fails to grow since it cannot overcome the old end's dominance. The daughter cell that inherits its parent cell's non-growing end typically displays precocious bipolar growth, suggesting that these cells do not contain a dominant end. Our data suggest that for a cell end to be dominant it needs to have grown in the previous generation. These results indicate that the new ends of *gef1* cells are not well-equipped to overcome old end dominance. Indeed, we find that in *gef1+* cells, 97% of daughter cells derived from a growing end display a normal growth pattern in which new end take-off occurs only after the old end initiates growth (Figure 5C). In *gef1Δ* cells, only 9% of daughter cells derived from a growing end display the same pattern; instead, 81% of daughter cells derived from a growing end failed to initiate growth at their new end and were thus monopolar (Figure 5C). These observations indicate that to promote bipolar growth, Gef1 enables the new end to overcome old end dominance.

Taken together, our findings suggest that Gef1 helps promote bipolar growth by enabling Scd1 localization to the new end. This is further supported by our previous observation that the hyperactive *gef1* mutant allele *gef1S112A* shows premature bipolar growth (Das et al., 2015). Gef1 sparsely localizes to cell ends and instead remains mainly cytoplasmic. Gef1 is phosphorylated by the NDR kinase Orb6 (Das et al., 2009), resulting in a 14-3-3 binding site (Das et al., 2015). Interaction with a 14-3-3 protein sequesters Gef1 to the cytoplasm and away from the cortex. The *gef1S112A* mutation eliminates the Orb6 phosphorylation site, thus enabling excessive Gef1S112A localization to both cell ends, where it activates Cdc42 to promote premature bipolar growth (Das et al., 2015). Consistent with these findings, we report that Scd1 is significantly more bipolar in *gef1S112A* mutants. 53% of *gef1S112A* cells exhibit Scd1-3xGFP localization at both ends, compared to 32% in *gef1+* cells (Figure 5D-E, p<0.0001). These findings support the idea that Gef1 overcomes old end dominance by recruiting Scd1 to the new end to establish a nascent growth site.

Scd1 is required to restrict Gef1 localization to the cell ends

Next, we asked whether Scd1 and actin similarly regulate Gef1 at sites of polarized growth. Cells lacking *scd1* are round, and under a cell cycle arrest, these cells show polarized growth with increased cell width (Chang et al., 1994; Kelly and Nurse, 2011). We find that active Cdc42 appears depolarized in *scd1Δ* mutants during interphase. While CRIB-3xGFP remains restricted to the ends in *scd1+* cells, in *scd1Δ* mutants its localization appears depolarized with random patches all over the cortex (Figure 6B i and iii). We find that in *scd1+* cells, Gef1-mNG displayed sparse but polarized localization at cell ends (Figure 6A i). In *scd1Δ* mutants, Gef1-mNG showed better cortical localization when compared to *scd1+* cells (Fig. 6A iii). Further, Gef1-mNG showed depolarized cortical localization in *scd1Δ* mutants with random patches all over the cortex. This suggests that Scd1 is required to restrict Gef1 localization to the cell ends, thus maintaining polarized growth.

Since we find that actin plays a role in the removal of Gef1 from the division site (Figure 3C), we ask whether actin also regulates Gef1 localization at sites of polarized growth. We treated cells expressing Gef1-mNG with DMSO or LatA. Gef1-mNG localizes to the ends of control cells treated with DMSO. Upon LatA treatment, Gef1-mNG localizes to ectopic patches at the cortex (Figure 6A i and ii). Next, we analyzed Gef1-mNG localization in LatA-treated *scd1Δ* mutants. In *scd1Δ* mutants treated with either DMSO or LatA, we find that Gef1-mNG localizes to broad patches along the cortex (Figure 6A iii and iv). To determine if ectopic Gef1 at the cortex in *scd1Δ* mutants or LatA-treatment of cells results in ectopic Cdc42 activation, we analyzed CRIB-3xGFP localization in these cells. We find that in cells treated with LatA, CRIB-3xGFP localizes randomly to the cortex, signifying ectopic Cdc42 activation, similar to previous reports (Mutavchiev et al., 2016). In mock DMSO-treated control cells, CRIB-3xGFP forms caps at the growing ends (Fig. 6B, i). Upon treatment with LatA, CRIB-3xGFP localizes ectopically to diffuse cortical patches (Fig. 6B, ii). Similarly, CRIB-3xGFP localization appears as diffuse cortical patches in *scd1Δ* mutants (Fig. 6B, iii). If ectopic Cdc42 activation in LatA-treated cells occurs due to ectopic Gef1 localization, then loss of *gef1* should restore polarized Cdc42 activation in these cells. Indeed, in *gef1Δ* mutants CRIB-3xGFP remains polarized upon LatA treatment (Fig.

6B, vi). In *scd1Δ* mutants, CRIB-3xGFP localization appears ectopic in cells treated with either DMSO or LatA (Figure 6, iii and iv). Together, these data indicate that Scd1 and actin are required to prevent ectopic Gef1 localization and Cdc42 activation to maintain proper cell shape.

DISCUSSION

While Cdc42 is a major regulator of polarized cell growth, its regulation is not well understood, largely due to the presence of multiple activators and inhibitors that often function in a redundant manner. In fission yeast, Cdc42 is activated by only two GEFs, Gef1 and Scd1 (Chang et al., 1994; Coll et al., 2003). While these GEFs are partially redundant (Coll et al., 2003; Hirota et al., 2003), they display distinct phenotypes and it is unclear why the cell requires two Cdc42 GEFs. We have recently shown that Gef1 and Scd1 localize sequentially to the division site to activate Cdc42 during cytokinesis (Wei et al., 2016). Here we take advantage of the temporal difference between Gef1 and Scd1 localization at the division site to determine the significance of these two GEFs in Cdc42 regulation. We uncover a novel interplay between the Cdc42 GEFs that functions in both cytokinesis and polarized cell growth (Figure 7A). Given the conserved nature of Cdc42 and its regulators, we posit that this interplay between the GEFs is a common feature of Cdc42 regulation.

Crosstalk between Gef1 and Scd1 during cytokinesis

We have previously reported that Gef1 recruitment precedes Scd1 localization to the division site (Wei et al., 2016). Given that Scd1 appears to be the primary GEF, we asked whether the role of Gef1 is to recruit Scd1. Indeed, we report that Scd1 localizes to the division site in a Gef1-dependent manner (Figure 7B). We report that Scd1 is recruited by its scaffold Scd2, which is in turn recruited by Gef1. Furthermore, we show that while Scd1 localization is dependent on Scd2, the reciprocal is not true. Unlike the cell ends, the division site has no prior history of Cdc42 activation or Scd1 localization. Thus, it is possible that Scd1 recruitment to the division site requires assistance from Gef1, as is also seen at the nascent growth site at the new end.

Misregulation of Cdc42 has been reported to result in cytokinesis failure in many organisms. Specifically, failure to inactivate Cdc42 leads to failed cell abscission in budding yeast and HeLa cells, and prevents cellularization in *Drosophila* embryos (Atkins et al., 2013; Crawford et al., 1998; Dutartre et al., 1996; Onishi et al., 2013). The mechanism by which Cdc42 is inactivated prior to cell abscission has not been investigated in fission yeast. We have previously reported that Gef1 localization to the division site is lost after ring constriction (Wei et al., 2016). Here, we show that Scd1 promotes the clearance of Gef1 from the division site after ring disassembly (Figure 7A). This suggests that Scd1 ensures that Gef1 does not persist at the division site in the final stages of cytokinesis, preventing inappropriate Cdc42 activation. Our data also show that Gef1 removal depends on the presence of actin cables and the formin For3. Actin cytoskeleton organization is primarily regulated by Cdc42 (Sit and Manser, 2011). We find that *scd1Δ* mutants show depolarized actin cables and patches likely due to mis-regulation of Cdc42. We posit that Scd1-dependent actin cytoskeleton organization promotes Gef1 removal from the division site after ring disassembly.

Gef1 and Scd1 cooperate to drive polarized cell growth

Since we observed that Gef1 recruits Scd1 at the division site, we addressed whether this crosstalk also instructs Cdc42 at sites of polarized growth. Indeed, we find that Gef1 is necessary for bipolar localization of Scd1 and Scd2. Cells lacking *gef1* are mostly monopolar, with polarized growth occurring only at the old end (Coll et al., 2003; Das et al., 2012). This suggests that Gef1 promotes bipolar growth by recruiting Scd1 and Scd2. We report that while constitutively active Cdc42 itself is bipolar, it is not sufficient to restore bipolar localization of Scd1 in *gef1* mutants. This suggests that active Cdc42 alone does not feed into a positive feedback pathway to promote bipolar Scd1 localization. Our findings highlight a requirement for Gef1 in this process. The two ends in fission yeast compete for active Cdc42; the old end is the dominant end and initially wins this competition (Das et al., 2012). Bipolarity is established when the new end overcomes the dominance of the old end and can initiate growth. Thus, the new end is a nascent growth site that must activate Cdc42 in the absence of pre-established cues. Analysis of the growth pattern of *gef1Δ* mutants indicates that new ends frequently fail to overcome old end dominance, resulting in monopolar growth in these cells. Bipolar growth in *gef1Δ* mutants is typically observed in cells that do not contain a dominant old end. Taken together, our findings indicate that Gef1 allows the new end to overcome old end dominance through Scd1 recruitment and Cdc42 activation, leading to bipolar growth (Figure 7C). *gef1S112A* mutants (Das et al., 2015) and constitutively active Cdc42 mutants, both display bipolar growth. However, only *gef1S112A* mutants display bipolar Scd1 localization in which both the old and the new end recruit Scd1 and initiate growth almost immediately after completion of division. This provides further evidence that Gef1 promotes Scd1 recruitment to initiate bipolar growth.

In fission yeast, Scd1 is the primary GEF that promotes polarized growth (Chang et al., 1994). Cells lacking *scd1* are depolarized due to ectopic Cdc42 activation. We find that ectopic Cdc42 activation in these mutants is most likely due to mislocalized Gef1. In the presence of *scd1*, Gef1 shows sparse localization and is restricted to the cell ends. Cells lacking *scd1*, fail to restrict Gef1 localization to the ends. We find that Gef1 is mislocalized in the absence of the actin cytoskeleton, leading to ectopic Cdc42 activation. Furthermore, ectopic Cdc42 activation in LatA-treated cells is abolished in *gef1Δ* mutants. This suggests that the ectopic Cdc42 activation observed in LatA-treated cells is Gef1-dependent. Since *scd1* mutants display defects in actin organization, we posit that Scd1 promotes polarized Gef1 localization via the actin cytoskeleton. A recent report shows that ectopic Cdc42 activation in LatA-treated cells depends on the stress-activated MAP kinase Sty1 (Mutavchiev et al., 2016). We found that fission yeast cells treated with LatA did not display ectopic Cdc42 activation in the absence of *sty1*. It is possible that in the absence of actin the cells elicit a stress response, leading to Sty1 activation that results in the mislocalization of Gef1. Further analysis will be necessary to test this hypothesis.

Multiple GEFs combinatorially regulate Cdc42 during complex processes

The presence of multiple regulators generates combinatorial control that allows for the fine-tuning of a system in different conditions. Furthermore, multiple regulators may interact to instruct each other. Here we report an interesting interplay between Gef1 and Scd1, in which

Gef1 promotes Scd1 localization while Scd1 restricts Gef1 localization. This regulatory interaction between the GEFs facilitates precise Cdc42 activation. Cdc42 is an essential protein and is the major regulator of polarized cell growth and membrane trafficking (Das and Verde, 2013; Estravis et al., 2012; Estravis et al., 2011; Etienne-Manneville, 2004; Harris and Tepass, 2010; Johnson, 1999; Miller and Johnson, 1994). We have shown that Cdc42 is also involved in septum formation during cytokinesis (Wei et al., 2016). Given the essential nature of Cdc42, it is not surprising that it has multiple regulators that can function in an at least partially redundant manner. While *gef1Δ* and *scd1Δ* mutants are viable, the *gef1Δscd1Δ* double mutant is inviable (Coll et al., 2003). However, these mutants display distinct phenotypes in which *gef1Δ* mutants are monopolar while *scd1Δ* mutants are depolarized (Chang et al., 1994; Coll et al., 2003). This indicates that Gef1 and Scd1 function in different ways in fission yeast cells.

Here we show that Gef1 and Scd1 crosstalk with each other to establish and maintain polarized growth. Polarized cell growth requires symmetry breaking, and several models have indicated a need for positive feedback loops in this process (Irazoqui et al., 2003; Kozubowski et al., 2008; Slaughter et al., 2009b). Cdc42 is able to break symmetry and establish polarization through positive feedback (Kozubowski et al., 2008; Slaughter et al., 2009b). Elegant experiments in budding yeast demonstrate that local activation of Cdc42 establishes positive feedback through the recruitment of additional GEFs to amplify the conversion of Cdc42-GDP to Cdc42-GTP (Butty et al., 2002; Kozubowski et al., 2008). We expected Gef1-mediated recruitment of Scd1 to function via local activation of Cdc42, which would recruit Scd1 through the establishment of positive feedback. Surprisingly, we report that active Cdc42 is not sufficient to recruit Scd1 in the absence of Gef1. While it has been proposed that Cdc42 establishes positive feedback through the formation of the ternary complex consisting of the Cdc42 effector PAK (p21-activated kinases) and its associated scaffold protein (Scd2 or Bem1) (Butty et al., 2002; Kozubowski et al., 2008), studies in *S. pombe* and *S. cerevisiae* suggest that Pak1 kinase activity antagonizes either the Cdc42 scaffold or the GEF, rather than establishing a positive feedback (Das et al., 2012; Gulli et al., 2000; Kuo et al., 2014; Rapali et al., 2017). In support of this antagonistic role of Pak1, we find that more Scd1 accumulates at cell ends and at the division site in the *pak1* switch-off mutant. This indicates that Pak1 does not drive positive feedback, but rather serves to limit the level of Cdc42 activation. Our findings that active Cdc42 failed to recruit Scd1 in the absence of *gef1* and that Pak kinase antagonizes Scd1 do not agree with current models of Cdc42-mediated positive feedback. This may be due, in part, to the fact that most of these models are based on studies in budding yeast, and it highlights that the mechanisms that generate feedbacks critical to many biological processes are not well understood in fission yeast.

Cdc42 activation undergoes an oscillatory pattern at the cell ends that promotes bipolarity. Current models to explain these oscillations indicate the presence of a positive feedback, time-delayed negative feedback, and competition between the two ends for active Cdc42. Since Scd1 is the Cdc42 GEF that establishes polarized growth, we posit that Scd1 activates Cdc42 through positive feedback at the dominant old end. Dominance at the old end ensures that Scd1 localization is mainly restricted to this end at the expense of the new end. A previous model suggests that as the cell reaches a certain size, a corresponding increase in Scd1 levels would allow the new end to overcome old end dominance to initiate growth and promote bipolarity. Competition for Scd1 alone cannot explain our finding that bipolarity ensues when old end dominance is overcome through Gef1-mediated recruitment of Scd1 to the new end. Furthermore, *gef1S112A* cells display bipolar growth at a smaller cell size. Although an increase

in Scd1 levels may promote bipolarity, our data suggest that Gef1-mediated Scd1 recruitment is more important to establish bipolar growth.

Our finding that active Cdc42 alone does not promote localization of Scd1 to nascent sites may provide an advantage to the cell. A caveat of a positive feedback model driven solely by active Cdc42 is that any stochastic activation of Cdc42 at the cell cortex may generate random Scd1-mediated growth sites. Our data indicate that apart from the dominant old end, Scd1 cannot be localized to additional sites of growth merely by the presence of active Cdc42. Instead, in a cell with a dominant old end, Gef1 must help recruit Scd1 to the new end to allow bipolar growth (Figure 7C). Given that Gef1 promotes Scd1-mediated polarized growth at the new end, it is conceivable that Gef1 itself is tightly regulated to prevent random Cdc42 activation. Indeed, Gef1 shows sparse localization to the cell ends and is mainly cytoplasmic (Das et al., 2015). The NDR kinase Orb6 prevents ectopic Gef1 localization via 14-3-3-mediated sequestration to the cytoplasm (Das et al., 2015; Das et al., 2009). Here we show that while Gef1 promotes Scd1 recruitment to a nascent site, Scd1 itself restricts Gef1 localization to the cell ends to precisely activate Cdc42 (Figure 7C). Together our findings describe an elegant system in which the two Cdc42 GEFs regulate each other to ensure proper cell polarization.

Significance of GEF coordination in other systems

In budding yeast, CDC24 is required for polarization during bud emergence and is essential for viability (Sloat et al., 1981; Sloat and Pringle, 1978), unlike Scd1 in fission yeast. Budding yeast also has a second GEF Bud3, which establishes a proper bud site (Kang et al., 2014). During G1 in budding yeast, bud emergence occurs via biphasic Cdc42 activation by the two GEFs: Bud3 helps select the bud site (Kang et al., 2014), and Cdc24 allows polarization (Sloat et al., 1981; Sloat and Pringle, 1978). This is analogous to new end growth in fission yeast, which requires Gef1-dependent recruitment of Scd1 for robust Cdc42 activation. It would be interesting to see if a crosstalk also exists between Bud3 and Cdc24.

The Rho family of GTPases includes Rho, Rac, and Cdc42. In certain mammalian cells, Cdc42 and Rac1 appear to activate cell growth in a biphasic manner (de Beco et al., 2018; Yang et al., 2016). For example, during motility, the GTPases, Rho, Rac, and Cdc42, regulate the actin cytoskeleton (Heasman and Ridley, 2008; Machacek et al., 2009). During cell migration, these GTPases form bands or 'zones' in the leading and trailing regions of the cell (Ridley, 2015). Their spatial separation is mediated by the organization of their GEFs and GAPs, as well as by regulatory signaling between these GTPases (Guilluy et al., 2011). Cdc42 and Rho are mutually antagonistic, explaining how such zones of GTPase activity can be established and maintained (Guilluy et al., 2011; Kutys and Yamada, 2014; Warner and Longmore, 2009). Similarly, Cdc42 can refine Rac activity (Guilluy et al., 2011). Cdc42 and Rac are activated by similar pathways and share the same effectors. Several recent experiments indicate that actin cytoskeleton reorganization leading to cell migration occurs in a biphasic manner, where Cdc42 activation at new sites sets the direction, while robust Rac activation determines the speed (de Beco et al., 2018; Yang et al., 2016). Unlike most eukaryotes, the genome of *S. pombe* does not contain a Rac GTPase. We speculate that the two Cdc42 GEFs of *S. pombe* allow it to fulfill the roles of both Cdc42 and Rac. Gef1 sets the direction of growth by establishing growth at a new site, while Scd1 promotes efficient growth through robust Cdc42 activation at the growth sites. It is possible that in addition to the crosstalk between different Rho GTPases, the crosstalk between

Cdc42 GEFs themselves as we report here is an intrinsic property of small GTPases and is necessary for fine-tuning their activity.

MATERIALS AND METHODS

Strains and cell culture

The *S. pombe* strains used in this study are listed in Supplemental Table S1. All strains are isogenic to the original strain PN567. Cells were cultured in yeast extract (YE) medium and grown exponentially at 25°C, unless specified otherwise. Standard techniques were used for genetic manipulation and analysis (Moreno et al., 1991). Cells were grown exponentially for at least 3 rounds of eight generations each before imaging.

Microscopy

Cells were imaged at room temperature (23–25°C) with an Olympus IX83 microscope equipped with a VTHawk two-dimensional array laser scanning confocal microscopy system (Visitech International, Sunderland, UK), electron-multiplying charge-coupled device digital camera (Hamamatsu, Hamamatsu City, Japan), and 100×/numerical aperture 1.49 UAPO lens (Olympus, Tokyo, Japan). Images were acquired with MetaMorph (Molecular Devices, Sunnyvale, CA) and analyzed by ImageJ (National Institutes of Health, Bethesda, MD).

Actin staining

The actin cytoskeleton was stained by Alexa Fluor phalloidin as described here (Das et al., 2009; Pelham and Chang, 2001). Briefly, exponentially growing cells were fixed with 3.5% formaldehyde for 10 mins at room temperature. the fixed cells were washed with PM buffer (35 mM KPO4, pH 6.8, 0.5 mM MgSO4) permeabilized with 1% triton X-100 and stained with Alexa Fluor phalloidin (Molecular Probes) for 30 mins.

Analysis of growth pattern

The growth pattern of *gef1+* and *gef1Δ* cells was observed by live imaging of cells through multiple generations. Cells were placed in 3.5-mm glass-bottom culture dishes (MatTek, Ashland, MA) and overlaid with YE medium plus 1% agar, and 100μM ascorbic acid to minimize photo-toxicity to the cell. A bright-field image was acquired every minute for 12 hours. Birth scars were used to distinguish between, as well as to measure, old end and new end growth.

Construction of fluorescently tagged Gef1 fusion proteins

The forward primer 5'-CCCGGGAACCCTCGCAGCTAAAGA-3' with a 5' BamHI site and the reverse primer 5'-GGATCCGTGTTTACCAAAGTTATGTAAGAC-3' with a 5' XmaI site were used to amplify a 3kb DNA fragment containing *gef1*, the 5' UTR, and the endogenous promoter. The fragment was then digested with BamHI and XmaI and ligated into the BamHI-XmaI site of pKS392 pFA6-tdTomato-kanMX and pKG6507 pFA6-mNeonGreen-kanMX. Constructs were linearized by digestion with XbaI and transformed into the *gef1* locus in *gef1Δ* cells.

Expression of constitutively active Cdc42

pjk148-nmt41x-leu1+ or *pjk148-nmt41x:cdc42G12V-leu1+* were linearized with NdeI and integrated into the leu1-32 loci in gef1+ and gef1Δ cells expressing either CRIB-3xGFP or Scd1-3xGFP. The empty vector *pjk148-nmt41x-leu1+* was used as control. Cells were grown in YE to promote minimal expression of *cdc42G12V*.

Latrunculin A treatment

Cells in YE were incubated at room temperature with 10μM Latrunculin A dissolved in dimethyl sulfoxide (DMSO) for 40 min prior to imaging. Control cells were treated with 1% DMSO and incubated for 40 min.

CK666 treatment
Cells in YE were incubated at room temperature with 100μM CK666 dissolved in dimethyl sulfoxide (DMSO) for 5 min prior to imaging. Control cells were treated with 1% DMSO and incubated for 5 min.

Analysis of fluorescent intensity
Mutants expressing fluorescent proteins were grown to OD 0.5 and imaged on slides. Cells in slides were imaged for no more than 3 mins to prevent any stress response as previously described (Das et al., 2015). Depending on the mutant and the fluorophore, 16-28 Z-planes were collected at a z-interval of 0.4μm for either or both the 488nm and 561nm channels. The respective controls were grown and imaged in an identical manner. ImageJ was used to generate sum projections from the z-series, and to measure the fluorescence intensity of a selected region (actomyosin ring, or growth cap at cell tip). The background fluorescence in a cell-free region of the image was subtracted to generate the normalized intensity. Mean normalized intensity was calculated for each image from all (n>5) measurable cells within each field. A Student's two-tailed t-test, assuming unequal variance, was used to determine significance through comparison of each strain's mean normalized intensities.

Acknowledgements

We thank J. Bembenek and T. Burch-Smith for critical review of our manuscript; K. Gould for supplying plasmids; and M. Balasubramanian, and S. Martin for providing strains. This work was supported by a grant from the National Science Foundation (1616495).

REFERENCES

Atkins, B.D., S. Yoshida, K. Saito, C.F. Wu, D.J. Lew, and D. Pellman. 2013. Inhibition of Cdc42 during mitotic exit is required for cytokinesis. *J Cell Biol*. 202:231-240.

Bos, J.L., H. Rehmann, and A. Wittinghofer. 2007. GEFs and GAPs: critical elements in the control of small G proteins. *Cell*. 129:865-877.

Butty, A.C., N. Perrinjaquet, A. Petit, M. Jaquenoud, J.E. Segall, K. Hofmann, C. Zwahlen, and M. Peter. 2002. A positive feedback loop stabilizes the guanine-nucleotide exchange factor Cdc24 at sites of polarization. *EMBO J*. 21:1565-1576.

Carlin, L.M., R. Evans, H. Milewicz, L. Fernandes, D.R. Matthews, M. Perani, J. Levitt, M.D. Keppler, J. Monypenny, T. Coolen, P.R. Barber, B. Vojnovic, K. Suhling, F. Fraternali, S. Ameer-Beg, P.J. Parker, N.S. Thomas, and T. Ng. 2011. A targeted siRNA screen identifies regulators of Cdc42 activity at the natural killer cell immunological synapse. *Sci Signal*. 4:ra81.

Chang, E.C., M. Barr, Y. Wang, V. Jung, H.P. Xu, and M.H. Wigler. 1994. Cooperative interaction of S. pombe proteins required for mating and morphogenesis. *Cell*. 79:131-141.

Coffman, V.C., J.A. Sees, D.R. Kovar, and J.Q. Wu. 2013. The formins Cdc12 and For3 cooperate during contractile ring assembly in cytokinesis. *J Cell Biol*. 203:101-114.

Coll, P.M., Y. Trillo, A. Ametzazurra, and P. Perez. 2003. Gef1p, a new guanine nucleotide exchange factor for Cdc42p, regulates polarity in Schizosaccharomyces pombe. *Molecular biology of the cell*. 14:313-323.

Crawford, J.M., N. Harden, T. Leung, L. Lim, and D.P. Kiehart. 1998. Cellularization in Drosophila melanogaster is disrupted by the inhibition of rho activity and the activation of Cdc42 function. *Dev Biol*. 204:151-164.

Das, M., T. Drake, D.J. Wiley, P. Buchwald, D. Vavylonis, and F. Verde. 2012. Oscillatory Dynamics of Cdc42 GTPase in the Control of Polarized Growth. *Science*.

Das, M., I. Nunez, M. Rodriguez, D.J. Wiley, J. Rodriguez, A. Sarkeshik, J.R. Yates, 3rd, P. Buchwald, and F. Verde. 2015. Phosphorylation-dependent inhibition of Cdc42 GEF Gef1 by 14-3-3 protein Rad24 spatially regulates Cdc42 GTPase activity and oscillatory dynamics during cell morphogenesis. *Mol Biol Cell*.

Das, M., and F. Verde. 2013. Role of Cdc42 dynamics in the control of fission yeast cell polarization. *Biochem Soc Trans*. 41:1745-1749.

Das, M., D.J. Wiley, X. Chen, K. Shah, and F. Verde. 2009. The conserved NDR kinase Orb6 controls polarized cell growth by spatial regulation of the small GTPase Cdc42. *Curr Biol*. 19:1314-1319.

de Beco, S., K. Vaidziulyte, J. Manzi, F. Dalier, F. Di Frederico, G. Cornilleau, M. Dahan, and M. Coppey. 2018. Optogenetic dissection of Rac1 and Cdc42 gradient shaping. *bioRxiv*.

Drubin, D.G., and W.J. Nelson. 1996. Origins of cell polarity. *Cell*. 84:335-344.

Dutartre, H., J. Davoust, J.P. Gorvel, and P. Chavrier. 1996. Cytokinesis arrest and redistribution of actin-cytoskeleton regulatory components in cells expressing the Rho GTPase CDC42Hs. *J Cell Sci*. 109 (Pt 2):367-377.

Endo, M., M. Shirouzu, and S. Yokoyama. 2003. The Cdc42 binding and scaffolding activities of the fission yeast adaptor protein Scd2. *J Biol Chem*. 278:843-852.

Estravis, M., S. Rincon, and P. Perez. 2012. Cdc42 regulation of polarized traffic in fission yeast. *Commun Integr Biol*. 5:370-373.

Estravis, M., S.A. Rincon, B. Santos, and P. Perez. 2011. Cdc42 regulates multiple membrane traffic events in fission yeast. *Traffic*. 12:1744-1758.

Etienne-Manneville, S. 2004. Cdc42--the centre of polarity. *Journal of cell science*. 117:1291-1300.

Feierbach, B., and F. Chang. 2001. Roles of the fission yeast formin for3p in cell polarity, actin cable formation and symmetric cell division. *Current biology : CB*. 11:1656-1665.

Feigin, M.E., and S.K. Muthuswamy. 2009. Polarity proteins regulate mammalian cell-cell junctions and cancer pathogenesis. *Curr Opin Cell Biol*. 21:694-700.

Gachet, Y., and J.S. Hyams. 2005. Endocytosis in fission yeast is spatially associated with the actin cytoskeleton during polarised cell growth and cytokinesis. *J Cell Sci*. 118:4231-4242.

Godde, N.J., R.C. Galea, I.A. Elsum, and P.O. Humbert. 2010. Cell polarity in motion: redefining mammary tissue organization through EMT and cell polarity transitions. *J Mammary Gland Biol Neoplasia*. 15:149-168.

Guilluy, C., R. Garcia-Mata, and K. Burridge. 2011. Rho protein crosstalk: another social network? *Trends Cell Biol*. 21:718-726.

Gulli, M.P., M. Jaquenoud, Y. Shimada, G. Niederhauser, P. Wiget, and M. Peter. 2000. Phosphorylation of the Cdc42 exchange factor Cdc24 by the PAK-like kinase Cla4 may regulate polarized growth in yeast. *Mol Cell*. 6:1155-1167.

Halaoui, R., and L. McCaffrey. 2014. Rewiring cell polarity signaling in cancer. *Oncogene*.

Harris, K.P., and U. Tepass. 2010. Cdc42 and vesicle trafficking in polarized cells. *Traffic*. 11:1272-1279.

Heasman, S.J., and A.J. Ridley. 2008. Mammalian Rho GTPases: new insights into their functions from in vivo studies. *Nat Rev Mol Cell Biol*. 9:690-701.

Hirota, K., K. Tanaka, K. Ohta, and M. Yamamoto. 2003. Gef1p and Scd1p, the Two GDP-GTP exchange factors for Cdc42p, form a ring structure that shrinks during cytokinesis in Schizosaccharomyces pombe. *Mol Biol Cell*. 14:3617-3627.

Howell, A.S., M. Jin, C.F. Wu, T.R. Zyla, T.C. Elston, and D.J. Lew. 2012. Negative feedback enhances robustness in the yeast polarity establishment circuit. *Cell*. 149:322-333.

Huang, J., Y. Huang, H. Yu, D. Subramanian, A. Padmanabhan, R. Thadani, Y. Tao, X. Tang, R. Wedlich-Soldner, and M.K. Balasubramanian. 2012. Nonmedially assembled F-actin cables incorporate into the actomyosin ring in fission yeast. *J Cell Biol*. 199:831-847.

Hwang, J.U., Y. Gu, Y.J. Lee, and Z. Yang. 2005. Oscillatory ROP GTPase activation leads the oscillatory polarized growth of pollen tubes. *Mol Biol Cell*. 16:5385-5399.

Irazoqui, J.E., A.S. Gladfelter, and D.J. Lew. 2003. Scaffold-mediated symmetry breaking by Cdc42p. *Nat Cell Biol*. 5:1062-1070.

Johnson, D.I. 1999. Cdc42: An essential Rho-type GTPase controlling eukaryotic cell polarity. *Microbiology and molecular biology reviews : MMBR*. 63:54-105.

Kang, P.J., M.E. Lee, and H.O. Park. 2014. Bud3 activates Cdc42 to establish a proper growth site in budding yeast. *J Cell Biol*. 206:19-28.

Kelly, F.D., and P. Nurse. 2011. Spatial control of Cdc42 activation determines cell width in fission yeast. *Mol Biol Cell*. 22:3801-3811.

Kozubowski, L., K. Saito, J.M. Johnson, A.S. Howell, T.R. Zyla, and D.J. Lew. 2008. Symmetry-breaking polarization driven by a Cdc42p GEF-PAK complex. *Current biology : CB*. 18:1719-1726.

Kuo, C.C., N.S. Savage, H. Chen, C.F. Wu, T.R. Zyla, and D.J. Lew. 2014. Inhibitory GEF phosphorylation provides negative feedback in the yeast polarity circuit. *Curr Biol*. 24:753-759.

Kutys, M.L., and K.M. Yamada. 2014. An extracellular-matrix-specific GEF-GAP interaction regulates Rho GTPase crosstalk for 3D collagen migration. *Nat Cell Biol*. 16:909-917.

Lauffenburger, D.A., and A.F. Horwitz. 1996. Cell migration: a physically integrated molecular process. *Cell*. 84:359-369.

Machacek, M., L. Hodgson, C. Welch, H. Elliott, O. Pertz, P. Nalbant, A. Abell, G.L. Johnson, K.M. Hahn, and G. Danuser. 2009. Coordination of Rho GTPase activities during cell protrusion. *Nature*. 461:99-103.

Miller, P.J., and D.I. Johnson. 1994. Cdc42p GTPase is involved in controlling polarized cell growth in Schizosaccharomyces pombe. *Mol Cell Biol*. 14:1075-1083.

Mitchison, J.M., and P. Nurse. 1985. Growth in cell length in the fission yeast Schizosaccharomyces pombe. *Journal of cell science*. 75:357-376.

Moreno, S., A. Klar, and P. Nurse. 1991. Molecular genetic analysis of fission yeast Schizosaccharomyces pombe. *Methods in enzymology*. 194:795-823.

Mutavchiev, D.R., M. Leda, and K.E. Sawin. 2016. Remodeling of the Fission Yeast Cdc42 Cell-Polarity Module via the Sty1 p38 Stress-Activated Protein Kinase Pathway. *Curr Biol*. 26:2921-2928.

Nabeshima, K., T. Nakagawa, A.F. Straight, A. Murray, Y. Chikashige, Y.M. Yamashita, Y. Hiraoka, and M. Yanagida. 1998. Dynamics of centromeres during metaphase-anaphase transition in fission yeast: Dis1 is implicated in force balance in metaphase bipolar spindle. *Mol Biol Cell*. 9:3211-3225.

Nance, J., and J.A. Zallen. 2011. Elaborating polarity: PAR proteins and the cytoskeleton. *Development*. 138:799-809.

Onishi, M., N. Ko, R. Nishihama, and J.R. Pringle. 2013. Distinct roles of Rho1, Cdc42, and Cyk3 in septum formation and abscission during yeast cytokinesis. *J Cell Biol*. 202:311-329.

Pelham, R.J., and F. Chang. 2001. Role of actin polymerization and actin cables in actin-patch movement in Schizosaccharomyces pombe. *Nature cell biology*. 3:235-244.

Pollard, T.D. 2010. Mechanics of cytokinesis in eukaryotes. *Curr Opin Cell Biol*. 22:50-56.

Rapali, P., R. Mitteau, C. Braun, A. Massoni-Laporte, C. Unlu, L. Bataille, F.S. Arramon, S.P. Gygi, and D. McCusker. 2017. Scaffold-mediated gating of Cdc42 signalling flux. *Elife*. 6.

Ridley, A.J. 2006. Rho GTPases and actin dynamics in membrane protrusions and vesicle trafficking. *Trends Cell Biol*. 16:522-529.

Ridley, A.J. 2015. Rho GTPase signalling in cell migration. *Curr Opin Cell Biol*. 36:103-112.

Sit, S.T., and E. Manser. 2011. Rho GTPases and their role in organizing the actin cytoskeleton. *J Cell Sci*. 124:679-683.

Slaughter, B.D., A. Das, J.W. Schwartz, B. Rubinstein, and R. Li. 2009a. Dual modes of cdc42 recycling fine-tune polarized morphogenesis. *Developmental cell*. 17:823-835.

Slaughter, B.D., S.E. Smith, and R. Li. 2009b. Symmetry breaking in the life cycle of the budding yeast. *Cold Spring Harbor Perspect Biol*. 1:a003384.

Sloat, B.F., A. Adams, and J.R. Pringle. 1981. Roles of the CDC24 gene product in cellular morphogenesis during the Saccharomyces cerevisiae cell cycle. *J Cell Biol*. 89:395-405.

Sloat, B.F., and J.R. Pringle. 1978. A mutant of yeast defective in cellular morphogenesis. *Science*. 200:1171-1173.

Spector, I., N.R. Shochet, Y. Kashman, and A. Groweiss. 1983. Latrunculins: novel marine toxins that disrupt microfilament organization in cultured cells. *Science*. 219:493-495.

Sun, S.C., Z.B. Wang, Y.N. Xu, S.E. Lee, X.S. Cui, and N.H. Kim. 2011. Arp2/3 complex regulates asymmetric division and cytokinesis in mouse oocytes. *PLoS One*. 6:e18392.

Wang, N., I.J. Lee, G. Rask, and J.Q. Wu. 2016. Roles of the TRAPP-II Complex and the Exocyst in Membrane Deposition during Fission Yeast Cytokinesis. *PLoS Biol*. 14:e1002437.

Warner, S.J., and G.D. Longmore. 2009. Cdc42 antagonizes Rho1 activity at adherens junctions to limit epithelial cell apical tension. *J Cell Biol*. 187:119-133.

Wedlich-Soldner, R., S.C. Wai, T. Schmidt, and R. Li. 2004. Robust cell polarity is a dynamic state established by coupling transport and GTPase signaling. *J Cell Biol*. 166:889-900.

Wei, B., B.S. Hercyk, N. Mattson, A. Mohammadi, J. Rich, E. DeBruyne, M.M. Clark, and M. Das. 2016. Unique spatiotemporal activation pattern of Cdc42 by Gef1 and Scd1 promotes different events during cytokinesis. *Mol Biol Cell*. 27:1235-1245.

Wu, J.Q., J.R. Kuhn, D.R. Kovar, and T.D. Pollard. 2003. Spatial and temporal pathway for assembly and constriction of the contractile ring in fission yeast cytokinesis. *Dev Cell*. 5:723-734.

Yang, H.W., S.R. Collins, and T. Meyer. 2016. Locally excitable Cdc42 signals steer cells during chemotaxis. *Nat Cell Biol*. 18:191-201.

Figure 1

Figure 1: Gef1 promotes Scd1 localization to the division site. (A) Scd1-3xGFP localization in *gef1+* and *gef1Δ* cells expressing the ring and SPB markers Rlc1-tdTomato and Sad1-mCherry respectively. Arrowheads label cells with Scd1-3xGFP localized to the division site, while arrows mark cells with constricting rings that lack Scd1-3xGFP at the division site. **(B and C)** Quantification of Scd1-3xGFP localization and intensity in the indicated genotypes (**, p<0.01). **(D)** CRIB-3xGFP, the active Cdc42 sensor, intensity at the division site in *gef1+* and *gef1Δ* cells transformed with the control vector *pJK148* or *cdc42G12V*. Arrows label cells with reduced Cdc42 activity at the division site and arrowheads indicate cells with increased Cdc42 activity. **(E)** Scd1-3xGFP localization at the division site in *gef1+* and *gef1Δ* cells transformed with the control vector *pJK148* or *cdc42G12V*. Arrows label cells with reduced Scd1-3xGFP at the division site. **(F and G)** Quantifications of *cdc42G12V*-mediated Cdc42 activity and Scd1 localization at the division site in the indicated genotypes (****, p<0.0001). Cell Division Site abbreviated as CDS. All data points are plotted in each graph, with black bars on top of data points that show the mean and standard deviation for each genotype. All images are inverted max projections. Scale bars = 5µm.

Figure 2

Figure 2: Gef1 promotes Scd1 localization to the division site via recruitment of Scd2. (A) Scd2 localization in *gef1+* and *gef1Δ* cells expressing the ring and SPB markers Rlc1-tdTomato and Sad1-mCherry. Arrowheads label cells with Scd2-GFP localized constricting rings, while arrows mark cells with assembled rings that lack Scd2-GFP at the division site. **(B and C)** Quantification of Scd2-GFP localization and intensity in the indicated genotypes (****, $p<0.0001$). **(D)** Scd1-3xGFP localization in *scd2+* and *scd2Δ* cells. Division site marked by black arrows in the bright field images. Scd1-3xGFP localization to the division site indicated by red arrowheads. Red arrows show absence of Scd1-3xGFP at the division site. **(E)** Scd2-GFP localization to the division site in *scd1+* and *scd1Δ* cells. **(F)** Representative images showing the localizations of Gef1-mNG, Scd2-GFP, and Scd1-3GFP (top panels) as a function of spindle pole body distance (bottom panels). The range of SPB distance is listed for each column. Green arrows indicate the earliest time point at which signal is visible. Red arrowheads indicate time points prior to localization. **(G)** Quantification of Gef1, Scd2, and Scd1 localization to the division site in a temporal manner, showing the means of the distance between spindle poles of the first 50^{th} percentile of early anaphase cells at which signal first appears (*, $p<0.05$). Cell Division Site abbreviated as CDS. All data points are plotted in each graph, with black bars on top of data points that show the mean and standard deviation for each genotype. All images are inverted max projections with the exception of bright field. Scale bars = 5µm.

Figure 3

Figure 3: Scd1 and actin promote Gef1 removal from the division site after ring constriction. (A) Gef1mNG localization to the division site after ring disassembly in *scd1+* and *scd1Δ* cells expressing the ring and SPB markers Rlc1-tdTomato and Sad1-mCherry. Black arrowheads mark the membrane barrier in cells post-ring disassembly. Red arrowheads mark cells post-ring disassembly that lack Gef1-mNG at the membrane barrier. Red arrows mark cells with Gef1-mNG localized to the membrane barrier post-ring assembly. **(B)** Quantification of Gef1 lingering at the division site in *scd1+* and *scd1Δ* cells (****, $p<0.0001$). **(C)** Gef1-mNG localization in septated cells expressing the ring and SPB markers Rlc1-tdTomato and Sad1-mCherry, treated with either DMSO, 10μM LatA, or 100μM CK666. Red arrowheads mark cells post-ring disassembly that lack Gef1-mNG at the membrane barrier. Red arrows indicate cells with Gef1-mNG localized to the membrane barrier post-ring assembly. Green arrowheads indicate cells with Gef1-mNG localized to the cortex orthogonal to the membrane barrier. **(D)** Quantification of Gef1 lingering at the division site in septated *scd1+* and *scd1Δ* cells treated with 10μM LatA or DMSO (*, $p<0.05$. **, $p<0.01$). **(E)** Gef1-3xYFP localization in *for3+* and *for3Δ* cells. Red arrowheads mark cells post-ring disassembly that lack Gef1-3xYFP at the membrane barrier. Red arrows mark cells with Gef1-3xYFP localized to the membrane barrier post-ring assembly. Cell Division Site abbreviated as CDS. All data points are plotted in each graph, with black bars on top of data points that show the mean and standard deviation for each genotype. All images are inverted max projections. Scale bars = 5μm.

Figure 4

Figure 4: Gef1 promotes Scd1 localization to the new end. (A) Scd2-GFP (top panel) and Scd1-3xGFP (bottom panel) localization to the sites of polarized growth in *gef1+* and *gef1Δ* cells. Red arrows indicate monopolar cells that do not recruit Scd2-GFP or Scd1-3xGFP to the new end. **(B)** Quantifications of bipolar Scd1-3xGFP and Scd2-GFP localization in the indicated genotypes (***, $p<0.001$, ****, $p<0.0001$). **(C)** CRIB-3xGFP and Scd1-3xGFP localization at cell tips, and restoration of bipolar growth in *gef1+* and *gef1Δ* cells transformed with the control vector *pJK148* or *cdc42G12V*. **(D and E)** Quantification of the percent of cells that exhibit bipolar CRIB-3xGFP and Scd1-3xGFP localization at cell tips in the indicated genotypes (**, $p<0.01$). **(F)** Scd2-GFP and Scd1-3xGFP localization to the cortex in *scd1Δ* and *scd2Δ* cells, respectively. Red arrows indicate cells with Scd2-GFP localized tot the cell cortex. All data points are plotted in each graph, with black bars on top of data points that show the mean and standard deviation for each genotype. All images are inverted max projections with the exception of bright field unless specified. Scale bars = 5µm.

C

Polarity of dominant end-containing cells

	gef1+	*gef1Δ*
monopolar	3%	81%
NETO	97%	9%
precociously bipolar	0%	9%

Figure 5

Figure 5: Gef1 promotes bipolar growth via new-end-take-off . (**A**) Wild-type cells predominately display old end growth followed by a delayed onset of new-end growth. (**B) i.** In *gef1Δ*, 68% of monopolar cells yield a monopolar cell from the end that grew in the previous generation and a bipolar cell from the end that failed to grow in the previous generation. **ii**.18% of monopolar cells yield two monopolar cells. **iii**. 6% of monopolar cells yield two monopolar cells. Circled numbers describe the order of growth. Arrows correspond to direction of growth. (**C**) Quantification of polarity of *gef1+* and *gef1Δ* cells that contain a dominant end. (**D**) Localization of Scd1-3xGFP to the cell poles in *gef1+* and *gef1S112A* cells. (**E**) Quantification of the percent of cells that exhibit bipolar Scd1-3xGFP localization at cell ends in the indicated genotypes (****, p<0.0001). Scale bar = 5μm.

Figure 6

Figure 6: Scd1 and actin prevent ectopic Gef1 localization to promote polarized growth. (A) Gef1-mNG localization in *gef1+* and *gef1Δ* cells treated with DMSO (top panel) or 10μM LatA (bottom panel). **(B)** CRIB-3xGFP localization in *gef1+* and *gef1Δ* cells treated with DMSO (top panel) or 10μM LatA (bottom panel). Arrowheads indicate cells with CRIB-3xGFP or Gef1-mNG localized to regions of polarized growth. Arrows indicate cells with CRIB-3xGFP or Gef1-mNG localized to non-polarized regions on the cell cortex. All images are inverted max projections of the medial 4-7 cell slices. Scale bar = 5μm.

A

B

D

Figure 7

Figure 7: Model of the crosstalk between Gef1 and Scd1 that promotes polarized bipolar growth. (A) Diagram of the crosstalk pathway between Gef1 and Scd1. Solid arrows indicate an activating or promoting relationship in the direction of the arrow. Red terminating arrow indicates inactivation or removal of the protein at the arrows terminus. Dashed arrows indicate that the mechanism that regulates the proteins to which these arrows point is not yet resolved. **(B)** Schematic depicting the sequential localization of Gef1, Scd2, and Scd1 to the division site during cytokinesis. At the division site Gef1 localizes first and promotes Scd2 localization. Scd2 at the division site then recruits Scd1. **(C)** Schematic illustrating the crosstalk between Gef1 and Scd1 that promotes bipolar growth and regulates cell shape. In wild type (WT) cells, Gef1 localizes Scd2 to the new end, which in turn recruits Scd1 thus enabling NETO. In *scd1Δ* cells Gef1 localization is no longer restricted to the cell ends leading to ectopic Cdc42 activation and loss of polarity.