CNNC: Convolutional Neural Networks for Co-Expression Analysis

Ye Yuan[1], Ziv Bar-Joseph[1,2*]

[1]Machine Learning Department, School of Computer Science, Carnegie Mellon

University, Pittsburgh, PA 15213, USA. [2]Computational Biology Department, School

of Computer Science, Carnegie Mellon University, Pittsburgh, PA 15213, USA.

*e-mail: zivbj@cs.cmu.edu

Abstract

Co-expression analysis has been extensively used in genomics studies and tools for

over two decades. To date, most methods for such analysis are unsupervised and

symmetric. Such methods cannot infer causality and are prone to both overfitting and

false negatives resulting from differences between cells in bulk studies. Here we

present a new, supervised method based on convolutional neural networks (CNNs)

for co-expression analysis. We use a normalized histogram image of gene pair

co-expression as the input to the CNN. Testing our method on several co-expression

prediction tasks we show that it outperforms prior methods and that scRNA-Seq data

leads to more accurate results when compared to bulk data. The method can be

directly extended to integrate sequence and epigenetic data and to infer causal

relationships.

Supporting website with software and data: https://github.com/xiaoyeye/CNNC.

Introduction

Co-expression analysis, which seeks to identify genes that are correlated or

anti-correlated across a large number of samples or time points, has been a key

research area of computational genomics for almost two decades[1-5]. In addition to the

identification of pairs of related genes, co-expression analysis serves as an initial step

in many of the most widely used computational methods for the analysis of genomics

data including various clustering methods[6], network inference and reconstruction

approaches[7-11], methods for classification based on genes expression[3] and many

more.

Given its centrality for several downstream applications, much work has focused on

improving the ability to infer correlated and anti-correlated genes. The most popular

method is based on Pearson correlation analysis[12]. Such analysis focuses on shared

trends rather than exact values. Other popular and widely used methods involve

mutual information (MI)[13-15], nonparametric methods, for example Spearman

correlation[16] methods based on alignment[17] and more[18-20].

While the above methods were shown to be useful in many applications, they also

suffer from serious drawbacks. The first major issue is overfitting. Given the large

number of genes that are profiled, and the often relatively small (at least in

comparison) number of samples, several genes that are determined to be

co-expressed may only reflect chance or noise in the data[21]. In addition, to date most

co-expression analysis utilized bulk gene expression data (either array or RNA-Seq).

In such data, correlations may be obscured by the different cell populations such that

even if two genes appear highly correlated, it may be that they are actually never

expressed in the same cell at the same time[22]. Finally, most of the widely used

co-expression analysis methods are symmetric which means that each pair has only

one co-expression value. While this is advantageous for some applications (for

example, clustering) it may be problematic for methods that aim at inferring causality

(for example, network reconstruction methods).

To address these issues we developed a new tool, CNNC which provides a

supervised way (that can be tailored to the condition / question of interest) to perform

co-expression analysis. The method utilizes both bulk and single cell RNA-Seq

(scRNA-Seq) data from tens of thousands of experiments, allowing us to overcome

cell population confounders. Our method utilizes CNNs which we tailor for the gene

expression analysis by representing input data for each pair of genes as an (image)

histogram. The network is trained with positive and negative examples for the specific

domain of interest (for example, known targets of a TF in a specific cell type, known

pathways in a specific biological process etc.). Depending on the input data the

training can be either symmetric or directed (for example, training the network to infer

that TF A regulate gene B but not vice versa). To reduce overfitting CNNC determines

specific thresholds based on the training for calling a pair correlated or anti-correlated

or for inferring causality.

We applied CNNC to data from tens of thousands of single cell and bulk experiments.

We noticed that scRNA-Seq data greatly improves performance when compared to

bulk RNA-Seq. Using the same expression data to learn different CNNs (by varying

the labels based on the specific domain the network was applied to) we show that

CNNC outperforms prior co-expression analysis methods both for directly inferring

interactions (including TF-gene and protein-protein interactions) and as a component

in algorithms for the reconstruction of known pathways and clustering. Finally, we

discuss the accuracy of the directionality predictions which are unique to CNNC and

shown that these predictions provide important information for determining missing

interactions in known pathways.

Results

We developed a general computational framework for supervised co-expression

analysis (**Fig. 1**). CNNC is based on CNN which is used to analyze summarized

co-expression histograms from pairs of genes from bulk and scRNA-Seq data. Given

a relatively small labeled set of positive pairs (with either negative or random pairs

serving as negative) the CNN learns to discriminate between interacting and / or

causal pairs and negative pairs. Once trained the CNN can be used to predict

co-expression scores for all gene pairs.

Learning a CNNC model

We used processed scRNA-Seq data that was collected from over 500 different

studies representing a wide range of cell types, conditions etc[23]. All raw data was

uniformly processed and assigned to a pre-determined set of more than 20,000

mouse genes (Methods). We also used bulk RNA-Seq RPKM data from Encyclopedia

of DNA Elements (ENCODE) project[24], which contains 58 mouse tissues or cell types.

For both datasets we first generated a normalized empirical probability distribution

function (NEPDF) for each gene pair (genes *a* and *b*) based on their expression in the

scRNA-Seq or bulk RNA-Seq data (**Fig. 1**, left). We calculated their normalized 2-

dimension (2D) histogram and fixed its size at 32X32, where columns represent gene

a expression levels and rows represent gene *b* such that entries in the matrix

represent the (normalized) co-occurrences of these values. See Methods for details.

Bulk and sc NEPDF were then either used separately or concatenated to form a

combined NEPDF with dimension of 32X64. Next, the histogram matrix is used as

input to a CNN which is trained using a N-dimension (ND) output label vector, where

N depends on specific tasks. In our case N can either be 1 (interacting or not) or 3 in

which case label 0 indicates that genes *a* and *b* are not interacting and label 1 (2)

indicates that gene *a* (*b*) regulates gene *b* (*a*). Because of the final 'softmax' layer

classification utilized by CNNs, for a three-label task CNNC's output is a vector

consisting of three respective probabilities, [*p0, p1, p2*], which sum to 1. In general,

our CNN model consists of one 32X32 or 32X64 input layer, ten intermediate layers

including six convolutional layers, three maxpooling layers, one flatten layer, and a

final ND 'softmax' layer or one scalar 'Sigmoid' layer (**Methods** and **Supplementary**

Fig. 1).

In addition to gene expression data, we can easily integrate other data types including

Dnase-seq, PWM, etc. For this, we concatenated the additional information as a

vector to the intermediate output of the gene expression data and continued with the

standard CNN architecture. See **Methods** and **Supplementary Fig. 1** for complete

details and **Supplementary Table 1** for information on training and run time.

Using CNNC to predict TF-gene interactions

Chromatin immunoprecipitation (ChIP)-seq has been widely used as a gold standard

for studying cell-specific protein-DNA interactions[25]. Here we first evaluated CNNC's

performance on regulator-target prediction based on data from the GTRD ChIP-seq

database[26].

We extracted data from GTRD for 41 TFs for which ChIP-seq experiments were

performed in mouse embryonic stem cell (mESC). To determine targets for each TF

based on the ChIP-seq data from GTRD, we followed ref 27 and 28 [27, 28] and defined

the promotor region as 10KB upstream to 1KB downstream from the transcription

start site (TSS) for each gene. If a TF X has at least one detected peak signal with

p-value smaller than 10^{-300} in or overlapping the promotor region of gene Y, we say

that TF X regulates gene Y. We also used this data to compare CNNC with the two

most popular co-expression analysis methods: Pearson correlation (PC) and mutual

information (MI) and to compare the accuracy of predictions based on the sc and bulk

RNA-Seq data. Since the prior methods used for comparison are symmetric, we

focused here on the two labels setting (interacting or not). We later discuss causality

inference on this data. To compare the methods and data types we performed

leave-one-TF-out cross validation analysis. For each of the 41TFs, we trained CNNC

with all other TFs and used the left out TF for testing (**Methods**).

Fig. 2 presents the results of this comparison analysis. First, we see that for all

methods scRNA-Seq data (left column) provides much more information when

compared to bulk (middle column). Note that while we had more scRNA-Seq profiles

in our training set when compared to bulk experiments, these actually represent much

fewer cells and conditions than those used in the bulk data. We have also tested the

performance when using the same number of bulk and scRNA profiles

(**Supplementary Fig. 2**). We found that even with this very small number of

scRNA-Seq profiles (with much fewer cells than the bulk) CNNC performs better when

using scRNA-Seq. These results support prior claims about convolution effects

resulting from population of cells that make target inference harder when using bulk

data[22, 29]. Still, bulk data did include some useful information and for all methods since

the joint sc and bulk data performed best when compared to individual data type on its

own. As for the methods themselves, for all types of input data, CNNC outperforms

the other two methods. This is especially noticeable when using the scRNA-Seq (and

combined) data where CNNC is 15% more accurate than MI and close to 20% more

accurate than PC. The difference is even more pronounced for the top ranked

predictions. Here, for CNNC we see almost no false negatives for the first 15% of

ranked pairs (inset, **Fig. 2i**).

Data Integration further improves TF target gene prediction

The above analysis was only based on using expression values. However, as noted

above, gene co-expression is often used as a component in more extensive

procedures that often integrate different types of genomics data. To test how the use

of the NN based method can aid such procedures we combined the co-expression

values obtained by our method and the other methods with sequence and DNase

hypersensitivity data. For sequence, we used PWMs for the TFs we tested from the

Jaspar website[30]. We have also used Dnase-seq data for the same cell line from the

mouse ENCODE project[24]. While there are several different methods for integrating

expression, sequence and DNase data, since our main focus here is on the

175 co-expression analysis methods we used a simple strategy for processing the PWM

176 and DNase data (**Methods**) which resulted in an additional 2D vector as input for

177 each pair which we embedded to create a 512D vector (**Fig. 1 and Methods**). We

178 next extended the CNN to utilize the additional data by concatenating it with the

179 NEPDF's 512D vector in the flatten layer to form a 1024D vector as shown in **Fig. 1**

180 and **Supplementary Fig. 1**.

181 Results, presented in **Fig. 2j**, show that these additional data sources indeed improve

182 the ability to predict TF-gene interactions. However, as before a combined framework

183 utilizing our CNNC method for co-expression analysis outperformed a method that

184 used both MI and PC. Thus, the NN based approach can successfully replace other

185 methods as a component in a more elaborated systems biology framework for

186 inferring interactions.

188 **CNNC can predict pathway regulator-target gene pairs**

189 While TFs usually directly impact the resulting expression profile of their target genes

190 (and so co-expression analysis seems like a natural option to study such interactions)

191 several methods have also utilized RNA-Seq data to infer pathways that combine

192 protein-protein and protein-DNA interactions[31-35]. To test whether CNNC can serve as

193 a component in pathway inference methods we selected two representative pathway

194 databases, KEGG[36] and Reactome[37] as gold standard and used these to train and

195 test our co-expression framework. Since we are interested in causal relationships we

196 only used directed edges with activation or inhabitation edge types and filtered out

197 cyclic gene pairs where genes regulate each other mutually (to allow for a unique

198 label for each pair). As for the negative data, here we limited the negative set to a

199 random set of pairs where both genes appear in pathways in the database but do not

200 interact. Here leave-one-gene-out cross validation strategy requires extremely large

201 computing resources due to the large number of genes with outgoing directed edges

202 (3,057 for KEGG and 2,519 for Reactome). Instead, we performed a three-fold cross

203 validation where we kept the set of genes for which we predicted interactions

completely separated (so a gene in the test set does not have any interaction in the

training set). The positive data was uniformly divided by the outgoing gene into three

equal sized outgoing subsets, CNNC was trained using any two subsets and

evaluated using the left subset, and then the test ROCs for each outgoing gene in the

three subsets were calculated (**Methods**). Results are presented in **Fig. 3**. As can be

seen, CNNC performs very well on the KEGG pathways (See **Supplementary Fig. 3**

for the different folds) and also performs quite well on Reactome pathways (see also

Supplementary Fig. 3). In contrast, the other co-expression analysis methods do not

perform as well on these datasets (**Supplementary Fig. 4**).

Using CNNC for casualty prediction

So far we focused on general interaction predictions, which is what most symmetric

co-expression analysis are aimed at. However, as discussed above CNNC can also

be used to infer directionality by changing the output of the NN.

We next used CNNC to infer causal edges for all three datasets we studied (for

TF-gene interactions causal relationships are clear, for the pathway database we only

analyzed directed edges and so had the ground truth for that data as well). As can be

seen in **Fig. 4**, when using the TF GTRD dataset, CNNC achieves a median AUROC

of 0.8227 (**Fig. 4a**), with 32 of the 41 TFs obtaining an AUROC of more than 0.5 on

this leave-one-TF-out classification task. Interestingly, as can be seen in

Supplementary Fig. 5, when only using bulk RNA-Seq data performance on the

GTRD data prediction is very weak. Thus, for the causality inference task scRNA-Seq

data is the only one that can provide enough information. For KEGG, CNNC is very

successful achieving a median AUROC of 0.9949 (**Fig. 4c**) (See **Supplementary Fig.**

6 for the different folds).   For Reactome (**Fig. 4e**) we see that the most confident

predictions are correct, but beyond the top predictions performance levels off (See

Supplementary Fig. 6 for the different folds). To try to understand the process used

by the NN to distinguish causal directions we plotted two NEPDF inputs to the NN

(**Figs. 4g and 4h**) which were correctly predicted as two different labels (1 for **4g** and

2 for **4h**). As can be seen, in both inputs the two genes display partial correlations and

there are places where both are up or down concurrently. However, the main

difference between the histograms in **4g** and **4h** are cases where one gene is up and

the other is not. In **4g** gene 2 is up while gene 1 is not indicating that the causal

relationship is likely g1 -> g2. The opposite holds for **4h** and so the method infers that

g2 -> g1 for that input. Thus, unlike the prior symmetric methods, the NEPDF that

serves as input provides important clues that the NN can utilize to infer both

interactions and causality.

Pathway application

Given the results for KEGG we asked whether we can use the CNNC method to infer

missing edges in current pathways. There have been several attempts to utilize

expression and other data to further refine known pathways and many of these are

based on co-expression analysis[9, 19-21, 32, 33, 38, 39]. Since our method provides both

direction and score we can extract all predicted directed edges above a certain score

and compare the resulting pathway to the database pathway to see if any additional

edges, that do not appear in the database, are predicted by our method. For this we

focused on the interleukin 17 (IL-17) pathway from KEGG database, which plays

crucial roles in inflammatory responses. We extracted 6 proteins and 4 directed edges

from this pathway by only using directed edges with activation or inhabitation edge

types and filtering out cyclic gene pairs (**Fig. 5a**). We applied CNNC trained on all

database pathway edges that do not contain any of these 6 proteins. As can be seen

in **Supplementary Fig. 7**, CNNC predicted 8 of the possible 30 edges for this path

(15 pairs with two possible directions each). 4 of these 8 were the original 4 directed

edges annotated in the database itself. The other 4 edges were not present in the

KEGG as causal interactions for this pathway. However, all are either supported by

their presence in other KEGG pathways or by recent publications (we reiterate that

interactions for all six proteins in other KEGG pathways were excluded from the

training data as mentioned above so these predictions are not contaminated by their

presence in other pathways in KEGG). Specifically, (traf6, nfkb1) and (map3k7, nfkb1)

have been annotated as causal pairs in KEGG's 'Pertussis pathway' and 'RIG-I-like

receptor signaling pathway'[36], respectively. (rela, nfkb1) is the known p50/p65

heterodimer of NF-κB[40]. As for the (nfkbia, traf6) pair, it was found that traf6's binding

to MAP3K7 activates ikbkb which in turn phosphorylates nfkbia[41].

CNNC output as a similarity matrix for clustering

To evaluate CNNC's performance on the downstream analysis, we used it to generate

a gene-gene similarity matrix. We next used this matrix as an input for a hierarchical

clustering algorithm.

We extracted the top 2,000 (top 1,000, see **Supplementary Fig. 8**) differentially

expressed genes based on the expression data used in this paper using fano factor

(FF) (**Supplementary Note**). Since we have trained CNNC using the KEGG database,

we removed KEGG genes from the test set. Next we performed hierarchical

clustering[42] using CNNC and PC based on all sc and bulk data (**Fig. 5b**, and **5c**). For

comparison, we selected the top 8 clusters for the resulting hierarchical clustering tree

for all inputs (see also **Supplementary Fig. 8**). Next, for each input we calculated the

significant GO terms (q-values < 0.05)[43] and plotted the results in **Fig. 5d**. As can be

seen, using CNNC as the input led to the identification of more significant GO terms

for the same set of genes indicating that the clustering obtained using this input is

more aligned with current biological knowledge.

Discussion and conclusion

Gene co-expression analysis has been a widely successful method for the analysis of gene expression data starting over two decades ago with the introduction of microarrays. Several methods have been suggested for this task and several other methods use co-expression analysis as a component in a more elaborate modeling framework.

While co-expression analysis performs well in some cases, it suffers from a number of drawbacks that often led to overfitting (false positives) or missing key relationships (false negatives). The former can be attributed to the unsupervised nature of most co-expression methods making it hard to 'train' them on a labeled dataset. The latter often resulted from the nature of the data used for co-expression analysis (bulk or population of cells data) which led to masking of relationships that existed in single cells. Moreover, while certain more sophisticated methods attempted to utilize gene expression to infer causality (for example, Bayesian network based methods [44]) these were only able to detect directed interactions, were based on very specific probabilistic modeling assumptions, and did not directly provide a confidence score for the resulting edges.

To address these issues we presented CNNC, a general framework for co-expression analysis which is based on convolutional NN (CNN). The key idea here is to convert the input data into a co-expression histogram which is very suitable for CNNs. Unlike most prior methods our method is supervised which allows the CNN to zoom in on subtle differences between positive and negative pairs. Supervision also helps fine tune the scoring function based on the different application. For example, different features may be important for analyzing TF-gene interactions when compared to inferring proteins in the same pathway. In addition to the supervised approach the fact that the network can utilize the large volumes of scRNA-Seq data allows it to better overcome masking issues reducing false negative.

Analysis of several different interaction prediction tasks indicates that CNNC can improve upon prior, unsupervised methods. It can also be naturally extended to

integrate complementary data including epigenetic and sequence information. Finally,

CNNC is easy to use either with general data or with condition specific data. For the

former, users can download the data and implementation from the supporting website

(**Supplementary Fig. 9**), provide a list of labels (positive and negative pairs for their

system of interest) and retrieve the scores for all possible gene pairs. These in turn

can be used for any downstream application including clustering, network analysis

etc.

In addition to comparing CNNC to prior methods we have also used it to evaluate the

advantages conferred by scRNA-Seq data. Models trained with scRNA-Seq data

outperformed those trained with bulk data for all systems we looked at. This supports

prior findings[45, 46] and addressed a key criticism of co-expression analysis – that many

interactions are observed or missed due to aggregation effects from the collection of

cells rather than because they truly represent specific molecular events. While the

scRNA-Seq data we used contained two orders of magnitude more samples, the total

number of cells profiled is smaller (each bulk experiment often profiles at least three

orders of magnitude more cells than a single scRNA-Seq profile[47]). In addition,

scRNA-Seq coverage is often two orders of magnitude less than bulk experiments so

that total number of reads in the two datasets is not very different. Even when

comparing with the same number of profiles for bulk and sc we find that CNNC

performs better when using scRNA-Seq data. This result seems to indicate that

despite the much greater noise associated with scRNA-Seq, such data can provide

more accurate models for the same overall costs, coverage and sample size.

CNNC is implemented in Python and both data and an open source version of the

software are available from the supporting website. We hope that this method would

become a useful component in future co-expression studies.

Online methods

Dataset sources and pre-process pipelines

We used mouse scRNA-Seq dataset collected by Alavi et al[23]. The dataset consists of

uniformly processed 43,261 expression profiles from over 500 different scRNA-Seq

studies. For each profile, expression values are available for the same set of 20,463

genes. Mouse bulk RNA-Seq dataset were downloaded from Mouse Encode project

[24]. That data included 249 samples and we only utilized genes that are present in the

scRNA-Seq dataset leading to the same number of genes for both datasets. mESC

Dnase data was also downloaded from Mouse Encode project[24]

(ENCFF096WRW.bed). Mouse TF motif information is from TRANSFAC database[48].

PWM values were calculated by Python package 'Biopython'[49].

For the DNase and PWM analysis we followed prior papers and defined the

transcription start site (TSS) region as 10KB upstream to 1KB downstream from the

TSS for each gene[27, 28]. For each TF and gene pair, using Biopython package we

calculated the score between the TF motif sequence and both the '+/-' sequences at

all possible positions along the TSS region of the gene, and then selected the

maximum one as the final PWM score. The maximum Dnase peak signal in the TSS

region was calculated as the scalar Dnase value for each gene.

Labeled data:

mESC ChIP-seq peak region data was downloaded from GTRD database, and we

used peaks with threshold p value $< 10^{-300}$. If one TF X has at least one ChIP-seq

peak signal in or partially in the TSS region of gene Y, as defined above, we say that X

regulates Y.

KEGG and Reactome pathway data were downloaded by the R package 'graphite'[50].

KEGG contains 290 pathways and Reactome contains 1581 pathways. For both, we

only select directed edges with either activation or inhabitation edge types and filter

out cyclic gene pairs where genes regulate each other mutually (to allow for a unique

label for each pair). In total, we have 3,057 proteins with outgoing directed edges in

KEGG and the total number of directed edges is 33,127. For Reacotome the

corresponding numbers are 2,519 and 33,641.

Constructing the input histogram

For any gene pair *a* and *b*, we first log transformed their expression, and then

uniformly divided the expression range of each gene to 32 bins. Next we created the

32X32 histogram by assigning each sample to an entry in the matrix and counting the

number of samples for each entry. Due to the very low expression levels and even

more so to dropouts in scRNA data, the value in zero-zero position is always very

large and often dominates the entire matrix. To overcome this, we added

pseudocounts to all entries. We combined bulk and scRNA-Seq NEPDFs by

concatenating them as a 32X64 matrix to achieve better performance.

CNN for RPKM data

We followed VGGnet[51] to build our convolutional neural networks (CNN) model

(**Supplementary Fig. 1**). The CNN consists of stacked layers of x 3X3 convolutional

filters (equation (1)) (x is a power of 2, ranging from 32 to 64 to 128) and interleaved

layers of 2X2 maxpooling (equation (2)). We used the constructed input data as input

to CNN. Each convolution layer computes the following function:

$\text{Convolution } (X)_{i,j}^k = \sum_{m=1}^{3} \sum_{n=1}^{3} W_{i,j}^k X_{i+m,j+n}$ (1)

Where *X* is the input from the previous layer, (*i,j*) is output position, *k* is convolutional

filter index and *W* is the filter matrix of size 3X3. In other words, each convolutional

layer computes a weighted average of the prior layer values where the weights are

determined based on training. The maxpooling layer computes the following function:

$\text{maxpooling } (X)_{i,j}^k = \max(\{X_{i,j}^k, X_{i+1,j}^k, X_{i,j+1}^k, X_{i+1,j+1}^k\})$ (2)

Where *X* is input, (*i,j*) is output position and *k* is the convolutional filter index. In other

words, the layer selects one of the values of the previous layer to move forward.

Overall structure

The overall structure of the CNN is presented in **Supplementary Fig. 1**. The input layer of the CNN is either 32X32 or 32X64 as discussed above. In addition, the CNN contains 10 intermediate layers and a single one or three-dimension output layer. The ten layers include both convolutional and maxpooling layers, and the exact dimensions of each layer are shown in **Supplementary Fig. 1**. Following ref 52 [52] we used rectified linear activation function (ReLU) as the activation function (equation (3)) across the whole network, except the final classification layers where 'sigmoid' function (equation (4)) was used for two categories classification and 'softmax' function (equation (5)) for multiple categories classification. These functions are defined below.

$$\text{ReLU}\,(x) = \begin{cases} x & if\ x \geq 0 \\ 0 & if\ x < 0 \end{cases} \tag{3}$$

$$\text{Sigmoid}_\theta\,(x) = {1}/{1 + e^{\theta x}} \tag{4}$$

$$\text{Softmax}_\theta\,(x) = \frac{1}{\sum_{j=1}^{k} e^{\theta_j x}} \begin{bmatrix} e^{\theta_1 x} \\ e^{\theta_2 x} \\ ... \\ e^{\theta_k x} \end{bmatrix} \tag{5}$$

Training and testing strategy

We evaluated the CNN using cross validation. In these, training and test datasets are strictly separated to avoid information leakage. See **Supplementary Note, Supplementary Fig. 10** and **Supplementary Table 1** for details. For the three labels (causality analysis) we did the following: for each gene, we generated (*a, b*) (label1) and (*b, a*)'s (label2) NEPDF matrices. For the 0 label we generated a (*a, N*) NEPDF matrices for GTRD where *N* was a random gene and *a* was the TF. 0 labels for KEGG and Reactome were generated from random (*M, N*) gene pairs. After training, we used *p*1(*a, b*) + *p*2(*a, b*) as the probability that *a* interacts *b*, *p*2(*a, b*) – *p*2(*b, a*) as the pseudo probability that *b* regulates *a*.

Integrating expression, sequence and DNase data

To integrate Dnase and PWM data with the processed RNA-Seq data, we first

computed the max value for a PWM scan and DNase accessibility for each promotor

region. We next generated a two-value vector from this data for each pair and

embedded it to a 512D vector using one fully connected layer containing 512 nodes.

Next these are concatenated with the expression processed data to form a 1024D

vector which serves as input to a fully connected 512-node plus 128-node layer neural

network classifier. See **Supplementary Fig. 1** for details. Early stopping strategy by

monitoring validation loss function is used to avoid overfitting.

Selection of edges for the IL-17 pathway analysis

We performed leave-one-pathway-out validation to evaluate CNNC' performance for

predicting edges for individual pathways. We selected a relatively small pathway

('IL-17' from KEGG) to improve our ability to present it visually. We discuss more

general results for KEGG as well (**Fig. 4**).   For this analysis we only selected directed

edges with either activation or inhabitation types and filtered out cyclic gene pairs

where genes regulate each other mutually to purify the edge types. In total, we had 6

nodes and 4 directed edges for the IL-17 pathway. Next, we trained CNNC with the

entire KEGG dataset excluding any interactions for the six 'IL-17' pathway proteins.

Hierarchical clustering and GO term enrichment analysis

We performed hierarchical clustering followed by GO term enrichment analysis to

evaluate CNNC' performance in downstream analysis. We selected the top 2,000 (or

1,000 (**Supplementary Fig. 8**)) genes with highest Fano factor (**Supplementary**

Note) We obtained the similarity matrices for the filtered gene list based on CNNC, sc

PC, bulk PC and sc&bulk PC respectively. We cut the tree at 8 clusters for all inputs.

Next, we performed GO term enrichment analysis using fisher's exact test and

counted the significant GO terms for each of the cluster result. Significance of

difference   between   different   inputs   was   computed   using   one-side

Wilcoxon-Mann-Whitney test for the q-values of the four strategies (**Fig. 5d**).

Acknowledgements

Work partially supported by NIH grant 1R01GM122096, US National Science

Foundation (DBI-1356505) to ZBJ. and a James S. McDonnell Foundation Scholars

Award in Studying Complex Systems to Z.B.-J.

Author contributions

Y.Y. and Z.B conceived the method. Y.Y. implemented CNNC and the support website.

Y.Y. and Z.B designed the experiments. Y.Y. and Z.B wrote the manuscript.

Competing interests

None

References

1.　De Smet, R. & Marchal, K. Advantages and limitations of current network inference

methods. *Nat Rev Microbiol* **8**, 717-729 (2010).

2.　Stuart, J.M., Segal, E., Koller, D. & Kim, S.K. A gene-coexpression network for global

discovery of conserved genetic modules. *Science* **302**, 249-255 (2003).

3.　van Dam, S., Vosa, U., van der Graaf, A., Franke, L. & de Magalhaes, J.P. Gene

co-expression analysis for functional classification and gene-disease predictions. *Brief*

Bioinform (2017).

4.　Butte, A.J. & Kohane, I.S. Unsupervised knowledge discovery in medical databases

using relevance networks. *Proc AMIA Symp*, 711-715 (1999).

5.　Eisen, M.B., Spellman, P.T., Brown, P.O. & Botstein, D. Cluster analysis and display

of genome-wide expression patterns. *Proc Natl Acad Sci U S A* **95**, 14863-14868

(1998).

6.　Newman, M.E. Spectral methods for community detection and graph partitioning.

Phys Rev E Stat Nonlin Soft Matter Phys **88**, 042822 (2013).

7.　Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation network

analysis. *BMC Bioinformatics* **9**, 559 (2008).

8.　Bartlett, T.E., Muller, S. & Diaz, A. Single-cell Co-expression Subnetwork Analysis. *Sci*

Rep **7**, 15066 (2017).

9.　Chan, T.E., Stumpf, M.P.H. & Babtie, A.C. Gene Regulatory Network Inference from

Single-Cell Data Using Multivariate Information Measures. *Cell Syst* **5**, 251-267 e253

(2017).

10.   Huynh-Thu, V.A., Irrthum, A., Wehenkel, L. & Geurts, P. Inferring regulatory networks

from expression data using tree-based methods. *PLoS One* **5** (2010).

11.   Faith, J.J. et al. Large-scale mapping and validation of Escherichia coli transcriptional

regulation from a compendium of expression profiles. *PLoS Biol* **5**, e8 (2007).

12.   Lee, H.K., Hsu, A.K., Sajdak, J., Qin, J. & Pavlidis, P. Coexpression analysis of human

genes across many microarray data sets. *Genome Res* **14**, 1085-1094 (2004).

13.   Song, L., Langfelder, P. & Horvath, S. Comparison of co-expression measures:

mutual information, correlation, and model based indices. *BMC Bioinformatics* **13**, 328

(2012).

14.   Daub, C.O., Steuer, R., Selbig, J. & Kloska, S. Estimating mutual information using

B-spline functions--an improved similarity measure for analysing gene expression

data. *BMC Bioinformatics* **5**, 118 (2004).

15.   Basso, K. et al. Reverse engineering of regulatory networks in human B cells. *Nat*

Genet **37**, 382-390 (2005).

16.   Kotlyar, M., Fuhrman, S., Ableson, A. & Somogyi, R. Spearman correlation identifies

statistically significant gene expression clusters in spinal cord development and injury.

Neurochem Res **27**, 1133-1140 (2002).

17.   Qian, J., Dolled-Filhart, M., Lin, J., Yu, H. & Gerstein, M. Beyond synexpression

relationships: local clustering of time-shifted and inverted gene expression profiles

identifies new, biologically relevant interactions. *J Mol Biol* **314**, 1053-1066 (2001).

18.   Reverter, A. & Chan, E.K. Combining partial correlation and an information theory

approach to the reversed engineering of gene co-expression networks. *Bioinformatics*

24, 2491-2497 (2008).

19. Kumari, S. et al. Evaluation of gene association methods for coexpression network

construction and biological knowledge discovery. *PLoS One* **7**, e50411 (2012).

20. Wang, Y.X., Waterman, M.S. & Huang, H. Gene coexpression measures in large

heterogeneous samples using count statistics. *Proc Natl Acad Sci U S A* **111**,

16371-16376 (2014).

21. Freytag, S., Gagnon-Bartsch, J., Speed, T.P. & Bahlo, M. Systematic noise degrades

gene co-expression signals but can be corrected. *BMC Bioinformatics* **16**, 309 (2015).

22. Crow, M., Paul, A., Ballouz, S., Huang, Z.J. & Gillis, J. Exploiting single-cell

expression to characterize co-expression replicability. *Genome Biol* **17**, 101 (2016).

23. Amir Alavi, M.R., Aiyappa Parvangada, Zhilin Huang, Ziv Bar-Joseph scQuery: a web

server for comparative analysis of single-cell RNA-seq data.   (2018).

24. Yue, F. et al. A comparative encyclopedia of DNA elements in the mouse genome.

Nature **515**, 355-364 (2014).

25. Johnson, D.S., Mortazavi, A., Myers, R.M. & Wold, B. Genome-wide mapping of in

vivo protein-DNA interactions. *Science* **316**, 1497-1502 (2007).

26. Yevshin, I., Sharipov, R., Valeev, T., Kel, A. & Kolpakov, F. GTRD: a database of

transcription factor binding sites identified by ChIP-seq experiments. *Nucleic Acids*

Res **45**, D61-D67 (2017).

27. Schulz, M.H. et al. Reconstructing dynamic microRNA-regulated interaction networks.

Proc Natl Acad Sci U S A **110**, 15686-15691 (2013).

28. Schulz, M.H. et al. DREM 2.0: Improved reconstruction of dynamic regulatory

networks from time-series expression data. *BMC Syst Biol* **6**, 104 (2012).

29. Wang, J. et al. Single-Cell Co-expression Analysis Reveals Distinct Functional

Modules, Co-regulation Mechanisms and Clinical Outcomes. *PLoS Comput Biol* **12**,

e1004892 (2016).

30. Khan, A. et al. JASPAR 2018: update of the open-access database of transcription

factor binding profiles and its web framework. *Nucleic Acids Res* **46**, D260-D266

(2018).

31. Gitter, A., Carmi, M., Barkai, N. & Bar-Joseph, Z. Linking the signaling cascades and

dynamic regulatory networks controlling stress responses. *Genome Res* **23**, 365-376

(2013).

32. van Noort, V., Snel, B. & Huynen, M.A. Predicting gene function by conserved

co-expression. *Trends Genet* **19**, 238-242 (2003).

33. Parikshak, N.N. et al. Integrative functional genomic analyses implicate specific

molecular pathways and circuits in autism. *Cell* **155**, 1008-1021 (2013).

34. Willsey, A.J. et al. Coexpression networks implicate human midfetal deep cortical

projection neurons in the pathogenesis of autism. *Cell* **155**, 997-1007 (2013).

35. Young, J.H. & Marcotte, E.M. Predictability of Genetic Interactions from Functional

Gene Modules. *G3 (Bethesda)* **7**, 617-624 (2017).

36. Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y. & Morishima, K. KEGG: new

perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res* **45**,

D353-D361 (2017).

37.   Fabregat, A. et al. The Reactome Pathway Knowledgebase. *Nucleic Acids Res* **46**,

582   D649-D655 (2018).

583   38.   Angelini, C. & Costa, V. Understanding gene regulatory mechanisms by integrating

584   ChIP-seq and RNA-seq data: statistical solutions to biological problems. *Front Cell*

585   *Dev Biol* **2**, 51 (2014).

586   39.   D'Haeseleer, P., Liang, S. & Somogyi, R. Genetic network inference: from

587   co-expression clustering to reverse engineering. *Bioinformatics* **16**, 707-726 (2000).

588   40.   Chen, F.E., Huang, D.B., Chen, Y.Q. & Ghosh, G. Crystal structure of p50/p65

589   heterodimer of transcription factor NF-kappaB bound to DNA. *Nature* **391**, 410-413

590   (1998).

591   41.   Kramer, I. Signal Transduction. (Elsevier, 2015).

592   42.   Bar-Joseph, Z., Gifford, D.K. & Jaakkola, T.S. Fast optimal leaf ordering for

593   hierarchical clustering. *Bioinformatics* **17 Suppl 1**, S22-29 (2001).

594   43.   Ashburner, M. et al. Gene ontology: tool for the unification of biology. The Gene

595   Ontology Consortium. *Nat Genet* **25**, 25-29 (2000).

596   44.   Spirtes, P. & Zhang, K. Causal discovery and inference: concepts and recent

597   methodological advances. *Appl Inform (Berl)* **3**, 3 (2016).

598   45.   Wagner, A., Regev, A. & Yosef, N. Revealing the vectors of cellular identity with

599   single-cell genomics. *Nat Biotechnol* **34**, 1145-1160 (2016).

600   46.   Wang, Y. & Navin, N.E. Advances and applications of single-cell sequencing

601   technologies. *Mol Cell* **58**, 598-609 (2015).

47.   Ozsolak, F. & Milos, P.M. RNA sequencing: advances, challenges and opportunities.

Nat Rev Genet **12**, 87-98 (2011).

48.   Wingender, E., Dietze, P., Karas, H. & Knuppel, R. TRANSFAC: a database on

transcription factors and their DNA binding sites. *Nucleic Acids Res* **24**, 238-241

(1996).

49.   Cock, P.J. et al. Biopython: freely available Python tools for computational molecular

biology and bioinformatics. *Bioinformatics* **25**, 1422-1423 (2009).

50.   Sales, G., Calura, E., Cavalieri, D. & Romualdi, C. graphite - a Bioconductor package

to convert pathway topology to gene network. *BMC Bioinformatics* **13**, 20 (2012).

51.   Karen Simonyan, A.Z. Very Deep Convolutional Networks for Large-Scale Image

Recognition. (2014).

52.   Xavier Glorot, A.B., Yoshua Bengio Deep Sparse Rectifier Neural Networks.

Proceedings of the Fourteenth International Conference on Artificial Intelligence and

Statistics, PMLR, 315-323 (2011).

Figure legends

Figure 1 The CNNC architecture

For each gene pair, expression levels from bulk and sc RNA-Seq are transformed into two 32X32 normalized empirical probability function (NEPDF) matrices, and the two are concatenated to form a combined 32X64 NEPDF matrix (left). The combined NEPDF serves as an input to a convolutional neural network (CNN). The intermediate layer of the CNN can be further concatenated with input vectors representing Dnase-seq and PWM data (top). The output layer contains three probability nodes where *p0* is the probability that genes *a* and *b* are not interacting, *p1* encodes the case that gene *a* regulates gene *b*, and *p2* is the probability that gene *b* regulates gene *a*.

Figure 2 GTRD TF-target prediction

(**a**) ROCs for Pearson Correlation (PC) based on scRNA-Seq. Light gray lines represent the performance for each TF. Black line represents the median ROC, and light green region represents the 25~75 quantile. (**b**) PC for bulk RNA-Seq. (**c**) PC for combined bulk and scRNA-Seq. (**d**) ROCs for Mutual Information (MI) when using scRNA-Seq. (**e**) MI when using bulk RNA-Seq. (**f**) MI using the combined bulk and scRNA-Seq. (**g**) ROCs for CNNC using scRNA-Seq. (**h**) CNNC using bulk RNA-Seq. (**i**) CNNC for combined data. Inset – Top ranking CNNC pairs are much more likely to be correct pairs when compared to other methods. (**j**) Comparison of TF-target predictions with additional data. Columns 1-3 show median AUROC of PC, MI, and CNNC using scRNA-Seq, bulk and the combine data, respectively. 4^{th} column shows prediction of TF-gene interactions using only PWM or Dnase. 5^{th} column shows performance when integrating expression, sequence and DNase data.

Figure 3 Predicting undirected pathway edges

(**a**) Overall ROCs for CNNC performance on KEGG pathway undirected edge prediction with bulk and scRNA-Seq. (**b**) The Area Under the Receiver Operating Characteristic curve (AUROC) histogram for (**a**). (**c**) The overall ROCs for performance of CNNC on Reactome pathway undirected edge prediction with bulk and scRNA-Seq. (**d**) The AUROC histogram for (**c**).

Figure 4 Directed (causal) edge prediction

(**a**) Overall ROCs for performance of CNNC on GTRD directed prediction with bulk and scRNA-Seq. (**b**) The AUROC histogram for (**a**). (**c**) Overall ROCs for performance of CNNC on KEGG pathway directed edge prediction with bulk and scRNA-Seq. (**d**) The AUROC histogram for (**c**). (**e**) Overall ROCs for performance of CNNC on Reactome pathway directed edge prediction with bulk and scRNA-Seq. (**f**) The

AUROC histogram for (**e**). (**g**) A typical NEPDF sample from a KEGG interaction that

is correctly predicted as label 1. (**h**) A typical NEPDF sample that is correctly predicted

as label 2.

Figure 5 Downstream applications using CNNC

CNNC can be used as a component in downstream analysis algorithms including for

pathway analysis and clustering. (**a**) Top: Directed edges annotated in KEGG for the

IL-17 pathway Bottom: predicted directed edges for the pathway. (**b**) Hierarchical

clustering of top 2,000 DE genes based on CNNC similarity matrix score. The number

under the horizontal line represents the distance between the two groups, and the

black horizontal line shows the resulting eight clusters groups. (**c**) Hierarchical

clustering based using PC as the input. (**d**) GO term analysis of the clustering results

from (**Figs. 5b, 5c, Supplementary Figs. 8a** and **8b**).

a
KEGG
0.9714

c
Reactome
0.8487

b

d