Thy1 (CD90) expression is regulated by DNA methylation during adipogenesis

E'Lissa M. Flores[1], Collynn F. Woeller*[2], Megan L. Falsetta[2], Martha Susiarjo[2], Richard P. Phipps[1,2,3]

Clinical & Translational Science Institute[1], Department of Environmental Medicine[2], Department of Microbiology and Immunology[3], University of Rochester School of Medicine and Dentistry. Rochester, NY. *Corresponding author, Collynn_Woeller@urmc.rochester.edu

Corresponding author (to whom all correspondence should be addressed):
Collynn Woeller, Ph.D.
Research Associate Professor
University of Rochester
Department of Environmental Medicine
601 Elmwood Ave, Box 850, Room 3-11001
Rochester, NY 14642
Phone: 585-275-7836

The authors have no conflict of interest.

Work funded by F31ES027767, TL1-TR000096, ES001247, and ES0023032

Running title: Thy1 methylation and adipogenesis
Key words: adipogenesis, Thy1 (CD90), methylation, 5-aza-dC, epigenetics

Abstract

The obesity epidemic is developing into the most costly health problem facing the world. Obesity, characterized by excessive adipogenesis and enlarged adipocytes, promotes morbidities such as diabetes, cardiovascular disease and cancer. Regulation of adipogenesis is critical to our understanding of how fat cell formation causes obesity and associated health problems. Thy1 (also called CD90), a widely used stem cell marker, blocks adipogenesis and reduces lipid accumulation. Thy1 knockout-mice are prone to diet-induced obesity. While the importance of Thy1 in adipogenesis and obesity is now evident, how its expression is regulated is not. We hypothesized that DNA methylation plays a role in promoting adipogenesis and affects *Thy1* expression. Using the methylation inhibitor 5-aza-2'-deoxycytidine (5-aza-dC), we investigated whether DNA methylation alters *Thy1* expression during adipogenesis in both mouse 3T3-L1 pre-adipocytes and mouse mesenchymal stem cells. Thy1 protein and mRNA levels were decreased dramatically during adipogenesis. However, 5-aza-dC treatment prevented this phenomenon. Pyrosequencing analysis shows that the CpG sites at the *Thy1* locus are methylated during adipogenesis. These new findings highlight the potential role of *Thy1* and DNA methylation in adipogenesis and obesity.

Introduction

Obesity rates have risen markedly in the last 30 years. More than 700 million people worldwide are clinically obese[1,2]. Obesity promotes type 2 diabetes, fatty liver disease, and cardiovascular disease, and is linked with certain cancers[3-5]. Health care costs associated with obesity and its comorbidities are enormous and will continue to

rise as currents trends continue and the population ages[6]. Thus, a further understanding

of obesity and its underlying mechanisms and causes are urgently needed.

Obesity results from a positive energy balance when more calories are

consumed than are used. The surplus energy is packaged into lipid-based storage

molecules and sent to fat storage cells called adipocytes. In obesity, there is both an

increase in adipocyte size and an increase in adipocyte number to accommodate the

lipid[7]. Adipocytes are formed during the process of adipogenesis and arise from stem

cells, fibroblasts, or other progenitor cells when appropriately programmed[8].

Adipogenesis is a highly regulated process that requires the activation of several key

signaling pathways, including STAT5 and Fyn and activation of the transcription factors

PPARγ and C/EBPα. Numerous genes involved in fatty acid transport and storage, such

as fatty acid binding protein 4 (Fabp4) are induced during adipogenesis to promote lipid

accumulation in adipocytes.

Several proteins including Pref-1, Wnt, TGFβ, and Thy1 (formally called CD90)

have been shown to inhibit adipogenesis by blocking pro-adipogenic signaling[9-13]. Our

recent study showed that Thy1 blocked the activity of the Src family kinase, Fyn, in pre-

adipocytes[14]. Thy1 mediated inhibition of Fyn activity prevented adipocyte formation.

Interestingly, while pre-adipocytes expressed high levels of Thy1, its expression was

lost during adipogenesis, and mature adipocytes expressed almost no Thy1. Thy1 is a

member of the immunoglobulin supergene family and is a glycophosphatidyl inositol

linked surface protein. While Thy1 is expressed on pre-adipocytes and subsets of

fibroblasts, neurons, and stem cells, little is known about how its expression is

controlled. We recently showed that Thy1 levels can be regulated by microRNAs.

Specifically, the miR-103/107 family of miRNAs can target Thy1 mRNA and reduce its

expression[15]. Furthermore, the *Thy1* gene contains several CpG rich elements termed

CpG islands, which are hotspots for cytosine methylation and gene regulation[16,17]. To

date, there have been no reports studying DNA methylation of *Thy1* during

adipogenesis. However, *Thy1* methylation has been studied in context of fibrosis and in

T cells, where they show an increase in *Thy1* methylation correlated with a decrease in

Thy1 expression[18-20]. Therefore, we investigated the same CG rich region within intron 1

(termed Thy1-CGI1), which is part of the promoter[17,20-23].

While changes in DNA methylation patterns at the *Thy1* locus have not been

characterized in the context of adipogenesis, recent reports have shown that DNA

methylation changes are an integral part of adipocyte formation[24,25]. In the most widely

used and well accepted model of adipogenesis, the murine 3T3-L1 pre-adipocyte line,

global DNA methylation has been shown to increase during adipogenesis[26,27].

Interestingly, addition of the DNA methylation inhibitor 5-aza-2'-deoxycytidine (5-aza-

dC) at the onset of adipogenic differentiation can completely block adipocyte formation,

suggesting that DNA methylation changes are essential for adipogenesis[27,28]. Specific

changes in regional DNA methylation are also critical for adipogenesis, and blocking

these patterns can alter the differentiation pathway of precursor cells towards

osteoblastogenesis rather than adipogenesis[24,29,30]. For example, when blocking DNA

methylation using 5-aza-dC in 3T3-L1 cells, Wnt10a expression increases through the

hypomethylation of several of its CpG sites. The increase in Wnt10a expression helps

steer the differentiation program towards osteoblastogenesis and away from

adipogenesis[30].

Because Thy1 protein and mRNA levels are rapidly suppressed during

adipogenic differentiation, we hypothesized that its reduced expression is linked to

hypermethylation of the *Thy1* locus. Excitingly, we are the first to show methylation

sensitive pyrosequencing data of the Thy1-CGI1 region and report herein that DNA

methylation is one of the essential regulators of *Thy1* expression during adipogenesis.

Results

Reduced Thy1 expression during adipogenesis is partially attenuated by 5-aza-dC in

bone marrow-derived mouse mesenchymal stem cells.

Previously, we demonstrated that pre-adipocytes and mesenchymal stem cells

must down-regulate *Thy1* expression to differentiate into adipocytes[14]. To determine if

DNA methylation regulates *Thy1* expression during adipogenesis, we first

pharmacologically inhibited DNA methyltransferase in bone-marrow derived mouse

mesenchymal stem cells (mMSCs) before differentiation into adipocytes[28]. mMSCs

were treated daily with either DMSO (vehicle) or the DNA methyltransferase inhibitor, 5-

aza-dC for 12 days and were concurrently treated with either media alone or with the

adipogenic cocktail medium (ACT), starting at day 3 (Fig. 1A). mMSCs typically

differentiate into adipocytes in 2 weeks when exposed to ACT[31]. Therefore, we

examined an intermediate time point to examine Thy1 levels. As expected, when given

ACT, Thy1 levels decreased while fatty acid binding protein 4 (Fabp4) levels were

increased compared to media alone (Fig. 1B-D); Fabp4 is highly expressed by

adipocytes and serves as an adipogenic marker. However, inhibiting DNA methylation

with 5-aza-dC increased Thy1 protein and mRNA levels versus media alone and

attenuated the down-regulation of Thy1 in ACT samples (Fig. 1B-C). We also

determined that protein and mRNA levels of Fabp4, a protein highly expressed by

adipocytes that serves as an adipogenic marker, were decreased in mMSCs treated

with 5-aza-dC (Fig. 1B,D). Immunofluorescence staining revealed that compared to

cells not treated with ACT, 5-aza-dC-treated mMSCs had increased surface expression

of Thy1 and decreased Fabp4 levels (Fig. 1E). These results indicate that inhibition of

DNA methylation impairs adipogenesis.

Reduced Thy1 expression, which is necessary for adipogenesis, is prevented by

inhibition of global DNA methylation in 3T3-L1 cells

Primary mesenchymal stem cells are heterogeneous with only some cells fully

differentiating into adipocytes[32]. Thus, we next used the well-established pre-adipocyte

murine cell line, 3T3-L1. Previously, we showed that fully differentiated adipocytes

treated with ACT no longer expressed Thy1 protein and mRNA[33]. To observe changes

in Thy1 expression, we examined an intermediate point in adipogenesis. At only four

days of treatment with ACT the cells are only partially differentiated, and Fabp4

expression increases. Cells were treated daily for 7 days with either DMSO or 5-aza-dC

and starting at day 3 were either given ACT or continued with media alone (Fig. 2A). As

expected, Thy1 decreased at both the protein and mRNA level in cells given ACT

compared to media alone (Fig 2B-C). However, we show that treatment with 5-aza-dC

prevents the decrease in Thy1 mRNA levels in ACT-treated cells, while media alone

with 5-aza-dC causes no significant changes (Fig 2C). 5-aza-dC also caused a

decrease in Fabp4 levels in ACT-treated cells, indicative of decreased adipogenesis

(Fig 2B,D). Inhibiting global methylation blunts adipogenesis, which is reflected by the

decreased levels of Fabp4 expression and in part sustains Thy1 levels.

5-aza-dC partially restores Thy1 cell surface expression in 3T3-L1 cells when exposed

to adipogenic cocktail

We next examined Thy1 cell surface expression on 3T3-L1 cells, since Thy1 is a

known cell marker on pre-adipocytes and is readily detected via flow cytometry and

immunofluorescence. Cells were treated for 7 days daily with either DMSO or 5-aza-dC,

while ACT samples were given the adipogenic cocktail starting at day 3, as previously

described. The representative histogram in Figure 3A shows cells treated with ACT shift

out of the Thy1+ gate into Thy1- during differentiation, while cells treated with ACT and

5-aza-dC mostly remain in the Thy1+ gate. As expected, pre-adipocytes treated with

ACT expressed significantly less surface Thy1 than cells with media alone, as

evidenced by a lower mean fluorescence intensity (MFI) (Fig 3B). However, pre-

adipocytes cultured with ACT and 5-aza-dC showed a significant increase in Thy1 MFI

compared to ACT alone, which occurred in tandem with an increase in the percentage

of Thy1-positive cells (Fig 3B-C). Using immunofluorescence, we confirmed that Thy1

expression was sustained in ACT-treated pre-adipocytes also treated with 5-aza-dC,

while Fabp4 expression decreased compared to cells treated with ACT alone (Fig. 3D).

Thy1(CD90) gene expression is regulated by DNA methylation during adipogenesis

We went on to examine *Thy1* DNA methylation, which typically occurs in cytosine

(CG) rich regions and is commonly associated with gene silencing [34,35]. As the

transcriptional activation of the *Thy1* gene involves both the promoter and intron 1 in

some cell types[21,23], and previous publications have referred to intron 1 as part of the

promoter[17,19-23], we focused on a CpG rich region within intron 1 that we termed Thy1-

CGI1 (Fig 4A). Using a pyrosequencer, we analyzed methylation levels of 5 consecutive

CpG sites within Thy1-CGI1. 3T3-L1 pre-adipocytes were treated as described

previously. We found that Thy1-CGI1 is hypermethylated during differentiation

comparing the average DNA methylation of ACT-treated samples to those treated with

media alone. Treatment with 5-aza-dC resulted in reduced methylation of these CpG

sites in the *Thy1* gene in both media alone and ACT-treated samples (Fig. 4B).

Furthermore, individual CpG positions within Thy1-CGI1 showed an increase in DNA

methylation when treated with the adipogenic cocktail versus media alone, with a

significant increase at CpG position 2 (Fig 4C). Methylation decreased across all 5 CpG

sites when treated with 5-aza-dC, whether the cells were treated with media alone or

with ACT (Fig 4D-E). Four of the five CpG sites examined had significantly reduced

methylation levels when cells were treated with ACT and 5-aza-dC. Our data indicates

that this region is methylation sensitive and can influence *Thy1* expression during

adipogenesis.

To test the methylation status of *Thy1* over time during adipocyte differentiation,

we examined methylation levels at these five sites at days 0 (prior to ACT), 2, 4, and 6.

Cells were pretreated with DMSO or given 5-aza-dC for 24 h, then either harvested at

day 0 or given ACT medium and with continued treatment with DMSO (vehicle) or 5-

aza-dC daily. Overall, average DNA methylation percentages increased in a time-

dependent manner when exposed to the cocktail, with the highest degree of methylation

occurring on day 6 (Fig 5A). Day 0 samples showed lower methylation levels compared

to other time points in which cells had been exposed to ACT. Furthermore, there were

no significant changes at any of the 5 CpG sites when given a single dose of 5-aza-dC

(Fig. 5B). However, day 6 samples showed a significant two-fold increase in methylation

compared to day 0, along with a significant decrease in methylation when cells were

exposed daily to 5-aza-dC at all five CpG sites (Fig. 5C). This implies that during the

normal adipogenic process, CpG sites at the *Thy1* locus become hypermethylated,

which may blunt *Thy1* expression and allow for differentiation to occur.

Discussion

Excessive adipogenesis can lead to weight gain and obesity, which affects over

700 million people worldwide. The consequences of obesity can be dire, including the

development of cardiovascular or liver disease, diabetes, and other comorbidities, which

result in significant morbidity and mortality. Therefore, understanding the adipogenic

pathway(s) and molecular changes that foster adipocyte differentiation will elucidate the

mechanisms contributing to the pathogenesis of obesity. New understanding should

lead to better solutions for this growing problem, as lifestyle changes (e.g. improved diet

and exercise) are often insufficient. Although genome-wide studies have shown that

changes in histone methylation/acetylation occur during adipogenesis[25,36,37], few

specific adipogenesis-relevant genes have been identified as influenced by epigenetic

changes (e.g. DNA methylation). In this study, we identify *Thy1* as a methylation

sensitive gene and demonstrate that DNA methylation plays an active role in

adipogenesis. Ultimately, we found that inhibiting DNA methylation blunts adipogenesis

and sustains Thy1 at levels that may retard or suppress adipogenesis.

Thy1 is a cell surface protein that is expressed on mouse thymocytes and on

both mouse and human pre-adipocytes[38,39]. We have previously shown in mouse 3T3-

L1 cells that Thy1 is down-regulated in a time dependent manner during adipocyte

differentiation, while cells overexpressing Thy1 cells no longer differentiate, even when

given an adipogenic cocktail, which typically causes 3T3-L1 cells to differentiate into

adipocytes after 6-8 days of exposure. However, here we show that 5-aza-dC blunts

adipogenesis, even in the presence of ACT. These findings correlate with reduced

levels of Fabp4 expression at both the protein and mRNA levels, along with partially

attenuated levels of Thy1 when exposed to a global methylation inhibitor. Fabp4 levels

inversely correlate with Thy1 levels; Fabp4 expression increases while Thy1 is

suppressed during adipogenesis[14]. Therefore, blunted levels of Fabp4 are likely another

contributing factor that impedes adipogenesis. We also demonstrated that during

adipogenesis, Thy1-CGI1 methylation increases, which likely contributes to reduced

Thy1 expression. While we saw significant changes at CpG position 2 in ACT-treated

samples relative to untreated (media alone) cells (Fig. 4C), we observed similar trends

for the other CpG positions. These results are consistent with previous studies showing

alterations in methylation status at CpG sites can cause significant changes in gene

expression[40,41]. Treatment with the methylation inhibitor, 5-aza-dC, resulted in

hypomethylation of Thy1-CGI1, which in part may contribute to the attenuation of *Thy1*

expression. Since 5-aza-dC is a global DNA methylation inhibitor, it can affect other

genes involved in adipogenesis in addition to *Thy1*. Our data suggest that DNA

methylation is a necessary regulatory mechanism for pre-adipocytes to differentiate into

fat cells, consistent with recent reports implying DNA methylation is involved in lineage-

specific adipocyte development[24,42].

Adipogenesis is a complex process that is controlled by many factors, which

include epigenetic and post-transcriptional modifications. Previous studies have

established that the expression/activity of microRNAs, small RNAs ~20-22 bp in length

that bind to and block the transcription of specific targeted genes, play a role

adipogenesis[43]. Furthermore, in obesity, it has been shown that there are significant

changes in the expression of microRNAs (miR), such as an increase in miR-103

levels[44,45]. We have recently shown that miR-103 levels increase during adipogenesis

and can bind the 3' UTR of Thy1 to blunt its expression[15]. Here, we confirmed that the

levels of miR-103 increase during adipogenesis, and that these levels were unchanged

by treatment with 5-aza-dC, suggesting that the regulatory effects of *Thy1* methylation

are separate from changes in microRNA expression (Figure S3). This represents

another potential facet of *Thy1* regulation. Therefore, several, possibly overlapping

mechanisms are involved in the adipogenic pathway. This has important implications for

developing therapeutic interventions to combat adipogenesis and obesity; more than

one aspect of this regulatory mechanism may need to be targeted to cause a significant

effect.

Stem cells, such as, mesenchymal stem cells (MSCs) are distinguished and

defined by various cell surface markers, including Thy1[46]. Mesenchymal stem cells can

also differentiate into osteocytes and chondrocytes[39], where Thy1 is expressed

heterogeneously in each subset[47]. Recent studies have shown 5-aza-dC prevents

adipogenesis and promotes osteoblastogenesis through the activation of Wnt10a[30].

Wnt10a is known to be upregulated and essential for bone formation[48-50]. However,

Thy1 also plays a critical role[51,52]. It was recently shown that Thy1 is upregulated during

osteoblastogenesis and that Thy1[-] cells (knockdown and knockout) cannot differentiate

into osteoblasts[53]. However, changes in *Thy1* DNA methylation during osteoblast

formation have not been analyzed. In our present study, we show that Thy1 levels are

sustained, while FABP4 levels are lowered in mMSCs treated with 5-aza-dC in the

presence of ACT. While many epigenetic factors aid in stem cell maintenance,[54]

changes in the methylation status of genes, such as *Thy1*, may alter stem cell state.

Since *Thy1* is highly expressed in MSCs and pre-adipocytes, maintaining *Thy1*

expression may be the key to remaining a precursor cell. However, further investigation

is needed. While we saw an increase in *Thy1* methylation in mouse pre-adipocyte cells,

a crucial next step would be to determine basal *Thy1* expression levels and the

methylation status of MSCs derived from adipose tissues of obese and non-obese

individuals. Testing the ability of these cells to differentiate into fat cells may correlate

with their *Thy1* expression profiles. Such investigations are likely to provide additional

evidence of Thy1's critical role in adipogenesis and would further underline the

importance of our findings.

While we show that 5-aza-dC treatment decreased methylation in the Thy1-CGI1

region and blunted adipogenesis, it is possible that hypomethylating the *Thy1* gene at

the same time could affect other cell types, such as, fibroblasts (involved in fibrosis); it

has been established that *Thy1* expression is up-regulated during and involved in

myofibroblast differentiation[55]. Future studies could examine other CpG islands in the

traditional promoter region and downstream regions to investigate whether methylation

of these sites is also essential for *Thy1's* involvement in adipogenesis, fibrosis, and

other functions. While the *Thy1* gene may be a key target for methylation during

adipogenesis, there are undoubtedly other genes regulated by methylation during

differentiation. Further investigation is necessary to identify other essential genes, which

is fundamental to understanding the adipogenic pathway.

In summary, our work shows for the first time that *Thy1* has increased DNA

methylation during adipogenesis. We demonstrate herein that inhibiting DNA

methylation attenuates the loss of *Thy1* when cells are stimulated to differentiate into

adipocytes. Blocking methylation leads to sustained *Thy1* expression and prevents

adipogenesis. These studies further highlight the role of genomic methylation and

Thy1's involvement in adipogenesis, which suggests these pathways may be

dysregulated in metabolic diseases in which adipogenesis is elevated, such as obesity.

Materials and Methods

Chemicals

5-aza-2′-deoxycytidine (5-aza-dC), 3-isobutyl-1-methylxanthine (IBMX),

dexamethasone, and human recombinant insulin were all purchased from Sigma-

Aldrich (St. Louis, MO).

Cell Culture

All cells were incubated at 37°C with 7% humidified CO_2. 3T3-L1 cells were maintained

in 10% calf serum supplemented with DMEM media. C57BL/6 Mouse Bone Marrow

Mesenchymal Stem Cells were purchased from Cell Biologics (Chicago, IL) and

maintained in 10% mesenchymal stem cell-qualified fetal bovine serum in

supplemented MEM media from Thermo Fisher. Cells were plated at 60% confluence

and treated at 80% confluency. To induce adipogenesis, media containing an

adipogenic cocktail (ACT) was added to confluent cells, which consists of 0.5 mM IBMX,

0.5 µM dexamethasone, and 2 µg/ml insulin. Fresh ACT was added every 2 days. To

inhibit methylation, cells were treated daily with 0.5uM 5-aza-dC or DMSO as a control.

Cells were then harvested on days indicated per experiment.

Quantitative real-time PCR (qPCR) detection of mRNA

RNA was extracted with a Qiagen miRNeasy Kit and quantified using a NanoDrop 1000

spectrophotometer (Thermo Scientific, Wilmington, DE). A BioRad iScript reverse

transcription kit was used to make cDNA from 150 ng RNA. RT-qPCR assays were then

performed using SsoAdvanced Universal SYBR Green Supermix (Bio-Rad), according

to the manufacturer's instructions. All genes of interest were normalized to 18S rRNA,

and the relative percentages were normalized to 100% to media + DMSO for Thy1

mRNA or 100% to ACT + DMSO for Fabp4 mRNA levels. Primer sequences were as

follows Thy1: Fwd 5'-CCTTACCCTAGCCAACTTCAC and Rv 5'-

AGGATGTGTTCTGAACCAGC; Fabp4: Fwd 5'-ATGTGTGATGCCTTTGTGGGAAC

and Rv5'-TCATGTTGGGCTTGGCCATG; 18s rRNA: Fwd 5'-

GTAACCCGTTGAACCCCATT and Rv 5'-CCATCCAATCGGTAGTAGCG.

Western blot analysis

Cells were lysed with 60 mM Tris, 2% SDS, and protease inhibitor cocktail (Sigma-

Aldrich). Ten µg of protein was loaded per lane and run on SDS-PAGE gels. Protein

gels were transferred to 0.45 um Immobilon-PVDF membranes (Millipore, Temecula,

CA) and blocked with 5% BSA in 0.1% Tween 20 in PBS. Primary antibodies, sheep

anti-mouse Thy1 (R&D), rabbit anti-mouse Fabp4 (Cell Signaling), and rabbit anti-

mouse β-tubulin (Cell Signaling) were diluted 1:5000, 1:500, and 1:5000, respectively,

and incubated for 1 h. Membranes were washed in 0.1% Tween 20 in PBS then

incubated in anti-sheep or anti-rabbit HRP-conjugated secondary antibodies at 1:5000

or 1:20,000 dilution, respectively. Protein was visualized using Immobilon Western

chemiluminescent horseradish peroxidase substrate (Millipore). MagicMark XP protein

standard protocol used for ladder (Novex). Blots were developed by X-ray film. All blots

are provided as uncropped images in the supplementary data.

Flow cytometry

Cells were trypsinized and washed in PBS, then fixed with 2% PFA and blocked with

1:50 human Fc receptor blocker (Miltenyi Biotech Inc., San Diego, CA) in PBS. The

cells were then incubated with anti-mouse Thy1.2-PE conjugated antibody, 1:500, (BD

Biosciences, San Jose, CA) for 1 h on ice. Cells were washed and resuspended in PBS.

Cells were analyzed on a LSR II flow cytometer running FACSDIVA software (BD

Biosciences). Analysis of fluorescence data was performed using FlowJo software

v10.1 (FlowJo, LLC, Ashland, Oregon).

 Immunofluorescent staining

 Cells treated in 12-well plates were washed with 1X PBS and fixed with 2% PFA for 10

 min and washed three times with PBS. Cells were blocked in 1% BSA and 0.1% Triton

 X-100 in PBS with normal donkey serum (Jackson Immunoresearch) and Fc-blocker

 1:50 (BD Biosciences). The primary antibodies used were Thy1.2-PE conjugated

 antibody, (BD Biosciences, San Jose, CA) and Fabp4 (Cell Signaling), which were

 diluted 1:500 in 1% BSA and incubated for 2 h at room temperature in the dark. After

 removal of primary antibody and three washes, secondary antibody (donkey anti-rabbit

 AF647) was applied at a 1:2000 dilution for an hour. Cells were then washed and

 visualized on an EVOS-FL Cell Imaging System (Thermo Fisher).

 DNA extraction and Bisulfite conversion

 Genomic DNA was isolated from cells using a DNeasy DNA extraction kit (Qiagen,

 Valencia, CA) and quantified using the NanoDrop 1000 spectrophotometer (Thermo

 Scientific, Wilmington, DE). 1000 ng of genomic DNA was then bisulfite converted using

 an Epitect Plus Bisulfite Conversion Kit (Qiagen) to be analyzed by a pyrosequencer.

 Pyrosequencing assays

 Bisulfite-treated DNA was amplified using the PyroMark PCR kit (Qiagen, Valencia, CA)

 with the conditions of, 95°C for 5 min, 45 cycles of (95°C for 30 s, annealing

 temperature of 58°C for 30 s, 72°C for 30 s), 72°C for 10 min. The PCR product sizes

 were then verified with electrophoresis on a 2% agarose gel. Ten µl of the biotinylated

 PCR products were mixed with 1 µl streptavidin-coated Sepharose beads, 40 µl

PyroMark binding buffer (Qiagen), and 29 µl RNAase-free water for a total volume of 80

µl. This mixture was then run on a PyroMark Vacuum Workstation (Qiagen). The

purified PCR products were then added to the annealing buffer, which contained the

corresponding sequencing primer. After annealing, the plate was loaded into the

PyroMark Q96 MD instrument (Qiagen). PyroMark-CpG software automatically

generates a dispensation order of dNTPs and control dispensations, based on the

sequence to analyze. Controls are included in the dispensation order to check the

performance of the reactions. All runs also included a no template control. We analyzed

the data with the PyroMark software for quantification of % DNA CpG methylation.

Pyrosequencing Primers

Methylation levels were measured in the first CpG island of intron 1, which is part of the

promoter[17,20-23] of mouse *Thy1* (chr9:44,043,384-44,048,579; GRCm38/mm10) (94bp-

349bp) using the pyrosequencing assay. Gene-specific primers for *Thy1* were designed

using the Pyro-Mark assay design software, version 2.0 (Qiagen, Valencia, CA). The

program automatically generated primer sets that included both PCR and sequencing

primers, based on selected target sequences. One of the primers was biotinylated to

enable immobilization to streptavidin-coated beads. The sequences were as follows:

Forward primer: 5'-TTTAGTTATAGTTTTGGGAAAGGATAT

Reverse Biotinylated primer: 5'-CCACCTCCTCCCTCTATT

Sequencing primer: 5'-ATAGGGAFTTTTTATAT

Statistical analysis

All values are presented as mean ± SEM. Experiments were conducted in triplicate at

separate times. Two-way analysis of variance (ANOVA) were used for statistical

analysis using GraphPad Prism6. P-values < 0.05 were considered significant.

Acknowledgements

This project is funded by NIH grants F31ES027767, TL1-TR000096, ES001247, and

ES0023032.

Authors Contributions

E.F. performed all experiments. E.F. C.W. M.S. R.P. aided in experimental design and

data analysis. E.F. prepared figures and wrote main manuscript. All authors aided in

editing the manuscript. All authors have reviewed the manuscript.

Additional Information

Competing Interests: The authors declare no competing interests.

1.   Afshin A, Forouzanfar MH, Reitsma MB, et al. Health Effects of Overweight and Obesity in 195

Countries over 25 Years. *The New England journal of medicine.* 2017;377(1):13-27.

2.   Flegal KM, Kruszon-Moran D, Carroll MD, Fryar CD, Ogden CL. Trends in Obesity Among Adults in

the United States, 2005 to 2014. *Jama.* 2016;315(21):2284-2291.

3.   Ahima RS. Connecting obesity, aging and diabetes. *Nature medicine.* 2009;15(9):996-997.

4.   Smyth S, Heron A. Diabetes and obesity: the twin epidemics. *Nature medicine.* 2006;12(1):75-80.

5.   Heymsfield SB, Wadden TA. Mechanisms, Pathophysiology, and Management of Obesity. *The*

New England journal of medicine. 2017;376(3):254-266.

6.   Cawley J, Meyerhoefer C. The medical care costs of obesity: an instrumental variables approach.

Journal of health economics. 2012;31(1):219-230.

7.   Jo J, Gavrilova O, Pack S, et al. Hypertrophy and/or Hyperplasia: Dynamics of Adipose Tissue

Growth. *PLoS computational biology.* 2009;5(3):e1000324.

8.   Joe AW, Yi L, Even Y, Vogl AW, Rossi FM. Depot-specific differences in adipogenic progenitor

abundance and proliferative response to high-fat diet. *Stem cells (Dayton, Ohio).*

2009;27(10):2563-2570.

9.   Jing K, Heo JY, Song KS, et al. Expression regulation and function of Pref-1 during adipogenesis of

human mesenchymal stem cells (MSCs). *Biochimica et biophysica acta.* 2009;1791(8):816-826.

10.   Hudak CS, Sul HS. Pref-1, a gatekeeper of adipogenesis. *Frontiers in endocrinology.* 2013;4:79.

11.   Kim CY, Kim GN, Wiacek JL, Chen CY, Kim KH. Selenate inhibits adipogenesis through induction

of transforming growth factor-beta1 (TGF-beta1) signaling. *Biochemical and biophysical research*

communications. 2012;426(4):551-557.

12.   Kumar A, Ruan M, Clifton K, Syed F, Khosla S, Oursler MJ. TGF-beta mediates suppression of

adipogenesis by estradiol through connective tissue growth factor induction. *Endocrinology.*

2012;153(1):254-263.

13.   Cawthorn WP, Bree AJ, Yao Y, et al. Wnt6, Wnt10a and Wnt10b inhibit adipogenesis and

stimulate osteoblastogenesis through a beta-catenin-dependent mechanism. *Bone.*

2012;50(2):477-489.

14.   Woeller CF, O'Loughlin CW, Pollock SJ, Thatcher TH, Feldon SE, Phipps RP. Thy1 (CD90) controls

adipogenesis by regulating activity of the Src family kinase, Fyn. *FASEB journal : official*

publication of the Federation of American Societies for Experimental Biology. 2015;29(3):920-

931.

15.   Woeller CF, Flores E, Pollock SJ, Phipps RP. Editor's Highlight: Thy1 (CD90) Expression is Reduced

by the Environmental Chemical Tetrabromobisphenol-A to Promote Adipogenesis Through

Induction of microRNA-103. *Toxicological sciences : an official journal of the Society of*

Toxicology. 2017;157(2):305-319.

16.   Robinson CM, Neary R, Levendale A, Watson CJ, Baugh JA. Hypoxia-induced DNA

hypermethylation in human pulmonary fibroblasts is associated with Thy-1 promoter

methylation and the development of a pro-fibrotic phenotype. *Respiratory research.*

2012;13:74.

17.   Kolsto AB, Kollias G, Giguere V, Isobe KI, Prydz H, Grosveld F. The maintenance of methylation-

free islands in transgenic mice. *Nucleic acids research.* 1986;14(24):9667-9678.

18.   Mann V, Szyf M, Razin A, Chriqui-Zeira E, Kedar E. Characterization of a tumorigenic murine T-

lymphoid-cell line spontaneously derived from an IL-2-dependent T-cell line. *International*

journal of cancer. 1986;37(5):781-786.

19.   Neveu WA, Mills ST, Staitieh BS, Sueblinvong V. TGF-beta1 epigenetically modifies Thy-1

expression in primary lung fibroblasts. *American journal of physiology Cell physiology.*

2015;309(9):C616-626.

20.   Sanders YY, Pardo A, Selman M, et al. Thy-1 promoter hypermethylation: a novel epigenetic

pathogenic mechanism in pulmonary fibrosis. *American journal of respiratory cell and molecular*

biology. 2008;39(5):610-618.

21.   Giguere V, Isobe K, Grosveld F. Structure of the murine Thy-1 gene. *The EMBO journal.*

1985;4(8):2017-2024.

22.   Gundersen G, Kolsto AB, Larsen F, Prydz H. Tissue-specific methylation of a CpG island in

transgenic mice. *Gene.* 1992;113(2):207-214.

23.   Spanopoulou E, Giguere V, Grosveld F. The functional domains of the murine Thy-1 gene

promoter. *Molecular and cellular biology.* 1991;11(4):2216-2228.

24.   Lim YC, Chia SY, Jin S, Han W, Ding C, Sun L. Dynamic DNA methylation landscape defines brown

and white cell specificity during adipogenesis. *Molecular metabolism.* 2016;5(10):1033-1041.

25.   Broholm C, Olsson AH, Perfilyev A, et al. Human adipogenesis is associated with genome-wide

DNA methylation and gene-expression changes. *Epigenomics.* 2016;8(12):1601-1617.

26.   Guo W, Chen J, Yang Y, Zhu J, Wu J. Epigenetic programming of Dnmt3a mediated by AP2alpha is

required for granting preadipocyte the ability to differentiate. *Cell death & disease.*

2016;7(12):e2496.

27.   Sakamoto H, Kogo Y, Ohgane J, et al. Sequential changes in genome-wide DNA methylation

status during adipocyte differentiation. *Biochemical and biophysical research communications.*

2008;366(2):360-366.

28.   Yang X, Wu R, Shan W, Yu L, Xue B, Shi H. DNA Methylation Biphasically Regulates 3T3-L1

Preadipocyte Differentiation. *Molecular endocrinology (Baltimore, Md).* 2016;30(6):677-687.

29.   Zhao QH, Wang SG, Liu SX, et al. PPARgamma forms a bridge between DNA methylation and

histone acetylation at the C/EBPalpha gene promoter to regulate the balance between

osteogenesis and adipogenesis of bone marrow stromal cells. *The FEBS journal.*

2013;280(22):5801-5814.

30.   Chen YS, Wu R, Yang X, et al. Inhibiting DNA methylation switches adipogenesis to

osteoblastogenesis by activating Wnt10a. *Scientific reports.* 2016;6:25283.

31.   Xu S, De Becker A, Van Camp B, Vanderkerken K, Van Riet I. An improved harvest and in vitro

expansion protocol for murine bone marrow-derived mesenchymal stem cells. *Journal of*

biomedicine & biotechnology. 2010;2010:105940.

32.   Qian SW, Li X, Zhang YY, et al. Characterization of adipocyte differentiation from human

mesenchymal stem cells in bone marrow. *BMC developmental biology.* 2010;10:47.

33.   Lehmann GM, Woeller CF, Pollock SJ, et al. Novel anti-adipogenic activity produced by human

fibroblasts. *American journal of physiology Cell physiology.* 2010;299(3):C672-681.

34.   Suzuki MM, Bird A. DNA methylation landscapes: provocative insights from epigenomics. *Nature*

reviews Genetics. 2008;9(6):465-476.

35.   Hsieh CL. Dependence of transcriptional repression on CpG methylation density. *Molecular and*

cellular biology. 1994;14(8):5487-5494.

36.   Keller M, Hopp L, Liu X, et al. Genome-wide DNA promoter methylation and transcriptome

analysis in human adipose tissue unravels novel candidate genes for obesity. *Molecular*

metabolism. 2017;6(1):86-100.

37.   Mikkelsen TS, Xu Z, Zhang X, et al. Comparative epigenomic analysis of murine and human

adipogenesis. *Cell.* 2010;143(1):156-169.

38.   Haeryfar SM, Hoskin DW. Thy-1: more than a mouse pan-T cell marker. *Journal of immunology*

(Baltimore, Md : 1950). 2004;173(6):3581-3588.

39.   Orbay H, Tobita M, Mizuno H. Mesenchymal stem cells isolated from adipose and other tissues:

basic biological properties and clinical applications. *Stem cells international.* 2012;2012:461718.

40.   Attwood JT, Yung RL, Richardson BC. DNA methylation and the regulation of gene transcription.

478   *Cellular and molecular life sciences : CMLS.* 2002;59(2):241-257.

479   41.   Baylin SB, Herman JG, Graff JR, Vertino PM, Issa JP. Alterations in DNA methylation: a

480   fundamental aspect of neoplasia. *Advances in cancer research.* 1998;72:141-196.

481   42.   Bowers RR, Kim JW, Otto TC, Lane MD. Stable stem cell commitment to the adipocyte lineage by

482   inhibition of DNA methylation: role of the BMP-4 gene. *Proceedings of the National Academy of*

483   *Sciences of the United States of America.* 2006;103(35):13022-13027.

484   43.   Du J, Cheng X, Shen L, et al. Methylation of miR-145a-5p promoter mediates adipocytes

485   differentiation. *Biochemical and biophysical research communications.* 2016;475(1):140-148.

486   44.   Li M, Liu Z, Zhang Z, Liu G, Sun S, Sun C. miR-103 promotes 3T3-L1 cell adipogenesis through

487   AKT/mTOR signal pathway with its target being MEF2D. *Biological chemistry.* 2015;396(3):235-

488   244.

489   45.   Perri R, Nares S, Zhang S, Barros SP, Offenbacher S. MicroRNA modulation in obesity and

490   periodontitis. *Journal of dental research.* 2012;91(1):33-38.

491   46.   Alipour R, Sadeghi F, Hashemi-Beni B, et al. Phenotypic characterizations and comparison of

492   adult dental stem cells with adipose-derived stem cells. *International journal of preventive*

493   *medicine.* 2010;1(3):164-171.

494   47.   Yoshimura H, Muneta T, Nimura A, Yokoyama A, Koga H, Sekiya I. Comparison of rat

495   mesenchymal stem cells derived from bone marrow, synovium, periosteum, adipose tissue, and

496   muscle. *Cell and tissue research.* 2007;327(3):449-462.

497   48.   Kasaai B, Moffatt P, Al-Salmi L, Lauzier D, Lessard L, Hamdy RC. Spatial and temporal localization

498   of WNT signaling proteins in a mouse model of distraction osteogenesis. *The journal of*

499   *histochemistry and cytochemistry : official journal of the Histochemistry Society.* 2012;60(3):219-

500   228.

49.   Rudnicki MA, Williams BO. Wnt signaling in bone and muscle. *Bone.* 2015;80:60-66.

50.   Yin X, Wang X, Hu X, Chen Y, Zeng K, Zhang H. ERbeta induces the differentiation of cultured

osteoblasts by both Wnt/beta-catenin signaling pathway and estrogen signaling pathways.

Experimental cell research. 2015;335(1):107-114.

51.   Chung MT, Liu C, Hyun JS, et al. CD90 (Thy-1)-positive selection enhances osteogenic capacity of

human adipose-derived stromal cells. *Tissue engineering Part A.* 2013;19(7-8):989-997.

52.   Bearden RN, Huggins SS, Cummings KJ, Smith R, Gregory CA, Saunders WB. In-vitro

characterization of canine multipotent stromal cells isolated from synovium, bone marrow, and

adipose tissue: a donor-matched comparative study. *Stem cell research & therapy.*

2017;8(1):218.

53.   Paine A, Woeller CF, Zhang H, et al. Thy1 is a positive regulator of osteoblast differentiation and

modulates bone homeostasis in obese mice. *FASEB journal : official publication of the Federation*

of American Societies for Experimental Biology. 2018:fj201701379R.

54.   Fernandez-Arroyo S, Cuyas E, Bosch-Barrera J, Alarcon T, Joven J, Menendez JA. Activation of the

methylation cycle in cells reprogrammed into a stem cell-like state. *Oncoscience.*

2015;2(12):958-967.

55.   Hansen TC, Woeller CF, Lacy SH, Koltz PF, Langstein HN, Phipps RP. Thy1 (CD90) Expression Is

Elevated in Radiation-Induced Periprosthetic Capsular Contracture: Implication for Novel

Therapeutics. *Plastic and reconstructive surgery.* 2017;140(2):316-326.

Figure Captions

Figure 1: The reduction of *Thy1* expression during adipogenesis is attenuated by 5-aza-dC in mouse mesenchymal stem cells. mMSCs were treated daily with either DMSO or 5-aza-dC for 12 days continually with basal media alone or with the introduction of the adipogenic cocktail (ACT) starting at day 3 A) Timeline diagram of treatment for mMSCs. B) Western blot shows treatment with 5-aza-dC increases Thy1 protein levels and decreases protein levels of the adipogenic marker, Fabp4 versus cells only receiving the adipogenic cocktail (ACT). C-D) RT-qPCR results show that treatment with 5-aza-dC significantly increases Thy1 mRNA and decreases Fabp4 mRNA levels. Relative percentages were normalized to Media DMSO for Thy1 mRNA and ACT DMSO for Fabp4 mRNA levels. E) Immunofluorescent images depict reduced FAPB4 (Green) expression and increased Thy1 (Red) expression in cells receiving 5-aza-dC versus those receiving ACT alone. Cell nuclei are stained with DAPI and depicted in blue. Scale bars in white represent 200μm. *$p<0.5$, **$p<.01$, ***$p<.001$.

Figure 2: Inhibiting methylation increases *Thy1* expression in 3T3-L1 during adipogenesis. 3T3-L1 cells were treated daily with either DMSO or 5-aza-dC for 7 days continually with basal media alone or with the introduction of the adipogenic cocktail (ACT) starting at day 3. A) Timeline diagram of treatment for 3T3-L1s. B) Western blot showing treatment with 5-aza-dC resulted in an increase in Thy1 total protein levels in both media alone and adipogenic cocktail exposed samples. ACT samples also had a decrease in Fabp4 protein levels when treated with 5-aza-dC, shown by a

representative western blot. C-D) RT-qPCR shows treatment with 5-aza-dC increases

Thy1 mRNA and decreases Fabp4 mRNA levels in ACT samples. Relative percentages

were normalized to 100% to media DMSO for Thy1 mRNA or 100% of ACT DMSO for

Fabp4 mRNA levels. *p<05, **p<.01, ****p<.0001

Figure 3: Thy1 surface protein levels attenuated by 5-aza-dC. 3T3-L1 cells were

treated daily with either DMSO or 5-aza-dC for 7 days with the introduction of the

adipogenic cocktail (ACT) starting at day 3. A) Representative histogram of Thy1

surface levels measured by flow cytometry using a CD90.2-PE conjugated antibody. B)

ACT samples had a decrease in Thy1 mean fluorescence intensity (MFI) that was

partially attenuated by 5-aza-dC treatment. C) ACT samples had a decrease in Thy1+

cells compared to Media alone, while ACT cells treated with 5-aza-dC had an increase

in the Thy1+ population. D) Immunofluorescent images show decreased staining of

Thy1 (Red) with ACT compared to pre-adipocytes (media alone) and increased Fabp4

(Green). Thy1 staining was sustained when ACT samples were treated with 5-aza-dC,

along with a decreased staining of Fabp4. Cell nuclei are stained with DAPI and

depicted in blue. Scale bars in white represent 200μm. **p<.01, ***p<.001, ****p<.0001

Figure 4: *Thy1* is hypermethylated during adipogenesis and demethylated by 5-

aza-dC. 3T3-L1 cells were treated daily with either DMSO or 5-aza-dC for 7 days with

the introduction of the adipogenic cocktail (ACT) starting at day 3. A) Schematic of *Thy1*

gene showing Thy1-CGI1 and CpG sites between exon 1 and 2. B) Average percent

DNA methylation of Thy1-CGI1 showed an increase in methylation in ACT cells

compared to media alone, while treatment with the methylation inhibitor, 5-aza-dC

resulted in a decrease of overall methylation in both groups. C) Individual CpG position

sites were measured across Thy1-CGI1 and showed increases in methylation with ACT

compared to media alone. D-E) Both media alone and ACT groups had decreases in

methylation in all individual CpG sites when treated 5-aza-dC. * $p<.05$, ***$p<.001$

Figure 5: *Thy1* is hypermethylated in a time dependent manner during adipocyte

differentiation and attenuated by 5-aza-dC. 3T3-L1 cells were pretreated for 24 h with

either DMSO or 5-aza-dC, along with continuous daily treatment and then harvested

corresponding to days with adipogenic cocktail (ACT) (Day 0= no ACT). A) Average

percent DNA methylation of Thy1-CGI1 showed an increase in methylation in normal

adipogenesis differentiation process, while treatment with the methylation inhibitor, 5-

aza-dC resulted in a decrease in overall methylation. B-C) Individual CpG position sites

were measured across Thy1-CGI1 and day 0 showed no relevant changes, while day 6

had a notable decrease in methylation when treated with 5-aza-dC. *= $p<.05$

582
583 **Figure 1**

584
585 **Figure 2**
586

587
588   **Figure 3**

589
590 **Figure 4**

A

B

C

591
592   **Figure 5**