Unbiased whole genomes from mammalian feces using fluorescence-activated cell sorting

Joseph D. Orkin[1,2]*, Marc de Manuel[3], Roman Krawetz[4], Javier del Campo[5], Claudia Fontsere[3], Lukas F. K. Kuderna[3], Ester Lizano[3], Jia Tang[6], Tomas Marques-Bonet[3,7,8,9], Amanda D. Melin[1,2]

1.   Department of Anthropology & Archaeology, University of Calgary, Calgary, Alberta, Canada

2.   Alberta Children's Hospital Research Institute, University of Calgary, Calgary, Alberta, Canada

3.   Institute of Evolutionary Biology (UPF-CSIC), PRBB, Dr. Aiguader 88, 08003 Barcelona, Spain.

4.   Department of Cell Biology and Anatomy, University of Calgary, Calgary, Alberta, Canada

5.   Department of Marine Biology and Oceanography, Institut de Ciències del Mar (Consejo Superior de Investigaciones Científicas), Barcelona, Catalonia, Spain

6.   Department of Biological Sciences, University of Calgary, Calgary, Alberta, Canada

7.   Catalan Institution of Research and Advanced Studies (ICREA), Passeig de Lluís Companys, 23, 08010, Barcelona, Spain

8.   CNAG-CRG, Centre for Genomic Regulation (CRG), Barcelona Institute of Science and Technology (BIST), Baldiri i Reixac

9.   Institut Català de Paleontologia Miquel Crusafont, Universitat Autònoma de Barcelona, Edifici ICTA-ICP, c/ Columnes s/n, 08193 Cerdanyola del Vallès, Barcelona, Spain

*Corresponding Author:
     Joseph D. Orkin (joseph.orkin@ucalgary.ca)

ABSTRACT

35
36

Non-invasive genomic research on free-ranging mammals typically relies on the use of
fecal DNA. This requires the isolation and enrichment of endogenous DNA, given its small
proportion compared to bacterial DNA. Current approaches for acquiring large-scale genomic
data from feces rely on bait-and-capture techniques. While this technique has greatly improved
our understanding of mammalian population genomics, it is limited by biases inherent to the
capture process, including allele dropout, low mapping rates, PCR duplication artifacts, and
structural biases. We report here a new method for generating whole mammalian genomes from
feces using fluorescence-activated cell sorting (FACS). Instead of enriching endogenous DNA
from extracted fecal DNA, we isolated mammalian cells directly from feces. We then built
fragment libraries with low input material from commercially available kits, which we
sequenced at high and low coverage. We validated this method on feces collected from primates
stored in RNAlater for up to three years. We sequenced one fecal genome at high coverage
(12X) and 15 additional fecal genomes at low coverage (0.1X - 4X). For comparative purposes,
we also sequenced DNA from nine blood or tissue samples opportunistically collected from
capuchin monkeys that died of natural causes or were treated in a local rehabilitation center.
Across all fecal samples, we achieved median mapping and duplication rates of 82% and 6%,
respectively. Our high-depth fecal genome did not differ in the distribution of coverage,
heterozygosity, or GC content from those derived from blood or tissue origin. As a practical
application of our new approach with low coverage fecal genomes, we were able to resolve the
population genetic structure of capuchin monkeys from four sites in Costa Rica.

INTRODUCTION

Advances in DNA sequencing technology have allowed for great strides to be made in comparative genomics (Arandjelovic & Vigilant, 2018; Corlett, 2017; Perry, 2014). It is now commonplace in a single study of a non-model organism to sequence partial genomes from multiple individuals. If fresh tissue or blood samples can be acquired from a handful of individuals, sequencing a *de novo* reference genome or generating a panel of single nucleotide variants is relatively straightforward. However, answering population level questions typically depends upon the non-invasive collection of fecal samples from free-ranging animals. Unfortunately, less than 5% of the extracted DNA from a fecal sample typically originates from an endogenous source (i.e. the host animal) (Hernandez-Rodriguez et al., 2017; Snyder-Mackler et al., 2016), while the remaining 95% comes from microorganisms and dietary items. For many species, this combination of factors makes is unfeasible to sequence whole genomes at high coverage from a large number of individuals. The resulting dearth of population-wide high-coverage genomes limits the scope of questions that can be asked and answered in genomics, ecology, and conservation.

Thanks to recent advances in non-invasive genomics, it has become possible to sequence partial genomes by enriching the proportion of endogenous DNA in feces (Chiou & Bergey, 2018; Perry, Marioni, Melsted, & Gilad, 2010; Snyder-Mackler et al., 2016). Through the use of targeted bait-and-capture and reduced representation libraries, this approach allows for the sequencing of single nucleotide variant (SNV) sets, which has begun to provide important insights into population structure and local adaptation of free-ranging mammals (Chiou, 2017; de Manuel et al., 2016; Wall et al., 2016). Despite the promise of this approach, DNA enrichment suffers from biases and impracticalities that limit its ability to uniformly cover a genome. Current bait-and-capture techniques are subject to inherent biases in the type of DNA captured (e.g. non-repetitive elements, GC content, reduced representation libraries, inconsistent hybridization); requires the costly and time consuming generation of RNA or DNA baits; have limited ability to enrich endogenous DNA (mean: ~57% of mapped reads (Snyder-Mackler et al., 2016)); and have high average PCR duplication rates (mean: ~38% of mapped reads (Snyder-Mackler et al., 2016)). Methylation-based enrichment offers a promising and cost-effective alternative to bait-and-capture for SNV generation, although it also suffers from inherent bias in the composition of the enriched libraries, and has limited enrichment capacity (mean: <50% of mapped reads (Chiou & Bergey, 2018)). While both approaches are viable for partial data, neither offers the realistic possibility of truly unbiased, cost-effective whole genome sequencing.

Through a novel application of fluorescence-activated cell sorting (FACS), we present a rapid, cost-effective method of isolating a host animal's intestinal epithelial cells for DNA extraction and genome sequencing. With this approach, we have routinely mapped more than 80% of reads to the host genome, strongly suggesting they are from endogenous DNA. This method requires no targeted enrichment of DNA, RNA baits, or methylation. It allows for DNA to be extracted and libraries built with commercially available kits, removing many of the challenges of enrichment-based techniques. Furthermore, our method allows for the long-term, room-temperature stabilization of samples, making it possible for field workers to collect samples with ease from remote areas without the need for temperature sensitive storage.

Here, we propose a novel protocol to isolate intestinal epithelial cells from the feces of white-faced capuchin monkeys (*Cebus imitator*) up to three years after initial collection. From these cells, we generated low coverage genomes from 17 fecal samples and selected one of them

104 for deeper sequencing (targeting ~10X - 15X coverage). In so doing, we have generated the first
105 uniformly-distributed, high-coverage, whole genome of a mammal from its feces. To
106 demonstrate the breadth of fecal FACS, we also conducted an analysis of population genetic
107 structure in two Costa Rican forest reserves using DNA derived from both fecal FACS and
108 traditional blood/tissue extractions.
109
110   **METHODS**

2.1 Sample Collection
We collected fecal samples from free-ranging white-faced capuchin monkeys (*Cebus*
imitator) at Sector Santa Rosa (SSR), part of the Área de Conservación Guanacaste in
northwestern Costa Rica, which is a 163,000 hectare tropical dry forest nature reserve (Figure 1).
Behavioral research of free-ranging white-faced capuchins has been ongoing at SSR since the
1980's which allows for the reliable identification of known individuals from facial features and
bodily scars (Fedigan & Rose-Wiles, 1996). We collected 14 fresh fecal samples from 12 white-
faced capuchin monkeys immediately following defecation (Table 1). We placed 1 mL of feces
into conical 15 mL tubes pre-filled with 5 mL of RNAlater. RNAlater preserved fecal samples
were sent to the University of Calgary, where they were stored at room temperature for up to
three years. To evaluate other preservation methods, we also collected two additional capuchin
monkey fecal samples (SSR-FL and a section of SSR-ML) and one spider monkey (*Ateles*
geoffroyi) fecal sample, which we stored in 1X PBS buffer and then froze in liquid nitrogen with
a betaine cryopreservative (Rinke et al., 2014). Given the logistical challenges of carrying liquid
nitrogen to remote field sites, we prioritized evaluation of samples stored in RNAlater.
Finally, we took tissue and blood samples opportunistically. During the course of our
study, 4 individual capuchin monkeys died of natural causes at SSR, from whom we were able to
collect tissue samples, which we stored in RNAlater. By collaborating with *Kids Saving the*
Rainforest veterinary rehabilitation clinic in Quepos, Costa Rica, we acquired blood samples
from 5 more Costa Rican white-faced capuchins who were undergoing treatment at the facility
(although we were unable to collect paired fecal samples). Samples were collected with
permission from the Area de Conservacion Guanacaste (ACG-PI-033-2016) and CONAGEBIO
(R-025-2014-OT-CONAGEBIO). Samples were exported from Costa Rica under permits from
CITES and Area de Conservacion Guanacaste (2016-CR2392/SJ #S 2477, 2016-CR2393/SJ #S
2477, DGVS-030-2016-ACG-PI-002-2016; 012706) and imported with permission from the
Canadian Food and Inspection agency (A-2016-03992-4).
137
2.2 FACS
Before isolating cells by Fluorescence-activated cell sorting (FACS), fecal samples were
prepared using a series of washes and filtration steps. Fecal samples were vortexed for 30 s and
centrifuged for 30 s at 2,500 g. Then the supernatant was passed through a 70 um filter into a 50
mL tube and washed with DPBS. After transferring the resultant filtrate to a 15 mL tube, it was
centrifuged at 1,500 RPM for 5 minutes to pellet the cells. Then we twice washed the cells with
13 mL of DPBS. We added 500 uL of DPBS to the pellet and re-filtered through a 35 um filter
into a 5 mL FACS tube. We prepared a negative control (to control for auto-fluorescence) with
500 uL of DPBS and one drop of the cell solution. To the remaining solution, we added 1 uL of
AE1/AE3 Anti-Pan Cytokeratin Alexa Fluor® 488 antibody or TOTO-3 DNA stain, which we
allowed to incubate at 4°C for at least 30 minutes.

We isolated cells using a BD FACSAria™ Fusion (BD Biosciences) flow cytometer at the
University of Calgary Flow Cytometry Core. To sterilize the cytometer's fluidics before
processing each sample, we ran a 3% bleach solution through the system for four minutes at
maximum pressure. We assessed background fluorescence and cellular integrity, by processing
the negative control sample prior to all prepared fecal samples. For each sample we first gated
our target population by forward and side scatter characteristics that were likely to minimize
bacteria and cellular debris (Figure 2). Secondary and tertiary gates were implemented to remove
cellular agglomerations. Finally, we selected cells with antibody or DNA fluorescence greater
than background levels. In cases when staining was not effective, we sorted solely on the first
three gates. Cells were pelleted and frozen at -20°C.
159
2.3 DNA Extraction and Shotgun Sequencing
We extracted fecal DNA (fDNA) with the QIAGEN DNA Micro kit, following the
"Small volumes of blood" protocol. To improve DNA yield, we increased the lysis time to three
hours, and incubated 50 µL of 56°C elution buffer on the spin column membrane for 10 minutes.
DNA concentration was measured with a Qubit fluorometer. Additionally, to calculate
endogenous DNA enrichment, we extracted DNA directly from five fecal samples prior to their
having undergone FACS. We extracted DNA from the nine tissue and blood samples using the
QIAGEN Gentra Puregene Tissue kit and DNeasy blood and tissue kit, respectively.
For the fecal samples, DNA was fragmented to 350 bp with a Covaris sonicator. We built
whole genomic sequencing libraries with the NEB Next Ultra 2 kit using 10-11 PCR cycles.
Fecal genomic libraries were sequenced on an Illumina NextSeq (2x150 PE) at the University of
Calgary genome sequencing core. We resequenced one fecal sample at high coverage on an
Illumina HighSeq 4000 at the McDonnell Genome Institute at Washington University in St.
Louis (MGI). High-coverage, whole genomic shotgun libraries were prepared for the blood and
tissue DNA samples and sequenced on an Illumina X-10 at MGI.
175
2.3 Mapping and SNV Generation
Reads were trimmed of sequencing adaptors with Trimmomatic (Bolger, Lohse, &
Usadel, 2014). Subsequently, we mapped the *Cebus* reads to the *Cebus imitator* 1.0 reference
genome (GCF_001604975.1) with BWA mem (Li & Durbin, 2009) and removed duplicates with
Picard Tools (http://broadinstitute.github.io/picard/). We called SNVs for each sample
independently using the *Cebus* genome and the GATK UnifiedGenotyper pipeline (*-out_mode
EMIT_ALL_SITES*) (McKenna et al., 2010). Genomic VCFs were then combined using GATK's
CombineVariants restricting to positions with a depth of coverage between 3 and 100, mapping
quality above 30, no reads with mapping quality zero and variant PHRED scores above 30.
Sequencing reads from one of the high coverage fecal samples (SSR-FL) bore a strong signature
of human contamination (16%), and were thus excluded from SNV generation. We included
reads from nine tissue/blood samples and one frozen fecal sample with high coverage (SSR-ML).
In total, we generated 4,184,363 SNVs for downstream analyses.
To remove potential human contamination from sequenced libraries, we mapped trimmed
reads to the *Cebus imitator* 1.0 and human (hg38) genomes simultaneously with BBsplit
(Bushnell, 2016). Using default BBsplit parameters, we binned separately reads that mapped
unambiguously to either genome. Ambiguously mapping reads (i.e. those mapping equally well
to both genomes) were assigned to both genomic bins, and unmapped reads were assigned to a
third bin. We calculated the amount of human genomic contamination as the percentage of total

reads unambiguously mapping to the human genome (Table 2). After removing contaminant
reads, all libraries with at least 0.5X genomic coverage were used for population analysis.
In order to test the effect of fecal FACS on mapping rates, we selected five samples at
random (SSR-CH, SSR-NM, SSR-LE, SSR-PR, SSR-SN) to compare pre- and post-FACS
mapping rates. To test for an increase in mapping percentage, we ran a one-sample paired
Wilcoxon signed-rank test on the percentages of reads that mapped exclusively to the *Cebus*
genome before and after flow FACS. Additionally, we ran Pearson's product moment
correlations to test for an effect of the number of cells (log10 transformed) on rates of mapping,
read duplication, and ng of input DNA. The above tests were all performed in R.
204
2.5 High coverage fecal genome comparison
We made several comparisons between our high-coverage feces-derived genome and the
blood/tissue-derived genomes using window-based approaches. For each test, the feces-derived
genome should fall within the range of variation for members of its population of origin (SSR).
Deviations from this, for examples all fecal genomes clustering together, would indicate biases
in our DNA isolation methods. To assess this, we constructed 10 KB / 4KB sliding windows
along the largest scaffold (21,314,911 bp) in the *C. imitator* reference genome. From these
windows, we constructed plots of coverage density and the distribution of window coverage
along the scaffold. Secondly, we assessed the level of heterozygosity in 1 MB / 200 KB sliding
windows throughout the ten largest scaffolds. For each high-coverage genome, we plotted the
density distribution of window heterozygosity. We measured genome-wide GC content with the
Picard Tools CollectGcBiasMetrics function. The percentage of GC content was assessed against
the distribution of normalized coverage and the number of reads in 100 bp windows per the
number reads aligned to the windows.
219
2.6 Population genomic analysis
Given the large degree of difference in coverage among our samples, (less than 1X to
greater than 50X), we performed pseudodiploid allele calling on all samples using custom
scripts. For each library, at each position in the SNV set, we selected a single, random read from
the sequenced library. From that read, we called the variant information at the respective SNV
site for the given library. In so doing, we generated a VCF with a representative degree of
variation and error for all samples.
To assess population structure and infer splits between northern and southern groups of
Costa Rican white-faced capuchins, we constructed principal components plots with
EIGENSTRAT (Price et al., 2006) and built population trees with TreeMix (Pickrell & Pritchard,
2012). Because we ascertained variants predominantly with libraries that were of tissue/blood
origin, we built principal components solely with SNVs from these libraries and projected the
remaining fecal libraries onto the principal components. For our maximum likelihood trees, we
used three outgroups (*Ateles geoffroyi*, *Saimiri sciureus*, and *Cebus albifrons*), with *A. geoffroyi*
serving as the root of the tree. Given the geographic distance and anthropogenic deforestation
between northern and southern populations, we assumed no migration. To account for linkage
disequilibrium, we grouped SNVs into windows of 1,000 SNVs.
237
238   **RESULTS**
239
240 *3.1 Isolation of intestinal epithelial cells using Fluorescence-activated cell sorting (FACS)*

Flow cytometry can be used to discriminate among categories of cells by examining the manner in which light scatters in response to cellular properties. We interpreted forward scatter (FSC) and side scatter (SSC) as measures of cellular size and granularity (complexity), respectively. When cells are intact, free of agglomerations, and of limited variety, they form easily identifiable clusters, particularly when bound with fluorescently labeled antibodies. In contrast to this idealized schema, abundant cellular debris prevented us from observing distinct cellular populations when assessing the relationship between FSC and SSC (Figure 2) of fecal samples. The vast majority of events were usually clustered in lower range of FSC, and likely of bacterial origin. To exclude bacteria insofar as possible, we implemented a FSC gate that only included events above this cluster, typically the top ½ to ⅔ of the FSC range. From the 14 RNAlater preserved capuchin fecal samples, we isolated a median of 1,739 cells, with a range of 129 - 62,201 (Table 2). Typically, we collected a few hundred or thousand cells, but in two cases of poor fluorescent staining (SSR-FN and the RNAlater preserved SSR-ML sample), we sorted the larger gated populations, irrespective of fluorescent intensity. From the frozen samples, SSR-FL and SSR-ML, we collected 4,405 and 2,546 cells, respectively. Similarly, from the spider monkey sample, which we split into two separate FACS runs, we isolated 4,026 and 602 cells.

3.2 Mapping of genomic libraries

From each cellular population, we successfully extracted DNA and prepared sequencing libraries. Among the RNAlater preserved capuchin samples, the total amount of DNA per sample was low, ranging from 2.96 to 21.50 ng, with a median value of 7.85 ng (Table 2). A relationship between the number of cells was not significantly correlated with the amount of extracted DNA (R=0.227; 95% CI (-0.345, 0.676); t=0.808, p > 0.05) or mapping rate (R = -0.204; 95% CI (-0.663, 0.367); t = -0.721; p > 0.05). Median mapping rates reached 93% (range: 55 - 98%) with BWA-MEM and 82% (range: 11 - 95%) with the more stringent BBsplit settings (Figures 3A, 3B, Table 2). Read duplication levels were low, with a median value of 9% (range: 2 - 40%) resulting in 63% (range: 8 - 92%) of reads being unique and mapping to the *Cebus imitator* 1.0 genome. The amount of duplicate reads was distributed bimodally across individuals, with reads from five samples having substantially higher duplication rates than the remaining nine. The rate of duplication was significantly correlated (R = -0.751; 95% CI (-0.917, -0.366); t = -3.94; p < 0.01) with the number of cells (log10 transformed), decreasing sharply above a threshold of about 1,000 cells (Figure 3D).

The samples frozen in liquid nitrogen mapped at comparable rates to those preserved in RNAlater. From the two frozen capuchin samples, SSR-ML and SSR-FL, respectively, we extracted 10.50 and 6.72 ng of DNA. These two samples mapped at 96% and 80.4% with BWA-MEM and 90% and 42% with BBsplit (5% and 3% duplicates), respectively. We extracted 6.96 and 4.50 ng of DNA from the two runs of the spider monkey sample, which mapped at a substantially lower rate of 54% and 49% with BWA-MEM and 12% with BBsplit for both (1% duplicates for both).

We observed little to no human contamination in the RNAlater preserved samples. For nine of the 14 samples, BBsplit mapped between 0.61 and 1.25% of reads to hg38 (median 0.96%); however, in four cases 2.86 - 5.80% of reads were binned to the human genome. Human mapped reads were also low for the frozen SSR-ML (1.25%) and spider monkey (2.83% and 1.82%) samples. However, SSR-FL appeared to have substantial human contamination (15.77% of reads). This may be due to initial processing of these three samples, which were stored using the cryopreservation method, at the field site. We conducted the initial vortexing, centrifugation,

and collection of supernatant (see section 2.2) at the SSR field station, which is likely where
SSR-FL was contaminated. Due to this, we examine the mapping rates using only the RNAlater
preserved samples. However, we were able to decontaminate reads bioinformatically, and
include the decontaminated reads in downstream analyses where appropriate.
By sorting fecal samples with FACS, we substantially increased the percentage of reads
mapping to the target genome. We selected five samples at random (SSR-CH, SSR-NM, SSR-
LE, SSR-PR, SSR-SN) to compare pre- and post-FACS mapping rates. The mapping rates of
unsorted feces ranged from 10 - 42%, with a median of 14% (Figure 3C). After flow sorting
aliquots of these fecal samples, we obtained significantly higher mapping rates (V = 15, p <
0.05) for each sample, ranging from 64 - 95%, with a median of 85%, resulting in a median 6.07
fold enrichment.
298
3.3 High coverage fecal genome
Given that the sample SSR-ML had a high mapping percentage, a low rate of duplication,
and was effectively free of human-specific mapping, we selected it for sequencing at high
coverage. Using ½ of one HiSeq 4000 lane, we achieved an average coverage of ~12X across the
Cebus imitator 1.0 genome.
When comparing the high coverage fecal and tissue genomes from the Santa Rosa site,
we observed no substantial difference in quality, coverage, heterozygosity, or GC content
(Figures 3 and 4). For each genome, the distribution of per site coverage followed a roughly
normal distribution with a small number of positions uncovered (~2%) (Figure 3A). Coverage
along the largest scaffold from the *Cebus* genome was uniform in both tissue and fecal samples
(Figure 3B). No obvious area of excessively high or low coverages is apparent in the fecal
genome compared to that of the tissue derived genomes. Importantly, the fecal genome does not
have any obvious gaps in coverage. Likewise, levels of heterozygosity were comparable between
fecal and tissue genomes (Figure 3C, D). The fluctuating levels of heterozygosity across the
largest genomic scaffold in 100 KB windows is highly similar for SSR-ML and SSR-CR (Figure
3D), indicative of their close familial relationship. Finally, the distribution of GC content across
the genome does not suffer from substantial bias (Figure 5B). Although the normalized coverage
at the extremes of the GC distribution is on the higher end of the capuchin samples (Figure 5A),
it falls well within the range of other samples for the vast majority of the genome where GC
content ranges from ~20 - 75% (Figure 5B).
319
3.4 Population structure
We observed likely population subdivision between the northern and southern groups of
white-faced capuchins in our SNV set. This separation corresponds to the ecological division of
the season tropical dry forests in the north from the non-seasonal tropical wet forests in the
south. Given the limitations of the available sampling sites, it is possible that the appearance of
an ecological divide is actually evidence of isolation by distance.
All individuals from the north and the south are sharply discriminated by the first
principle component of the PCA (Figure 6A). The second component indicates a higher degree
of genetic variation within the southern individuals. All the northern individuals form a tight
cluster on the PCA plot, in contrast to those from the south, which are more widely dispersed
along PC 2. Furthermore, the single individual from the northern site of Cañas clusters closely
with the individuals from Santa Rosa, despite a geographic distance of more than 100 km, which
suggests that isolation by distance might not be the sole reason for population differentiation. No

clustering was observed within the four individuals from the southern sites of Manuel Antonio and Quepos, apart from their separation from the northern individuals along PC 1. Because we generated the principal components with samples from the primary SNV set and projected the remaining samples (fecal flow FACS and tissue-based outgroups), the outgroup taxa are expected to fall in between the two main sampling clusters of white-faced capuchins. As expected, the three outgroup taxa (*C. albifrons*, *S. sciureus*, and *A. geoffroyi*) fall in the center of the PCA plot.

The pattern of clustering generated by our maximum likelihood SNV tree recapitulates the expected patterns of geographic distance and ecological separation in our sample (Figure 6). Among the white-faced capuchin monkeys, the northern and southern clades represent the main split in the tree. Each clade is subdivided according to the two sampling sites within the geographic/ecological regions. Furthermore, the three outgroup taxa split by the expected degree of evolutionary distance. These relationships are not perturbed by the fact that samples were a mixture of traditional tissue-based genomic libraries and libraries generated by fecal flow-FACS. This pattern is evident both within the northern sites and outgroup taxa. Additionally, depth of coverage does not appear to affect the pattern of clustering. Our sample ranged in coverage from less than 1X to greater than 50X. In spite of this, the pattern of geographic/ecological subdivision held.

DISCUSSION

In this manuscript, we describe a novel use of FACS to isolate cells from the feces of free-ranging mammals for population and comparative genomics. We have demonstrated that fecal FACS is an effective means for: 1) the enrichment of endogenous DNA from non-invasive primate samples; 2) the generation of unbiased whole genomes at high coverage or low coverage sequencing libraries suitable for population genomic analysis. Isolating genome-scale information from non-invasively collected samples remains a major challenge in molecular ecology. Although DNA can be extracted readily from museum specimens and captive individuals (Guschanski et al., 2013; Prado-Martinez et al., 2013; van der Valk, Lona Durazo, Dalén, & Guschanski, 2017), the vast majority of the world's mammalian genomic diversity remains in free-ranging individuals. Our results indicate that fecal FACS has the potential for widespread application in molecular ecology and the broadening of non-invasive genomics for threatened and cryptic mammals.

4.1 Performance and cost-effectiveness

Current techniques to isolate whole genomic information from fecal samples depend upon the enrichment of endogenous DNA from extracted fDNA (Chiou & Bergey, 2018; Perry et al., 2010; Snyder-Mackler et al., 2016). While these methods have proven effective for SNV analyses, particularly at low coverage (Chiou, 2017; de Manuel et al., 2016; Wall et al., 2016), they remain of limited genomic scope. The total mapping rate of endogenous reads from the highest performing enrichment protocol is 57%, with a non-duplicate mapping rate of 38% (Snyder-Mackler et al., 2016). The median non-duplicate rate that we generated through FACS is 63% (82% when including duplicates), substantially outperforming that of enrichment-based approaches. While sequencing costs have fallen dramatically in recent years, maximizing the proportion of non-duplicate reads in sequencing libraries remains a critical factor in determining the feasibility of sampling schemes. Studies that aim to sequence tens or hundreds of fecal

individuals at high coverage are simply not practical for most labs, given the current cost
structure. We were able to isolate primate cells from feces for roughly $40 per sample. Given
that each sample required about 30 minutes of FACS time and three hours of wet lab preparation
time (per batch of samples), a trained lab worker could prepare five to ten samples per day,
presuming the availability of FACS resources. Although these costs of time and money are not
negligible, this may be a justifiable expense for projects where the increased mapping rate and
genomic coverage are desired.
While our fecal FACS method is effective in white-faced capuchin monkeys and
Geoffroy's spider monkeys, we acknowledge that further validation in other species is warranted.
Given the disparity in mapping rates between the capuchin and spider monkey samples, it is
possible that cytometry protocols would need to be optimized toward the particularities of a
given species' feces and conditions. Consistent with this notion is the fact that the fecal sample
(SSR-SB1) with low mapping success, was substantially darker than the other capuchin samples,
which, depending on the dietary items consumed, typically have a green, brown, or rust
coloration. Mapping was substantially improved in the replicate sample (SSR-SB2), which was
collected on a different day. Curiously, we did not observe a relationship between the number of
sorted cells and the concentration of extracted DNA. However, this is likely explained by
residual intercalating dyes used in FACS process remaining in the sorted cells and interfering
with Qubit quantification (Kuderna et al., 2018). Additionally, it is peculiar that the mapping
rates of the libraries we built from unsorted fDNA were so high (median 14%). Typically, less
than 5% of fDNA is of an endogenous source, although some chimpanzee samples have been
reported to have up to 25% endogenous reads (Hernandez-Rodriguez et al., 2017). Further
testing of capuchin fecal samples with lower endogenous DNA concentration is worth pursuing,
as mapping rates for endogenous DNA from unprocessed and enriched fDNA are often
correlated (Chiou & Bergey, 2018; Hernandez-Rodriguez et al., 2017; Snyder-Mackler et al.,
2016). However, because cell sorting is not a targeted DNA enrichment process, we find it
unlikely that post-FACS mapping rates should depend on the concentration of endogenous
fDNA; accordingly, we did not observe any such relationship among the five samples we tested
for enrichment (Figure 3C). Furthermore, we did not observe a correlation between the number
of isolated cells and the mapping rate; in one case, we obtained a 94% mapping rate with only
140 cells. Presuming that the flow cytometer is sorting cells correctly, and that those cells
contain viable DNA, the mapping rate should only be contingent upon the accuracy of the cell
sorting process.
We have demonstrated that RNAlater is an effective, long-term, room-temperature
cellular storage medium for fecal FACS. In the great majority of cases, FACS involves the
sorting of living cellular populations, and attempts to sort dead cells are often met with
skepticism (Sasaki, Dumas, & Engleman, 1987). Dead cells are typically distorted and
fragmented, yielding populations that are difficult to discriminate. We attempted to freeze fresh
feces with liquid nitrogen and a betaine cryopreservative, following the single-cell protocol of
Rinke et al. (2014). Unfortunately, many of these samples contained extremely large amounts of
cellular debris, likely from improper cryopreservation in field conditions. Additionally,
cryopreservation of samples required a non-trivial amount of laboratory preparation in non-
sterile field conditions that we believe introduced substantial human contamination to SSR-FL.
While we were able to sequence one of the frozen samples (SSR-ML) at high coverage and
replicate it with RNAlater, we cannot presently recommend in-field cryopreservation of fecal
samples for FACS. RNAlater is often commonly used in molecular field primatology, because it

offers long-term, stable preservation of host DNA at room temperature. For our purposes, it also
offered the distinct advantage of not requiring any in-field laboratory preparation, which
minimized human contamination of our cellular populations. Attempts to flow sort RNAlater
preserved cells of any origin are extremely scant, and we only found two such studies in the
literature (Barrett et al., 2002; Zaitoun, Erickson, Schell, & Epstein, 2010). We observed a
substantial improvement in the cellular integrity in the RNAlater preserved samples. Although
cells preserved in RNAlater are dead, they suffer minimal histological disruption, and maintain
cellular epitopes critical for antibody binding (Florell et al., 2001). Given this array of benefits,
we recommend preservation of fresh fecal samples in RNAlater when collected in field
conditions.
435
4.2 Quality and feasibility of high-coverage fecal genomics
We have presented the first high-coverage, unbiased mammalian genome, derived
exclusively from feces. While traditional bait-and-capture approaches to non-invasive genomics
have allowed for broad sampling of the mammalian genome from feces, such methods remain
limited by genomic bias. When compared to tissue-derived capuchin genomes, our FACS-
derived fecal genome indicates no such biases. SSR-ML consistently fell within or immediately
adjacent to the observable range of variation of the other tissue samples collected from Sector
Santa Rosa. While we acknowledge that it would have been optimal to compare high-coverage
whole genomes generated from the blood and feces of the same individual, this was not possible,
because of our non-invasive sampling strategy. Nonetheless, we are able to infer such a
comparison through the use of pedigree data in our SSR samples. The scaffold-wide pattern of
heterozygosity observable in SSR-ML (Figure 4D) is nearly identical to that of SSR-CR (tissue),
who was his sibling. This relationship is further supported by the population clustering results
(section 4.3). Furthermore, the SSR-ML sample we used in SNV calling did not bear any
indication of human contamination. In order to remain consistent in comparison with the tissue
and blood derived samples, we did not remove reads mapping to the hg38 with BBsplit during
SNV calling. Because the *Cebus* genome is less complete than hg38, it is likely that the majority
of human-specific mapping from this and other samples is artifactual. Given the consistency
similarity of the SSR-ML sample to the others from SSR, we and suggest that FACS is a viable
approach to expand the horizons of non-invasive population and conservation genomics.
Prior to selecting libraries for high-coverage sequencing, we suggest that multiple
libraries should be run on a lower throughput sequencing platform (e.g. MiSeq). Given the
variability in sequencing outcomes inherent in our technique, it would be prudent to avoid
wasting sequencing capacity on libraries that lack the requisite diversity for high-depth
sequencing. Working with extremely low numbers of cells, which is sometimes the result of the
FACS process, can result non-trivial duplication rates and the potential for the introduction of
human contaminants. Given that our FACS protocol only requires a small amount of fecal slurry,
processing two or three aliquots from the same fecal sample would increase the number of cells
and, presumably, the available diversity in cases where it was deemed necessary.
465
4.3 Population structure of white-faced capuchin monkeys
By successfully discriminating among two populations of white-faced capuchins in Costa
Rica, we have demonstrated that fecal FACS is effective for low-coverage applications of
population and conservation genomics. While bait-and-capture approaches remain a valuable

tool for the assessment of population genetic structure from real-world distributions of free-
ranging mammals, fecal FACS provides a simple alternative approach.
The clustering patterns in our trees and PCA plots do not reveal any samples that deviate
from their expected geographic or ecological origin. These relationships are robust to both the
coverage levels (< 1X to > 50X) and biological origins (feces, tissue, and blood) of the samples.
The tight geographic clustering of individuals within the SSR sampling locale provides
reasonable evidence that there is no substantial effect from fecal FACS on population structure.
Were it the case that fecal FACS introduced substantial bias, we would have expected the fecal
samples from SSR to plot in a separate cluster from those of tissue origin. As fecal and tissue
samples fall in the same general cluster, this is no evidence of such an effect. Furthermore,
known pedigree information from SSR corresponds to the genetic relationships observed in our
SNV tree. SSR-ML (fecal) and SSR-CR (tissue) form an internal clade in the tree (sixth and
seventh points from the top). These two individuals also cluster adjacent to each other on the
PCA plot.
484
4.4 Summary
Through a novel use of flow cytometry/FACS, we have developed a new method for the
isolation of epithelial cells from mammalian feces for population genomics. We generated the
first high-coverage, unbiased mammalian genome solely from feces. Additionally, we have
demonstrated that fecal FACS can be used to generate low coverage SNP datasets that function
well in population assignment and clustering algorithms. Fecal FACS is cost-effective and free
of the biases that commonly occur in traditional bait-and-capture approaches to the enrichment
of endogenous DNA from feces. Furthermore, fecal FACS does not require costly impractical
preservation of biomaterial in liquid nitrogen; rather, we rely on room-temperature stable storage
in RNAlater. Fecal FACS offers great benefits to the field of mammalian conservation and
population genomics.
496
497   **AUTHOR CONTRIBUTIONS**
498
499 Research was designed by JDO, ADM, RK and JC; performed by JDO, RK, CF, LFKK, EL, and
500 JT; analyzed by JDO and MM; and written by JDO, ADM, and TMB.
501
502   **DATA ACCESSIBILITY STATEMENT**
503
504 Sequencing data will be archived on NCBI SRA and made publically available.
505
506   **ACKNOWLEDGEMENTS**
507

We would like to thank Laurie Kennedy, Yiping Liu form the University of Calgary Flow
Cytometry Core for their patience and assistance with developing this protocol. Additionally, we
thank Jene Weatherhead, Shelley Wegener, Frank Visser, Gwen Duytschaever for molecular
laboratory assistance. We acknowledge Oscar Fornas for helpful discussion. Thanks to Wes
Warren, Pat Minx, Mike Montague, Shoji Kawamura, and J. Pedro Magalhaes for their
involvement with the development of the *Cebus imitator* reference genome. PJ Perry, Shasta
Webb, Rachel Williamson, and Saul Cheves Hernandez assisted with sample acquisition. This
research was supported by the National Sciences and Engineering Research Council of Canada

(NSERC), and the Canada Research Chairs program to ADM. TMB is supported by BFU2017-
86471-P (MINECO/FEDER, UE), Howard Hughes International Early Career, Obra Social "La
Caixa" and Secretaria d'Universitats i Recerca del Departament d'Economia i Coneixement de la
Generalitat de Catalunya. JDO is supported by the Alberta Children's Hospital Research Institute
(ACHRI). CF is supported by "La Caixa" PhD fellowship. L.F.K.K. is supported by an FPI
fellowship associated with BFU2014-55090-P (MINECO/FEDER, UE)
522
523

524 **FIGURES**
525

526
527
Figure 1: Map of sampling sites. Sector Santa Rosa (SSR) and Cañas are situated in the northern
dry forest and samples from Quepos and Manuel Antonio are from the southern wet forest. Map
courtesy of Eric Gaba—Wikimedia Commons user: Sting.
531
532

533
534
Figure 2: FACS gating strategy. Cells were gated first by size and complexity to avoid bacteria
and cellular debris (A), followed by discrimination of cellular agglomerations (B and C).
Fluorescence of AE1/AE3 Anti-Pan Cytokeratin Alexa Fluor® 488 antibody (FITC-A) is
depicted in unstained (D) and stained (E) cellular populations. Epithelial cells were identified as
those fluorescing beyond background levels, as depicted in the P4 gate.
540

541
542
Figure 3: Mapping percentages of sequencing reads from RNAlater preserved fDNA libraries
prepared with FACS for A) all samples, and B) individual libraries. C) Increase in mapping rate
for RNAlater preserved samples. D) Relationship between mapped read duplication and number
of cells with LOESS smoothing. The duplicate rate decreases sharply once a threshold of about
1,000 cells is reached.
548

Figure 4: A) Density of genomic coverage of high coverage genomes from Santa Rosa. B)
Average coverage per 100 KB window along the largest scaffold of the *C. imitator* 1.0 reference
genome. C) Density of 1 MB windows at varying levels of heterozygosity along the entire
genome. D) Heterozygosity of 100 KB windows along the largest scaffold of the *C. imitator* 1.0
reference genome. The top two genomes (SSR-CR and SSR-ML) are from siblings. The order of
individuals in figures B and D correspond to that of figure A.

558
559
560 **Figure 5:** Percent of GC content across the genome for the four tissue (red) and one fecal (blue)
561 samples from Sector Santa Rosa. GC content does not substantially differ for each type of
562 sample. A) Average normalized coverage at each percentage of GC. B) Number of reads per 100
563 bp window (scaled by the number aligned reads) at each percentage of GC.
564
565

566
567
568 **Figure 6:** Left: Principal components of 14 fecal and 10 blood/tissue libraries from white faced
569 capuchin and three outgroups. Right: Maximum likelihood tree of 9 fecal and 10 blood/tissue
570 libraries. Samples with less 0.5X coverage were excluded. Among the white-faced capuchin
571 samples, individuals from northern (dry forest) and southern (wet forest) regions form the
572 primary split; secondary splits reflect the individuals from different sites within regions.
573

574 **TABLES**:
575
576 **Table 1:** Origins and preservation information for *Cebus imitator* samples.
577

Sample	Region	Site	Sample Type	Preservation
SSR-NM	North	Sector Santa Rosa	Feces	RNA*later*
SSR-TY	North	Sector Santa Rosa	Feces	RNA*later*
SSR-FN	North	Sector Santa Rosa	Feces	RNA*later*
SSR-LE	North	Sector Santa Rosa	Feces	RNA*later*
SSR-CH	North	Sector Santa Rosa	Feces	RNA*later*
SSR-FG	North	Sector Santa Rosa	Feces	RNA*later*
SSR-LU	North	Sector Santa Rosa	Feces	RNA*later*
SSR-SN	North	Sector Santa Rosa	Feces	RNA*later*
SSR-KI	North	Sector Santa Rosa	Feces	RNA*later*
SSR-RF	North	Sector Santa Rosa	Feces	RNA*later*
SSR-ML	North	Sector Santa Rosa	Feces	RNA*later*
SSR-PR	North	Sector Santa Rosa	Feces	RNA*later*
SSR-SB1	North	Sector Santa Rosa	Feces	RNA*later*
SSR-SB2	North	Sector Santa Rosa	Feces	RNA*later*
SSR-ML	North	Sector Santa Rosa	Feces	Frozen
SSR-FL	North	Sector Santa Rosa	Feces	Frozen
SSR-CR	North	Sector Santa Rosa	Tissue	Frozen
SSR-FL	North	Sector Santa Rosa	Tissue	Frozen
SSR-TH	North	Sector Santa Rosa	Tissue	Frozen
SSR-T5-1	North	Sector Santa Rosa	Tissue	Frozen
SSR-RM08	North	Sector Santa Rosa	Tissue	Frozen
CNS-HE	North	Cañas	Blood	Frozen
KSTR29	South	Manuel Antonio	Blood	Frozen
KSTR116	South	Manuel Antonio	Blood	Frozen
KSTR159	South	Manuel Antonio	Blood	Frozen
KSTR64	South	Quepos	Blood	Frozen

578
579

580   **Table 2**: FACS and mapping results from *Cebus* and *Ateles* fecal samples
581

Monkey	Library	Cells	PCR Cycles	Total DNA (ng)	% Mapping					X Coverage
					BWA mem	BBsplit *Cebus*	Unique *Cebus*	Duplicate *Cebus*	BBsplit Human	
SSR-ML	SSR-ML Frozen	2546	11	10.50	96	90	85	5	1.25	11.7
SSR-FL	SSR-FL	4405	12	6.72	80	42	40	3	15.77	4.4
SSR-FN	SSR-FN	62601	8	21.50	97	93	86	6	0.81	2.8
SSR-FG	SSR-FG	580	10	9.75	94	84	60	24	2.86	2.0
SSR-LU	SSR-LU	8998	10	8.00	93	84	72	11	0.89	2.0
SSR-ML	SSR-ML RNAlater	42837	10	8.26	88	67	63	4	1.08	1.9
SSR-TY	SSR-TY	140	10	7.70	98	94	64	30	1.24	1.5
SSR-SB	SSR-SB 2	129	10	9.00	79	60	39	22	1.00	1.1
SSR-SB	SSR-SB 1	11944	10	6.25	55	11	8	2	0.61	
SSR-KI	SSR-KI	612	10	9.00	93	78	45	33	5.80	1.0
SSR-RF	SSR-RF	257	10	10.00	92	78	38	40	5.18	0.7
SSR-NM	SSR-NM	3336	11	3.38	98	95	92	3	0.66	0.4
SSR-CH	SSR-CH	957	11	4.06	93	85	80	5	0.74	0.4
SSR-LE	SSR-LE	1612	11	2.96	96	91	81	11	0.91	0.3
SSR-SN	SSR-SN	1866	11	3.96	92	79	74	6	3.07	0.2
SSR-PR	SSR-PR	12316	11	3.13	78	64	58	6	0.68	0.1
Spider	Spider 1	4026	12	6.96	54	12	11	1	2.83	0.4
Spider	Spider 2	602	11	4.50	49	12	10	1	1.82	
	Median (*Cebus*)*	2079		7.85	93	82	63	6	1	

582
583
584
585

REFERENCES

Arandjelovic, M., & Vigilant, L. (2018). Non-invasive genetic censusing and monitoring of
primate populations. *American Journal of Primatology*. doi:10.1002/ajp.22743

Barrett, M. T., Glogovac, J., Prevo, L. J., Reid, B. J., Porter, P., & Rabinovitch, P. S. (2002).
High-quality RNA and DNA from flow cytometrically sorted human epithelial cells and
tissues. *BioTechniques*, *32*(4), 888–90, 892, 894, 896.

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* , *30*(15), 2114–2120.

Bushnell, B. (2016). BBMap short read aligner. *University of California, Berkeley, California.*
URL Http://sourceforge.net/projects/bbmap.

Chiou, K. L. (2017). *Population Genomics of a Baboon Hybrid Zone in Zambia* (PhD Thesis).
Washington University in St. Louis. Retrieved from https://doi.org/10.7936/K7348HS3

Chiou, K. L., & Bergey, C. M. (2018). Methylation-based enrichment facilitates low-cost,
noninvasive genomic scale sequencing of populations from feces. *Scientific Reports*, *8*(1),
1975.

Corlett, R. T. (2017). A Bigger Toolbox: Biotechnology in Biodiversity Conservation. *Trends in*
Biotechnology, *35*(1), 55–65.

de Manuel, M., Kuhlwilm, M., Frandsen, P., Sousa, V. C., Desai, T., Prado-Martinez, J., …
Marques-Bonet, T. (2016). Chimpanzee genomic diversity reveals ancient admixture with
bonobos. *Science*, *354*(6311), 477–481.

Fedigan, L., & Rose-Wiles, L. (1996). See how they grow: Tracking capuchin monkey
populations in a regenerating Costa Rican dry forest. In M. A. Norconk, A. L. Rosenberger,
& P. A. Garber (Eds.), *Adaptive radiations of Neotropical primates* (pp. 289–307). Springer.

Florell, S. R., Coffin, C. M., Holden, J. A., Zimmermann, J. W., Gerwels, J. W., Summers, B.

K., … Leachman, S. A. (2001). Preservation of RNA for functional genomic studies: a

multidisciplinary tumor bank protocol. *Modern Pathology: An Official Journal of the United*

States and Canadian Academy of Pathology, Inc, *14*(2), 116–128.

Guschanski, K., Krause, J., Sawyer, S., Valente, L. M., Bailey, S., Finstermeier, K., …

Savolainen, V. (2013). Next-generation museomics disentangles one of the largest primate

radiations. *Systematic Biology*, *62*(4), 539–554.

Hernandez-Rodriguez, J., Arandjelovic, M., Lester, J., de Filippo, C., Weihmann, A., Meyer,

M., … Marques-Bonet, T. (2017). The impact of endogenous content, replicates and pooling

on genome capture from faecal samples. *Molecular Ecology Resources*. doi:10.1111/1755-

0998.12728

Kuderna, L. F. K., Lizano, E., Julia, E., Gomez-Garrido, J., Serres-Armero, A., Kuhlwilm, M., …

Marques-Bonet, T. (2018, June 13). *Selective single molecule sequencing and assembly of a*

human Y chromosome of African origin. *bioRxiv*. doi:10.1101/342667

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics* , *25*(14), 1754–1760.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., … DePristo,

M. A. (2010). The Genome Analysis Toolkit: a MapReduce framework for analyzing next-

generation DNA sequencing data. *Genome Research*, *20*(9), 1297–1303.

Perry, G. H. (2014). The Promise and Practicality of Population Genomics Research with

Endangered Species. *International Journal of Primatology*, *35*(1), 55–70.

Perry, G. H., Marioni, J. C., Melsted, P., & Gilad, Y. (2010). Genomic-scale capture and

sequencing of endogenous DNA from feces. *Molecular Ecology*, *19*(24), 5332–5344.

Pickrell, J. K., & Pritchard, J. K. (2012). Inference of population splits and mixtures from

genome-wide allele frequency data. *PLoS Genetics*, *8*(11), e1002967.

Prado-Martinez, J., Sudmant, P. H., Kidd, J. M., Li, H., Kelley, J. L., Lorente-Galdos, B., …

Marques-Bonet, T. (2013). Great ape genetic diversity and population history. *Nature*,

499(7459), 471–475.

Price, A. L., Patterson, N. J., Plenge, R. M., Weinblatt, M. E., Shadick, N. A., & Reich, D.

(2006). Principal components analysis corrects for stratification in genome-wide association

studies. *Nature Genetics*, *38*(8), 904–909.

Rinke, C., Lee, J., Nath, N., Goudeau, D., Thompson, B., Poulton, N., … Woyke, T. (2014).

Obtaining genomes from uncultivated environmental microorganisms using FACS-based

single-cell genomics. *Nature Protocols*, *9*(5), 1038–1048.

Sasaki, D. T., Dumas, S. E., & Engleman, E. G. (1987). Discrimination of Viable and Non-

Viable Cells Using Propidium Iodide in Two Color Immunofluorescencel. *Alan R. Liss, Inc.*

Cytometry, *8*, 413–420.

Snyder-Mackler, N., Majoros, W. H., Yuan, M. L., Shaver, A. O., Gordon, J. B., Kopp, G. H., …

Tung, J. (2016). Efficient Genome-Wide Sequencing and Low-Coverage Pedigree Analysis

from Noninvasively Collected Samples. *Genetics*, *203*(2), 699–714.

van der Valk, T., Lona Durazo, F., Dalén, L., & Guschanski, K. (2017). Whole mitochondrial

genome capture from faecal samples and museum-preserved specimens. *Molecular Ecology*

Resources, *17*(6), e111–e121.

Wall, J. D., Schlebusch, S. A., Alberts, S. C., Cox, L. A., Snyder-Mackler, N., Nevonen, K.

A., … Tung, J. (2016). Genomewide ancestry and divergence patterns from low-coverage

sequencing data reveal a complex history of admixture in wild baboons. *Molecular Ecology*,

25(14), 3469–3483.

Zaitoun, I., Erickson, C. S., Schell, K., & Epstein, M. L. (2010). Use of RNAlater in

fluorescence-activated cell sorting (FACS) reduces the fluorescence from GFP but not from

DsRed. *BMC Research Notes*, *3*, 328.