Title: **Shared Song Object Detector Neurons in *Drosophila* Male and Female Brains**

Drive Divergent, Sex-Specific Behaviors

Authors: Jan Clemens[1,2]*, David Deutsch[1]*, Stephan Y. Thiberge[1,3], and Mala Murthy[1,3,4,#]

Affiliations:

[1] Princeton Neuroscience Institute, Princeton University, Princeton 08540, NJ, USA

[2] current address: European Neuroscience Institute, Grisebachstrasse 5, Göttingen,

Germany

[3] Bezos Center for Neural Circuit Dynamics, Princeton Neuroscience Institute, Princeton

University, Princeton, NJ, USA

[4] Department of Molecular Biology, Princeton University, Princeton, NJ, USA

* These authors contributed equally

Lead contact

Correspondence: mmurthy@princeton.edu

Running Title: Shared Song Detector Neurons Drive Sex-Specific Behaviors

Abstract:

Males and females often produce distinct responses to the same sensory stimuli. How such

differences arise – at the level of sensory processing or in the circuits that generate behavior

– remains largely unresolved across sensory modalities. We address this issue in the

acoustic communication system of *Drosophila*. During courtship, males generate time-

varying songs, and each sex responds with specific behaviors. We characterize male and

female behavioral tuning for all aspects of song, and show that feature tuning is similar

between sexes, suggesting sex-shared song detectors drive divergent behaviors. We then

identify higher-order neurons in the *Drosophila* brain, called pC2, that are tuned for multiple

temporal aspects of one mode of the male's song, drive sex-specific behaviors, and show a

mirrored correspondence between sensory and motor tuning. We thus uncover acoustic

object detector neurons at the sensory-motor interface that flexibly link auditory perception

with sex-specific behavioral responses to communication signals.

Introduction

Across animals, males and females produce distinct, dimorphic behaviors in response to common sensory stimuli (e.g., pheromones, visual cues, or acoustic signals), and these differences are critical for social and reproductive behaviors (Billeter and Levine, 2013; Dulac and Wagner, 2006; Kelley, 2003; Yamamoto et al., 2013; Yang and Shah, 2014). The molecular dissection of sexual dimorphisms in the nervous system of flies and mice in particular (Cachero et al., 2010; Dulac and Wagner, 2006; Rideout et al., 2010; Stowers and Logan, 2010; Yang and Shah, 2014; Yu et al., 2010) has identified neurons involved in either processing important social cues or driving social behaviors, but it remains open as to how sex-specific behaviors to common sensory signals emerge along sensorimotor pathways. It could be that males and females process sensory information differently, leading to different behavioral outcomes, or that males and females process sensory information identically, but drive different behaviors downstream of common detectors.

This issue has been most heavily investigated for pheromone processing, and these studies point to differences along sensory pathways. For instance, in *Drosophila*, the male pheromone cVA induces either aggression in males (Wang and Anderson, 2010) or receptivity in females (Billeter et al., 2009; Kurtovic et al., 2007). The pheromone is detected by shared circuits in males and females and the similarly processed sensory information (Datta et al., 2008) is then routed to sex-specific higher-order olfactory neurons (Kohl et al., 2013; Ruta et al., 2010) that likely exert different effects on behavior, although this hypothesis has not yet been tested. In the mouse, the male pheromone ESP1 triggers lordosis in females, but has no effect on male behavior. This pheromone activates V2Rp5 sensory neurons in both sexes but, analogous to cVA processing in flies, these neurons exhibit sex-specific projection patterns in the hypothalamus that drive sex-specific behavioral responses (Haga et al., 2010; Ishii et al., 2017). For pheromone processing then, the rule appears to be that early olfactory processing is largely shared between the sexes and then common percepts are routed to separate higher-order neurons or circuits for control of differential behaviors. But does this rule apply for other modalities, or for stimuli that can be defined by multiple temporal or spatial scales (e.g. visual objects or complex sounds). For such stimuli, selectivity typically emerges in higher-order neurons (Dicarlo et al., 2012; Gentner, 2008; Tsao and Livingstone, 2008) and we do not yet know if such neurons are shared between males and females, and therefore if dimorphic responses emerge in downstream circuits.

Here, we investigate this issue in the auditory system in *Drosophila*. Similar to birds (Fortune et al., 2011; Konishi, 1985), frogs (Gerhardt and Huber, 2002), or other insects (Ronacher et al., 2014), acoustic communication in *Drosophila* involves different behaviors in males and females relative to the courtship song. During courtship, males chase females and produce a species-specific song that comprises two major modes – pulse song consists of trains of brief pulses and sine song consists of a sustained harmonic oscillation (Bennet-Clark and Ewing, 1967). In contrast with males, females are silent but arbitrate mating decisions (Bennet-Clark and Ewing, 1969). Males use visual feedback cues from the female (rapid changes in her walking speed and her distance relative to him) to determine which song mode (sine or pulse) to produce over time (Clemens et al., 2017; Coen et al., 2014; 2016) – this gives rise to the variable structure of song bouts (Fig. 1A). Receptive females slow in response to song (Aranha et al., 2017; Bussell et al., 2014; Clemens et al., 2015; Coen et al., 2014; Cook, 1973; Crossley et al., 1995; F. Von Schilcher, 1976; Tompkins et al., 1982), while playback of courtship song to males can induce them to increase their walking speed (Crossley et al., 1995; F. Von Schilcher, 1976; Vaughan et al., 2014), and to display courtship-like behaviors (Eberl et al., 1997; Li et al., 2018; Yoon et al., 2013; Zhou et al., 2015). These behavioral differences surrounding song production and perception between *Drosophila* males and females, combined with the wealth of genetic and neural circuit tools, make the *Drosophila* acoustic communication system an excellent one in which to investigate whether males and females share common sensory detection strategies for their courtship song, and how divergent behaviors arise.

Each major mode of *Drosophila* courtship song, sine or pulse, contains patterns on multiple temporal scales (Arthur et al., 2013; Bennet-Clark and Ewing, 1967) (Fig. 1A) – neurons that represent either the pulse or sine mode should in theory bind all of the temporal features of each mode, similar to object detectors in other systems (Bizley and Cohen, 2013; Dicarlo et al., 2012; Gentner and Margoliash, 2003; Griffiths and Warren, 2004), and their tuning should match behavioral tuning. Historically, behaviorally relevant song features have been defined based on the parameters of the species' own song (Bennet-Clark and Ewing, 1969). However, there is now ample evidence that the preferred song can diverge from the conspecific song (Amézquita et al., 2011; Blankers et al., 2015; Ryan et al., 2001) – for instance if females prefer exaggerated song features (Rosenthal and Ryan, 2011; Ryan and Cummings, 2013) or respond to signal parameters not normally produced by their male

conspecifics (Hennig et al., 2016). Moreover, the job of the female nervous system is not

only to tell one species apart from another, but also to select a particular mate from within

the species distribution. It is therefore important to define song modes by the acoustic tuning

of specific behavioral outputs. This has been done for other insects (e.g. (Clemens and

Hennig, 2013; D. von Helversen and O. von Helversen, 1997)) but never for flies in a

systematic way that also permits a direct comparison between sexes.

To that end, we developed a behavioral assay for assessing dynamic changes in walking

speed in response to sound playback in both sexes, and we then measured locomotor

tuning for all features of either pulse or sine song. This reveals that males and females have

similar tuning but different behavioral responses and that they are tuned for every major

feature of the song. We then identified a small set of sexually dimorphic neurons, termed

pC2 (Kimura et al., 2015; Rideout et al., 2010; Zhou et al., 2014) , that serve as shared

pulse song detectors in both sexes: the tuning of pC2 neurons is matched to behavioral

tuning for pulse song – but not for sine song – across a wide range of temporal scales. We

find that optogenetic activation of pC2 is sufficient to drive sex-specific behaviors produced

in response to pulse song – changes in locomotion with sex-specific dynamics as well as

singing in males. pC2 is also part of a sensorimotor loop, since it drives and is driven by

pulse song. Finally, we establish the importance of pC2 neurons by showing that early social

experience changes both the tuning of these neurons and the tuning of the behavior. Our

results indicate that the fly brain contains common song object detectors in males and

females which control sex-specific behavioral responses to song via downstream circuits.

Results

Comprehensive characterization of behavioral tuning for courtship song features

We designed a single-fly playback assay in which individual males or females receive

acoustic stimuli in the absence of any confounding social interactions, and we implemented

an automated tracker to analyze changes in locomotion relative to acoustic playback (Fig.

1B and Movie S1). The assay (which we refer to as FLyTRAP (**F**ly **L**ocomotor **TR**acking and

Acoustic **P**layback)) monitors dynamic changes in walking speed, which provides a readout

that can be directly compared between both males and females, as opposed to slower

readouts of sex-specific behaviors such as the female time to copulation (Bennet-Clark and

Ewing, 1969; Zhou et al., 2014) or male-male chaining (Yoon et al., 2013; Zhou et al., 2015).

Because of the high-throughput nature of our assay combined with automated tracking, we

can easily test a large number of flies and song parameters, including those only rarely

produced by conspecifics but to which animals might be sensitive (Coen et al. 2014; Aranha

et al., 2017; Bussell et al., 2014; Crossley et al., 1995; Eberl et al., 1997; Rybak et al.,

2002a; 2002b; F. Von Schilcher, 1976; Yoon et al., 2013; Zhou et al., 2015).

Previous studies that assayed either male-female copulation rates or male-male chaining

often focused on behavioral selectivity for the interval between pulses in a pulse train (inter-

pulse interval (IPI), Fig. 1A) (Bennet-Clark and Ewing, 1969; Rybak et al., 2002b; F. Von

Schilcher, 1976; Zhou et al., 2015). Using FLyTRAP, we systematically compared male and

female locomotor tuning to 82 acoustic stimuli that span the features and timescales present

in courtship song (see Supplemental Table 1). Typically, each stimulus was presented 23

times to 120 females and 120 males, generating >2500 responses per stimulus and sex (see

Methods).

We started by examining behavioral tuning for IPI using the wild type strain NM91. We

generated artificial pulse trains and varied only the IPI (between 16 and 96ms) – we chose a

stimulus intensity of 5 mm/s since varying intensity had minimal effect on pulse song

responses in females (Fig. S1A, B). Observed changes in speed were stimulus-locked, sex-

specific and tuned to IPI (Fig. 1C). While females slow down to pulse trains, males exhibit

transient slowing at pulse train onset followed by a long-lasting acceleration. The transient

component of the locomotor response was present for all stimuli (Fig. S1C, D, S2A-C) and

may correspond to an unspecific startle response to sound onset (Lehnert et al., 2013). The

transient was also present in females but masked by the stimulus-dependent slowing that

followed (Fig. 1C). Due to the briefness of the transient response, the integral change in

speed following stimulus onset reflects mostly the speed during the sustained phase (Fig.

S1C, D). For simplicity, we therefore used the full integral as an overall measure of

behavioral tuning. We found that in FLyTRAP, female IPI tuning is a band-pass filter

matched to the statistics of male song (Fig. 1D): the mode of the distribution of *Drosophila*

melanogaster IPIs is centered between 30 and 50 ms and females decrease their speed

most for the same IPI range, and less for shorter or longer IPIs. Males produced a similar

band-pass tuning curve peaked at the same IPI range - but their locomotor response was

opposite in sign (males accelerated, females decelerated). This is consistent with the results

of other assays that have found band-pass tuning for IPI in both sexes (Bennet-Clark and

Ewing, 1969; Rybak et al., 2002b; F. Von Schilcher, 1976; Zhou et al., 2015) and a sex-
specific sign of locomotor responses (Crossley et al., 1995; F. Von Schilcher, 1976).
170
We next systematically varied parameters that characterize pulse song to cover (and extend
beyond) the distribution of each parameter within *D. melanogaster* male song (just as we did
for IPI) (see Supp. Fig. 2). We examined behavioral tuning in both sexes for parameters that
varied on timescales of milliseconds (carrier frequency, pulse duration and IPI) to seconds
(pulse train duration) (Fig. 1A). We found that male and female tuning curves are of opposite
sign but similar shape for all pulse song features tested across time scales (Fig. 2A, B, S2D-
F, see Fig. S2A-C for speed traces), and that the behavioral tuning for pulse parameters
often overlapped the distribution found in natural song (Fig. 2C). While the behavioral tuning
curves for all pulse song features on short time scales are band-pass with a well-defined
peak, we found that tuning for pulse train duration was monotonous: both females and males
increase their locomotor response with increasing pulse train duration up to four seconds
(Fig. 2A, B). During natural courtship, pulse trains longer than four seconds are rarely
produced (Coen et al., 2014). Males also produce two distinct types of pulses (Clemens et
al., 2017) – we find that while females appear to be broadly tuned for both types of pulses in
the FLyTRAP assay, males respond preferentially to higher frequency pulses. Finally, we
found that both males and females are more selective for the pulse duration versus the
pulse pause, the two components of the IPI (Fig. S2D-F) – this is in contrast to other insects
that produce and process song pulses (e.g. crickets, grasshoppers, katydids), and that are
preferentially tuned to pulse pause, pulse period or pulse train duty cycle (Hennig et al.,
2014; Ronacher et al., 2014).
191
We next tested locomotor tuning for the parameters that characterize sine stimuli – carrier
frequency and the duration of sine trains (Fig. 1A). Both males and females slow for sine
tones of different frequencies, with very low and very high frequencies eliciting the strongest
responses (Fig. 2A, B and Fig. S2A-C). Notably, the frequencies inducing the strongest
slowing (100Hz) are not typically produced by males (Fig. 2C). As for sine train duration
tuning, we observed sustained responses that increased with duration and saturated only
weakly, possibly because of the weak response magnitude.
199
Pulse and sine song usually co-occur within a single bout but it is not known why males
produce two different modes (although females respond to both during natural courtship

(Clemens et al., 2017; Coen et al., 2014)). One possibility is that one mode exerts a priming
effect on the other (F. V. Schilcher, 1976). To test interactions between the two song modes,
we presented sequences in which a 2-second pulse train was followed by a 2-second sine
tone or in which a sine tone was followed by a pulse train and compared the responses for
these sequences to the responses to an individual pulse train or sine tone (Fig. S2G). The
responses are well explained by a linear combination of the responses to individual sine or
pulse trains. Deviations from linearity occur due to sound onset responses, but otherwise
responses do not strongly depend on the order of presentation in a bout (see also (Talyn
and Dowse, 2004)). This suggests that these stimuli are processed in independent
pathways.
212
To summarize, we compared behavioral responses in males and females for all features that
define the courtship song. We found that male and female speed changes were strongly
correlated for both song modes but that the sign of the correlation was negative for pulse
stimuli and positive for sine stimuli (Fig. 2E). The opposite sign of the correlations along with
the independence of responses to sine and pulse stimuli (Fig. S2G) indicates that sine and
pulse song are processed by different circuits. The large magnitude of the correlations
implies that feature tuning of the behavioral responses is similar between sexes and
suggests that detector neurons for each song mode should be shared between sexes.
221
Hearing pulse song drives wing extension in males, but not in females
Another sex-specific aspect of song responses is courtship: playback of conspecific song
induces courtship-like behavior in males – this can even be directed towards other males,
leading to the male chaining response, in which males follow other males, chasing and
extending their wings (Eberl et al., 1997; F. Von Schilcher, 1976; Yoon et al., 2013). In our
single-fly assay, males lack a target for courtship and the song-induced arousal likely
manifests as an increase in speed. Since FLyTRAP does not permit simultaneous recording
of fly acoustic signals during playback, we quantified wing extension as a proxy for singing,
and examined whether song playback alone drives singing in solitary males. We found that
solitary males extend their wings in response to pulse song stimuli specifically (Fig. 2F, G,
Movie S2). This behavior is tuned for the inter-pulse interval (similar to the locomotor
response, Fig. 1D) – the conspecific IPI of 36ms drives the most wing extension, and shorter
and longer IPIs evoke fewer wing extensions. By contrast, conspecific sine song (150Hz)
does not induce wing extension (Fig. 2F) (see also (Eberl et al., 1997; Yoon et al., 2013)).

We also found that playback of pulse does not elicit wing extension in females, even though
females have been shown to possess functional circuitry for singing (Clyne and Miesenböck,
2008; Rezával et al., 2016) – wing extension in response to pulse song is thus sex-specific.
239
These results are consistent with those for locomotor tuning: pulse song, but not sine song,
generates sex-specific differences in behavior. The identical tuning of the two behavioral
responses in males (locomotion (Fig. 1C) and song production (Fig. 2G)) suggests that the
behavioral responses are driven by a common circuit. Finally, that playback of pulse song
alone is sufficient to drive singing in males implies the existence of neurons with
sensorimotor correspondences, similar to "mirror neurons" (Mooney, 2014; Prather et al.,
2008; Rizzolatti and Arbib, 1998) – we explore this hypothesis below.
247

***Drosophila* male and female brains share pulse song detector neurons**
Our systematic exploration of song stimulus space using the FLyTRAP assay revealed
behavioral tuning for song parameters across temporal scales (from the carrier frequencies
of sine and pulse lasting milliseconds to the duration of sine and pulse trains lasting
seconds). We next searched for neurons with tuning across temporal scales that could serve
as object detectors of either the pulse or sine mode of courtship song. We focused on
neurons expressing the Doublesex (Dsx) transcription factor that regulates sexual
dimorphism in cell number and neuronal morphology between males and females. In the
central brain there are ~70 Dsx+ neurons per hemisphere in females and ~140 Dsx+
neurons per hemisphere in males (Kimura et al., 2015; Rideout et al., 2010). Previous
studies found calcium responses to both song-like stimuli and pheromones in Dsx+ neuron
projections in females (Zhou et al., 2014) and tuning for the inter-pulse interval in males
(Zhou et al., 2015). In addition, silencing subsets of Dsx+ neurons in females affected
receptivity (Zhou et al., 2014). These data suggest that Dsx+ neurons could serve as the
common pulse song detectors in males and females. To test this possibility, we recorded
auditory responses in Dsx+ neurons and examined tuning for song features across
timescales, in both males and females, to compare with our behavioral results.
265

266 We imaged neural activity using the calcium sensor GCaMP6m (Chen et al., 2013)
267 expressed only in Dsx+ neurons and focused on the lateral junction (LJ) (Cachero et al.,
268 2010; Ito et al., 2014; Yu et al., 2010), a site of convergence for the majority of Dsx+ neuron
269 projections (Fig. 3A, B, S5B, C, Movie S3, S4). We found that male and female Dsx+

projections in the LJ are driven strongly by pulse, but not by sine, stimuli (Fig. 3C) similar to previous results (Zhou et al., 2014). While males overall produce weaker responses to auditory stimuli compared with females (Fig. 3C), the normalized LJ responses are highly correlated between sexes – stimuli that evoked the strongest responses in females also evoked the strongest responses in males (Fig. 3D).

In order to connect behavioral tuning to neural tuning, we validated that the genetic strain in which we imaged calcium responses possessed the same behavioral tuning as the strain used in FLyTRAP (Fig. S3A). The sex-specific sign and shape of the IPI tuning curves of flies from these two strains matches acoustic tuning measured in other playback assays: females slow for pulse song (Clemens et al., 2017; Coen et al., 2014; Crossley et al., 1995; F. Von Schilcher, 1976) and copulate most when exposed to conspecific IPIs and less for shorter or longer IPIs (Bennet-Clark and Ewing, 1969; Li et al., 2018); males in groups increase their speed (Crossley et al., 1995; F. Von Schilcher, 1976; Vaughan et al., 2014) and court other females or males most when exposed to pulse song with the conspecific IPI (Crossley et al., 1995; Yoon et al., 2013; Zhou et al., 2015). The two strains chosen for the behavioral and calcium imaging experiments thus display the species-typical acoustic tuning likely expressed during natural courtship encounters. Interestingly, the behavioral tuning for IPI in seven additional wild type strains is still sex-specific but deviates from the species-typical tuning (Fig. S3B) when assayed in FLyTRAP, even though the same wild type strains – including NM91 – display virtually identical responses to song in a courtship assay in which a male courts a female (Clemens et al., 2017; 2015; Coen et al., 2016; 2014). This indicates that these strains require additional cues (e.g., pheromones or visual cues) to fully express their preference for conspecific song features.

When examining neuronal tuning curves, we found a good match between Dsx+ LJ responses and the magnitude of changes in speed across all timescales of pulse song in both sexes (Fig. 3E, F). For example, the Dsx+ LJ tuning curve for IPI is similar in females and males with the strongest responses at 36ms, matching the behavioral tuning curves (compare with Fig. 2A,B) – moreover, Dsx+ neurons are also preferentially tuned for the pulse duration over the pulse pause (Fig. S4D, E – compare with Fig. S2E, F). At longer timescales, LJ tuning curves also match behavioral tuning curves for pulse train duration – while the integral calcium continues to increase with train duration, the peak fluorescence saturates in both sexes, similar to the behavioral response. Apart from this difference, peak

and integral ΔF/F are similar (Fig. S4F, G). Overall, LJ responses are selective for the

features found in conspecific pulse song: LJ responses were strongest for stimuli with a

carrier frequency of 250Hz, an inter-pulse interval of 36ms, and a pulse duration of 16ms

(Fig. S4A-C). A match in only two of these three features was not sufficient to maximally

drive Dsx+ neurons. Sine stimuli have lower carrier frequencies, long durations, and no

pauses (they are by definition continuous) – which explains the weak responses of Dsx+

neurons to all sine stimuli (Fig. 3C).

We next directly compared Dsx+ LJ responses and behavioral responses for all stimuli and

found strong correlations for pulse stimuli in both sexes – although with a sex-specific sign.

That is, male neural and behavioral tuning for pulse stimuli are positively correlated (r=0.55,

p=9x10^{-6}) – high Dsx+ neuron activity correlates with the most acceleration (Fig. 3G).

Female neural and behavioral tuning for pulse stimuli are negatively correlated (r=-0.62,

p=2x10^{-7}) – high Dsx+ neuron activity correlates with the most slowing (Fig. 3H). This is

consistent with these neurons controlling the magnitude, but not the direction of speed

changes. We observed no statistically significant correlation for sine stimuli (male: r=-0.15,

p=0.56; female: r=0.27, p=0.30), as Dsx+ LJ responses to sine stimuli were weak. Note that

Dsx+ LJ activity only accounts for roughly 1/3 of the variability in behavioral responses for

pulse song. This suggests that the behavior is driven and modulated by additional pathways

outside of the Dsx+ neurons in the LJ. Nonetheless, Dsx+ neurons that innervate the LJ

have tuning properties expected for pulse song object detectors – they prefer pulse over

sine stimuli, are similarly tuned in males and females, and their feature tuning matches the

behavioral tuning for all pulse, but not sine, stimuli across timescales.

Dsx+ pC2l neurons are pulse song detectors

The Dsx+ neurons of the central brain form a morphologically heterogeneous population with

several distinct, anatomical clusters many of which project to the LJ (Kimura et al., 2015;

Rideout et al., 2010; Zhou et al., 2015; 2014) (Fig. 3A). Previous studies that examined

auditory responses in Dsx+ neurons (Zhou et al., 2015; 2014) did not resolve which subtype

carried the response. Using a stochastic labeling approach (Nern et al., 2015), we confirmed

that only five out of eight Dsx+ cell types in the female brain project to the LJ (Kimura et al.,

2015): pC1, pC2l/m, pMN1, and pMN2, but not pCd1/2 and aDN (Fig. 4A and Fig. S5D, E).

We next imaged calcium responses to pulse and sine stimuli in the somata of all five Dsx+

cell types that innervate the LJ and found that a subset of neurons in the pC1 and pC2

clusters possess auditory responses, in addition to cell type pMN2 (a female-specific neuron (Kimura et al., 2015) comprising only one cell body per hemisphere) (Fig. 4B, C, Movie S5, S6). All responsive cells preferred pulse over sine stimuli (Fig. 4D). We did not observe auditory responses in pMN1 neurons (not shown), although we cannot rule out that this neuron class has responses that are below the level of detection by the calcium indicator GCaMP6m.

The pC1 cluster contained a small number of auditory cells (2-3 cells in the female brain; we found none in the male brain) (Fig. 4C). Since pC1 and pNM2 contain only few auditory-responsive neurons and/or are present only in females, we focused on pC2 as the putative pulse song detector common to both sexes. Although there are more pC2 neurons in males versus females (~67 vs. ~26, (Kimura et al., 2015)) the number of auditory ROIs is similar in both sexes (~15). pC2 neurons can be subdivided into a lateral and a medial type, termed pC2l and pC2m (Robinett et al., 2010), and each type projects to the lateral junction via a distinct process (see Fig. 3A, S5B, C). Most auditory ROIs were lateral in the pC2 cluster (Fig. S5A), and all somata as well as the pC2l process produced strong auditory responses that were highly correlated with those we recorded in the LJ (Fig. 4F, G). From this we conclude that LJ responses reflect the tuning of pC2l neurons. Importantly, the tuning of the pC2l process matches the behavioral tuning (Fig. 4H), indicating that pC2l neurons are pulse song object detectors.

Optogenetic activation of pC2l neurons drives sex-specific behaviors

If pC2l neurons serve as pulse song detectors, then their activation should also be sufficient to drive the sex-specific behaviors observed for pulse song – changes in locomotion and singing that are distinct between males and females. To test this hypothesis, we used a driver line (Zhou et al., 2014; Rezával et al., 2016) that labels 11/22 female and 22/36 male pC2l neurons, in addition to 5-6 pCd neurons, but no pC2m or pC1 neurons (Fig. S6A). At least 5 of the pC2l cells in this driver line responded to song (Fig. S6B), which corresponds to ~1/3 of the auditory pC2l neurons. We expressed CsChrimson, a red-shifted channelrhodopsin (Klapoetke et al., 2014), in these neurons and optogenetically activated them in both males and females.

We first recorded behavior in a chamber tiled with microphones (Coen et al., 2014) to test whether pC2 activation was sufficient to induce singing, as we previously showed that pulse

song playback alone drives wing extension in males (Fig. 2F, G). Upon red light activation,

males produced pulse song, while sine song was produced transiently after stimulus offset

(Fig. 5A, Movie S7), and the amount of pulse song produced scaled with the strength of

activation (Fig. 5B). The evoked pulse and sine songs were virtually indistinguishable from

natural song (Fig. S6C, D). In *Drosophila*, retinal (the channelrhodopsin cofactor) must be

supplied via feeding, and red light stimulation drove singing significantly more in males fed

with retinal versus those fed regular food (Fig. S6E). Activation of a control line that only

labels pCd neurons (Zhou et al., 2014) did not drive any singing (Fig. S6E), implying that

song production results from the activation of the pC2 neurons in our driver. Importantly, we

never observed song production upon pC2 activation in females (Fig. S6E) – pC2 neurons

thus drive song in a sex-specific manner. These results also establish pC2 neurons as

serving a dual sensory and motor role, analogous to "mirror neurons" in other systems

(Mooney, 2014): they respond selectively to the pulse song (Fig. 3C, F) and also bias the

song pathway towards producing the same song mode (Fig. 5A, B).

We next evaluated changes in locomotor speed in both sexes for multiple levels of

optogenetic activation by placing flies in the FLyTRAP assay but using red light for activation

(instead of sound). To account for innate visual responses to the light stimulus, we

subtracted the responses of normally fed flies from retinal fed flies (Fig. S6F). Because the

behavioral responses to acoustic stimuli can depend on genetic background (Fig. S3), we

assessed responses to playback of pulse stimuli with varying IPI for the genotype used in

the optogenetic activation experiments (see Methods for list of genotypes). For this particular

strain, both males and females decreased their walking speed in response to pulse stimuli.

Nonetheless, we observed sex-specific differences in both the magnitude and dynamics of

the behavioral responses to sound (Fig. 5D-G), and optogenetic activation of pC2 neurons

reproduced many of these sex-specific differences (Fig. 5 H-K). For instance, male

responses outlasted the stimulus for both sound and optogenetic activation, while female

responses were multiphasic for both conditions. To confirm that the response dynamics

were sex-specific, we performed principal component analysis (PCA) on the speed traces of

males and females for sound playback and optogenetic activation experiments (Fig. 5L). The

first two principal components were sufficient to explain 84% of the variance in the speed

traces, and the responses varied along sex-specific axes: the PCA scores for males varied

most in the direction of the first principal component and female PCA scores align with the

orthogonal, second principal component. In each sex, responses for acoustic and

optogenetic stimulation were co-aligned, suggesting that pC2 neuron activation recapitulates

the sex-specific locomotor dynamics observed for acoustic playback in this strain.

Finally, we used the same pC2-specific driver to constitutively suppress the synaptic output

of pC2 (via expression of TNT (Sweeney et al., 1995)) in females and paired them with wild

type virgin males (see Methods). We quantified female song responses as the correlation

between different song features and female speed (Clemens et al., 2015; Coen et al., 2014)

(Fig. 5M-O). Because male song is structured via sensory feedback cues from the female

(Coen et al., 2014), silencing pC2 neurons in females could affect the content of male song

– however, the statistics of male song were unchanged by the female manipulation (Fig.

S6G-H). pC2 inactivation specifically affected the correlation between female speed and the

pulse song IPI, which changed from ~0 to 0.3 (Fig. 5M-O). While control – and wild type

(Clemens et al., 2015) – females do not change their speed relative to the range of natural

IPIs produced by conspecific males, females with pC2 neurons silenced accelerate more

with increasing IPI. This indicates that pC2 neurons are required for the proper response to

pulse song, and that other neurons that represent pulse song are tuned for longer IPIs.

While female locomotor responses to courtship song were affected by pC2 inactivation,

copulation rates were not significantly reduced (Fig. S6I), consistent with previous studies

(Zhou et al., 2014). In conjunction with the match between behavioral tuning and pC2 tuning,

these results add to the evidence that pC2 neurons serve as pulse song object detectors

and play a critical role at the sensorimotor interface – they relay information about pulse

song to sex-specific downstream circuits that control either singing or locomotion, and

thereby mediate acoustic communication behaviors.

pC2 neural activity is modulated by social experience

Many sexual behaviors change with social experience (Keleman et al., 2012; Li et al., 2018;

Marlin et al., 2015; Remedios et al., 2017). This plasticity could be mediated by modulating

the selectivity and the gain of object detectors or that of downstream circuits. Social

experience is known to affect courtship behavior in Drosophila (Ellis and Kessler, 1975;

Kohatsu and Yamamoto, 2015; F. Von Schilcher, 1976), and prolonged exposure to

conspecific song sharpens the IPI selectivity of the female mating decision and the male

chaining response (Li et al., 2018). We first tested whether locomotor responses in

FLyTRAP are also modulated by social experience. We found that males that were exposed

to social cues (including the song of other males) via group-housing exhibited stronger and

more selective changes in speed than individually housed males (Fig. 6A). Females do not

sing to other females when group housed and accordingly, their locomotor responses were

unaffected by housing conditions (Fig. 6A). These sex-specific effects of housing were

reflected in changes in the neuronal responses of pC2 neurons: calcium responses in pC2

(measured via the LJ) (Fig. 4F,G) do not change strongly with housing conditions in females,

but become more selective for IPI in group-housed males (Fig. 6B). Notably, sine song

responses and responses to pulse trains with different durations are not affected by housing

conditions (Fig. S7).

Discussion

Using a quantitative behavioral assay, we characterized locomotor responses in both males

and females to the features that define the *Drosophila melanogaster* courtship song. Males

and females showed similar tuning for pulse song stimuli, but nonetheless produced distinct

responses (for example, males accelerate while females decelerate; males sing while

females do not) (Fig. 1, 2). Both males and females were responsive to all features of pulse

song, across timescales, and tuning was matched to the distribution of each parameter in

the male's song. We then identified Dsx+ pC2 neurons in the brain that respond selectively

to pulse song stimuli, and whose tuning is matched to behavioral tuning (Fig. 3, 4). The

activation of pC2 neurons elicited sex-specific behavioral responses to pulse song (Fig. 5),

and social experience sharpened both behavioral feature selectivity and pC2 tuning (Fig. 6).

We thus conclude that Dsx+ pC2 neurons are pulse song object detectors that couple song

detection with the execution of sex-specific behaviors.

In other systems, object detectors are typically higher-order neurons that bind the multiple

low-level features that compose an object across spatial and temporal scales (Brincat and

Connor, 2004; Dicarlo et al., 2012): visual object detectors match visual features across

spatial scales (Freiwald and Tsao, 2010; Quiroga, 2012; Tsao et al., 2006) – for instance, a

face is recognized based on individual elements and their spatial relation (Freiwald et al.,

2009). Likewise, acoustic object detectors match acoustic features across temporal scales

(Gentner, 2008; Griffiths and Warren, 2004) – song detection neurons in the bird brain

recognize the bird's own song by both syllable identity and sequence (Doupe and Konishi,

1991; Gentner and Margoliash, 2003; Margoliash, 1983). pC2 neurons also bind different

properties of the pulse song to selectively signal the presence of conspecific pulse song –

pC2 is tuned to several features of pulse song like pulse carrier frequency, pulse duration,

inter-pulse interval, and pulse train duration and a match in only one feature is not sufficient

to strongly drive these neurons (Fig. 3, S4A-C). They can therefore be considered acoustic

object detectors.

Matches between behavioral tuning and conspecific song

Behavioral selectivity for species-specific signals is thought to serve species separation. In

FLyTRAP, locomotor tuning of *D. melanogaster* females overlaps with the conspecific song

– females slow to conspecific song (Fig. 2A) and do not change their speed or may even

accelerate for deviant pulse parameters (Fig. S2E). However, the tuning for any single song

feature is not sufficiently narrow to serve as an effective filter for conspecific song. For

instance, females also slow for IPIs produced by a sibling species *D. simulans* (50-65 ms)

(Bennet-Clark and Ewing, 1969). However, *D. simulans* pulses would be rejected based on

a mismatch in other song features – *D. simulans* pulses are too short and of too high

frequency to be accepted by females (Clemens et al., 2017; Riabinina et al., 2011).

Selectivity for multiple song features may thus enable species discrimination with relatively

broad single-feature tuning (Amézquita et al., 2011). In addition, males and females are

exposed to additional non-acoustic cues during courtship that may further sharpen

behavioral tuning. For instance, chemical cues prevent males from courting heterospecific

females (Fan et al., 2013) and likely also contribute to female rejection (Billeter et al., 2009;

Rybak et al., 2002b) – it will be interesting to explore how non-auditory cues (Keleman et al.,

2012; Zhang et al., 2016) modulate locomotor responses to song and whether multi-modal

integration occurs in pC2 neurons or elsewhere. The absence of non-acoustic cues may

also explain the diversity of locomotor responses across strains in the FLyTRAP assay (Fig.

S3). Using a naturalistic courtship assay, previous studies show that these same strains

exhibit similar behaviors – males pattern their song in response to the female behavior and

females change their locomotor speed to the natural courtship song similarly across all

strains (Clemens et al., 2017; 2015; Coen et al., 2016; 2014).

In contrast to pulse song responses, the locomotor and singing responses for sine song

were less sex-specific (Fig. 2E) and the behavioral tuning did not match well the conspecific

song – very low frequencies never produced by males slowed females the most (Fig. 2A, B).

This implies divergent roles for the two song modes and is consistent with previous studies

(Eberl et al., 1997; F. V. Schilcher, 1976) – for instance sine song does not induce male-

male courtship (Yoon et al., 2013). It has been suggested that pulse song may modulate

sine song responses (F. V. Schilcher, 1976) but we did not detect strong serial interactions

between the two song modes (Fig. S2G). Alternatively, responses to sine song may depend

more strongly on the presence of male chemical cues (Billeter et al., 2009; Kurtovic et al.,

2007) that are absent in the FLyTRAP assay. This is consistent with sine song being

produced when the male is near the female (Coen et al., 2014) – that is, when these

chemical cues are particularly strong.

Pathways for detecting sine and pulse

Our behavioral and neuronal results suggest that pulse and sine song are processed in

parallel pathways (Fig. 2E, 3C, F-H) but it is unclear as of yet how and where sounds are

split into different streams. Sine and pulse can be separated based on spectral and temporal

properties (Fig. S5). In fact, the frequency tuning in auditory receptor neurons (JON) and

first-order auditory brain neurons (AMMC) may already be sufficient to separate the lower-

frequency sine (150Hz) from the higher-frequency pulse (>220Hz) (Azevedo and Wilson,

2017; Ishikawa et al., 2017; Kamikouchi et al., 2009; Patella and Wilson, 2018; Yorozu et al.,

2009). Temporal pattern could further discriminate pulse from sine by either suppressing

responses to the sustained sine via adaptation or by tuning temporal integration such that

the brief pulse stimuli fail to drive neuronal spiking. A complete mapping of auditory

pathways and auditory activity throughout the *Drosophila* brain is required to identify where

and how the neural selectivity for the different song modes arises.

Here, we have identified pC2 as one of the pathways driving responses to pulse song – pC2

tuning matches the behavioral tuning for pulse song (Fig. 3G, H), pC2 activation drives sex-

specific responses to song (Fig. 5), and experience-dependent modulation of pC2 tuning

matches the behavioral tuning (Fig. 6). But, our data also indicate that it is not the only

pathway used to detect pulse song – for example, we observed locomotor responses to

pulse song even when pC2 neurons were silenced in females (Fig. 5M). Interestingly,

previous studies have implied pC1 as a pulse song detector (Zhou et al., 2015; 2014). Like

pC2, pC1 exists in males and females (Rideout et al., 2010), and activation drives several

courtship-related behaviors in males – including singing, male-male courtship, or aggression

(Koganezawa et al., 2016; Kohatsu et al., 2011; Pan et al., 2012; Philipsborn et al., 2011;

Zhou et al., 2015) – and also in females (Li et al., 2018; Rezával et al., 2016; Zhou et al.,

2014). All previous studies have relied on imaging activity in the lateral junction (LJ) to show

that pC1 preferentially responds to pulse song (Zhou et al., 2015; 2014). However, we show

here that calcium responses of Dsx+ neurons in the LJ reflect the auditory activity of multiple

Dsx+ cell types – and we detected auditory responses in the somata of pC2, pC1 (only in

females) and pMN2 (only in females) (Fig. 4). Because the number of auditory neurons

within the pC2 cluster is much larger than for pC1 or pMN2 (Fig. 4C), and because tuning in

pC2 somas matches the tuning in the LJ (Fig. 4E-H), we conclude that the LJ activity largely

reflects pC2 responses. Nonetheless, we have not exhaustively assessed the match

between the neuronal responses of female pC1 and pMN2 neurons and behavior. Those

neurons may also be critical for the female's response to pulse song, including behaviors not

investigated here (such as oviposition (Kimura et al., 2015)).

Inputs and outputs of pC2 neurons

How the selectivity of pC2 for pulse song arises is as of yet unclear since systematic studies

of tuning for multiple pulse song features in the early auditory pathway are missing.

However, existing evidence suggests that pC2 may acquire its feature selectivity in a serial

manner – via a cumulative sharpening of tuning for song features at successive stages of

auditory processing (Kamikouchi et al., 2009; Yamada et al., 2018; Zhou et al., 2015; 2014).

Auditory receptor neurons display diverse and specific band-pass tuning for carrier

frequency (Ishikawa et al., 2017; Kamikouchi et al., 2009; Patella and Wilson, 2018; Yorozu

et al., 2009) and first order auditory B1 neurons further sharpen frequency tuning via

resonant conductances (Azevedo and Wilson, 2017). Likewise, peripheral responses are

already weakly tuned for IPI (Clemens et al., 2018; Ishikawa et al., 2017) and this tuning is

further sharpened in downstream neurons (Vaughan et al., 2014; Zhou et al., 2015) through

the interplay of excitation and inhibition (Yamada et al., 2018). This serial sharpening is

similar to how selectivity for pulse song arises in crickets, in which a delay-line and

coincidence detector mechanism produces broad selectivity for pulse duration and pulse

pause which is subsequently sharpened in a downstream neuron (Schöneich et al., 2015).

More direct readouts of the membrane voltage of auditory neurons in the fly brain are

required to determine the biophysical mechanisms that generate song selectivity in pC2.

Similarly, the circuits downstream of pC2 neurons that control the diverse and sex-specific

behaviors reported here remain to be identified. pC2 neurons may connect directly with

descending interneurons (DNs) (Cande et al., 2017; Namiki et al., 2017) that control motor

behaviors. For example, pC2 activation in males drives pulse song production, followed by

sine song production at stimulus offset (Fig. 5A). This behavior resembles that caused by

pIP10 activation (Clemens et al., 2017) – pIP10 is a sex-specific descending neuron

(Philipsborn et al., 2011), but we don't yet know if it directly connects with pC2 neurons. The

fact that song responses are bi-directional – pulse song can induce both slowing and

acceleration within each sex (Fig. 2A, B, S2) – implies that the sex-specificity of motor

control is more than a simple re-routing from accelerating DNs in males to slowing DNs in

females. Notably, song also promotes copulation, but we did not detect a significant effect of

pC2 inactivation on copulation rates (Fig. S6I). This could be because our driver only labeled

1/3 of the auditory pC2 neurons or because pC2 activity does not inform the decision to

mate. That is, there may exist parallel pathways that control song responses on different

timescales: one pathway that accumulates song information over timescales of minutes

(Clemens et al., 2015; Ratcliff et al., 2016) and ultimately controls the mating decision while

another, independent pathway controls behavioral responses to song on sub-second

timescales, such as dynamic adjustments in locomotion and the production of courtship

song.

Modularity facilitates plasticity of behavioral responses to song

Our behavioral data strongly suggest that the sex-specificity of behavior arises after feature

tuning – that is, the shared object detector – pC2 – determines the magnitude of behavioral

responses in both sexes (Fig. 3G, H), and sex-specific aspects, such as the sign and

dynamics of locomotor responses or singing in males, are driven by sex-specific circuits

downstream of pC2 (Fig. 5). This is reminiscent of how sex-specific behaviors are driven to

the male pheromone cVA in flies: shared detector neurons – olfactory receptor neurons and

projection neurons in the antennal lobe – detect cVA in both sexes, and this information is

then routed to sex-specific higher-order neurons in the lateral horn, which are thought to

drive the different behaviors (Datta et al., 2008; Kohl et al., 2013; Ruta et al., 2010). This

modular architecture with object detectors being flexibly routed to different behavioral

outputs is beneficial if these routes are plastic. For instance, here we show that social

experience can shape male responses to song (similar to (Li et al., 2018)), along with tuning

at the level of the object detector neurons (Fig. 6). During mating, males transfer a sex

peptide to females (Yapici et al., 2008) that alters female behavioral responses to song from

slowing to acceleration (Coen et al., 2014) – these effects may be mediated at the level of

the motor circuits downstream of pC2, shifting pulse song responses in females to resemble

those of males. Modularity also facilitates behavioral plasticity on evolutionary time scales

since only one element – the feature detector – needs to change for behavioral tuning in

both sexes to adapt to new songs that evolve during speciation (Capranica et al., 1973;

Kostarakos et al., 2009). The identification of pC2 neurons as the pulse object detectors is

therefore likely to benefit future studies of the evolution of song recognition.

pC2 neurons have a dual sensory and motor role

Strikingly, our results imply a dual sensory and motor role of pC2 neurons: they drive the

production of the sensory object they detect (Fig. 3F, G, 5A-C). This dual role may guide

social interactions and communication via imitation. In *Drosophila melanogaster*, hearing the

song of other males induces a male to court and sing to other females and even males

(Eberl et al., 1997; Yoon et al., 2013). This behavior may have originated because the song

of another male indicates the presence of a female nearby. pC2 is thus reminiscent of

sensorimotor correspondence neurons found in vertebrates (Mooney, 2014; Prather et al.,

2008; Rizzolatti and Fogassi, 2014) – neurons that are active during the production as well

as the observation of a behavior. Such neurons are hypothesized to play a role in learning

(Mooney, 2014) or – as is probably the case in *Drosophila* – imitation and communication

between conspecifics (Rizzolatti and Arbib, 1998). However, pC2 differs crucially from these

instances of "mirror" neurons in that it directly drives the production of the acoustic object it

detects (Fig. 5A-C), while other sensorimotor correspondence neurons are activated by both

sensory and motor-like inputs, but do not drive the object production itself. Because we

recorded pC2 activity in passively listening males, we do not yet know whether pC2 is

activated by sound in an actively singing male. If so, hearing its own song could induce self-

stimulation and form a positive feedback loop to maintain courtship behavior by mediating

persistent behavioral state-changes (Hoopfer et al., 2016). Alternatively, auditory inputs

could be suppressed during singing via a corollary discharge (Poulet and Hedwig, 2003;

Schneider et al., 2014), which would allow pC2 to maintain sensitivity to the song of other

males to coordinate inter-male competition during singing. Additional studies of pC2 activity

in behaving animals are required to fully understand how these pulse song detector neurons

integrate into the acoustic communication behavior.

In summary, we show how the circuits that recognize song to drive diverse and sex-specific

behavioral responses are organized in *Drosophila*: common detector neurons – pC2 –

recognize pulse song in both males and females, and this identically processed information

is then routed to drive multiple sex-specific behaviors. Similar principles may underlie the

production of sex-specific behavioral responses to acoustic communication signals in other
insects, song birds or mammals.
645

Acknowledgements: Isabel D'Allesandro for help with playback behavioral experiments,
Alex Hammons and Nofar Ozeri-Engelhard for help with dissections for immunostaining,
Georgia Guan for help with crosses, immunostaining, and general technical assistance,
Diego Pacheco for help with aligning the volumetric GCaMP scans, Robert Court and Doug
Armstrong (http://www.virtualflybrain.org) for help with brain registration, David Stern, Ben
Arthur and Barry Dickson for discussions during the development of the FLyTRAP assay,
and Kai Feng and Barry Dickson for sharing the design of their playback assay chamber,
Bruce Baker, Stephen Goodwin, Gerry Rubin, Peter Andolfatto for gifts of flies, and Kristin
Scott, Asif Ghazanfar, Tim Buschman, as well as Christa Baker, Frederic Römschied and
other members of the Murthy lab for feedback on the manuscript.
656

657 **Funding:** JC was funded by a Princeton Sloan-Swartz postdoctoral fellowship and by an
658 Emmy-Noether grant from the German Research Foundation (DFG CL 596/1-1). MM was
659 funded by an NIH Director's New Innovator (DP2) award, an NSF CAREER award, and an
660 HHMI Faculty Scholar award.
661

662 **Author contributions:** JC, DD, and MM designed the study; JC primarily built the acoustic
663 playback assay and collected behavioral data, with contributions from DD, and DD primarily
664 collected anatomical and calcium imaging data, with contributions from JC; ST designed and
665 built the two-photon imaging microscope; JC and DD analyzed data; JC and MM wrote the
666 manuscript, with contributions from DD.

References

Amézquita, A., Flechas, S.V., Lima, A.P., Gasser, H., Hödl, W., 2011. Acoustic interference and recognition space within a complex assemblage of dendrobatid frogs. Proc Natl Acad Sci U S A 108, 17058–17063. doi:10.1073/pnas.1104773108

Aranha, M.M., Herrmann, D., Cachitas, H., Neto-Silva, R.M., Dias, S., Vasconcelos, M.L., 2017. apterous Brain Neurons Control Receptivity to Male Courtship in Drosophila Melanogaster Females. Sci Rep 7, 46242. doi:10.1038/srep46242

Arthur, B.J., Sunayama-Morita, T., Coen, P., Murthy, M., Stern, D.L., 2013. Multi-channel acoustic recording and automated analysis of Drosophila courtship songs. BMC Biol 11, 11. doi:10.1186/1741-7007-11-11

Azevedo, A.W., Wilson, R.I., 2017. Active Mechanisms of Vibration Encoding and Frequency Filtering in Central Mechanosensory Neurons. Neuron 1–25. doi:10.1016/j.neuron.2017.09.004

Bennet-Clark, H.C., Ewing, A.W., 1969. Pulse interval as a critical parameter in the courtship song of Drosophila melanogaster. Animal Behaviour 17, 755–759. doi:10.1016/S0003-3472(69)80023-0

Bennet-Clark, H.C., Ewing, A.W., 1967. Stimuli provided by Courtship of Male Drosophila melanogaster. Nature 215, 669–671. doi:10.1038/215669a0

Billeter, J.-C., Atallah, J., Krupp, J.J., Millar, J.G., Levine, J.D., 2009. Specialized cells tag sexual and species identity in Drosophila melanogaster. Nature 461, 987–991. doi:10.1038/nature08495

Billeter, J.-C., Levine, J.D., 2013. Who is he and what is he to you? Recognition in Drosophila melanogaster. Current Opinion in Neurobiology 23, 17–23. doi:10.1016/j.conb.2012.08.009

Bizley, J.K., Cohen, Y.E., 2013. The what, where and how of auditory-object perception. Nature Reviews Neuroscience 14, 693–707. doi:10.1038/nrn3565

Blankers, T., Hennig, R.M., Gray, D.A., 2015. Conservation of multivariate female preference functions and preference mechanisms in three species of trilling field crickets. Journal of Evolutionary Biology n/a–n/a. doi:10.1111/jeb.12599

Brincat, S.L., Connor, C.E., 2004. Underlying principles of visual shape selectivity in posterior inferotemporal cortex. Nature neuroscience 7, 880–886. doi:10.1038/nn1278

Bussell, J.J., Yapici, N., Zhang, S.X., Dickson, B.J., Vosshall, L.B., 2014. Abdominal-B Neurons Control Drosophila Virgin Female Receptivity. Current Biology 24, 1584–1595. doi:10.1016/j.cub.2014.06.011

Cachero, S., Ostrovsky, A.D., Yu, J.Y., Dickson, B.J., Jefferis, G.S.X.E., 2010. Sexual dimorphism in the fly brain. Current biology : CB 20, 1589–1601. doi:10.1016/j.cub.2010.07.045

Cande, J., Berman, G.J., Namiki, S., Qiu, J., Korff, W., Card, G., Shaevitz, J.W., Stern, D.L., 2017. Optogenetic dissection of descending behavioral control in Drosophila. bioRxiv 1–50. doi:10.1101/230128

Capranica, R.R., Frishkopf, L.S., Nevo, E., 1973. Encoding of Geographic Dialects in the Auditory System of the Cricket Frog. Science 182, 1272–1275. doi:10.1126/science.182.4118.1272

Chen, T.-W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R., Kerr, R.A., Orger, M.B., Jayaraman, V., Looger, L.L., Svoboda, K., Kim, D.S., 2013. Ultrasensitive fluorescent proteins for imaging neuronal activity. Nature 499, 295–300. doi:10.1038/nature12354

Clemens, J., Coen, P., Roemschied, F., Pereira, T., Mazumder, D., Pacheco, D., Murthy, M., 2017. Discovery of a new song mode in Drosophila reveals hidden structure in the sensory and neural drivers of behavior. bioRxiv 221044. doi:10.1101/221044

Clemens, J., Girardin, C.C., Coen, P., Guan, X.-J., Dickson, B.J., Murthy, M., 2015. Connecting Neural Codes with Behavior in the Auditory System of Drosophila. Neuron 87, 1332–1343. doi:10.1016/j.neuron.2015.08.014

Clemens, J., Hennig, R.M., 2013. Computational principles underlying the recognition of acoustic signals in insects. Journal of Computational Neuroscience 35, 75–85. doi:10.1007/s10827-013-0441-0

Clemens, J., Ozeri-Engelhard, N., Murthy, M., 2018. Fast intensity adaptation enhances the encoding of sound in Drosophila. Nat Commun 9, 134. doi:10.1038/s41467-017-02453-9

Clyne, J.D., Miesenböck, G., 2008. Sex-specific control and tuning of the pattern generator for courtship song in Drosophila. Cell 133, 354–363. doi:10.1016/j.cell.2008.01.050

Coen, P., Clemens, J., Weinstein, A.J., Pacheco, D.A., Deng, Y., Murthy, M., 2014. Dynamic sensory cues shape song structure in Drosophila. Nature 507, 233–237. doi:10.1038/nature13131

Coen, P., Xie, M., Clemens, J., Murthy, M., 2016. Sensorimotor Transformations Underlying Variability in Song Intensity during Drosophila Courtship. Neuron 89, 629–644. doi:10.1016/j.neuron.2015.12.035

Cook, R.M., 1973. Courtship processing in Drosophila melanogaster. II. An adaptation to selection for receptivity to wingless males. Animal Behaviour 21, 349–358. doi:10.1016/S0003-3472(73)80077-6

Crossley, S.A., Bennet-Clark, H.C., Evert, H.T., 1995. Courtship song components affect male and female Drosophila differently. Animal Behaviour 50, 827–839. doi:10.1016/0003-3472(95)80142-1

Datta, S.R., Vasconcelos, M.L., Ruta, V., Luo, S., Wong, A., Demir, E., Flores, J., Balonze, K., Dickson, B.J., Axel, R., 2008. The Drosophila pheromone cVA activates a sexually dimorphic neural circuit. Nature 452, 473–477. doi:10.1038/nature06808

Dicarlo, J.J., Zoccolan, D., Rust, N.C., 2012. How Does the Brain Solve Visual Object Recognition? Neuron 73, 415–434. doi:10.1016/j.neuron.2012.01.010

Doupe, A.J., Konishi, M., 1991. Song-selective auditory circuits in the vocal control system of the zebra finch. Proceedings of the National Academy of Sciences of the United States of America 88, 11339–11343.

Dulac, C., Wagner, S., 2006. Genetic Analysis of Brain Circuits Underlying Pheromone Signaling. Annu. Rev. Genet. 40, 449–467. doi:10.1146/annurev.genet.39.073003.093937

Eberl, D.F., Duyk, G.M., Perrimon, N., 1997. A genetic screen for mutations that disrupt an auditory response in Drosophila melanogaster. Proceedings of the National Academy of Sciences of the United States of America 94, 14837–14842. doi:10.1146/annurev.neuro.20.1.567

Ellis, L.B., Kessler, S., 1975. Differential posteclosion housing experiences and reproduction in Drosophila. Animal Behaviour 23, 949–952. doi:10.1016/0003-3472(75)90119-0

Fan, P., Manoli, D.S., Ahmed, O.M., Chen, Y., Agarwal, N., Kwong, S., Cai, A.G., Neitz, J., Renslo, A., Baker, B.S., Shah, N.M., 2013. Genetic and Neural Mechanisms that Inhibit Drosophila from Mating with Other Species. Cell 154, 89–102. doi:10.1016/j.cell.2013.06.008

Fortune, E.S., Rodríguez, C., Li, D., Ball, G.F., Coleman, M.J., 2011. Neural mechanisms for the coordination of duet singing in wrens. Science 334, 666–670. doi:10.1126/science.1209867

Freiwald, W.A., Tsao, D.Y., 2010. Functional Compartmentalization and Viewpoint Generalization Within the Macaque Face-Processing System. Science 330, 845–851. doi:10.1126/science.1194908

Freiwald, W.A., Tsao, D.Y., Livingstone, M.S., 2009. A face feature space in the macaque temporal lobe. Nature neuroscience 12, 1187–1196. doi:10.1038/nn.2363

Gentner, T.Q., 2008. Temporal scales of auditory objects underlying birdsong vocal
recognition. The Journal of the Acoustical Society of America 124, 1350–1359.
doi:10.1121/1.2945705
Gentner, T.Q., Margoliash, D., 2003. Neuronal populations and single cells representing
learned auditory objects. Nature 424, 669–674. doi:10.1038/nature01731
Gerhardt, C.H., Huber, F., 2002. Acoustic Communication in Insects and Anurans. University
Of Chicago Press.
Griffiths, T.D., Warren, J.D., 2004. What is an auditory object? Nature Reviews
Neuroscience 5, 887–892. doi:10.1038/nrn1538
Haga, S., Hattori, T., Sato, T., Sato, K., Matsuda, S., Kobayakawa, R., Sakano, H.,
Yoshihara, Y., Kikusui, T., Touhara, K., 2010. The male mouse pheromone ESP1
enhances female sexual receptive behaviour through a specific vomeronasal receptor.
Nature 466, 118–122. doi:10.1038/nature09142
Helversen, von, D., Helversen, von, O., 1997. Recognition of sex in the acoustic
communication of the grasshopper Chorthippus biguttulus (Orthoptera, Acrididae).
Journal of Comparative Physiology A: Neuroethology, Sensory, Neural, and Behavioral
Physiology 180, 373–386. doi:10.1007/s003590050056
Hennig, R.M., Blankers, T., Gray, D.A., 2016. Divergence in male cricket song and female
preference functions in three allopatric sister species. Journal of Comparative
Physiology A: Neuroethology, Sensory, Neural, and Behavioral Physiology 202, 347–
360. doi:10.1007/s00359-016-1083-2
Hennig, R.M., Heller, K.-G., Clemens, J., 2014. Time and timing in the acoustic recognition
system of crickets. Frontiers in Physiology 5. doi:10.3389/fphys.2014.00286
Hoopfer, E.D., Jung, Y., Inagaki, H.K., Rubin, G.M., Anderson, D.J., Ramaswami, M., 2016.
P1 interneurons promote a persistent internal state that enhances inter-male aggression
in Drosophila. eLife 4, e11346. doi:10.7554/eLife.11346
Ishii, K.K., Osakada, T., Mori, H., Miyasaka, N., Yoshihara, Y., Miyamichi, K., Touhara, K.,
2017. A Labeled-Line Neural Circuit for Pheromone-Mediated Sexual Behaviors in Mice.
Neuron 95, 123–137.e8. doi:10.1016/j.neuron.2017.05.038
Ishikawa, Y., Okamoto, N., Nakamura, M., Kim, H., Kamikouchi, A., 2017. Anatomic and
Physiologic Heterogeneity of Subgroup-A Auditory Sensory Neurons in Fruit Flies. Front.
Neural Circuits 11, 46. doi:10.3389/fncir.2017.00046
Ito, K., Shinomiya, K., Ito, M., Armstrong, J.D., Boyan, G., Hartenstein, V., Harzsch, S.,
Heisenberg, M., Homberg, U., Jenett, A., Keshishian, H., Restifo, L.L., Rössler, W.,
Simpson, J.H., Strausfeld, N.J., Strauss, R., Vosshall, L.B., 2014. A Systematic
Nomenclature for the Insect Brain. Neuron 81, 755–765.
doi:10.1016/j.neuron.2013.12.017
Kamikouchi, A., Inagaki, H.K., Effertz, T., Hendrich, O., Fiala, A., Göpfert, M.C., Ito, K.,
2009. The neural basis of Drosophila gravity-sensing and hearing. Nature 458, 165–171.
doi:10.1038/nature07810
Keleman, K., Vrontou, E., Krüttner, S., Yu, J.Y., Kurtovic-Kozaric, A., Dickson, B.J., 2012.
Dopamine neurons modulate pheromone responses in Drosophila courtship learning.
Nature. doi:10.1038/nature11345
Kelley, D.B., 2003. Sexually Dimorphic Behaviors.
http://dx.doi.org/10.1146/annurev.ne.11.030188.001301 11, 225–251.
Kimura, K.-I., Sato, C., Koganezawa, M., Yamamoto, D., 2015. Drosophila Ovipositor
Extension in Mating Behavior and Egg Deposition Involves Distinct Sets of Brain
Interneurons. PLoS ONE 10, e0126445. doi:10.1371/journal.pone.0126445
Klapoetke, N.C., Murata, Y., Kim, S.S., Pulver, S.R., Birdsey-Benson, A., Cho, Y.K.,
Morimoto, T.K., Chuong, A.S., Carpenter, E.J., Tian, Z., Wang, J., Xie, Y., Yan, Z.,
Zhang, Y., Chow, B.Y., Surek, B., Melkonian, M., Jayaraman, V., Constantine-Paton, M.,

Wong, G.K.-S., Boyden, E.S., 2014. Independent optical excitation of distinct neural
populations. Nat. Methods 11, 338–346. doi:10.1038/nmeth.2836
Koganezawa, M., Kimura, K.-I., Yamamoto, D., 2016. The Neural Circuitry that Functions as
a Switch for Courtship versus Aggression in Drosophila Males. Current Biology 26,
1395–1403. doi:10.1016/j.cub.2016.04.017
Kohatsu, S., Koganezawa, M., Yamamoto, D., 2011. Female contact activates male-specific
interneurons that trigger stereotypic courtship behavior in Drosophila. Neuron 69, 498–
508. doi:10.1016/j.neuron.2010.12.017
Kohatsu, S., Yamamoto, D., 2015. Visually induced initiation of Drosophila innate courtship-
like following pursuit is mediated by central excitatory state. Nat Commun 6, 6457.
doi:10.1038/ncomms7457
Kohl, J., Ostrovsky, A.D., Frechter, S., Jefferis, G.S.X.E., 2013. A Bidirectional Circuit Switch
Reroutes Pheromone Signals in Male and Female Brains. Cell 155, 1610–1623.
doi:10.1016/j.cell.2013.11.025
Konishi, M., 1985. Birdsong: From Behavior to Neuron. Annu. Rev. Neurosci. 8, 125–170.
doi:10.1146/annurev.neuro.8.1.125
Kostarakos, K., Hennig, M.R., Römer, H., 2009. Two matched filters and the evolution of
mating signals in four species of cricket. Frontiers in Zoology 6, 22. doi:10.1186/1742-
9994-6-22
Kurtovic, A., Widmer, A., Dickson, B.J., 2007. A single class of olfactory neurons mediates
behavioural responses to a Drosophila sex pheromone. Nature 446, 542–546.
doi:10.1038/nature05672
Lehnert, B.P., Baker, A.E., Gaudry, Q., Chiang, A.-S., Wilson, R.I., 2013. Distinct Roles of
TRP Channels in Auditory Transduction and Amplification in Drosophila. Neuron 77,
115–128. doi:10.1016/j.neuron.2012.11.030
Li, X., Ishimoto, H., Kamikouchi, A., 2018. Auditory experience controls the maturation of
song discrimination and sexual response in Drosophila. eLife 7, e34348.
doi:10.7554/eLife.34348
Margoliash, D., 1983. Acoustic parameters underlying the responses of song-specific
neurons in the white-crowned sparrow. J. Neurosci. 3, 1039–1057.
Marlin, B.J., Mitre, M., D'Amour, J.A., Chao, M.V., Froemke, R.C., 2015. Oxytocin enables
maternal behaviour by balancing cortical inhibition. Nature 520, 499–504.
doi:10.1038/nature14402
Mooney, R., 2014. Auditory–vocal mirroring in songbirds. Philosophical Transactions of the
Royal Society B: Biological Sciences 369, 20130179–399. doi:10.1098/rstb.2013.0179
Namiki, S., Dickinson, M.H., Wong, A.M., Korff, W., Card, G.M., 2017. The functional
organization of descending sensory-motor pathways in Drosophila. bioRxiv 231696.
doi:10.1101/231696
Nern, A., Pfeiffer, B.D., Rubin, G.M., 2015. Optimized tools for multicolor stochastic labeling
reveal diverse stereotyped cell arrangements in the fly visual system. Proc Natl Acad Sci
U S A 112, E2967–76. doi:10.1073/pnas.1506763112
Pan, Y., Meissner, G.W., Baker, B.S., 2012. Joint control of Drosophila male courtship
behavior by motion cues and activation of male-specific P1 neurons. Proc Natl Acad Sci
U S A 109, 10065–10070. doi:10.1073/pnas.1207107109
Patella, P., Wilson, R.I., 2018. Functional Maps of Mechanosensory Features in the
Drosophila Brain. Current Biology 0. doi:10.1016/j.cub.2018.02.074
Philipsborn, von, A.C., Liu, T., Yu, J.Y., Masser, C., Bidaye, S.S., Dickson, B.J., 2011.
Neuronal control of Drosophila courtship song. Neuron 69, 509–522.
doi:10.1016/j.neuron.2011.01.011

Poulet, J.F.A., Hedwig, B., 2003. Corollary discharge inhibition of ascending auditory
neurons in the stridulating cricket. The Journal of neuroscience : the official journal of
the Society for Neuroscience 23, 4717–4725.

Prather, J.F., Peters, S., Nowicki, S., Mooney, R., 2008. Precise auditory–vocal mirroring in
neurons for learned vocal communication. Nature 451, 305–310.
doi:10.1038/nature06492

Quiroga, R.Q., 2012. Concept cells: the building blocks of declarative memory functions.
Nature Reviews Neuroscience 13, 587–597. doi:10.1038/nrn3251

Ratcliff, R., Smith, P.L., Brown, S.D., McKoon, G., 2016. Diffusion Decision Model: Current
Issues and History. Trends Cogn. Sci. (Regul. Ed.) 20, 260–281.
doi:10.1016/j.tics.2016.01.007

Remedios, R., Kennedy, A., Zelikowsky, M., Grewe, B.F., Schnitzer, M.J., Anderson, D.J.,
2017. Social behaviour shapes hypothalamic neural ensemble representations of
conspecific sex. Nature 550, 388–392. doi:10.1038/nature23885

Rezával, C., Pattnaik, S., Pavlou, H.J., Nojima, T., Brüggemeier, B., D'Souza, L.A.D.,
Dweck, H.K.M., Goodwin, S.F., 2016. Activation of Latent Courtship Circuitry in the
Brain of Drosophila Females Induces Male-like Behaviors. Current Biology 26, 2508–
2515. doi:10.1016/j.cub.2016.07.021

Riabinina, O., Dai, M., Duke, T., Albert, J.T., 2011. Active process mediates species-specific
tuning of Drosophila ears. Current biology : CB 21, 658–664.
doi:10.1016/j.cub.2011.03.001

Rideout, E.J., Dornan, A.J., Neville, M.C., Eadie, S., Goodwin, S.F., 2010. Control of sexual
differentiation and behavior by the doublesex gene in Drosophila melanogaster. Nature
neuroscience 13, 458–466. doi:10.1038/nn.2515

Rizzolatti, G., Arbib, M.A., 1998. Language within our grasp. Trends in Neurosciences 21,
188–194. doi:10.1016/S0166-2236(98)01260-0

Rizzolatti, G., Fogassi, L., 2014. The mirror mechanism: recent findings and perspectives.
Philosophical Transactions of the Royal Society B: Biological Sciences 369, 20130420–
20130420. doi:10.1098/rstb.2013.0420

Robinett, C.C., Vaughan, A.G., Knapp, J.M., Baker, B.S., 2010. Sex and the single cell. II.
There is a time and place for sex. PLoS Biology 8, e1000365.
doi:10.1371/journal.pbio.1000365.g009

Ronacher, B., Ronacher, B., Hennig, R.M., Clemens, J., 2014. Computational principles
underlying recognition of acoustic signals in grasshoppers and crickets. J Comp Physiol
A 201, 61–71. doi:10.1007/s00359-014-0946-7

Rosenthal, G.G., Ryan, M.J., 2011. Conflicting preferences within females: sexual selection
versus species recognition. Biol. Lett. 7, 525–527. doi:10.1098/rsbl.2011.0027

Ruta, V., Datta, S.R., Vasconcelos, M.L., Freeland, J., Looger, L.L., Axel, R., 2010. A
dimorphic pheromone circuit in Drosophila from sensory input to descending output.
Nature 468, 686–690. doi:10.1038/nature09554

Ryan, M.J., Cummings, M.E., 2013. Perceptual Biases and Mate Choice. Annu. Rev. Ecol.
Evol. Syst. 44, 437–459. doi:10.1146/annurev-ecolsys-110512-135901

Ryan, M.J., Phelps, S.M., Rand, A.S., 2001. How evolutionary history shapes recognition
mechanisms. Trends Cogn. Sci. (Regul. Ed.) 5, 143–148.

Rybak, F., Aubin, T., Moulin, B., Jallon, J.-M., 2002a. Acoustic communication in Drosophila
melanogaster courtship: Are pulse-and sine-song frequencies important for courtship
success? Canadian journal of zoology 80, 987–996.

Rybak, F., Sureau, G., Aubin, T., 2002b. Functional coupling of acoustic and chemical
signals in the courtship behaviour of the male Drosophila melanogaster. Proc. R. Soc. B
269, 695–701. doi:10.1098/rspb.2001.1919

Schilcher, F.V., 1976. The function of pulse song and sine song in the courtship of
Drosophila melanogaster. Animal Behaviour 24, 622–625.
Schilcher, Von, F., 1976. The role of auditory stimuli in the courtship of Drosophila
melanogaster. Animal Behaviour 24, 18–26. doi:10.1016/S0003-3472(76)80095-4
Schneider, D.M., Nelson, A., Mooney, R., 2014. A synaptic and circuit basis for corollary
discharge in the auditory cortex. Nature. doi:10.1038/nature13724
Schöneich, S., Kostarakos, K., Hedwig, B., 2015. An auditory feature detection circuit for
sound pattern recognition. Science Advances 1, e1500325–e1500325.
doi:10.1126/sciadv.1500325
Stowers, L., Logan, D.W., 2010. Sexual dimorphism in olfactory signaling. Current Opinion in
Neurobiology 20, 770–775. doi:10.1016/j.conb.2010.08.015
Sweeney, S.T., Broadie, K., Keane, J., Niemann, H., O'Kane, C.J., 1995. Targeted
expression of tetanus toxin light chain in Drosophila specifically eliminates synaptic
transmission and causes behavioral defects. Neuron 14, 341–351. doi:10.1016/0896-
6273(95)90290-2
Talyn, B.C., Dowse, H.B., 2004. The role of courtship song in sexual selection and species
recognition by female Drosophila melanogaster. Animal Behaviour 68, 1165–1180.
doi:10.1016/j.anbehav.2003.11.023
Tompkins, L., Gross, A.C., Hall, J.C., Gailey, D.A., Siegel, R.W., 1982. The role of female
movement in the sexual behavior ofDrosophila melanogaster. Behav Genet 12, 295–
307. doi:10.1007/BF01067849
Tsao, D.Y., Freiwald, W.A., Tootell, R.B.H., Livingstone, M.S., 2006. A cortical region
consisting entirely of face-selective cells. Science (New York, N.Y.) 311, 670–674.
doi:10.1126/science.1119983
Tsao, D.Y., Livingstone, M.S., 2008. Mechanisms of Face Perception. Annu. Rev. Neurosci.
31, 411–437. doi:10.1146/annurev.neuro.30.051606.094238
Vaughan, A.G., Zhou, C., Manoli, D.S., Baker, B.S., 2014. Neural Pathways for the
Detection and Discrimination of Conspecific Song in D. melanogaster. Current Biology.
doi:10.1016/j.cub.2014.03.048
Wang, L., Anderson, D.J., 2010. Identification of an aggression-promoting pheromone and
its receptor neurons in Drosophila. Nature 463, 227–231. doi:10.1038/nature08678
Yamada, D., Ishimoto, H., Li, X., Kohashi, T., Ishikawa, Y., Kamikouchi, A., 2018.
GABAergic Local Interneurons Shape Female Fruit Fly Response to Mating Songs. J.
Neurosci. 38, 4329–4347. doi:10.1523/JNEUROSCI.3644-17.2018
Yamamoto, D., Yamamoto, D., Koganezawa, M., Koganezawa, M., 2013. Genes and circuits
of courtship behaviour in Drosophila males. Nature Reviews Neuroscience 14, 681–692.
doi:10.1038/nrn3567
Yang, C.F., Shah, N.M., 2014. Representing Sex in the Brain, One Module at a Time.
Neuron 82, 261–278. doi:10.1016/j.neuron.2014.03.029
Yapici, N., Kim, Y.J., Ribeiro, C., Dickson, B.J., 2008. A receptor that mediates the post-
mating switch in Drosophila reproductive behaviour. Nature.
Yoon, J., Matsuo, E., Yamada, D., Mizuno, H., Morimoto, T., Miyakawa, H., Kinoshita, S.,
Ishimoto, H., Kamikouchi, A., 2013. Selectivity and Plasticity in a Sound-Evoked Male-
Male Interaction in Drosophila. PLoS ONE 8, e74289. doi:10.1371/journal.pone.0074289
Yorozu, S., Wong, A., Fischer, B.J., Dankert, H., Kernan, M.J., Kamikouchi, A., Ito, K.,
Anderson, D.J., 2009. Distinct sensory representations of wind and near-field sound in
the Drosophila brain. Nature 458, 201–205. doi:10.1038/nature07843
Yu, J.Y., Kanai, M.I., Demir, E., Jefferis, G.S.X.E., Dickson, B.J., 2010. Cellular organization
of the neural circuit that drives Drosophila courtship behavior. Current biology : CB 20,
1602–1614. doi:10.1016/j.cub.2010.08.025

Zhang, S.X., Rogulja, D., Crickmore, M.A., 2016. Dopaminergic Circuitry Underlying Mating
Drive. Neuron 91, 168–181. doi:10.1016/j.neuron.2016.05.020
Zhou, C., Franconville, R., Vaughan, A.G., Robinett, C.C., Jayaraman, V., Baker, B.S., 2015.
Central neural circuitry mediating courtship song perception in male Drosophila. eLife 4,
11. doi:10.7554/eLife.08477
Zhou, C., Pan, Y., Robinett, C.C., Meissner, G.W., Baker, B.S., 2014. Central Brain Neurons
Expressing doublesex Regulate Female Receptivity in Drosophila. Neuron 83, 149–163.
doi:10.1016/j.neuron.2014.05.038
977

Methods

Flies

The following fly lines were used in our study:

Genotype	Figures	Source/comment
D. melanogaster NM91	1, 2, 3G,H,I, 4H, S1, S2, S3A, S6B,C	gift from Peter Andolfatto
8 *D. melanogaster* strains CM07, CarM03, N30, NM91, TZ58, ZH23, ZW109, and Canton S (lab stock)	S3B	Canton S is a lab stock, the 7 other strains are a gift from Peter Andolfatto
UAS-20X-GCaMP6m,UAS-tdTomato;dsx-Gal4 (Chen et al., 2013; Rideout et al., 2010)	3B-I 4B-D, F-H S3A , S4,S5A-C, S6, S7	dsx-Gal4 is a gift from Stephan Goodwin
UAS-eGFPX2;dsx-Gal4 (Rideout et al., 2010)	3A	
UAS>STOP>CsChrimson.mVenus/LexAop-flp; dsx-LexA, 8LexAop2-flp/R42B01-Gal4 (Klapoetke et al., 2014; Zhou et al., 2015; 2014)	5A-L S6A-E	R42B01-Gal4 and dsx-LexA are gifts from Bruce Baker. Drives expression of csChrimson in pC2 neurons and in a few pCd1/pCd2 neurons
UAS-GCaMP6m, UAS-TdTom/+;R42B01-Gal4/+ (Chen et al., 2013; Zhou et al., 2015; 2014)	S6A, right	R42B01-Gal4 is a gift from Bruce Baker
UAS>STOP>csChrimson/LexAop-flp; dsx-LexA, 8LexAop-Flp/R41A01-Gal4 (Klapoetke et al., 2014; Zhou et al., 2014)	S6D	Used to control for pCd1 neurons in the R42B01-Dsx intersection
UAS>STOP>TNT/LexAop-flp; dsx-LexA/R42B01-Gal4 (Sweeney et al., 1995; Zhou et al., 2015; 2014)	5M-O S6F-H	Inhibit synaptic output of pC2 neurons in females during courtship
+/LexAop-flp; dsx-LexA/R42B01-Gal4 (Zhou et al., 2015; 2014)	5M-O S6F-H	Control for pC2 TNT females
R71G01.AD/UAS-myrGFP;dsx.DBD/+ (Pan et al., 2012)	4A (pC1)	R71G01.AD is a gift from Gerald Rubin, dsx.DBD is a gift from Stephen Goodwin
R57G10-flpG5/+; dsx-Gal4/10UAS>STOP>HA, 10UAS>STOP>V5,10UAS>STOP>FLAG (Nern et al., 2015; Rideout et al., 2010)	4A (pMN2), 4E, S5E	Bloomington #64088 crossed with ;;dsx-Gal4
R57G10-flpl/+; dsx-Gal4/10UAS>STOP>HA, 10UAS>STOP>V5,10UAS>STOP>FLAG (Nern et al., 2015; Rideout et al., 2010)	4A (pC2l), S5D	Bloomington #64087 crossed with ;;dsx-Gal4

FLyTRAP

Fly behavior was recorded with PointGrey cameras (FL3-U3-13Y3M-C or FL3-U3-13E4C-C). Grey color frames with a resolution of 1280x960 pixels were acquired at 30 frames per second using custom written software in python and saved as compressed videos. Sound representation was controlled using custom software written in Matlab. The sound stimuli were converted to an analog voltage signal using a National Instruments DAQ card (PCIe-6343). The signal was then amplified by a Samson s-amp headphone amp and used to drive a speaker (HiVi F6 6-1/2" Bass/Midrange). Sound intensity was calibrated as in (Clemens et al., 2015) by converting the voltage of a calibrated microphone (placed where the fly chambers would be during an experiment) to sound intensity and adjusting the sound amplification to match the target intensity. Sound and video where synchronized by placing into the camera's field-of-view a 650nm LED whose brightness was controlled using a copy of the sound signal. The chamber consisted of an array of 12 small arenas (7 by 46 mm, made from red plastic) that was placed in front of the loudspeaker (Movie S1). The arena floor consisted of plastic mesh to let sound into the chamber and the top was covered with a thin, translucent plastic sheet. Flies were illuminated using a white LED back light from below and a desk lamp from above.

Playback experiments

Virgin male and female flies were isolated within 6 hours of eclosion and aged for 3-7 days prior to the experiments. Flies were raised at low density on a 12:12 dark:light cycle, at 25°C and 60% humidity. Flies were introduced gently into the chamber using an aspirator. Recordings were performed at 25°C and timed to start within 60 minutes of the incubator lights switching on to catch the morning activity peak. Stimulus playback was block-randomized to ensure that all stimuli within a set occur at the same overall rate throughout the stimulus. The stimulus set (e.g. five pulse trains with different IPIs, see Supplemental Table 1 for a list of all stimulus sets) was repeated for the duration of the experiment (2 hours). Stimuli were interleaved by 60 seconds of silence to reduce crosstalk between responses to subsequent stimulus presentations.

Stimulus design

Sound was generated at a sampling frequency of 10 kHz using custom Matlab scripts. Sine song stimuli were created as pure tones of the specified frequency and intensity (typically 5mm/s). Pulse song was generated by arranging Gabor wavelets in trains interleaved by a specified pause. The Gabor wavelets were built by modulating the amplitude of a short sinusoidal using a Gaussian: $exp(-t^2/(2\sigma^2)) \, sin(2\pi f * t + \phi)$, where f is the pulse carrier frequency, ϕ is the phase of carrier, and σ is proportional to the pulse duration. The parameters for all stimuli used along with the behavioral responses obtained in FLyTRAP are listed in Supplemental Table 1.

Analysis of FLyTRAP data

Fly positions where tracked using custom-written software. Briefly, the image background was estimated as the median of 500 frames spaced to cover the full video. Foreground pixels (corresponding to the fly body) were identified by thresholding the absolute values of the

difference between each frame and the background estimate. The fly center position was then taken as the median of the position of all foreground pixels in each chamber. The sequence of fly positions across video frames was then converted into a time series using the light onset frames of the synchronization LED (indicating sound onset) as a reference. From the position time series fly speed was calculated and the speed traces where then aligned to stimulus onset for each trial. Base line speed was calculated as the average of the speed over an interval starting 30 seconds and ending 2 seconds before stimulus onset. Test speed was calculated over an interval starting at stimulus onset and ending 2 seconds after stimulus offset. Tuning curves were calculated as the difference between baseline speed and test speed for each trial, averaged over trials for each stimulus and animal. Speed traces were obtained by subtracting the baseline speed from the trace for each trial and averaging over trials for each stimulus and animal. All data (tuning curves, speed traces) are presented as mean +/- s.e.m. over flies.

Manual scoring of wing extension in FLyTRAP

To evaluate the number of flies that extend their wings upon playback of pulse or sine song, we manually scored wing extension in the videos using the VirtualDub software. For pulse song (see Movie S2), we scored 25 stimuli/fly, choosing trials randomly but ensuring that each IPI (16/36/56/76/96 ms) was scored 5 times/fly. To avoid bias, the scorer was blind to the IPI presented to the fly in each trial. A total of 120 male flies and 36 female flies were scored (3000 and 900 single-fly responses total for pulse song). We scored wing extension only when the wing was extended in the first 1/3 second following stimulus onset, and only when the wings where not extended during the 1 second before stimulus onset. For sine song (150Hz carrier frequency), 60 males fly were scored.

Joint tuning for pulse duration and pulse pause

To visualize locomotor (Fig. S2E, F) and calcium (Fig. S4D,E) responses to pulse trains with different combinations of pulse duration and pulse pause we generated smooth surface plots using Matlab's "scatteredInterpolant" function with the interpolation mode set to "natural". The boundaries of the plots were set as follows: Pulse duration of zero corresponds to silence and the speed values were set to 0 since all speed traces are always base line subtracted. A pulse pause of zero corresponds to a continuous oscillation and we set the corresponding speed values to those obtained for a 4 second pure tone with a frequency of 250 Hz.

Measurement of song features from natural song data

The inter-pulse interval (IPI) is given by the interval between the peaks of subsequent pulses in a pulse train. Pulse trains correspond to continuous sequences of pulses with IPIs smaller than 200ms. Measuring the pulse durations from natural song data is non-trivial since pulses vary in their shape and can be embedded in background noise. We quantified pulse duration by 1) calculating the envelope of each pulse using the Hilbert transform, 2) smoothing that envelope using a Gaussian window with a standard deviation of 2 ms, and 3) taking as the pulse duration the full width of the smoothed envelope at 20% of the maximum amplitude of the pulse. Pulse durations for artificial stimuli used in our pulse train were defined to be consistent with this method. Pulse carrier frequency is given by the center of mass of the amplitude spectrum of

each pulse (Clemens et al., 2017). Sine carrier frequency was calculated as the peak frequency of the power spectrum of individual sine tones.

PCA of speed traces
For the PCA of sex-specific responses to sound and optogenetic activation of pC2 (Fig. 5L) we collected male and female speed traces for all IPIs (Fig. 5D, F) and optogenetic activation levels (Fig. 5H, J) into a large matrix. Each speed trace was cut to include only the 12 seconds after sound onset and then normalized to have zero mean and unit variance. The first two principal components explain 84% of the total data variance.

Optogenetic experiments
CsChrimson was expressed in pC2 neurons using the intersection between R42B01-Gal4 and dsx-LexA. 655nm light was emitted from a ring of 6 Tri-Star LEDs (LuxeonStar, SinkPAD-II 20mm Tri-Star Base) in FLyTRAP (Fig. 5D-K). Flies were fed with food that contained all-trans retinal for a minimum of three days post eclosion. Control were raised on regular fly food after eclosion. The LED was on for four seconds with 60 seconds pause between stimuli, similar to the temporal pattern used for auditory stimulus delivery in FLyTRAP (1-5mW/cm^2, 100Hz, duty cycle 0.5). To measure the amount of song driven by pC2 activation in solitary flies, we used a chamber whose floor was tiled with 16 microphones to allow recording of the song (Fig. 5A-C, Movie S7; see (Clemens et al., 2017)). The LED (627nm LEDs, LuxeonStar) was on for four seconds (frequency 25 Hz, duty cycle 0.5) and off for 60 seconds. We tested three different light intensities (1.8, 9, and 13 mW/cm^2) that were presented in 3 blocks of 18 trials. The order of the three blocks (light intensities) was randomized for each fly. Fly song was segmented as described previously (Arthur et al., 2013; Coen et al., 2014).

pC2 inactivation in females during courtship
Tetanus neurotoxin light chain (TNT) (Sweeney et al., 1995). was used to block synaptic transmission in pC2 neurons in females. 3-7 days old virgin males (wild type, NM91) and females (pC2-TNT: UAS>STOP>TNT/LexAop-flp; dsx-LexA/R42B01-Gal4, pC2-control: +/LexAop-flp; dsx-LexA/R42B01-Gal4) were paired in a custom-built chamber, designed to record fly song (~25 mm diameter, tiled with 16 microphones). Flies were allowed to interact for 30 minutes, and the percent of flies copulated as a function of time was scored (Fig. S6H). A monochrome camera (Point Grey, FL3-U3-13Y3M) was used to record the fly behavior at 60 frames per second. Fly position was tracked offline and song was segmented as previously described (Arthur et al., 2013; Coen et al., 2014). We then calculated song statistics (e.g. amount of song or number of pulses per window) and female locomotion (average female speed) in windows of 60s with 30s overlap (Clemens et al., 2015). For the rank correlations between male song features and female speed (Fig. 5M-O), we binned the female speed values into 16 bins with the bin edges chosen such that each bin was populated by an equal amount of samples (see Fig. 5M) and calculated the rank correlation between the binned female speed and the average male song feature per bin. Changes in correlation between control and experimental flies (Fig. 5O) were analyzed using an ANCOVA model with independent slopes

and intercepts. Significance was determined based on the p-value of the interaction term (model's genotype by song-feature) after Bonferroni correction.

Calcium imaging

Imaging experiments were performed on a custom built two-photon laser scanning microscope equipped with 5mm galvanometer mirrors (Cambridge Technology), an electro-optic modulator (M350-80LA-02 KD*P, Conoptics) to control the laser intensity, a piezoelectric focusing device (P-725, Physik Instrumente), a Chameleon Ultra II Ti:Sapphire laser (Coherent) and a water immersion objective (Olympus XLPlan 25X, NA=1.05). The fluorescence signal collected by the objective was reflected by a dichroic mirror (FF685 Dio2, Semrock), filtered using a multiphoton short-pass emission filter (FF01-680/sp-25, Semrock), split by a dichroic mirror (FF555 Dio3, Semrock) into two channels, green (FF02-525/40-25, Semrock) and red (FF01-593/40-25, Semrock), and detected by GaAsP photo-multiplier tubes (H10770PA-40, Hamamatsu). Laser power (measured at the sample plane) was restricted to 15 mW. The microscope was controlled in Matlab using ScanImage 5.1 (Vidrio). Single plane calcium signals (Fig. 3C-I, 4F,G and pMN2 neuron in Fig 4C-E) were scanned at 8.5 Hz (256X256 pixels). Pixel size was ~0.5µmX0.5µm when imaging the lateral junction or pC2l process and ~0.25µmX0.25µm when imaging cell bodies in a single plan (Fig 4G and pMN2 in Fig. 4B-D). For volumetric scanning of cell bodies (Figs 4B-D, S5A), volumes were acquired at 0.5Hz (256*216, 20 planes, voxel size ~ 0.34µm X 0.4µm X 1.5µm), scanning one group of cells at a time (pC1, pC2, pCd).

After surgery (opening of the head capsule to reveal the brain), flies were placed beneath the objective and perfusion saline was continuously delivered directly to the meniscus. Sound playback was controlled using custom written Matlab software (Clemens et al., 2018). The software also stopped and started the calcium imaging via a TTL pulse sent to ScanImage ("external hardware trigger" mode). The sound stimulus was generated at a sampling rate of 10kHz and sent through an amplifier (Crown, D-75A) to a set of head phones (Koss, 'The Plug'). A single ear plug was connected to one side of a plastic tube (outer-inner diameters 1/8''-1/16'') and the other tube tip was positioned 2 mm away from the fly arista. Sound intensity was calibrated by measuring the sound intensity 2 mm away from the tube tip with a pre-calibrated microphone at a range of frequencies (100Hz-800Hz) and the output signal was corrected according to the measured intensities. The pause between stimulus representation was 25 seconds. A stimulus set (26-36 stimuli) was presented to each fly for 3 times in block-randomized order as in the playback experiments. If the response decayed in the middle of a repetition (possibly because of drift in the z-axis), the whole repetition was discarded from the analysis. Typically, two full repetitions per fly were used for analysis.

Regions of interest (ROIs) for calcium response measurements (in the LJ, pC2 process and in single Dsx+ somata) were selected manually based on a z-projection of the tdTomato channel. $\Delta F/F$ of the GCaMP signal was calculated as $(F(t)-F_0)/F_0$, where F_0 is the mean fluorescence in the ROI in the 10 seconds preceding stimulus onset. Integral $\Delta F/F$ (Fig. 3D, F-I) and peak $\Delta F/F$

(Fig. 3F, inset) values were calculated in a window starting at sound stimulus onset and ending 25 seconds after sound stimulus offset. To compensate for differences in overall responsiveness across flies, we normalized ΔF/F values of each fly by dividing the integral or peak ΔF/F by the maximal value (of integral or peak ΔF/F) across all stimuli for that fly. For volumetric scanning (Fig. 4), each time series was first motion corrected using the rigid motion correction algorithm NoRMCorre (Pnevmatikakis and Giovannucci, 2017) using the tdTomato signal as the reference image.

Light microscopy
Flies expressing GFP in Dsx+ neurons (UAS-eGFP2X; dsx-Gal4; Fig. 3A) and flies expressing CsChrimson.mVenus in pC2 neurons (R42B01-Gal4 intersected with dsx-LexA; Fig. S6) were immunostained and scanned in a confocal microscope. 2-4 day old flies were cold-anesthetized on ice, dissected in cold S2 insect medium (Sigma Aldrich, #S0146) and fixed for 30-40 minutes on a rotator at room temperature in 4% PFA in 0.3% PBTS (0.3% Triton in PBSX1), followed by 4x15 minutes washes in 0.3% PBTS and 30 minutes in blocking solution (5% normal goat serum in 0.3%PBTS). Brains were incubated over two nights at 4°C with primary antibody, washed with 0.3%PBT and incubated for two more nights at 4°C in secondary antibody, followed by washing (4x15 minutes in 0.3%PBTS and 4x20 minutes in PBS), and mounting with Vestashield for 2-7 days before imaging. Antibodies were diluted in blocking solution at the following concentrations: rabbit anti-GFP (Invitrogen #1828014; used against GFP and mVenus) 1:1000, mouse anti-Bruchpilot (nc82, DSHB AB2314866) 1:20, goat anti-rabbit Alexa Flour 488 (Invitrogen #1853312) 1:200, goat anti-mouse Alexa Flour 633 (Invitrogen #1906490) 1:200.

Stochastic labeling of Dsx+ neurons in the female brain (Fig 4A, E) was done using multi-color-flip-out (MCFO, (Nern et al., 2015)) with three different epitope tags (HA,V5,FLAG). We followed the JFRC FlyLight Protocol 'IHC-MCFO' (https://www.janelia.org/project-team/flylight/protocols) for the preparation of brains. Flp was induced using R5710C10 promotor-coding sequence fusions of the flpG5 and flpl. Flies were 4-7 days old when dissected. Flies were stored at 25˚C. Confocal stacks were acquired with a white light laser confocal microscope (Leica TCS SP8 X) and a Leica objective (HC PL APO 20x/0.75 CS2). A high-resolution scan of a pC2 cell (Fig 4E) was performed with an oil immersion Leica objective (HC PL APO 63x/1.40 Oil CS2, fig 4E). Images were registered to the Janelia brain template (JFRC2) (Jenett et al., 2012) using vfbaligner (http://vfbaligner.inf.ed.ac.uk), which internally uses CMTK for registration (Rohlfing and Maurer, 2003). The images of the fly brain in Figs 4A and S5D were deposited by G. Jefferis (Jefferis, 2014). Image processing was performed in FIJI (Schindelin et al., 2012).

References

Arthur, B.J., Sunayama-Morita, T., Coen, P., Murthy, M., Stern, D.L., 2013. Multi-channel acoustic recording and automated analysis of Drosophila courtship songs. BMC Biol 11, 11. doi:10.1186/1741-7007-11-11

Chen, T.-W., Wardill, T.J., Sun, Y., Pulver, S.R., Renninger, S.L., Baohan, A., Schreiter, E.R., Kerr, R.A., Orger, M.B., Jayaraman, V., Looger, L.L., Svoboda, K., Kim, D.S., 2013. Ultrasensitive fluorescent proteins for imaging neuronal activity. Nature 499, 295–300. doi:10.1038/nature12354

Clemens, J., Coen, P., Roemschied, F., Pereira, T., Mazumder, D., Pacheco, D., Murthy, M., 2017. Discovery of a new song mode in Drosophila reveals hidden structure in the sensory and neural drivers of behavior. bioRxiv 221044. doi:10.1101/221044

Clemens, J., Girardin, C.C., Coen, P., Guan, X.-J., Dickson, B.J., Murthy, M., 2015. Connecting Neural Codes with Behavior in the Auditory System of Drosophila. Neuron 87, 1332–1343. doi:10.1016/j.neuron.2015.08.014

Clemens, J., Ozeri-Engelhard, N., Murthy, M., 2018. Fast intensity adaptation enhances the encoding of sound in Drosophila. Nat Commun 9, 134. doi:10.1038/s41467-017-02453-9

Coen, P., Clemens, J., Weinstein, A.J., Pacheco, D.A., Deng, Y., Murthy, M., 2014. Dynamic sensory cues shape song structure in Drosophila. Nature 507, 233–237. doi:10.1038/nature13131

Jefferis, G.S.X.E., 2014. JFRC2 Template Brain. doi:10.5281/zenodo.10567

Jenett, A., Rubin, G.M., Ngo, T.-T.B., Shepherd, D., Murphy, C., Dionne, H., Pfeiffer, B.D., Cavallaro, A., Hall, D., Jeter, J., Iyer, N., Fetter, D., Hausenfluck, J.H., Peng, H., Trautman, E.T., Svirskas, R.R., Myers, E.W., Iwinski, Z.R., Aso, Y., DePasquale, G.M., Enos, A., Hulamm, P., Lam, S.C.B., Li, H.-H., Laverty, T.R., Long, F., Qu, L., Murphy, S.D., Rokicki, K., Safford, T., Shaw, K., Simpson, J.H., Sowell, A., Tae, S., Yu, Y., Zugates, C.T., 2012. A GAL4-Driver Line Resource for Drosophila Neurobiology. Cell Reports 2, 991–1001. doi:10.1016/j.celrep.2012.09.011

Klapoetke, N.C., Murata, Y., Kim, S.S., Pulver, S.R., Birdsey-Benson, A., Cho, Y.K., Morimoto, T.K., Chuong, A.S., Carpenter, E.J., Tian, Z., Wang, J., Xie, Y., Yan, Z., Zhang, Y., Chow, B.Y., Surek, B., Melkonian, M., Jayaraman, V., Constantine-Paton, M., Wong, G.K.-S., Boyden, E.S., 2014. Independent optical excitation of distinct neural populations. Nat. Methods 11, 338–346. doi:10.1038/nmeth.2836

Nern, A., Pfeiffer, B.D., Rubin, G.M., 2015. Optimized tools for multicolor stochastic labeling reveal diverse stereotyped cell arrangements in the fly visual system. Proc Natl Acad Sci U S A 112, E2967–76. doi:10.1073/pnas.1506763112

Pan, Y., Meissner, G.W., Baker, B.S., 2012. Joint control of Drosophila male courtship behavior by motion cues and activation of male-specific P1 neurons. Proc Natl Acad Sci U S A 109, 10065–10070. doi:10.1073/pnas.1207107109

Pnevmatikakis, E.A., Giovannucci, A., 2017. NoRMCorre: An online algorithm for piecewise rigid motion correction of calcium imaging data. Journal of Neuroscience Methods 291, 83–94. doi:10.1016/j.jneumeth.2017.07.031

Rideout, E.J., Dornan, A.J., Neville, M.C., Eadie, S., Goodwin, S.F., 2010. Control of sexual differentiation and behavior by the doublesex gene in Drosophila melanogaster. Nature neuroscience 13, 458–466. doi:10.1038/nn.2515

Rohlfing, T., Maurer, C.R., 2003. Nonrigid image registration in shared-memory multiprocessor environments with application to brains, breasts, and bees. IEEE Trans Inf Technol Biomed 7, 16–25.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri, K., Tomancak, P., Cardona, A., 2012. Fiji: an open-source platform for biological-image analysis. Nat. Methods 9, 676–682. doi:10.1038/nmeth.2019

Sweeney, S.T., Broadie, K., Keane, J., Niemann, H., O'Kane, C.J., 1995. Targeted expression of tetanus toxin light chain in Drosophila specifically eliminates synaptic transmission and causes behavioral defects. Neuron 14, 341–351. doi:10.1016/0896-6273(95)90290-2

Zhou, C., Franconville, R., Vaughan, A.G., Robinett, C.C., Jayaraman, V., Baker, B.S., 2015. Central neural circuitry mediating courtship song perception in male Drosophila. eLife 4, 11. doi:10.7554/eLife.08477

Zhou, C., Pan, Y., Robinett, C.C., Meissner, G.W., Baker, B.S., 2014. Central Brain Neurons Expressing doublesex Regulate Female Receptivity in Drosophila. Neuron 83, 149–163. doi:10.1016/j.neuron.2014.05.038

Figure 1

Figure 1 – FLyTRAP assay for comparing locomotor tuning for courtship song stimuli in males and females.

A *Drosophila melanogaster* produces song in bouts that can consist of two modes: Sine song corresponds to a weakly amplitude modulated oscillation with a species-specific carrier frequency (~150Hz) and pulse song corresponds to trains of Gabor-like wavelets each with a carrier frequency between 220 and 450Hz and a duration between 6 and 12 ms. These pulses are produced at an inter-pulse interval (IPI) of 30-45 ms.

B FLyTRAP consists of behavioral chamber that is placed in front of a speaker through which sound is presented. Fly movement is tracked using a camera. Shown is a single video frame of females in the assay with fly tracks for the preceding 20 seconds overlaid in magenta. See Movie S1.

C Locomotor responses of females (magenta) and males (grey) for pulse trains with different IPIs (see legend). The gray shaded box indicates the duration of the sound stimulus. Red traces at the bottom of the plot show short snippets of the 5 IPI stimuli presented in this experiment. Baseline speed was subtracted before trial averaging.

D Speed tuning curves for different IPIs in females (magenta) and males (grey) are obtained by averaging the speed traces in the six seconds following stimulus onset. The histograms at bottom shows the IPI distribution found in male song (data from 47 males of NM91 wild type strain totaling 82643 pulses).

Lines and shaded areas or error bars in C and D correspond to the mean ± s.e.m. across 112 male and 112 female flies.

See also Figure S1 and Movie S1.

Figure 2

Figure 2 – Responses to song playback are sex-specific and tuned for multiple features of pulse and sine song.

A, B Locomotor tuning curves for females (A, magenta) and males (B, grey) for 6 different features of pulse and sine song. Lines and error bars correspond to the mean±s.e.m. across flies (see Table S1 for a description of all stimuli and N flies).

C Distribution of the six different song features tested in A, B in the natural courtship song of *Drosophila melanogaster* males (data from 47 males of NM91 wild type strain totaling 82643 pulses and 51 minutes of sine song from 5269 song bouts). Histograms are normalized to a maximum of 1.0.

D Pictograms (not to scale) illustrating each song feature examined in A-C. Pulse and sine song features are marked red and blue, respectively.

E Changes in speed for males and females for all pulse (red) and sine (blue) stimuli tested (data from A, B, S2, see Table S1). Responses to sine stimuli are strongly and positively correlated between sexes (r=0.86, p=4×10^{-7}). Pulse responses are also strongly but negatively correlated (r=-0.79, p=7×10^{-18}). Blue and red lines correspond to linear fits to the responses to sine and pulse song, respectively.

F Fraction of trials for which male and female flies extended their wings during the playback of pulse song (five different IPIs as in 1C, D) and sine song (150Hz, quantified only for males). Solitary males (grey) frequently extend their wings in response to pulse but not to sine song. Solitary females (magenta) do not extend wings for pulse song. See also Movie S2.

G Fraction of trials that evoke wing extension in males (grey) and females (magenta) as a function of IPI. In males, wing extension and locomotor behavior (Figure 1D) exhibit strikingly similar tuning with a peak at the conspecific IPI. Females almost never extend their wing for any IPI.

See also Figure S2 and Movie S2.

Figure 3

Figure 3: Neuronal tuning in the LJ matches behavioral tuning for pulse stimuli in males and females.

A Anatomy of Dsx+ neurons in the female brain. Max z-projection of a confocal stack of a fly brain in which all Dsx+ are labeled with GFP. 5/8 cell types (pC1, pC2l (yellow), pC2m (blue), pMN1, pMN2) project to the lateral junction (LJ), while 3 cell types (pCd1, pCd2, aDN (movie S11)) do not. Yellow and blue arrows point to the processes that connect pC2l and pC2m to the LJ. See also Fig. S5B,C.

B Grayscale image (see color bar) of calcium responses ($\Delta F/F$) to a pulse train (IPI 36ms) in an ROI centered around the LJ (red) and the pC2l process (yellow) in a female. Shown are snapshots of the recording at three different time points relative to stimulus onset - before (T=-10s), during (T=1.2s), and after (T=20s) the stimulus. Flies express GCaMP6m in all Dsx+ cells. Conspecific-like pulse elicits strong increases in fluorescence in the LJ and the pC2 process.

C LJ responses to sine (blue) and pulses (red) stimuli in females (left) and males (right). Individual dots correspond to integral $\Delta F/F$ responses for individual stimuli averaged over the 3-12 individuals tested for each stimulus. Many pulse stimuli evoke much stronger responses than the most effective sine stimulus (p=8x10^{-11} for females and p=2x10^{-11} for males, two-sided rank sum comparison of sine and pulse responses).

D Comparison of male and female LJ responses to sine (blue) and pulse (red) stimuli. Responses to both song modes are correlated strongly for pulse (r^2=0.74, p=3x10^{-15}) and weakly for sine (r^2=0.31, p=0.002) stimuli. Individual dots correspond to the integral $\Delta F/F$ for individual stimuli averaged across animals. Before averaging, the responses of each animal were normalized to compensate for inter-individual differences in calcium levels (see methods for details).

E Fluorescence traces from the LJ in females (top, magenta) and males (bottom, grey) for pulse trains with three different IPIs (see legend, average over 6 individuals for each sex). In both sexes, the LJ responds most strongly to the conspecific IPI of 36ms (Fig. 1D). Responses are much weaker for shorter (16ms) and longer (76ms) IPIs. Calcium responses in the LJ are smaller in males than in females (compare C). See also Supp. Movie S3, S4.

F Tuning curves of calcium responses in the female (magenta) and the male (gray) LJ for features of pulse and sine song (compare to behavioral tuning in Fig. 2A, B). Lines and error bars correspond to the mean±s.e.m. across flies. Integral $\Delta F/F$ normalized as in D. Insets show peak $\Delta F/F$ tuning curves for pulse train duration. In contrast to the integral $\Delta F/F$, the peak $\Delta F/F$ saturates for long pulse trains, indicating that the LJ stops accumulating calcium for pulse trains not normally produced by males. Apart from this difference, integral $\Delta F/F$ and peak $\Delta F/F$ are similar (see Fig. S4F, G).

G, H Comparison of behavioral and neuronal tuning in males (G) and females (H). Dots correspond to the average normalized integral $\Delta F/F$ over individuals, lines indicate linear fits. In males (G), behavioral and neuronal responses are strongly and *positively* correlated for pulse (red, r=0.55, p=9x10^{-6}) but not for sine stimuli (blue, r=-0.15, p=0.06). In females (H), behavioral and neuronal responses are strongly and *negatively* correlated for pulse (red, r=-0.62, p=2x10^{-7}) but not for sine stimuli (blue, r=0.27, p=0.30). All $\Delta F/F$ values are from flies expressing GCamp6m under the control of Dsx-Gal4. See also Figure S3 and S4, and Movie S3 and S4.

Figure 4

Figure 4 - pC2 neurons are pulse object detectors common to both sexes.
A Individual Dsx+ neuron types (black) with somas in the female central brain in which we detected calcium responses for pulse or sine song, registered to a common template brain (gray) (see Methods for details). Of the 8 Dsx+ cell types in the central brain, pC2l, pC2m, the single female-only neuron pMN2 and a small number of pC1 neurons (and only in some individuals) respond to courtship sounds. The lateral junction (LJ) is marked in magenta and somata are marked with golden arrow heads. See also Supp. Movie S10, S11.
B Example somatic fluorescence traces from single somata of the pC1, pC2, and pMN2 cells in response to pulse trains (IPI=36ms, single trial responses). Fluorescence trace from the LJ (magenta) shown for comparison. The gray box marks the duration of the sound stimulus. In each panel. Horizontal and vertical scale bars correspond to 6 seconds and 0.25 ΔF/F, respectively. Horizontal black line marks ΔF/F=0.
C Fraction of cells in Dsx+ clusters with detectable somatic calcium responses to pulse or sine song (females, light grey dots; males, dark grey squares). Complete clusters were imaged using volumetric scan for pC1, pC2 and single plane scans for pMN2. We did not distinguish between pC2l/m, since in most flies both groups are spatially intermingled at the level of cell bodies. Note that all flies included showed calcium responses to sound in the LJ, even when we did not detect responses in specific somata.
D Peak somatic ΔF/F for pulse (red) and sine (stimuli). Lines connect responses to pulse (36ms IPI) and sine (150Hz) recorded in the same animal. Note that responses are plotted on a log scale – the average of the ratio between sine and pulse for all cells is ~2.6. 36/38 pC2, 4/5 pC1 and 2/2 pMN2 prefer pulse over sine. See also Supp. Movie S5, S6.
E High resolution confocal scan of a single pC2l neuron (obtained via a stochastic labelling technique, see Methods for details). Only the side ipsilateral to the cell body is shown. The neurites in the lateral junction appear varicose, indicating that they contain pre-synaptic sites.
F Normalized integral ΔF/F values recorded simultaneously in the LJ and the process that connects the LJ with the somata of pC2l (and no other Dsx+ cell type) are highly correlated (r^2=0.99, p=6x10^{-39}). Each point corresponds to an individual stimulus (pulse or sine) averaged over flies (N=10-24 flies per stimulus). The high correlation means that calcium responses in the LJ reflect responses in pC2l neurons.
G Normalized ΔF/F recorded first in the LJ and then in single pC2l somata in the same fly are highly correlated (r^2=0.81, p=2x10^{-14}, N=8 flies per stimulus), demonstrating that calcium responses in the LJ represent the responses of individual pC2l cells, with some variability across individual cells and animals.
H Comparison of calcium responses pC2l process and female speed for the same stimuli. pC2l and behavioral responses are highly correlated for pulse but not for sine stimuli (pulse: r=-0.74, p=3x10^{-7}, sine: r=0.44, p=0.46), just as for the LJ (compare Fig. 3H). The match between neuronal and behavioral tuning for pulse song indicates that pC2l neurons are pulse song detectors. Each point corresponds to an individual stimulus (Δspeed: N~120 flies per stimulus, ΔF/F: N=10-24 flies per stimulus).
All ΔF/F values correspond to the integral fluorescence and are from flies expressing GCamp6m under the control of Dsx-Gal4. See also Figure S5 and Movie S5 and S6.

Figure 5

Figure 5 – pC2 activation generates sex-specific behaviors.
A Song evoked in males by optogenetic activation (627nm LEDs, intensity 13 mW/cm^2) of a driver line that labels pC2l and pCd neurons (R42B01∩Dsx). Top trace shows a song recording marking pulse and sine song in red and blue, respectively. The grey area indicates the duration (4 seconds) of optogenetic activation. Pulse song is evoked during activation while sine song occurs immediately following activation. Bottom plots show pulse rate (red) and sine song probability (blue) averaged over 7 flies (18 stimulation epochs per animals). See also Movie S7.
B, C Average pulse rate (B) and sine song probability (C) evoked in the 6 seconds following LED light onset (LED duration is 4 seconds). Dose-response curves for individuals are shown as thin lines, population averages (mean±s.e.m.) are shown as thick lines with error bars. P-values result from two-sided sign tests and are adjusted for multiple comparisons using Bonferroni's method. Same data as in A. See also Supp. Movie S7, S8.
D-G Changes in speed in FLyTRAP for pulse trains (duration indicated by grey boxes) with different IPIs (see legend) in the R42B01∩Dsx strain. Female behavioral responses (magenta) were weak with multi-phasic dynamics (D, E). Males (grey) slowed for all pulse stimuli tested and responses outlast the stimulus duration (F, G). Data represent averages over 112 female and 113 male flies.
H-K Optogenetic activation of R42B01∩Dsx using csChrimson evokes locomotor responses with sex-specific dynamics. Changes in speed (grey box) and tuning curves were corrected for intrinsic light responses by subtracting the responses of control flies with the same genotype that were not fed retinal (see Fig. S6E). Females slow for weak and speed for strong activation with multi-phasic dynamics as for sound (H, I, compare D). Males decrease their speed and responses outlast the optogenetic stimulus as for sound (J, K, compare F). See also Supp. Movie S9. See S6E for N flies. The x scales of panels I, K are inverted to match the tuning for sound since the direction of the mapping between LED intensity and IPI is arbitrary.
L Principal component (PC) analysis of male and female locomotor speed traces (12s following stimulus LED or sound onset, traces taken from D, F, H, J). Shown are first and second principal component (PC) scores of females (magenta) and males (grey) for sound (squares) and optogenetic stimulation (circles). Female responses to different LED intensities and to different IPIs spread along the second PC while male responses to sound and optogenetic stimulation largely spread along the first PC.
M Locomotor tuning for IPI during natural courtship obtained from single females that were courted by a wild-type NM91 male. pC2l synaptic output in the females was inhibited using TNT using the R42B01∩Dsx driver. Lines and error bars correspond to the mean±s.e.m speed of N females per genotype tested (pC2l TNT– orange, pC2l control – blue, N=48 females for each genotype, see methods for details on how the tuning curves were computed). pC2l control females (blue) do not change their speed with IPI within the range commonly produced by males (rank correlation 0.02, p=0.59, compare Fig. 1D). pC2l TNT females (orange) accelerate for longer IPIs (rank correlation 0.31, p=3x10^{-30}).
N Rank correlation between female speed and different song features during natural courtship (pC2l control – blue, pC2l TNT – orange).
O Difference between the rank correlations for control (blue) and pC2l TNT (orange) flies in N.

pC2l inactivation specifically changes the correlation between female speed and IPI (dark gray, $p=6\times10^{-8}$). All other changes in correlation are much smaller and not significant ($p>0.18$). P-values were obtained by fitting an ANCOVA model (see methods for details) and were corrected for multiple comparisons using the Bonferroni method.

See also Figure S6 and Movie S7.

Figure 6

A

B

C

Figure 6 – Locomotor and pC2 responses are similarly modulated by social experience.
A Changes in speed for pulse trains measured using FLyTRAP with different IPIs in single-housed (solid line) or group-housed (dashed lines) female (left, magenta) and male flies (right, grey). Plots show mean ± s.e.m. across 112 group-housed and 119 single-housed female or male flies. Female IPI tuning is not strongly affected by housing conditions. By contrast, males change their speed more strongly and more selectively when group-housed.
B Calcium responses from the LJ for pulse trains with different IPIs in single-housed (solid line) or group-housed (dashed lines) female (left, magenta) and male flies (right, grey). Plots show mean ± s.e.m. across 5-6 female or male flies in each condition. In females, group housing only weakly suppresses LJ responses for some IPIs. By contrast, male LJ responses are selectively suppressed for long IPIs, which sharpens the IPI tuning.
C Ratio of Calcium responses to 36 and 56 ms IPIs in single-housed or group-housed female (left, magenta) and male flies (right, grey). Individual dots correspond to individual flies, the solid lines connect the population average ratios. P-values were obtained from a two-sided rank sum test.
All ΔF/F values correspond to the integral fluorescence and are from flies expressing GCamp6m under the control of Dsx-Gal4.
See also Figure S7.

Figure S1

Figure S1. Related to Figure 1.

A Locomotor responses of females for pulse trains (36ms IPI stimulus) and sine tones (100Hz tones) of different intensities (color coded, see legend). Intensity is given in mm/s since flies are sensitive to the particle velocity of sound, not sound pressure.

B Speed tuning curves for the traces in A obtained by averaging the speed in the six seconds following stimulus onset. For pulse song, responses are weak for intensities <1mm/s and don't change much beyond that. While sine responses are weak, there is a tendency for speed to change more for louder sine tones. Lines and error bars indicate mean±s.e.m. over ~120 flies.

C Excluding the transient response component only negligibly affects behavioral tuning curves. Shown are IPI tuning curves for males (gray) and females (magenta) obtained by averaging different epochs of the speed traces. The full response (solid lines) corresponds to the average, base-line subtracted speed in the 6 seconds following sound onset. For the sustained response (dashed lines), we start integration of the speed traces not at sound onset but 500ms into the sound. Tuning curves for the full and sustained phases are very similar – the negative transient response component adds only a weak negative bias to the tuning curves.

D Full vs. sustained responses for all stimuli tested in males (gray) and females (magenta). Both measures yield highly correlated responses (r^2=1.00, p=2x10^{-284}). The purely negative transient response component in the full responses adds a negligible negative bias of -0.05mm/s.

Figure S2

Figure S2. Related to Figure 2.
A, B Locomotor response traces for all stimuli in Fig. 2A, B for females (A) and males (B). Stimulus values are color coded (see legends). Gray shaded areas mark the duration of the sound stimulus. Vertical black lines indicate sound onset for stimulus sets with varying durations (pulse train duration, sine tone duration). Lines correspond to the mean over typically ~120 animals (see supp. table 1 for exact values). Error bars are similar to those in 1C and omitted for clarity.
C Pictograms (not to scale) illustrating each song feature examined in A-B. Pulse and sine song features are marked red and blue, respectively.
D Three principal types of tuning are observable when testing behavioral responses for stimuli with different pulse durations and pauses. Black ellipses indicate the range of stimulus parameters that evoke strong behavioral responses, dots correspond to the three (gray) stimuli shown below each tuning field. Horizontal, vertical, and anti-diagonal lines mark stimuli with constant pause, duration and period, respectively. Pulse duration tuning (left) corresponds to high selectivity (narrow tuning) for pulse duration and higher tolerance for pulse pauses. Pause duration tuning (middle), corresponds to high selectivity for pulse pause and high tolerance for pulse pause. Note that for both types, the tuning for pulse duration and pulse pause does not interact – e.g. the preferred pause does not change with pulse duration. Pulse period (a.k.a. inter-pulse interval) tuning (right) corresponds to a preference for stimuli with a constant pulse period, given by the sum of pulse duration and pause. For this type of tuning, pulse duration and pulse pause interact – the preferred pulse pause increases with decreasing pulse duration.
E, F Locomotor responses of females (E) and males (F) for pulse trains with different pulse durations and pulse pauses. Speed values are color coded (see color bar in F). Black dots mark the parameter combinations of the stimuli tested in FLyTRAP. Intermediate values were obtained using interpolation (see methods). Male and female response fields are similar except for the sign – were females tend to slow (blue colors), males tend to accelerate (red colors), and vice versa (compare Fig. 2C). Responses are more selective for pulse duration than for pulse pause and pulse duration tuning is relatively independent of the pulse duration.
G Responses to sequences of sine tones (blue) and pulse trains (red). First and third row correspond to 2s sine followed by 2s pulse for females and males, respectively. For the second and fourth the order is reversed – stimuli start with 2s pulse song and transition into 2s sine song. The top row shows responses to the individual components of the sequence aligned to the onset of the component in the combined stimulus. The bottom row shows responses for both sexes to the compound stimuli (black) and the linear prediction obtained by summing the responses to the individual components (green). The linear prediction matches the measured responses well except at the transition due to transient response to sound onset only present in individual component responses. Lines and shaded areas correspond to the mean ± s.e.m. over 189 female and 217 male flies.

Figure S3

Figure S3. Related to Figure 3.
A Male (left) and female (right) locomotor tuning for pulse trains with different IPIs for NM91 – the wild type strain used in FLyTRAP – and flies expressing GCaMP6m in all Dsx+ neurons used for calcium imaging (see legend). Dsx-GCaMP6m and NM91 flies exhibit similar and species-typical IPI tuning that also matches the tuning found using alternative assays (see text for details).
B IPI tuning in seven additional wild type strains is diverse and not consistent with the species-typical tuning (male left, female right, see legend for strains used). Note, however, that most strains still respond sex-specifically to pulse song sex, similar to NM91 and Dsx-GCaMP6m. These strains produce similar responses to song in a natural courtship assay, suggesting that these strains require sensory cues that are missing in FLyTRAP for expressing their song preference.
Graphs show mean ± s.e.m. over individuals (100-150 flies per strain and sex).

Figure S4

Figure S4. Related to Figure 3.

A Calcium response of the female LJ as a function of the pulse duration for all stimuli in our data set with a duration of 4 seconds. Blue circles correspond to pulse stimuli and red triangles mark sine stimuli, which by definition do not have pauses and can be thought of as very long pulses. Note that besides pulse duration, all stimuli also differ in pulse pause duration and carrier frequency – for instance all sine stimuli (red triangles) differ in carrier frequency (see C). This explains why stimuli with the same pulse duration evoke diverse calcium responses (integral ΔF/F) - they differ in these other stimulus features. The black line connects stimuli that have the optimal pulse carrier frequency (250 Hz) and pulse pause (20ms). To account for differences in ΔF/F across individuals, values were normalized by the maximal ΔF/F for each individual.

B Same calcium response data as in A but now plotted as a function of pulse pause. Sine stimuli correspond to pulse trains with no pauses – they are by definition continuous oscillations. The black line connects stimuli with the optimal pulse duration (12ms) and pulse carrier frequency (250Hz).

C Same calcium response data as in A but now plotted as a function of pulse carrier frequency. Sine stimuli differed in their carrier frequencies. The black line connects stimuli with the optimal pulse duration (12ms) and pulse pause duration (24 ms).

D, E Calcium responses from the female (D) or male (E) LJ for pulse trains with different combinations of pulse pauses and pulse durations. The stimuli constitute a subset of those measured in the behavior. LJ tuning for pulse trains with different pulse pauses and pulse durations recapitulates the behavioral tuning (compare Fig. S2E, F): LJ responses are more selective for pulse duration and the preferred pulse duration does not change with pause duration.

F, G Comparison of peak and integral ΔF/F values from the LJ for all stimuli tested in females (F) and males (G). Pulse and sine song are marked with red and blue, respectively. The black lines correspond to the best linear fit. Both measures of calcium responses are highly correlated (males: $r=0.79$, $p=4\times10^{-26}$, female: $r=0.80$, $p=5\times10^{-39}$).

Figure S5

individuals

▼ ■ ◆ ◀ ● response

▼ ■ ◆ ◀ ● no response

A

pC2 soma position along normalized **lateral-medial** axis

pC2 soma position along normalized **anterior-posterior** axis

pC2 soma position along normalized **dorsal-ventral** axis

B male

LJ

10um

pC1   pC2m   pC2l

C female

LJ

10um

pC1   pC2m&pC2l

D

LJ

pCd2

E

LJ

aDN

pC2l

Figure S5. Related to Figure 4.
A Locations of pC2 somata in the female brain with (pink) or without (grey) auditory responses along the lateral-medial and anterior-posterior (left) or dorsal-ventral (right) axis. Soma positions were normalized to between 0 and 1 for each animal. Individual symbols (see legends) correspond to somata from different animals. Responsive pC2 somata are concentrated in the lateral proportion of the cell cluster (lower half of each plot) and largely correspond to the pC2l subcluster, although we also observed auditory responses in pC2m somate (see Movie S5).
B, C Max z-projection of baseline fluorescence values from a two-photon volumetric scan of a male (B) and female (C) brain expressing GCaMP6m in all Dsx+ neurons. Each of the three clusters (pC1 - white, pC2m – blue, pC2l - yellow) connects to the LJ (red) via unique processes (arrows). pC2l and pC2m are hard to distinguish by soma location in many flies (mostly females) since they are intermingled.
D A single pCd2 neuron labeled using MCFO (black) registered to a template brain (JFRC2, gray). The lateral junction (LJ) is marked in magenta. pCd1 (not shown, {Kimura:2015ff}) and pCd2 do not project to the LJ.
E Max z-projection of a confocal stack in which aDN (green) and pC2l (red) were labelled with different colors using MCFO. pC2l but not aDN projects to the LJ (blue).

Figure S6

Figure S6. Related to Figure 5.

A Expression pattern of CsChrimson.mVenus in the intersection of R42B01 and Dsx (green) for females (left) and males (right). Neuropil is labelled with nc82 (magenta). The intersection labels 11/22 female and 22/36 male pC2l neurons, as well as 5-6 pCd1 and 2 pCd2 in either sex

B Calcium responses from a female of this line for pulse trains (IPI 36ms) in pC2l somata (blue), the pC2l process (yellow) and the LJ (red). We detected auditory response in all 5 pC2l cells visible in the imaging plane, as well as in the LJ and in the pC2l process.

C, D Song evoked by 655nm activation of R42B01∩Dsx neurons (black, N=7 flies) resembles the natural song produced by wild type flies (NM91, N=47 flies) during courtship (red). Shown are distributions of IPIs, pulse carrier frequencies, and sine carrier frequencies (C) as well as average pulse shapes (D).

E Pulse rates and sine song probability upon optogenetic activation of males expressing CsChrimson (red-shifted channel rhodopsin) in R42B01∩Dsx (see A) on food without added retinal (top, solid lines, N=3 flies) or on food with retinal (top, dashed lines, N=7 flies), males expressing CsChrimson in R41A01∩Dsx (labels pCd1) fed retinal food (middle, N=2 flies), and females expressing CsChrimson in R42B01∩Dsx neurons fed retinal food (bottom, N=3 flies). Optogenetic activation of pCd1 in the male (middle) and of pC2 (and pCd) in the female (bottom) does not evoke song, demonstrating that the singing evoked by R42B01∩Dsx activation in males is 1) due to pC2 and not the pCd1 also labeled in this line, and 2) sex-specific. Males expressing CsChrimson in R42B01∩Dsx that were kept on normal food produced some song upon activation, though much less than males fed retinal (see dashed lines). This residual activation likely stems from small amounts of retinal in normal food.

F Speed traces for female (magenta, top) and male (grey, bottom) flies expressing CsChrimson in R42B01∩Dsx. Shown are responses for flies fed retinal (left), normal food (middle) and the difference between the traces of retinal and normally fed flies (right). Colors correspond to different driving voltages of an array of 655nm LEDs. Lines correspond to averages over 70 females and 83 males fed retinal food, and 29 female and 34 males fed normal food.

G Statistics of male song do not change when they court pC2l TNT females (using the R42B01∩Dsx driver). Shown are distributions of 11 song parameters from NM91 males courting pC2 control (blue) or pC2 TNT females (orange).

H Female genotype does not change correlation statistics (correlations between pairs of song parameters) in the song of NM91 males. Shown are all unique pair-wise correlations between the 11 song parameters in F for the song of NM91 males courting pC2l control (x values) or pC2l TNT females (y values).

I Cumulative fraction of copulated pairs of NM91 males courting pC2l control (blue) or pC2l TNT females (orange). There is weak but statistically not significant effect of female genotype on copulation rates (p=0.19, Cox's proportional hazards regression model).

F-H: Data from 48 pC2 control and 48 pC2 TNT pairs.

Figure S7

Figure S7. Related to Figure 6.

A Calcium responses in pC2 neurons (measured via the LJ) for pulse trains with different durations (IPI=36ms) in single- and group-housed (solid and dashed lines) females (left) and males (right). Data from 5-6 flies for each condition (single/group-housed, females/males).
B Same as A, but for sine carrier frequency. Data from 20/5 single/group-housed females and 12/6 single/group-housed males.

Responses to pulse trains with different durations and to sine tones with different carrier frequencies do not change substantially with housing conditions.

Lines and error bars correspond to mean ± s.e.m over flies.

Movie legends

Movie S1 – Female flies (wild type NM91) in FLyTRAP. Track histories are shown as colored trails. Movie speed corresponds to real time.

Movie S2 – Males extend their wings in response to pulse song in FLyTRAP. In this example, 10 out of 12 flies extended their wings after stimulus onset (4 second pulse train, inter-pulse interval = 36ms). Responding flies are marked with red circles. Movie is slowed down 4X. Some flies extend their wings spontaneously (not triggered by sound), see for example fly 5 in this movie.

Movie S3 – Two-photon Calcium imaging of the female lateral junction and pC2l process. GCaMP6m is expressed in all the Dsx+ neurons. The response in the lateral junction (LJ) and pC2l process are highly correlated (see also Fig 4F). Three responses are shown for a single fly – 4 second pulse trains with inter-pulse intervals of 16/36/76ms. The text in the corner appears when the sound stimulus is on. Movie speed corresponds to real time.

Movie S4 - Two-photon Calcium imaging of the male lateral junction and pC2l process (same as in S3).

Movie S5 – Two-photon Calcium imaging (single plane) of female pC2 cell bodies. The responses to a 4 second pulse train and to a 4 second sine tone (carrier frequency – 200Hz) are shown. In most females the pC2l and pC2m cell bodies can not easily be separated into distinct clusters. Movie speed corresponds to real time.

Movie S6 – Two-photon Calcium imaging (single plane) of male pC2 cell bodies (same stimuli as in movie S5). pC2l and pC2m cell bodies can be spatially organized in two distinct clusters. Movie speed corresponds to real time.

Movie S7 – Optogenetic activation of pC2l neurons in an isolated male. Three light intensities are shown (1.8, 9, 13 mW/cm2). Sound is recorded using microphones that tile the chamber floor. Video and audio are synced and played at real time.