RNASeq analysis of *Aedes albopictus* mosquitoes during chikungunya virus infection

Authors and Affiliations:

Ravikiran Vedururu[1,2], Matthew J. Neave[1], Mary Tachedjian[1], Paul R. Gorry[3], Jean-Bernard Duchemin[1#],

Prasad N. Paradkar[1*]

1. CSIRO Health & Biosecurity, Australian Animal Health Laboratory, Geelong, Australia

2. School of Applied Sciences, RMIT University, Bundoora, Australia

3. School of Health and Biomedical Science, RMIT University, Bundoora, Australia

[#]Current Address: Institut Pasteur de la Guyane, Cayenne, French Guiana

*Corresponding author: Prasad.Paradkar@csiro.au

Short Title: Chikungunya virus infection in *Aedes albopictus*

Abstract

Chikungunya virus (CHIKV), preferentially transmitted by *Aedes* mosquitoes, is an emerging pathogen around the world and causes significant morbidity in patients. A single amino acid mutation in the envelope protein of CHIKV has led to shift in vector preference towards *Aedes albopictus*, an invasive mosquito. Previous studies have shown that after infection, mosquitoes mount an antiviral immune response. However, molecular interactions during the course of infection at different tissues and time-points remain largely uncharacterised. Here we performed whole transcriptome analysis on dissected midguts and head/thorax of CHIKV (Indian Ocean strain) infected *Aedes albopictus* to identify differentially expressed genes compared with uninfected controls. For this, RNA was extracted at two days post-infection (D2) from pooled midguts and eight days post-infection (D8) from heads and the anterior 1/3rd of the thorax. We identified 25 and 96 differentially expressed genes from the D2 and D8 samples respectively (p-value <0.05). Custom *de novo* transcriptomes were assembled for the reads that did not align with the reference genome and an additional 225 and 4771 differentially expressed genes from D2 and D8, respectively, were identified. Twenty-two of the identified transcripts, possibly involved in immunity, were validated by qRT-PCR. Interestingly, we also detected changes in viral diversity, as shown by number of mutations in the viral genome, with increase in number of mutations in the midgut compared with mammalian host (Vero cell culture), followed by reduction in the number of mutations in head and thorax at D8, indicating a possible genomic bottleneck. Taken together, these results will help in understanding *Aedes Albopictus* interactions with CHIKV and can be utilised to reduce the impact of this viral infection.

Author Summary

Chikungunya virus has caused several outbreaks around the world in the last decade. Once a

relatively unknown virus, it now causes seasonal infections in tropical and some temperate

regions. This change in epidemiology is attributed to vector switch from *Aedes aegypti* to *Aedes*

albopictus, an invasive pest leading to spread and causing infections in temperate regions.

Although recent research has identified mosquito factors influencing infections, our

understanding of interaction between chikungunya virus and its vector is limited. Using whole

transcriptome sequencing of chikungunya infected mosquitoes, we identified differentially

expressed genes in the midgut and head and thorax, over the course of mosquito infection. We

also detected changes in the viral genome during mosquito infection and a possible genetic

bottleneck event with reduction in viral variants at the head and thorax region of mosquito in the

later stages of infection. These results will lead to improving our understanding of mosquito-virus

interactions with *Aedes albopictus* as a vector and in turn lead to development of novel disease

control strategies.

Introduction

Arboviruses, such as dengue, chikungunya and Zika viruses, cause a significant burden on public health systems worldwide. Transmitted by mosquitoes, these viruses can cause high morbidity and mortality, with dengue alone causing more than 300 million infections per year[1]. First identified and described in 1955 in a report on an outbreak at the border of Tanzania and Mozambique in 1952, chikungunya virus (CHIKV) is an enveloped, positive sense RNA virus belonging to the Alphavirus genus in *Togaviridae* family[2, 3]. CHIKV infection causes high febrile illness, cutaneous exanthema and debilitating and often prolonged arthralgia [4-6].

While *Aedes aegypti* is the traditional vector for CHIKV, since the Reunion Island outbreak in 2005-2006, *Aedes albopictus* is observed to be more involved in viral transmission[7]. A single amino acid change in codon 226 of the E1 gene, that codes for the envelope protein of the virus, has improved fitness of CHIKV in *Aedes albopictus*[8]. This genetic shift has been implicated for most of the recent outbreaks, where despite an absence of the traditional vector *Aedes aegypti*, CHIKV has successfully established infections through *Aedes albopictus*. As an invasive species, *Aedes albopictus* has been expanding its traditional habitat of tropical and sub-tropical regions to much cooler temperate regions. *Aedes albopictus* also survives in favourable microhabitats even in winter and freezing temperatures [9]. These factors have further increased the risk of CHIKV to cause outbreaks in newer areas where mosquito-borne viral diseases are uncommon, such as Northern America and temperate Europe [10, 11].

For a mosquito to become infective, the virus needs to cross 2 critical barrier tissues, the midgut and salivary glands. Infection barriers can be influenced by multiple factors, including viral factors

such as viral glycoproteins, or vector factors such as presence of a viral receptor, host replication

factors and the microbiome composition of the midgut. Inside mosquitoes, after feeding, the

blood meal moves down to the midgut where virus must contact epithelial cells before digestion

of the blood meal and formation of the peritrophic matrix takes place. Following the successful

infection of the midgut, the virus must overcome midgut escape barriers to disseminate to other

tissues such as the haemocoel, head and salivary glands[12]. When an adult female *Aedes*

albopictus mosquito is exposed to CHIKV in the process of blood feeding, the virus infects the

midgut usually in a matter of hours [13-15]. From haemocoel, the virus makes its way to the

salivary glands of the mosquito, which are present in the front 1/3rd thorax region. Once the virus

is detected in the saliva, the mosquito is considered to be infective and a competent vector [16,

17].

Although previous studies [18-23] have used transcriptome-based approaches to identify

mosquito-virus interactions, tissue-specific responses during the course of infection have not

been characterised. Here, using next-generation sequencing, we characterised the whole

transcriptome response at the midgut (MG) (the first barrier site) and head & thorax (HT)

(containing the salivary glands and viral dissemination sites), in *Aedes albopictus* in response to

CHIKV infection. Our results also showed that the virus undergoes mutations in its genome as it

passes through the mosquito and the number of mutations was significantly different at the two

sites and in the course of infection, indicating a possible genomic bottleneck.

Methods

Chikungunya virus

Chikungunya virus isolate 06113879 (Mauritius strain), isolated from a viraemic traveller who returned to Australia in 2006, was obtained from the Victorian Infectious Diseases Reference Laboratory (VIDRL), Melbourne [24]. The isolate was passaged in Vero cells (Cell culture facility, AAHL) four times, followed by once in C6/36 (*Aedes albopictus* larval cell line, obtained from Cell culture facility, AAHL) followed again by Vero cells, which was then used for the experiments. $TCID_{50}$ assay was performed on Vero cells to determine viral titer.

Infections of *Aedes albopictus* and sample processing

All experiments were performed under biosafety level 3 (BSL-3) conditions in the insectary at the Australian Animal Health Laboratory, CSIRO. Insectary conditions were maintained at 27.5°C and 70% in relative humidity with a 12hr light and dark cycle. Female mosquitoes (5-8 days old) were challenged with a chicken blood meal spiked with CHIKV (1 in 100 dilution of stock virus, $TCID_{50}$ 1.5×10^9/ml) through chicken skin membrane feeding (Hemotek membrane feeding system®). Uninfected chicken blood and skin were provided by the Small Animal Facility (Australian Animal Health Laboratory) from chicken bred in the laboratory without any arboviral infection. The procedure was conducted with approval from AAHL Animal Ethics Committee. After one hour, the mosquitoes were anesthetised with CO_2, blood fed females sorted and kept in 200 mL cardboard cup containers at 27.5°C, 70% humidity and 14:10 day:night photoperiod for 2 and 8 days with 10% sugar solution *ad libitum*. For controls, females were fed with blood mixed with media supernatant from uninfected Vero cell culture. Midguts were collected at 2 dpi (D2) and

114 head/thorax were collected at 8 dpi (D8), from infected and control mosquitoes. Dissected

115 tissues were stored in 50 µl of Qiagen RLTplus buffer with 5-10 silica beads (1 mm) at -80°C.

116 RNA extraction and cDNA preparation

117 Bead beating was performed on MP Biomedicals FastPrep -24™ homogeniser, 3 cycles, speed:

118 6.5 m/s, 45 seconds each cycle. RNA was extracted using the RNeasy™ kit (Qiagen Australia) and

119 cDNA was generated by using random hexamers and Superscript-III reverse transcriptase

120 (Thermo Fisher Scientific Inc. Australia) following the manufacturer's protocols.

121 qRT-PCR for infection check and RNASeq data validation

cDNA generated from the RNA extracted from the midguts of D2 pools and the carcasses of D8

pools were tested for CHIKV viral RNA using an in-house designed qRT-PCR (Table B in S1), using

primers specific for the E1 gene (Table A in S1). For RNASeq data validation, adult female

mosquitoes were infected with CHIKV as described before and RNA was extracted from the MGs

and HTs of 5 infected mosquitoes and cDNA was generated by protocols described previously.

cDNA from the corresponding tissue of 5 uninfected mosquitoes was used as controls. qPCR was

performed using gene-specific primers and 18s rRNA specific primers as internal controls (Table

A in S1).

qPCR was performed on an Applied Biosystems Quantstudio™ 6 using the Takara-Clonetech SYBR

Green Master Mix: SYBR Premix Ex Taq II (Tli RNase H Plus). The following cycling conditions were

used with a melt curve at the end. Cycling conditions were: 30 seconds at 95°C, 40 cycles of 5

seconds at 95°C and 30 seconds at 60°C followed by a melt curve. The baseline and Ct values

were calculated automatically using the supplied QuantStudio™ Software and the ΔΔCt values

were calculated using the average ΔCt value of controls and 18srRNA as reference.

RNASeq library construction and sequencing

Libraries for RNASeq were prepared using Nugen's Ovation Universal RNASeq kit, following

manufacturer's specification with a minor modification in the HL-dsDNAse treatment. During first

strand synthesis with DNase treatment, HL-dsDNase from Thermo Fisher Scientific was used in

our library preparation, along with the 10x buffer supplied and their protocol used. The libraries

were pooled and sequenced on a single lane of Hiseq-2500 (Macrogen Inc, South Korea) to

generate 2x100 bp reads. The fastq files were deposited in NCBI's Sequence Read Archive (SRA

Accession ID: SRP140387).

Data analysis-Differential gene expression and Gene ontology

Quality trimming of the raw sequences was performed using Trimmomatic v0.36. The reads were

aligned to the CHIKV reference sequence (GenBank ID: MH229986) to assess the infection status

using Hisat2 v2.0.5[25].

Following removal of *Gallus gallus* reads (due to chicken blood feeding) using SAMtools v1.3.1,

the remaining reads were aligned to the *Aedes albopictus* Foshan strain genome sequence

(AaloF1) from Vectorbase using Hisat2 and the resultant SAM file was sorted and converted into

a BAM file using SAMtools[26, 27].

On Galaxy virtual lab v1.4.6.p5, featureCounts v1.4.6-p5 was used to quantify aligned transcripts

from the sorted BAM files with default parameters for paired end reads, and DESeq2 v2.11.38

was used to obtain differentially expressed genes between the controls and infected samples by

using default parameters on featureCounts output files [28, 29].

Using Trinity v2.3.2, two custom De Novo transcriptomes were built by combining the unaligned

reads from MG (D2) and HT (D8) respectively[30, 31]. This transcriptome was used as a reference

genome and the reads aligned, transcript counts measured and differentially expressed genes

quantified using edgeR. The differentially expressed genes were annotated using BlastX and

BlastN [32, 33].

The gene ontology (GO) IDs of the differentially expressed genes were obtained using the Biomart

tool and topGO analysis was performed to identify the Molecular Functions (MF), Biological

Processes (BP) and Cellular Components (CC) that were either enriched or depleted in the

differentially expressed genes [34, 35]. Using topGO's classic algorithm and based on p-values

generated using Fisher's exact method, differentially expressed genes were grouped based on

their ontologies. Enrichment % was calculated as the ratio of the number of times particular

genes in the pathway were differentially expressed compared to the expected number by chance.

Whole genome sequencing of CHIKV

The Qiagen QIAseq FX Single Cell RNA Library kit was used for Illumina library preparation from

total RNA extracted from CHIKV infected Vero cell culture supernatant using the Qiagen

RNeasy^{TM} kit. The library was sequenced on the Illumina Miniseq, with the mid-output kit (300

cycles) generating 2x150 bp paired-end reads. The resultant fastq files were quality trimmed and

assembled to consensus sequence on CLC Genomics workbench v9.5.2. The sequence was

annotated and submitted to GenBank (MH229986).

Viral mutation calling and genetic diversity

From the infected D2 and D8 RNAseq libraries, the CHIKV reads were extracted using SAMtools. These reads were aligned to the reference sequence generated from the whole genome sequencing of the original cell culture isolate of the virus. Variant calling (SNPs and INDELs >3%) was performed using Varscan2 v2.3.9 on individual libraries as well as all D2 libraries merged together and all D8 libraries merged together [36, 37].

Results

RNASeq

For whole transcriptome analysis of *Aedes albopictus* mosquito tissues, during the course of CHIKV infection, dissected mosquito midguts at 2 dpi (as infection and first barrier site) and head/thorax at 8 dpi (as dissemination site containing salivary gland) were selected. The amount of RNA that could be extracted from a single *Aedes albopictus* midgut (MG) or head/thorax (HT) was below the lower limit of detection by the high-sensitivity kit on the Qubit (Thermo Fisher Scientific Inc. Australia). Hence, the MGs or HTs from 6 mosquitoes were pooled together for RNA extraction to obtain sufficient material. This was also required to avoid using a low-RNA input RNAseq kit, which would likely introduce bias during the PCR amplification stage. The pool size was kept as low as possible to retain information on biological variations.

To determine differentially expressed genes in MGs and HTs of *Aedes albopictus* mosquitoes infected with CHIKV, 5 pools of MGs collected at D2, and 5 pools of HTs collected at D8, from infected mosquitoes were created. For controls, 3 pools of MGs and 3 pools of HTs were used. Initially, to determine infection status of these tissues, qRT-PCR was performed using CHIKV-

specific primers. Based on these results (Table C in S1) and requirements of the Nugen Ovation

universal RNAseq kit, for D2, 2 control and 3 infected MG pools were used; while for D8, 1 control

and 2 infected HT pools were used to prepare libraries.

The sequencing of eight libraries resulted in between 37 million and 170 million reads each. After

quality trimming, reads mapping to the chicken genome were discarded to remove reads

originating from undigested chicken blood. The remaining reads were aligned to the *Aedes*

albopictus reference genome with the average alignment of 62.22% between the 8 libraries. The

results also confirmed that all five infected libraries (3 from D2 and 2 from D8) contained viral

reads, while the 3 control libraries (2 from D2 and 1 from D8) did not (Table 1).

Table 1: **RNASeq NGS data summary**.

	Total reads (millions)	% mapped to Refseq genome	% of CHIKV reads
D2_Infected MG 1	170.65	60.83%	0.01
D2_Infected MG 2	37.13	60.46%	0.02
D2_Infected MG 3	38.02	65.28%	0.10
D2_Control MG1	44.77	61.53%	0.00
D2_Control MG2	44.72	63.15%	0.00
D8_Infected HT1	47.15	63.28%	1.12
D8_Infected HT2	40.60	60.35%	2.09
D8_Control HT1	42.10	62.91%	0.00

Total reads obtained, their alignment percentage to the reference genome and the percentage of reads

aligned to the chikungunya genome.

Differential expression and TopGO analysis

Differentially expressed genes were identified using DESeq2 and edgeR and were plotted as

Volcano plots (Fig 1). The results showed a number of genes to be differentially expressed at the

two time points and tissue sites, using both methods (Table 2). The complete list of genes and

transcripts differentially expressed with p-values of less than 0.05 is provided in supplementary

information S2. The fasta files obtained as output for the custom transcriptome assembly are

provided in supplementary information S3.

Fig 1. Volcano plots from DESeq2 and edgeR for D2 and D8 DGE

Volcano plots from DESeq2 (top panels) and edgeR (bottom panels) of differentially expressed genes from

D2 (left panels) and D8 (right panels) samples. DESeq2 was performed by aligning reads to the *Aedes*

albopictus reference genome; while edgeR analysis was done on reads that did not align to the reference

genome and were aligned to the custom transcriptome.

Table 2: **Differential Gene Expression analysis**:

		No of Differentially expressed genes ($P < 0.05$)		
		Total	Up	Down
D2 (MG)	DESeq2	25	14	11
	edgeR	225	94	131
D8 (HT)	DESeq2	96	51	45
	edgeR	4771	3805	966

Number of genes found to be differentially expressed in D2 midguts and D8 HTs using Deseq2 and edgeR

analysis.

To determine the biological processes and molecular functions of the differentially expressed

genes, gene set enrichment analysis and ontology was performed using TopGO (Fig 2). As

expected, a number of biological and molecular processes were significantly affected during

CHIKV infection.

Fig 2. topGo enrichment comparison in differentially expressed genes

Enrichment analysis of down (left panel) and up (right panel) regulated genes in the midguts (D2) and

heads and thorax (D8) of *Aedes albopictus* mosquitoes in response to CHIKV at 2 and 8 dpi respectively.

Enrichment % is calculated as the ratio of 'significant' (Number of times the gene ontology number is

observed as differentially expressed) to 'expected' (Number of times the gene ontology number is

expected based on observation in control samples) gene numbers.

For D2 MGs, differentially regulated biological processes were metabolic (p-value 0.012) and

serine family amino acid biosynthetic process (p-value 0.026). The differentially regulated

molecular functions, likely of interest from an innate immune response point of view, were

lysozyme activity (down regulated, p-value 0.028), alkaline phosphatase (down regulated, p-

value 0.033) and carboxypeptidase activity (upregulated, p-value 0.042).

For D8 HTs, upregulated molecular functions were RNA (p-value 0.046) and mRNA binding (p-

value 0.049) and molecular functions down regulated were lysozyme activity (p-value 0.0038),

serine-type endopeptidase activity (p-value 0.0058), protein heterodimerisation (p-value

0.00181) and chitin binding (p-value 0.00532). Biological processes down regulated were defence

response (p-value 9.70E-05), proteolysis (p-value 0.0075) and chitin metabolic process (p-value

0.0106). Up regulated biological processes were homophilic cell adhesion via plasma membrane

(p-value 0.0056), UMP biosynthetic process (p-value 0.0245), regulation of transport (p-value

0.0245), spliceosomal complex assembly (p-value 0.0342).

Interestingly, genes involved in lysozymal activity were down regulated in both D2 MGs and D8

HTs, indicating significance of this pathway during infection as well as the dissemination process.

RNASeq data validation on qRT-PCR

We selected 22 differentially expressed genes for validation by qRT-PCR, based on their known

immune functions. This included 8 genes from D2 (1 aligned to reference genome and 7 from

custom transcriptome) and 14 genes from D8 (11 aligned to reference genome and 3 from

custom transcriptome) (see Table 3 for details).

Table 3: List of genes validated by qRT-PCR.

	GeneID	Time point/ site	Gene Annotation
1	AALF020406	D2/MG	Ankyrin repeat domain-containing protein 44
2	TRINITY_DN109663_c0_g2_i1	D2/MG	PREDICTED: mucin-22-like [*Aedes albopictus*]/ PREDICTED: flocculation protein FLO11-like, partial [*Aedes albopictus*]
3	TRINITY_DN110327_c1_g2_i1	D2/MG	PREDICTED: ATP-dependent RNA helicase dbp2 [*Aedes albopictus*]
4	TRINITY_DN46100_c0_g1_i1	D2/MG	PREDICTED: E3 ubiquitin-protein ligase MARCH6 [*Aedes albopictus*]
5	TRINITY_DN102975_c0_g1_i1	D2/MG	PREDICTED: *Aedes albopictus* ARF GTPase-activating protein GIT2
6	TRINITY_DN109582_c0_g2_i2	D2/MG	PREDICTED: protein NPC2 homolog [*Aedes albopictus*]
7	TRINITY_DN110186_c0_g1_i2	D2/MG	PREDICTED: translocon-associated protein subunit delta-like [*Aedes albopictus*]
8	TRINITY_DN110556_c0_g2_i1	D2/MG	PREDICTED: uncharacterized protein LOC109408932 [*Aedes albopictus*]
9	AALF011899	D8/HT	Quaking protein A
10	AALF004300	D8/HT	PREDICTED: PDZ and LIM domain protein 7-like isoform X1
11	AALF008354	D8/HT	Putative ecdysteroid-regulated 16 kDa protein [Source:UniProtKB/TrEMBL;Acc:A0A023EHE1]
12	AALF012634	D8/HT	PREDICTED: fat-like cadherin-related tumor suppressor homolog [*Aedes albopictus*]

13	AALF023547	D8/HT	Peptidylprolyl isomerase [Source:UniProtKB/TrEMBL;Acc:A0A023EQ01]
14	AALF021910	D8/HT	PREDICTED: fasciclin-2-like isoform X1 [*Aedes albopictus*]
15	AALF012324	D8/HT	glycine rich RNA binding protein, putative
16	AALF016505	D8/HT	leucine-rich immune protein (Long)
17	AALF025245	D8/HT	PREDICTED: inhibitor of Bruton tyrosine kinase [*Aedes albopictus*]
18	AALF016704	D8/HT	phosrestin i (arrestin b) (arrestin 2) [Source:Projected from *Aedes aegypti* (AAEL003116) VB Community Annotation]
19	AALF026574	D8/HT	Putative uncharactarised protein containing CCHC zinc finger domain
20	TRINITY_DN129476_c0_g1_i1	D8/HT	PREDICTED: protein no-on-transient A isoform X2 [*Aedes albopictus*]
21	TRINITY_DN131737_c0_g1_i2	D8/HT	PREDICTED: ficolin-3-like [*Aedes albopictus*]
22	TRINITY_DN131885_c0_g2_i2	D8/HT	PREDICTED: PDZ and LIM domain protein Zasp-like isoform X5 [*Aedes albopictus*]

List of genes selected for validation by qRT-PCR from the two time points and tissues and their

annotation based on BlastX and BlastN.

The qRT-PCR results, using gene-specific primers, were compared with the RNAseq data (Table

4). In D2 samples, among the 8 targets chosen for validation, 6 were concordant with one target

being discordant. In D8 samples, 7 out of 14 target genes were concordant. The ΔΔCt of

TRINITY_DN46100_c0_g1_i1 (D2) and AALF008354 (D8), were less than 0.1 and hence not

included in concordance calculation.

Table 4: Comparison of expression fold changes in chosen targets between qRT-PCR and RNASeq.

Target ID	Expression fold change	LogFC
AALF020406	0.5(↓)	-5.56
TRINITY_DN102975_c0_g1_i1	0.76(↓)	-8.7
TRINITY_DN109582_c0_g2_i2	5.35(↑)	6.29
TRINITY_DN109663_c0_g2_i1	0.16(↓)	-8.61
TRINITY_DN110186_c0_g1_i2	1.88(↑)	8.82

TRINITY_DN110327_c1_g2_i1	6.53(↑)	-9.07
TRINITY_DN110556_c0_g2_i1	1.51(↑)	8.84
TRINITY_DN46100_c0_g1_i1	1.03(↑)	-9.04
AALF011899	0.8(↓)	2.54
AALF004300	0.43(↓)	1.53
AALF008354	0.96(↓)	2.07
AALF012634	0.01(↓)	3.51
AALF023547	0.75(↓)	2.24
TRINITY_DN129476_c0_g1_i1	0.01(↓)	-5.08
TRINITY_DN131737_c0_g1_i2	0.07(↓)	-6.42
TRINITY_DN131885_c0_g2_i2	2.9(↑)	10
AALF021910	2.97(↑)	3.55
AALF012324	0(↓)	2.96
AALF016505	3.65(↑)	-1.65
AALF025245	17.54(↑)	2.49
AALF016704	1149.59(↑)	1.51
AALF026574	0.09(↓)	-3.12

263 Results of qRT-PCR on the 22 gene targets selected from RNAseq DGE analysis. Expression fold change

264 for qRT-PCR is calculated as $2^{-\Delta\Delta Ct}$ and the up or down regulation is indicated by arrows, while LogFC

265 was calculated by DESeq2 and edgeR

266 The mean ΔCt values of D2 and D8 targets in infected mosquitoes compared to uninfected

267 controls are shown in Table D in S1.

Chikungunya viral variants in mosquitoes

Previous studies have shown that significant viral genetic bottlenecks occur during arbovirus infection of mosquito vectors [24, 38]. Each bottleneck event may result in significant reduction in viral variant diversity and thus affect the viral variant ultimately transmitted via mosquito saliva. To determine the CHIKV variant diversity in MG and HT samples, we first sequenced the CHIKV isolate 06113879. After passage in Vero cells, whole genome sequencing was performed using the Illumina MiniSeq, resulting in about 8.5 million quality trimmed paired-end reads. The assembly resulted in an 11,929bp long consensus sequence, which perfectly matched a previously sequenced 559bp portion of the E1 gene from this isolate (GenBank ID: EU404186.1).

The viral consensus sequence was most homologous (Identity: 11705/11985 (97.7%), Similarity: 11705/11985 (97.7%), Gaps: 245/11985 (2.0%)) to the CHIKV strain LR2006_OPY1 (GenBank: KT449801.1) and had the E1-A226V mutation, indicating this isolate also belongs to the same lineage as the CHIKV which caused the *La Reunion* outbreak in the Indian ocean in 2006.

Reads from the Vero isolate, as well as from D2 MG and D8 HT, were compared with the consensus sequence to determine the number of mutations in each sample. The number of mutations in the coding region of the D2 midgut samples (50; SNPs: 43 and Indels: 7) were significantly higher compared with Vero (16; SNPs: 9 and Indels: 7), while the number of mutations in D8 HTs were significantly lower compared with other samples (7; SNPs: 4 and Indels: 3). Interestingly, while most of the non-coding mutations were in the AT-rich and variable 3` untranslated region (UTR), the Non-structural protein nsP3 gene was a hotspot for changes, particularly around position 5205 (Fig 3), indicating possible evolutionary pressures.

Fig 3. Variants detected in Vero cell culture isolate, D2 and D8 independent libraries plotted as per their location on the CHIKV genome

SNPs (red circles) and indels (blue circles) detected in the viral extract from Vero cell culture, three MG pools and two HT pools are plotted with reference to their position on the viral genome using VarScan2. The variants were called using Samtools and the figure was generated using R-Studio and annotated in Inkscape v0.91.

Discussion

Chikungunya virus is a re-emerging alphavirus causing a high morbidity with long term arthralgia. Previous studies have taken approaches to understand the interaction between chikungunya virus and the *Aedes aegypti* vector [39]. However, considering the switch in vector preference towards *Aedes albopictus* by Indian Ocean isolates and invasive nature of this mosquito species, it is paramount to characterize the interaction between CHIKV and the new vector.

Previous studies with whole transcriptome analysis in mosquito vectors have used either whole mosquitoes or cell culture [18-22, 40]. Our objective here was to study the vector-virus interaction specifically at midgut, the first barrier site, and head/thorax, which denote the dissemination sites and include salivary glands, to understand the factors that play a critical role in determining mosquito vector competence.

In the current study, unbiased transcriptional analysis was performed on tissues collected from lab reared adult female *Aedes albopictus* mosquitoes over the course of CHIKV infection. The gene expression patterns at these tissue sites compared to uninfected samples revealed the transcriptional changes that are likely to be in response to the viral infection.

Our analysis revealed that at 2 dpi in midgut, most of the transcriptional changes are related to

metabolism. Analysis of molecular functions revealed that while lysozyme activity and alkaline

phosphatase were down regulated, carboxypeptidase activity was upregulated. Indeed,

lysozymal and carboxypeptidase pathways are implicated in innate immune responses against

multiple arboviral infections [41-44].

At 8dpi in head/thorax, differential regulation of biological processes including RNA and mRNA

binding, lysosomal and serpin pathways and down regulation of defensin genes were observed.

These could be due to either mosquito immune responses to CHIKV or viral modulation of

immunity. Processes such as the regulation of transport and homophilic cell adhesion via plasma

membrane could be involved in viral assembly and export [45-48]. Functional studies to

determine whether these genes are pro or anti-viral are needed and could explain the role of

these genes in the infection process.

Interestingly, in D8 HTs, two odorant binding proteins (OBPs) were found to be differentially

expressed. Obp25 (AALF018602) was upregulated while D7 protein (AALF024478) was down

regulated. In *Aedes aegypti*, salivary glands infected with dengue virus also showed differential

expression of odorant binding proteins [49] and it was shown that down regulation of OBPs

reduced the chemosensory abilities of the mosquitoes and hence reduced exposure to virus via

feeding and thus hindered transmission capabilities. Similar mechanisms may be in play here as

well, although that remains to be verified.

Multiple prior publications have also shown that the RNAi pathway is one of the major pathways

involved in antiviral responses in insects [50-52]. In this study, we did not find any statistically

significant changes in expression of genes involved in RNAi pathways. It is possible that the

regulation of this pathway either does not occur at the transcriptional level or the proteins

involved are ubiquitously expressed and not differentially regulated. It is also possible that the

time points we selected did not coincide with RNAi activation.

Twenty-two genes were selected for validation by qRT-PCR, based on their possible involvement

in mosquito immunity. The concordance was lower in the D8 genes compared to D2 genes. This

could be due to the lower number of controls in D8 samples in the RNAseq analysis.

The incomplete and poorly annotated reference genome of *Aedes albopictus* was a hindrance in

performing data analysis and robust pathway analysis. We also used heads and anterior 1/3rd of

the thorax, which included salivary glands at 8 dpi. The results from D8 samples represent data

from heterogeneous tissue, and care needs to be taken before any broad conclusions are drawn.

Functional characterisation of the identified genes may help in deciphering the results and

understanding their role in mosquito-virus interactions.

RNA viruses, like CHIKV, have an inherent ability to rapidly mutate and generate variants for

adaptation in novel environments through an error-prone polymerase[53]. Viral diversification is

thought to be driven by the mosquito immune system leading to evolution of new genotypes.

Previous study has shown that West Nile virus exhibits stochastic reductions in genetic diversity,

which was recovered during intra-tissue population expansions [54]. Here, we showed that the

mutations that arose in MG and HT samples compared to the original Vero cell culture isolate

indicate possible changes in the viral sequence as it adapts from a mammalian host to different

mosquito tissues. Vero cells, derived from African green monkey kidney, are known to be type-1

interferon response deficient [55], which may result in a high amount of CHIKV viral diversity as

seen by high number of viral variants. Our results suggest that this diversity increases significantly

in the mosquito midgut samples at 2dpi. We hypothesise that this increased genomic diversity

represents CHIKV adapting and replicating in the mosquito midgut milieu. Interestingly, the

number of CHIKV viral variants decreased considerably in HT by 8 dpi, possibly indicating a genetic

bottleneck at this stage. These results are consistent with previous literature that show increase

in CHIKV fitness and its adaptability during host switch [56]. The CHIKV 3' UTR enhances viral

replication in mosquitoes by interacting with mosquito cell-specific factors. This region contains

highly conserved sequence elements for viral replication, binding sites for cellular miRNAs that

determine cell tropism, host range, and pathogenesis, and conserved binding regions for a

cellular protein that influences viral RNA stability [57]. The 3'-UTR of the CHIKV genome showed

the most variation in all our samples. Although significant, this may be due sequencing errors at

this AT-rich region. Non-structural protein, nSP3, is an important protein for viral pathogenesis

with immune modulatory functions [58]. We detected an apparent 'hotspot' of variability within

the coding sequence of this protein in all samples, suggesting an important functional effect in

mosquitoes that warrants further study.

Overall, our results showed significant changes in the transcriptome of *Aedes albopictus*

mosquitoes after CHIKV infection, with identified genes involved in multiple cellular processes.

This study, for the first time, examines differential gene expression at the midgut (the first critical

barrier site) and head and thorax (dissemination site containing salivary glands) in infected

mosquitoes. This study can be utilized in determining potential pro-viral and antiviral host factors

and in turn, will be helpful in reducing the high impact of CHIKV infections by targeting the vector,

Aedes albopictus.

Acknowledgments

The chikungunya virus isolate was obtained from the Victorian Infectious disease reference

laboratory, Melbourne. Thanks to Dr Kim Blasdell for helping with CLC Genomics workbench and

to Chris Freebairn for collection of *Aedes albopictus* eggs from Torres Strait islands. Thanks to

Adam Foord from AAHL-CSIRO for discussions about optimisation of qRT-PCR.

References

1.   Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The global

distribution and burden of dengue. Nature. 2013;496:504. doi: 10.1038/nature12060

2.   Lumsden WH. An epidemic of virus disease in Southern Province, Tanganyika Territory,

in 1952-53. II. General description and epidemiology. Transactions of the Royal Society of

Tropical Medicine and Hygiene. 1955;49(1):33-57. Epub 1955/01/01. PubMed PMID: 14373835.

3.   Robinson MC. An epidemic of virus disease in Southern Province, Tanganyika Territory,

in 1952-53. I. Clinical features. Transactions of the Royal Society of Tropical Medicine and

Hygiene. 1955;49(1):28-32. Epub 1955/01/01. PubMed PMID: 14373834.

4.   Kamath S, Das AK, Parikh FS. Chikungunya. J Assoc Physicians India. 2006;54.

5.   Higgs S, Vanlandingham D. Chikungunya virus and its mosquito vectors. Vector Borne

Zoonotic Dis. 2015;15(4):231-40. Epub 2015/02/13. doi: 10.1089/vbz.2014.1745. PubMed

PMID: 25674945.

6.   Simon F, Javelle E, Oliver M, Leparc-Goffart I, Marimoutou C. Chikungunya Virus

Infection. Current Infectious Disease Reports. 2011;13(3):218-28. doi: 10.1007/s11908-011-

0180-1. PubMed PMID: PMC3085104.

7.   Reiter P, Fontenille D, Paupy C. *Aedes albopictus* as an epidemic vector of chikungunya

virus: another emerging problem? The Lancet Infectious Diseases. 2006;6(8):463-4. doi:

10.1016/s1473-3099(06)70531-x.

8.   Tsetsarkin KA, Chen R, Weaver SC. Interspecies transmission and chikungunya virus

emergence. Current Opinion in Virology. 2016;16:143-50. doi:

http://dx.doi.org/10.1016/j.coviro.2016.02.007.

9.   Waldock J, Chandra NL, Lelieveld J, Proestos Y, Michael E, Christophides G, et al. The role

of environmental variables on Aedes albopictus biology and chikungunya epidemiology.

Pathogens and global health. 2013;107(5):224-41. doi: 10.1179/2047773213Y.0000000100.

PubMed PMID: PMC4001452.

10.   Rezza G, Nicoletti L, Angelini R, Romi R, Finarelli AC, Panning M, et al. Infection with

chikungunya virus in Italy: an outbreak in a temperate region. Lancet (London, England).

2007;370(9602):1840-6. Epub 2007/12/07. doi: 10.1016/s0140-6736(07)61779-6. PubMed

PMID: 18061059.

11.   Enserink M. Chikungunya: No Longer a Third World Disease. Science.

2007;318(5858):1860.

12.   Franz AWE, Kantor AM, Passarelli AL, Clem RJ. Tissue Barriers to Arbovirus Infection in

Mosquitoes. Viruses. 2015;7(7):3741-67. doi: 10.3390/v7072795. PubMed PMID: PMC4517124.

13.   Dong S, Kantor AM, Lin J, Passarelli AL, Clem RJ, Franz AWE. Infection pattern and

transmission potential of chikungunya virus in two New World laboratory-adapted Aedes

aegypti strains. Scientific reports. 2016;6:24729. doi: 10.1038/srep24729. PubMed PMID:

PMC4840389.

14.   Arias-Goeta C, Mousson L, Rougeon F, Failloux A-B. Dissemination and Transmission of

the E1-226V Variant of Chikungunya Virus in Aedes albopictus Are Controlled at the Midgut

Barrier Level. PLOS ONE. 2013;8(2):e57548. doi: 10.1371/journal.pone.0057548.

15.   Dubrulle M, Mousson L, Moutailler S, Vazeille M, Failloux A-B. Chikungunya Virus and

Aedes Mosquitoes: Saliva Is Infectious as soon as Two Days after Oral Infection. PLOS ONE.

2009;4(6):e5895. doi: 10.1371/journal.pone.0005895.

16.   Kuno G, Chang G-JJ. Biological Transmission of Arboviruses: Reexamination of and New

Insights into Components, Mechanisms, and Unique Traits as Well as Their Evolutionary Trends.

Clinical Microbiology Reviews. 2005;18(4):608-37. doi: 10.1128/CMR.18.4.608-637.2005.

17.   Chompoosri J, Thavara U, Tawatsin A, Boonserm R, Phumee A, Sangkitporn S, et al.

Vertical transmission of Indian Ocean Lineage of chikungunya virus in Aedes aegypti and Aedes

albopictus mosquitoes. (1756-3305 (Electronic)). doi: D - NLM: PMC4842298 OTO - NOTNLM.

18.   Dong S, Behura SK, Franz AWE. The midgut transcriptome of Aedes aegypti fed with

saline or protein meals containing chikungunya virus reveals genes potentially involved in viral

midgut escape. BMC Genomics. 2017;18(1):382. Epub 2017/05/17. doi: 10.1186/s12864-017-

3775-6. PubMed PMID: 28506207; PubMed Central PMCID: PMCPMC5433025.

19.   Etebari K, Hegde S, Saldaña MA, Widen SG, Wood TG, Asgari S, et al. Global

Transcriptome Analysis of Aedes aegypti Mosquitoes in Response to Zika Virus Infection.

mSphere. 2017;2(6):e00456-17. doi: 10.1128/mSphere.00456-17. PubMed PMID: PMC5700376.

20.   Shrinet J, Srivastava P, Sunil S. Transcriptome analysis of Aedes aegypti in response to

mono-infections and co-infections of dengue virus-2 and chikungunya virus. Biochemical and

Biophysical Research Communications. 2017;492(4):617-23. doi:

https://doi.org/10.1016/j.bbrc.2017.01.162.

21.   Bonizzoni M, Dunn WA, Campbell CL, Olson KE, Marinotti O, James AA. Complex

Modulation of the Aedes aegypti Transcriptome in Response to Dengue Virus Infection. PLOS

ONE. 2012;7(11):e50512. doi: 10.1371/journal.pone.0050512.

22.   Fragkoudis R, Chi Y, Siu RW, Barry G, Attarzadeh-Yazdi G, Merits A, et al. Semliki Forest

virus strongly reduces mosquito host defence signaling. Insect Mol Biol. 2008;17(6):647-56.

Epub 2008/09/25. doi: 10.1111/j.1365-2583.2008.00834.x. PubMed PMID: 18811601; PubMed

Central PMCID: PMCPmc2710796.

23.   Liu Y, Zhou Y, Wu J, Zheng P, Li Y, Zheng X, et al. The expression profile of Aedes

albopictus miRNAs is altered by dengue virus serotype-2 infection. Cell & bioscience. 2015;5:16.

Epub 2015/04/30. doi: 10.1186/s13578-015-0009-y. PubMed PMID: 25922662; PubMed Central

PMCID: PMCPMC4411651.

24.   Pyke AT. Partial E1 gene sequence of Mauritius 2006 isolate 06113879. Unpublished.

2008.

25.   Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory

requirements. Nature Methods. 2015;12:357. doi: 10.1038/nmeth.3317.

26.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence

Alignment/Map format and SAMtools. Bioinformatics (Oxford, England). 2009;25(16):2078-9.

Epub 2009/06/10. doi: 10.1093/bioinformatics/btp352. PubMed PMID: 19505943; PubMed

Central PMCID: PMCPMC2723002.

27.   Lawson D, Arensburger P, Atkinson P, Besansky NJ, Bruggner RV, Butler R, et al.

VectorBase: a data resource for invertebrate vector genomics. Nucleic acids research.

2009;37(suppl 1):D583-D7. doi: 10.1093/nar/gkn857.

28.   Afgan E, Sloggett C, Goonasekera N, Makunin I, Benson D, Crowe M, et al. Genomics

Virtual Laboratory: A Practical Bioinformatics Workbench for the Cloud. PLOS ONE.

2015;10(10):e0140829. doi: 10.1371/journal.pone.0140829.

29.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome biology. 2014;15(12):550. doi: 10.1186/s13059-014-0550-

8.

30.   Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, Bowden J, et al. De novo

transcript sequence reconstruction from RNA-Seq: reference generation and analysis with

Trinity. Nature protocols. 2013;8(8):10.1038/nprot.2013.084. doi: 10.1038/nprot.2013.084.

PubMed PMID: PMC3875132.

31.   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length

transcriptome assembly from RNA-Seq data without a reference genome. Nature

Biotechnology. 2011;29:644. doi: 10.1038/nbt.1883.

32.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:

architecture and applications. BMC Bioinformatics. 2009;10:421-. doi: 10.1186/1471-2105-10-

421. PubMed PMID: PMC2803857.

33.   Boratyn GM, Schäffer AA, Agarwala R, Altschul SF, Lipman DJ, Madden TL. Domain

enhanced lookup time accelerated BLAST. Biology Direct. 2012;7:12-. doi: 10.1186/1745-6150-

7-12. PubMed PMID: PMC3438057.

34.   Smedley D, Haider S, Durinck S, Pandini L, Provero P, Allen J, et al. The BioMart

community portal: an innovative alternative to large, centralized data repositories. Nucleic

acids research. 2015;43(W1):W589-W98. doi: 10.1093/nar/gkv350.

35.   Alexa A RJ. topGO: Enrichment Analysis for Gene Ontology. R package version 2330.

2016.

36.   Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, et al. VarScan 2: somatic

mutation and copy number alteration discovery in cancer by exome sequencing. Genome Res.

2012;22(3):568-76. Epub 2012/02/04. doi: 10.1101/gr.129684.111. PubMed PMID: 22300766;

PubMed Central PMCID: PMCPMC3290792.

37.   Koboldt DC, Zhang Q, Larson DE, Shen D, McLellan MD, Lin L, et al. VarScan 2: Somatic

mutation and copy number alteration discovery in cancer by exome sequencing. Genome

Research. 2012;22(3):568-76. doi: 10.1101/gr.129684.111. PubMed PMID: PMC3290792.

38.   Mudiganti U, Hernandez R, Brown DT. Insect response to alphavirus infection--

establishment of alphavirus persistence in insect cells involves inhibition of viral polyprotein

cleavage. Virus Res. 2010;150(1-2):73-84. Epub 2010/03/11. doi:

10.1016/j.virusres.2010.02.016. PubMed PMID: 20214937.

39.   McFarlane M, Arias-Goeta C, Martin E, O'Hara Z, Lulla A, Mousson L, et al.

Characterization of Aedes aegypti innate-immune pathways that limit Chikungunya virus

replication. PLoS neglected tropical diseases. 2014;8(7):e2994. Epub 2014/07/25. doi:

10.1371/journal.pntd.0002994. PubMed PMID: 25058001; PubMed Central PMCID:

PMCPmc4109886.

40.   Colpitts TM, Cox J, Vanlandingham DL, Feitosa FM, Cheng G, Kurscheid S, et al.

Alterations in the Aedes aegypti Transcriptome during Infection with West Nile, Dengue and

Yellow Fever Viruses. PLoS pathogens. 2011;7(9):e1002189. doi: 10.1371/journal.ppat.1002189.

41.   Johnston C, Jiang W, Chu T, Levine B. Identification of Genes Involved in the Host

Response to Neurovirulent Alphavirus Infection. Journal of Virology. 2001;75(21):10431-45. doi:

10.1128/JVI.75.21.10431-10445.2001. PubMed PMID: PMC114617.

42.   Merkling SH, van Rij RP. Beyond RNAi: Antiviral defense strategies in Drosophila and

mosquito. Journal of Insect Physiology. 2013;59(2):159-70. doi:

https://doi.org/10.1016/j.jinsphys.2012.07.004.

43.   Swevers L, Liu J, Smagghe G. Defense Mechanisms against Viral Infection in Drosophila:

RNAi and Non-RNAi. Viruses. 2018;10(5). Epub 2018/05/05. doi: 10.3390/v10050230. PubMed

PMID: 29723993; PubMed Central PMCID: PMCPMC5977223.

44.   Isoe J, Zamora J, Miesfeld RL. Molecular Analysis of the Aedes aegypti Carboxypeptidase

Gene Family. Insect biochemistry and molecular biology. 2009;39(1):68-73. doi:

10.1016/j.ibmb.2008.09.006. PubMed PMID: PMC2673731.

45.   Mateo M, Generous A, Sinn PL, Cattaneo R. Connections matter – how viruses use cell–

cell adhesion components. Journal of Cell Science. 2015;128(3):431-9. doi: 10.1242/jcs.159400.

PubMed PMID: PMC4311127.

46.   Bhella D. The role of cellular adhesion molecules in virus attachment and entry.

Philosophical Transactions of the Royal Society B: Biological Sciences.

2015;370(1661):20140035. doi: 10.1098/rstb.2014.0035. PubMed PMID: PMC4275905.

47.   Chisenhall DM, Christofferson RC, McCracken MK, Johnson A-MF, Londono-Renteria B,

Mores CN. Infection with dengue-2 virus alters proteins in naturally expectorated saliva of

Aedes aegypti mosquitoes. Parasites & vectors. 2014;7(1):252. doi: 10.1186/1756-3305-7-252.

48.   Gulley MM, Zhang X, Michel K. The roles of serpins in mosquito immunology and

physiology. Journal of Insect Physiology. 2013;59(2):138-47. doi:

https://doi.org/10.1016/j.jinsphys.2012.08.015.

49.   Sim S, Ramirez JL, Dimopoulos G. Dengue Virus Infection of the Aedes aegypti Salivary

Gland and Chemosensory Apparatus Induces Genes that Modulate Infection and Blood-Feeding

Behavior. PLoS pathogens. 2012;8(3):e1002631. doi: 10.1371/journal.ppat.1002631.

50.   Campbell CL, Keene KM, Brackney DE, Olson KE, Blair CD, Wilusz J, et al. Aedes

aegyptiuses RNA interference in defense against Sindbis virus infection. BMC Microbiology.

2008;8(1):47. doi: 10.1186/1471-2180-8-47.

51.   Cirimotich CM, Scott JC, Phillips AT, Geiss BJ, Olson KE. Suppression of RNA interference

increases alphavirus replication and virus-associated mortality in Aedes aegypti mosquitoes.

BMC Microbiology. 2009;9:49-. doi: 10.1186/1471-2180-9-49. PubMed PMID: PMC2660349.

52.   McFarlane M, Arias-Goeta C, Martin E, O'Hara Z, Lulla A, Mousson L, et al.

Characterization of Aedes aegypti Innate-Immune Pathways that Limit Chikungunya Virus

Replication. PLoS neglected tropical diseases. 2014;8(7):e2994. doi:

10.1371/journal.pntd.0002994. PubMed PMID: PMC4109886.

53.   Holland J, Spindler K, Horodyski F, Grabau E, Nichol S, VandePol S. Rapid evolution of

RNA genomes. Science. 1982;215(4540):1577.

54.   Grubaugh ND, Weger-Lucarelli J, Murrieta RA, Fauver JR, Garcia-Luna SM, Prasad AN, et

al. Genetic drift during systemic arbovirus infection of mosquito vectors leads to decreased

relative fitness during host switching. Cell Host & Microbe. 2016;19(4):481-92. doi:

10.1016/j.chom.2016.03.002. PubMed PMID: PMC4833525.

55.   Desmyter J, Melnick JL, Rawls WE. Defectiveness of Interferon Production and of Rubella

Virus Interference in a Line of African Green Monkey Kidney Cells (Vero). Journal of Virology.

1968;2(10):955-61. PubMed PMID: PMC375423.

56.   Coffey LL, Vignuzzi M. Host Alternation of Chikungunya Virus Increases Fitness while

Restricting Population Diversity and Adaptability to Novel Selective Pressures. Journal of

Virology. 2011;85(2):1025-35. doi: 10.1128/jvi.01918-10. PubMed PMID: PMC3020036.

57.   Hyde JL, Chen R, Trobaugh DW, Diamond MS, Weaver SC, Klimstra WB, et al. The 5' and

3' ends of alphavirus RNAs – non-coding is not non-functional. Virus research. 2015;206:99-107.

doi: 10.1016/j.virusres.2015.01.016. PubMed PMID: PMC4654126.

58.   Mathur K, Anand A, Dubey SK, Sanan-Mishra N, Bhatnagar RK, Sunil S. Analysis of

chikungunya virus proteins reveals that non-structural proteins nsP2 and nsP3 exhibit RNA

interference (RNAi) suppressor activity. Scientific reports. 2016;6:38065. doi:

10.1038/srep38065.

563
564

Supporting information Captions

S1 Supplementary information. Table A: List of primers, Table B: Ct values of 10-fold dilutions of CHIKV

RNA, Table C: Infection check of insect tissue pools and Table D: ΔΔCt of validated targets on qRT-PCR.

S2. List of differentially expressed genes.

S3. Custom transcriptome output.

Figure 1.

Figure 2.

Figure 3.