# New criteria for sympatric speciation in the genomic era

Emilie J Richards[1], Maria R. Servedio[1], Christopher H Martin[1]*

[1] Department of Biology, University of North Carolina at Chapel Hill, Chapel Hill NC

*corresponding author: chmartin@unc.edu

Main text: 3,139 words, 1 Figure, 3 Boxes

**Abstract**

Sympatric speciation illustrates how natural and sexual selection may create new species in

isolation without geographic barriers. However, so far, all genomic reanalyses of classic

examples of sympatric speciation indicate secondary gene flow occurred. Thus, there is a need to

revisit criteria for demonstrating sympatric speciation in the face of widespread gene flow. We

summarize theoretical differences between sympatric speciation and speciation-with-gene-flow

models and propose genomic criteria for sympatric speciation: 1) timing of fine-scale

introgression; 2) timing of selective sweeps and 3) functional annotation of this introgressed

variation; and 4) the absence of similar sweeps in outgroups. Monophyly is an insufficient

criterion for sympatric speciation; we must take a locus-specific approach to investigate whether

any introgression contributed to reproductive isolation.

**What is sympatric speciation?**

Sympatric speciation is the evolution of reproductive isolation within a single panmictic

population without the aid of any geographic isolation [1]. It represents the most extreme and

controversial endpoint on the divergence with gene flow continuum: panmictic gene flow and no

initial divergence at the start of speciation [2–5]. In the context of theoretical speciation models,

sympatric speciation is the most difficult process because the starting conditions involve no pre-

existing divergence, potentially tied with physical linkage, among loci involved in reproductive

isolation (i.e. 'barrier' loci [6,7]); instead, **linkage disequilibrium** (see Glossary) must build up

through time within a population through the action of disruptive natural selection and strong

assortative mating by ecotype, despite the countervailing eroding force of recombination [8–11].

We can thus distinguish different types of scenarios that will result in two sister species

being found in sympatry based on whether **secondary gene flow** aided population divergence: 1)

classic sympatric speciation without gene flow; 2) sympatric speciation in the presence of a)

neutral secondary gene flow or b) after differential sorting of an ancestral hybrid swarm. In the

latter case, it is important to distinguish whether the ancestral hybrid swarm population achieved

panmixia before later divergence (i.e. sympatric divergence); otherwise, differential sorting of

haplotypes within the hybrid swarm is better described by secondary contact speciation with

gene flow models. 3) Speciation may be aided by secondary gene flow that a) triggers initial

sympatric divergence or b) increases divergence after initial divergence in sympatry becomes

stalled, an outcome of many sympatric speciation models without sufficiently strong disruptive

selection [12–15]. Finally, 4) secondary contact after a period of allopatry between two

populations can result in coexistence or reinforcement, if there is not collapse into a single

admixed population [16–19]. We consider scenarios 1 and 2 to be examples of sympatric

speciation, whereas scenarios 3 and 4 would be examples of speciation aided by secondary gene

flow. Interestingly, hybrid swarm scenarios exist in a gray area, since substantial initial gene

flow from multiple sources may increase ecological or preference variation within a population

that is sufficient to trigger later sympatric divergence, even without segregating inversions or

genetic incompatibilities [20–22]. So far, we know of no examples of scenario 1 within any case

study of sympatric sister species examined using genomic tools; even long diverged species

show some evidence of **introgression** in their past (e.g. [23]). In contrast, sympatric speciation

with neutral gene flow (Scenario 2) and speciation aided by gene flow (Scenarios 3 and 4)

frequently appear to operate concurrently even within a single sympatric adaptive radiation (e.g.

[24–26]).

It is important to distinguish these scenarios because theoretical models predict that

sympatric divergence unaided by any form of secondary gene flow is substantially more difficult

than other speciation with gene flow scenarios (Box 1). Gene flow throughout the speciation

process allows recombination to break down linkage disequilibrium among alleles associated

with ecological divergence and assortative mating.  There are actually three different classes of

sympatric speciation models to consider: the most difficult process involves independently

segregating loci for ecotype, female preferences, and male traits within the population, whereas

sympatric divergence is much easier if any of these three types of traits are combined, such as

assortative mating based on phenotype matching instead of separate loci for preference and traits

[27,28] or "magic" traits (such as assortative mating based on microhabitat preference;

[9,10,29]). Sympatric speciation by sexual selection alone is also theoretically possible if there is

substantial preference variation either initially within the population or through secondary gene

flow [20,30,31].

Any form of linkage disequilibrium among ecological and mate choice loci formed in

allopatry, whether due to physical linkage, selection, or drift, can tend to shift the initial starting

conditions of panmixia in favor of sympatric divergence [17]. However, linkage disequilibrium

without physical linkage subsides within a few generations after secondary sympatry and thus

may not allow sufficient time for the evolution of assortative mating within the population. In

contrast, pre-existing physical linkage among ecological loci has been shown to increase the

probability of divergence, especially when it captures already divergent alleles as is more likely

after allopatric divergence [32,33]. Similarly, physical linkage can cause preference and trait

alleles to mimic phenotype matching, although even tight linkage can break down over long

timescales (shown in a model with population structure: [34]). Segregating inversions in the

ancestral population are now well-known empirical examples of physical linkage promoting

divergence in sympatry [35–37]. Sympatric divergence is also limited by many other restrictive

conditions regarding the costs of female choosiness and strengths of disruptive selection and

assortative mating (Box 1).

Despite extensive searches for examples of sympatric speciation in the wild, there are few

convincing case studies due to the difficulty of ruling out historical allopatric scenarios (see

below) and the new difficulty of ruling out a role of introgression in speciation. Furthermore, the

role of magic traits or matching vs. preference/trait mechanisms is not fully understood in any

existing case study. Thus, we still have very limited empirical tests of an extensive theoretical

literature and diverse competing models of the notoriously difficult process of sympatric

speciation [24,27,28,38–42].

**The classic problem of sympatric speciation**

There are four traditional criteria for demonstrating classic sympatric speciation (Scenario 1): 1)

sister species which are reproductively isolated, 2) form a **monophyletic** group, 3) largely

overlap in ranges, and 4) have biogeographic and evolutionary histories that make periods of

allopatric divergence highly unlikely [1]. Very few case studies have been able to meet these

rigorous criteria despite intense searches [1,5]. This has led to the prominent status of crater lake

cichlid radiations as some of the best examples of sympatric speciation in the wild due to the

uniform shape of isolated volcanic lakes which convincingly rule out phases of allopatry due to

water level changes (Box 2; [43–45]).

The monophyly criterion assumes that monophyly arises only when a single ancestral

population underlies the present-day daughter species. This is typically met by inferring a single

phylogeny from one or more loci. This single point-estimate view of evolutionary history is

problematic because it obscures the presence of non-bifurcating relationships among organisms

(e.g. sister species which derived ancestry from multiple source populations due to extensive

gene flow or hybrid speciation) and the real variation in evolutionary histories among genes

across the genome itself (e.g. [46]). Few regions of the genome may initially contribute to

reproductive isolation resulting in a heterogeneous genomic landscape of differentiation among

incipient species [47], a pattern now extensively supported across case studies [48–51].

Therefore, monophyletic relationships are consistent with, but not exclusive to a scenario of

sympatric speciation. Examining heterogeneous evolutionary histories across regions relevant to

speciation is thus crucial for understanding the processes and conditions under which sympatric

divergence can occur.

**The 'new' problem of sympatric speciation**

While genomics has increased our ability to resolve evolutionary relationships among organisms,

it has also revealed more complex evolutionary histories of multiple colonization and extensive

secondary gene flow in all examples of sympatric speciation that have been examined with

genomic data so far [52–68]. Indeed, only a handful of genes may directly contribute to the

speciation process whereas the rest of the genome is porous to gene flow while reproductive

isolation is incomplete [47,69]. Examples of 'classic' sympatric speciation without secondary

gene flow (Scenario 1) are now unknown after applying modern genomic tools to search for

introgression (and paleogenomics is therefore unnecessary to provide historical point estimates

of spatial isolation as recently suggested [70]). Instead, it is still possible that sympatric

speciation occurs in the face of secondary gene flow in nearly all these examples (Scenario 2;

[67]). Importantly, most evidence of secondary gene flow comes from genome-wide tests of

introgression from outgroup lineages, not gene flow between diverging populations in sympatry

(e.g. [64,65]). Therefore, introgression detected at the genome-wide level from lineages outside

the speciation event tells us little about the divergence process among incipient sympatric species

and how gene flow shaped the process of speciation.

The challenge of sympatric speciation in the genomic era is establishing or rejecting a

functional role for the ubiquitous secondary gene flow present during the speciation process, in

effect ruling out scenarios 3 and 4 in favor of scenario 2 (Fig. 1). Even if signatures of secondary

gene flow are detected, speciation could still have occurred solely via mechanisms of sympatric

speciation if that secondary gene flow did not play a causal role in divergence. Secondary gene

flow could play a causal role if it introduced novel genetic variation or physically linked alleles

(e.g. a segregating inversion) that promote speciation before the start of divergence, such as

**hybrid swarm (**Scenario 2b**;** ([21,22,25,71]), adaptive introgression (Scenario 3; [72–77]),

**transgressive segregation** (Scenarios 2-3; [78,79]), or hybrid speciation (Scenario 4; [80]). We

note a distinction between ancestral introgression of segregating haplotypes promoting speciation

(speciation with gene flow models apply) versus sufficient time for recombination to break down

these haplotypes after a hybrid swarm and create panmictic conditions before the start of

divergence (sympatric speciation models apply) versus simply inflated ecological and preference

variation within a population due to hybrid swarm (gray area; sympatric speciation more likely

due to this initial gene flow). Here we propose and discuss new genomic criteria to help establish

or reject a functional role of secondary gene flow in the speciation process (Fig. 1). This is

necessary to identify putative cases of the sympatric speciation process when gene flow appears

to be nearly universal in the wild, particularly among sympatric diverging populations.

**New criteria for sympatric speciation in the genomic era**

Although genome-wide analyses of introgression provide a starting point, ultimately

consideration of the time of arrival and functional role of each introgressed region within extant

sympatric sister species pairs will be necessary to distinguish between sympatric speciation with

gene flow (Scenario 2) or speciation aided by secondary gene flow (Scenario 3; e.g. segregating

inversions [35,81] or ancient balancing selection on regions containing multiple barrier loci

[82,83]). We suggest four major types of genomic analyses as new criteria to help identify

sympatric speciation with gene flow: 1) estimate the timing of introgression into sympatric sister

species relative to their divergence time, 2) infer the presence and timing of selective sweeps

within sympatric sister species, 3) annotate candidate adaptive introgression regions for

functional elements or trait associations that may be relevant to speciation, and 4) if closely

related non-speciating outgroups are available, confirm the lack of selective sweeps of these

regions in outgroups. Combining these statistics will aid in distinguishing where case studies fall

along the speciation with gene flow continuum and whether the starting conditions of panmixia

in sympatric speciation models will apply (Fig. 1).

*1) Secondary gene flow is constant across the speciation process or not concurrent with*

*divergence times*

Estimating the duration of gene flow and the timing of introgression relative to the timing of

divergence between sympatric sister species will help distinguish between scenarios of sympatric

speciation, speciation with gene flow, and secondary contact. If populations diverged in

sympatry independent of any concurrent secondary gene flow (Scenario 2), we might expect to

see weak concordance of the timing of gene flow with divergence times among species. This

discordance could be in the form of discrete gene flow events that date well before or after

divergence times among species (Fig 1A). In the case of continuous gene flow from the time of

colonization to the present, more information about the functional role and selection on

introgressed regions will be needed.

The timing of introgression is also useful in ruling out other evolutionary phenomena that

can leave similar genomic signatures. The random or biased assortment of ancestral variation

among lineages during the speciation process can create similar phylogenetic patterns to

introgression resulting from secondary gene flow [82,84]. Timing is important for differentiating

introgression from the sorting of ancient ancestral polymorphisms due to processes such as

balancing selection. For example, if genetic divergence in an introgressed region shared between

sister species is greater (e.g. elevated $D_{xy}$) than expected given divergence time between the

sister species, this pattern suggests differential sorting of ancestral variation and doesn't rule out

a scenario of sympatric speciation (Scenarios 1 & 2). If introgression after secondary contact did

occur, genetic divergence in these regions between recipient sister species should be lower than

expected given their divergence time. Increasingly sophisticated approaches for detecting fine-

scale patterns of introgression are available to estimate the timing and duration of gene flow

from genomic data (Box 3).

*2) Lack of selective sweeps or non-concurrent timing of sweeps in regions that have experience*

*gene flow*

We can use information about selective sweeps of introgressed variation to further characterize

the role of gene flow in sympatric divergence. When an allele is selectively favored in a

population, positive selection may cause it to increase in frequency and form a localized

selective sweep of reduced genetic variation surrounding the adaptive variant [85]. Such regions

of high differentiation in recently diverged species are often targeted as candidates for speciation

genes, although other processes not directly associated with speciation can lead to similar

patterns of high heterogeneity in differentiation across a genome (reviewed in [86–88]); indeed,

there is still no evidence that these regions are associated with reproductive isolation or reduced

gene flow and can also result from adaptive introgression [89,90]). If speciation was recent or

ongoing, there may be strong signatures of a selective sweep for particular haplotypes in at least

one of the sister species for regions involved in the divergence process (Fig. 1B). If secondary

gene flow was neutral with respect to speciation, we may find no signatures of selective sweeps

in those introgressed regions.

Importantly, a sweep of the same introgressed region in both sympatric sister species may

be interpreted as adaptation to the same new environment, which may not contribute to

reproductive isolation between the pair (dependent on their respective genetic backgrounds; e.g.

[90,91]). However, this pattern is also consistent with the sweep of a region contributing to a

'one-allele' mechanism of mate choice [27,28,92], such as increased female choosiness in both

sympatric sister species (e.g. [93]), which *would* contribute to reproductive isolation. Thus,

selective sweeps of an introgressed region in both sympatric sister species do not rule out its role

in aiding the speciation process.

Alternatively, if selective sweeps are detected, the timing of selective sweeps in the

regions affected by this gene flow can give indirect evidence about the selective pressure

underlying the sweep. It is challenging to infer something about the importance of an

introgressed region if the timing of introgression predates the timing of the selective sweep

because linkage disequilibrium among loci relevant to speciation may take time to build up, a

process involved in most speciation models [9,10]. However, the absence of selective sweeps or

introgression until long after population divergence would suggest that introgression was not

relevant to the speciation process (Scenario 2a). Introgressed regions that have undergone soft

selective sweeps and were important for divergence may easily be missed, but increasingly

sensitive methods [94,95] are making it easier to detect them.

*3) Weak support for casual role of secondary gene flow based on functional genetic analyses of*

*variants in the region*

Another potential source of evidence for the functional importance of gene flow can come from

associations between variants in introgressed regions with traits involved in ecological and

sexual isolation between sister species from genome-wide association studies (GWAS).

However, many complex traits are driven by a large number of variants of small effect and ruling

out a functional role for gene flow from any annotations is difficult (e.g. omnigenic model; [96]).

The conservation of sequences within introgressed regions across taxa may also provide strong

evidence of a functional role (e.g. PhastCons [97]). Finally, and most powerfully, genome editing

and gene expression reporter systems are increasingly tractable in non-model systems (e.g. [98–

101]. This is ultimately an asymmetric problem: finding evidence that an introgressed region

may have contributed to reproductive isolation is far easier than demonstrating that no

introgressed regions contributed to reproductive isolation in any way [67]. Finding evidence for

sympatric speciation in the wild is now the difficult problem of functional genetic analyses of

introgressed regions.

*4) Similar patterns of selection or divergence in the introgessed regions in closely related*

*outgroup populations*

Thorough investigations of these same regions in outgroups to the sympatric species gives added

power to distinguish whether secondary gene flow aided sympatric divergence. If non-

diversifying, closely related species exist in similar environments and haven't diversified in a

similar manner but share signatures of selective sweeps in the same regions, then the observed

introgression may have been neutral relative to speciation, e.g. due to adaptations to shared

changes in climate or pathogens or shared regions of reduced recombination or increased

background selection. For example, several studies comparing genomic landscapes of

differentiation across closely related taxa have found that high differentiation observed in the

same genomic regions across taxa reflects the action of linked selection across low-

recombination regions rather than selection against gene flow at barrier loci [102–110].

**Concluding Remarks**

Sympatric speciation remains among the most controversial evolutionary processes, beloved by

theorists and long sought after by empiricists. While evidence for this process appeared to be

mounting using traditional criteria [5], genomic data has now cast doubt on all these putative

examples due to the ubiquity of secondary gene flow. Furthermore, nearly all our existing case

studies involve some form of automatic magic trait, such as assortative mating by habitat

[35,111,112], along a depth gradient [63], or environment-induced phenology shifts [113].  We

think an outstanding remaining question is whether sympatric divergence can occur in nature

without the aid of some form of magic trait, as originally demonstrated to be possible in theory

[9,114].

Future fine-scale investigations of introgression will likely continue to paint a complex

picture of the role of secondary gene flow in nearly all speciation events. The highly polygenic

and multi-dimensional nature of adaptation and mate choice suggests that an 'all-of-the-above'

speciation scenario containing a mix of preference/trait, magic trait, and phenotype matching,

each spread across a wide distribution of allelic effect sizes with varying times of arrival, will be

the norm in nature.  In contrast, although numerous and diverse, most speciation models continue

to address these mechanisms in a piecemeal fashion with an assumption of large effect alleles. It

remains unclear how different mechanisms, effect sizes, and times of arrival will interact and

compete within a single model.

Interestingly, strict isolation of sympatric environments such as crater lakes may

ultimately become less important, since even these isolated environments are not isolated from

secondary gene flow [115,116]. Instead, recent sympatric divergence combined with well

characterized introgression and functional annotations may be the new limiting factor for

convincing case studies of sympatric divergence in the genomic era.

*Box 1*. **Why do we care whether speciation is sympatric?**

An increasingly common claim in the empirical speciation literature is that there is no difference

between 'speciation with gene flow' and 'sympatric speciation' (e.g. [70]). This contrasts sharply

with the rich theoretical literature differentiating models of sympatric speciation with models of

speciation with gene flow. Indeed, theory teaches us that we should care about the real

differences between the process of sympatric speciation (i.e. population divergence in sympatry

without the aid of introgression contributing to reproductive isolation) and other models of

speciation with gene flow. Sympatric speciation is uniquely and notoriously difficult [117], in

part because quite specific conditions of resource availability (e.g., [9,118]), mating traits and

preferences (e.g., [31,119]), and search costs (e.g., [120]) must be met for it to occur.

Inferences from theoretical models predict that, under a scenario of speciation with gene

flow (Scenario 3), introgression can make the process of speciation much easier in three ways.

First, by introducing additional variation in ecological traits into the population, introgression

could potentially facilitate a branching process due to competition for resources (although we are

not aware of a model that assesses this precise situation, it can be inferred from the dynamics of

[9]).  Second, introgression of novel alleles for mating preferences may provide a boost in

preference variation that could be an important trigger to aid the evolution of assortative mating

under a preference/trait mechanism, which requires preference variation to be large ([20,31]).

Indeed, we see exactly this pattern of secondary gene flow of olfactory alleles shortly before the

rapid divergence of a Cameroon cichlid radiation in Lake Ejagham [66]. Third, secondary

sympatry may lead to increased linkage disequilibrium between assortative mating and

ecological loci or among ecological loci.  It seems logical that this might facilitate sympatric

speciation as this metric is often described as progress along the speciation continuum. However,

initial linkage disequilibrium has been shown not to matter much in at least some scenarios [8],

because without physical linkage, linkage disequilibrium will break down quickly. However,

physical linkage may enable these alleles to remain in association for a sufficient time for

assortative mating to evolve within the population (e.g., [34]). Initial linkage disequilibrium may

also increase the probability of allelic capture by an inversion or for selection for new mutations

within the inversion that may affect both ecology and assortment [32]. Increased linkage

disequilibrium among ecological loci may also increase the probability of sympatric divergence,

but this is in effect similar to varying effect sizes of alleles at ecological loci (e.g. many small

effect alleles within a region resemble a large-effect locus [121–124]).

The fundamental difference between sympatric speciation and speciation with gene flow,

including secondary contact scenarios, lies in the fact that very often multiple equilibrium states

exist in speciation models, such that loss of divergence and maintenance of divergence in the

presence of gene flow are both possible outcomes, depending on the starting conditions of a

population (this is nicely illustrated for one measure of divergence by [17], Fig. I).  In such

cases, speciation is much more easily reached from starting conditions that match those of two

populations that have diverged largely in allopatry due to the large amount of allelic variation or

pre-existing phenotypic bimodality and assortative mating.  Even for scenarios of speciation with

gene flow that are much easier, such as geographic separation between two incipient species that

are undergoing gene flow, differentiation is much more difficult to reach or maintain from an

initially homogeneous population than from an initially differentiated one [125,126].

**Fig. I**

*Fig I.* *Two equilibrium cases exist for the linkage disequilibrium (LD), a proxy for*

*differentiation into two distinct "species" in this model, that can be maintained between two loci*

*that are under disruptive selection and determine assortative mating. With little initial LD, the*

*one-species equilibrium is likely to be reached even when the intensity of assortment is high.*

*When LD in the traits is initially large, as can be the case if there is initially divergence in*

*allopatry, the two-species equilibrium can be reached instead. Modified from* [17].

*Box 2.* **Evidence for sympatric speciation from crater lake cichlid radiations**

There are relatively few volcanic chains of crater lakes containing fishes in the tropics, notably

found only in Cameroon, Nicaragua, Tanzania, Uganda, Madagascar, and Papua New Guinea

[63,127,128]. Although sympatric radiations of endemic fishes are known from other isolated

saline, alkali, and ancient lakes [129–132], only three lineages of cichlids have radiated in the

world's crater lakes (Fig. II). The most diverse is Barombi Mbo, Cameroon with eleven endemic

species, followed by Lake Bermin, Cameroon with nine [43]. Nicaraguan crater lakes reach up to

five species [45,55,133,134], the East African craters never exceed two sympatric species

[63,68], and Madagascar's several crater lakes each contain only a single endemic cichlid [128].

It remains unknown why regional and lineage diversity varies so greatly because there appears to

be no relationship with crater lake size or age (up to approximately 5 km diameter and 2 million

years old) until reaching the much larger sizes of the East African rift lakes ([135], but also see

[134]).

In contrast to claims in a recent review [70], the evidence for sympatric speciation with

secondary gene flow is rather consistently in favor and remarkably similar across all crater lake

cichlid radiations. In all cases examined with genomic data so far, secondary gene flow was

detected, but there was little evidence it came from substantial divergence in allopatry followed

by secondary contact. Instead, nearly all studies have concluded sympatric divergence with

periodic or continuous gene flow, frequently from an initial hybrid swarm population (i.e.

introgression from multiple outgroup populations; [54,63–65,68]).

We think that the best evidence for secondary gene flow as a trigger of sympatric

divergence in cichlids comes from a radiation of three *Coptodon* species in Lake Ejagham:

demographic analyses of whole genomes suggest that this population did not diversify for 8,000

years in the face of frequent gene flow until an influx of olfactory receptor alleles coinciding

with the first sympatric divergence event in the lake [66]. Similarly in Lake Victoria, segregating

opsin alleles in riverine cichlid populations were differentially sorted among Lake Victorian

cichlids and may have triggered their diversification [25].

Evidence for sympatric divergence in crater lake cichlids without the aid of secondary

gene flow remains elusive. Malinsky et al. [63] showed that introgression and hybrid swarm

predated the divergence of a shallow/deep-water pair of cichlids in Lake Massoko, Tanzania;

however, these ancestral segregating haplotypes may have later aided sympatric divergence

(which admittedly is very difficult to rule out – see text). Very recent sympatric divergence in

some crater lakes on the order of thousands of years may also suggest that ongoing divergence

occurred in sympatry without the aid of gene flow [133,136]; however, it remains unclear if this

incipient divergence will become stalled as in other sympatric radiations [24]. Very rare

secondary gene flow without a clear functional role into the Barombi Mbo cichlid radiation ( <

1% introgressed regions in nearly all species) provides weak evidence of sympatric divergence,

but more functional characterization and timing of introgression is needed [67]. The recent

advent of transgenic reporters, CRISPR-Cas9, and *in situ* hybridization genetic tools within

Nicaraguan crater lake cichlids provides much promise for future investigations of the role of

introgression in sympatric divergence [99,137].

**Fig. II**

**Fig. II** Examples of volcanic crater lakes containing endemic cichlid radiations around the globe:

*a,d,f*) Barombi Mbo, Cameroon and its only outlet stream; *b*) Lake Apoyo, Nicaragua, *c*) Lake

Massoko, Tanzania, *e*) Lake Bermin, Cameroon. Satellite images (a-c) from Google Earth; other

images by CHM.

***Box 3.* Tools for detecting and dating local gene flow and selective sweeps**

*1) Detecting and dating local secondary gene flow*

While there are a variety of tests to detect gene flow on a local scale or within sliding genomic

windows [23,138–142], dating gene flow events is still accomplished mostly at the level of the

entire genome (although see [141] for dating introgression relative to divergence). Currently,

three types of demographic **coalescent** modeling approaches can infer local gene flow timing

based on different information from genomic data: 1) the distribution of allele frequencies from

genotype data (site frequency spectrum: [143,144]), 2) the distribution of haplotype block

lengths from phased genomes [145–148], and 3) variation in coalescent patterns among gene

trees [149]. Alternative approaches using **conditional random fields** [150] and **hidden Markov**

**models** [151,152] have also been used to detect and date gene flow events.

*2) Dating selective sweeps and ages of beneficial alleles:*

Methods for estimating the age of a sweep of a beneficial allele exploit several different aspects

about the pattern of variation surrounding the allele on its haplotypic background. These include

heuristic approaches that use point estimates of mean haplotype length or the number of derived

mutations within a chosen distance of the site [153–156] and model-based approaches that use

demographic information and summary statistics of allele frequencies and linkage disequilibrium

to model a distribution of ages that fit the observed data [157–160]. Alternatively, full sequence

data about haplotype structure on chromosomes and models that leverage the length of ancestral

haplotypes surrounding the beneficial allele and the accumulation of derived mutations can be

used to estimate the age of beneficial alleles [161–163].

3) *Functional genetic analyses of introgressed variants*

Functional assessments of introgressed regions minimally involve searching an annotated

reference genome for genes with relevant functions known for model organisms (or a pipeline

for assembling and annotating the organism of interest if an annotated reference is not already

available (reviewed in [164,165]). Introgressed regions that are unannotated can be searched for

evidence of potential functional importance based on strong sequence conservation across taxa

[97] or potential regulatory elements (reviewed in [166]). Additionally, genome wide association

studies (GWAS) can highlight variants in introgressed regions that may underlie complex traits

of interest, including novel variants previously unknown in model organisms (GWAS reviewed

in [167–169]). Functional validation of gene and regulatory element variants through genome-

editing experiments is also becoming increasingly tractable for non-model organisms (e.g.

[98,99,101,170]).

**Fig. 1**

**Figure 1: Genomic signatures of sympatric speciation and speciation with gene flow.**

Speciation scenarios are grouped into sympatric speciation scenarios (yellow box) and speciation

with gene flow scenarios (red box). Speciation from a hybrid swarm (orange box) can fall under

either category and additional information is necessary to determine what category of speciation

models best describe this process. **A)** The timing of gene flow relative to divergence can be used

to distinguish between speciation scenarios. The colored arrows represent gene flow events and

the colored lines within the tree represent a signature of introgression from that gene flow event

into the sympatric species. **B)** Venn diagrams illustrating the number of genomic windows across

the entire genome expected to have overlapping signatures of introgression (e.g. $f_d$ outliers),

genetic divergence (e.g. $F_{st}$ and $D_{xy}$ outliers), and selective sweeps (e.g. SweeD) for each

speciation scenario (e.g. see [77]). The highlighted sections of the Venn diagrams indicate the

key signature that can be used to distinguish between the scenarios. The scenarios that are

expected to leave very similar signatures of overlap are grouped by the bars colored with their

respective Venn diagram.

**Glossary Box**

**Coalescence**: The event of two sampled lineages from different populations merging back in

time in a shared ancestral lineage.

$D_{xy}$: A measure of absolute genetic divergence between populations calculated as the average

number of pairwise differences between sequences from two populations, excluding all the

comparisons between sequences within populations.

**Hybrid swarm**: A genetically diverse population with unique allele combinations derived from

the hybridization of multiple distinct taxa and subsequent backcrossing with hybrids and crossing

between hybrids themselves.

**Hidden Markov model**: A statistical modeling approach used to infer hidden states from

observed data along a sequence, where each hidden variable is independent of all others and

conditional only on the state of the previous hidden variable.

**Conditional random field**: A statistical modeling approach similar to hidden Markov models

except that each hidden variable can be conditional on regional hidden variables, not just the

immediately previous one.

**Incomplete lineage sorting**: The imperfect sorting of ancestral alleles between diverging

lineages that creates variable signatures about the evolutionary relationships among organisms.

**Introgression**: The movement and incorporation of genetic material from one distinct lineage

into another upon hybridization between the two and subsequent backcrossing with one of the

parent species.

**Linkage disequilibrium**: A non-random association of alleles at two or more loci.

**Monophyletic**: A group of lineages where the most recent ancestor of the group is not an

ancestor of any lineages outside the group.

**Secondary gene flow**: Any gene flow event from non-sympatric populations after the initial

colonization of the area that sympatric sister species diverged in. Introgression into the diverging

sister species following such events potentially brings in variation that has evolved in allopatry

that can aid the speciation process.

**Transgressive segregation**: The formation of extreme phenotypes in a segregating hybrid

population that are outside the range of phenotypes observed in parental species.

**Acknowledgements**

We thank J. McGirr, M. St. John, J. Poelstra, B. Reinhard, and J. Hermisson for valuable

discussion of content in this manuscript. This work was supported by the National Science

Foundation DEB CAREER Grant #1749764 and by the University of North Carolina to CHM.

**References**
539
540    1      Coyne, J.A. and Orr, H.A. (2004) *Speciation*, Sinauer Associates.
541    2      Mallet, J. *et al.* (2009) Space, sympatry and speciation. *J. Evol. Biol.* 22, 2332–41
542    3      Fitzpatrick, B.M. *et al.* (2009) Pattern, process and geographic modes of speciation. *J. Evol. Biol.*
543           22, 2342–2347
544    4      Fitzpatrick, B.M. *et al.* (2008) What, if anything, is sympatric speciation? *J. Evol. Biol.* 21, 1452–
545           1459
546    5      Bolnick, D.I. and Fitzpatrick, B.M. (2007) Sympatric Speciation: Models and Empirical Evidence.
547           *Annu. Rev. Ecol. Evol. Syst.* 38, 459–487
548    6      Nosil, P. and Schluter, D. (2011) The genes underlying the process of speciation. *Trends Ecol.*
549           *Evol.* 26, 160–167
550    7      Ravinet, M. *et al.* (2017) Interpreting the genomic landscape of speciation: a road map for finding
551           barriers to gene flow. *J. Evol. Biol.* 30, 1450–1477
552    8      Felsenstein, J. (1981) Skepticism towards Santa Rosalia, or why are there so few kinds of animals?
553           *Evolution (N. Y).* 35, 124–138
554    9      Dieckmann, U. and Doebeli, M. (1999) On the origin of species by sympatric speciation. *Nature*
555           400, 354
556    10     Gavrilets, S. (2004) *Fitness landscape and origin of species*, Princeton University Press.
557    11     Doebeli, M. *et al.* (2005) What we have also learned: adaptive speciation is theoretically plausible.
558           *Evolution (N. Y).* 59, 691–699
559    12     Matessi, C. *et al.* (2002) Long-term Buildup of Reproductive Isolation Promoted by Disruptive
560           Selection: How Far Does it Go? *Selection* 2, 41–64
561    13     Bolnick, D.I. and Doebeli, M. (2003) Sexual dimorphism: two sides of the same ecological coin.
562           *Evolution (N. Y).* 57, 2433–2449
563    14     Bolnick, D.I. (2006) Multi-species outcomes in a common model of sympatric speciation. *J.*
564           *Theor. Biol.* 241, 734–744
565    15     Bürger, R. and Schneider, K.A. (2006) Intraspecific Competitive Divergence and Convergence
566           under Assortative Mating. *Am. Nat.* 167, 190–205
567    16     Otto, S.P. *et al.* (2008) Frequency-dependent selection and the evolution of assortative mating.
568           *Genetics* 179, 2091–2112
569    17     Kirkpatrick, M. and Ravigné, V. (2002) Speciation by Natural and Sexual Selection: Models and
570           Experiments. *Am. Nat.* 159, S22–S35
571    18     Turelli, M. *et al.* (2001) Theory and speciation. *Trends Ecol. Evol.* 16, 330–343
572    19     Garner, A.G. *et al.* (2018) Genomic signatures of reinforcement. *Genes (Basel).* 9,
573    20     van Doorn, G.S. and Weissing, F.J. (2002) Ecological versus Sexual Selection Models of
574           Sympatric Speciation: A Synthesis. *Selection* 2, 17–40

21    Seehausen, O. (2004) Hybridization and adaptive radiation. *Trends Ecol. Evol.* 19, 198–207

22    Seehausen, O. (2013) Conditions when hybridization might predispose populations for adaptive
radiation. *J. Evol. Biol.* 26, 279–281

23    Turissini, D.A. and Matute, D.R. (2017) Fine scale mapping of genomic introgressions within the
Drosophila yakuba clade. *PLOS Genet.* 13, e1006971

24    Martin, C.H. (2012) Weak disruptive selection and incomplete phenotypic divergence in two
classic examples of sympatric speciation: Cameroon crater lake cichlids. *Am. Nat.* 180, E90–E109

25    Meier, J.I. *et al.* (2017) Ancient hybridization fuels rapid cichlid fish adaptive radiations. *Nat.*
*Commun.* 8, 1–11

26    Wagner, C.E. and McCune, A.R. (2009) Contrasting patterns of spatial genetic structure in
sympatric rock-dwelling cichlid fishes. *Evolution (N. Y).* 63, 1312–1326

27    Servedio, M.R. and Boughman, J.W. (2017) The Role of Sexual Selection in Local Adaptation and
Speciation. *Annu. Rev. Ecol. Evol. Syst.* 48, annurev-ecolsys-110316-022905

28    Kopp, M. *et al.* (2018) Mechanisms of Assortative Mating in Speciation with Gene Flow:
Connecting Theory and Empirical Research. *Am. Nat.* 191, 1–20

29    Servedio, M.R. *et al.* (2011) Magic traits in speciation: 'magic' but not rare? *Trends Ecol. Evol.*
26, 389–397

30    Turner, G.F. and Burrows, M.T. (1995) A model of sympatric speciation by sexual selection.
*Proc. R. Soc. London. Ser. B Biol. Sci.* 260, 287–292

31    Weissing, F.J. *et al.* (2011) Adaptive speciation theory: A conceptual review. *Behav. Ecol.*
*Sociobiol.* 65, 461–480

32    Kirkpatrick, M. and Barton, N.H. (2006) Chromosome inversions, local adaptation and speciation.
*Genetics* 173, 419–434

33    Feder, J.L. and Nosil, P. (2009) Chromosomal inversions and species differences: When are genes
affecting adaptive divergence and reproductive isolation expected to reside within inversions?
*Evolution (N. Y).* 63, 3061–3075

34    Servedio, M. and Bürger, R. The Effects on Parapatric Divergence of Linkage between Preference
and Trait Loci versus Pleiotropy. , *Genes* , 9. (2018)

35    Feder, J.L. *et al.* (2003) Allopatric genetic origins for sympatric host-plant shifts and race
formation in Rhagoletis. *Proc. Natl. Acad. Sci.* 100, 10314–10319

36    Fuller, Z. *et al.* The role of chromosomal inversions in speciation. *biorxiv.org* at
<https://www.biorxiv.org/content/early/2017/11/24/211771.abstract>

37    Kirkpatrick, M. *et al.* Chromosome inversions, local adaptation and speciation. *Genet. Soc Am.* at
<http://www.genetics.org/content/173/1/419.short>

38    Martin, C.H. (2013) Strong assortative mating by diet, color, size, and morphology but limited
progress toward sympatric speciation in a classic example: Cameroon crater lake cichlids.
*Evolution (N. Y).* 67, 2114–23

39    Bolnick, D.I. (2011) Sympatric Speciation in Threespine Stickleback: Why Not? *Int. J. Ecol.*
2011, 1–15

40    Gavrilets, S. *et al.* (2007) Case studies and mathematical models of ecological speciation. 1.
Cichlids in a crater lake. *Mol. Ecol.* 16, 2893–909

41    Gavrilets, S. and Vose, A. (2007) Case studies and mathematical models of ecological speciation.
2. Palms on an oceanic island. *Mol. Ecol.* 16, 2910–21

42    Gavrilets, S. (2014) Models of speciation: Where are we now? *J. Hered.* 105, 743–755

43    Schliewen, U.K. *et al.* (1994) Sympatric speciation suggested by monophyly of crater lake
cichlids. *Nature* 368, 629–631

44    Schliewen, U.K. and Klee, B. (2004) Reticulate sympatric speciation in Cameroonian crater lake
cichlids. *Front. Zool.* 1, 9–14

45    Barluenga, M. *et al.* (2006) Sympatric speciation in Nicaraguan crater lake cichlid fish. *Nature*
439, 719–723

46    Hahn, M.W. and Nakhleh, L. (2015) Irrational exuberance for resolved species trees. *Evolution (N.*

626              *Y).* DOI: 10.1111/evo.12832

627    47     Wu, C.-I. (2001) The genic view of the process of speciation. *J. Evol. Biol.* 14, 851–865

628    48     McGirr, J.A. and Martin, C.H. (2016) Novel candidate genes underlying extreme trophic
629              specialization in Caribbean pupfishes. *Mol. Biol. Evol.* 34, 873–888

630    49     Martin, S.H. *et al.* (2013) Genome-wide evidence for speciation with gene flow in Heliconius
631              butterflies. *Genome Res.* 23, 1817–1828

632    50     Fontaine, M.C. *et al.* (2015) Extensive introgression in a malaria vector species complex revealed
633              by phylogenomics. *Science (80-. ).* 347, 1258522–1258522

634    51     Campbell, C.R. *et al.* (2018) What is Speciation Genomics? The roles of ecology, gene flow, and
635              genomic architecture in the formation of species. *Biol. J. Linn. Soc.*

636    52     Papadopolus *et al.* (2011) Speciation with gene flow on Lord Howe Island. *Proc. Natl. Acad. Sci.*
637              *U. S. A.* 108, 13188–93

638    53     Geiger, M.F. *et al.* (2013) Crater Lake Apoyo Revisited - Population Genetics of an Emerging
639              Species Flock. *PLoS One* 8, 1–17

640    54     Elmer, K.R. *et al.* (2014) Parallel evolution of Nicaraguan crater lake cichlid fishes via non-
641              parallel routes. *Nat. Commun.* 5, 1–8

642    55     Barluenga, M. and Meyer, A. (2010) Phylogeography, colonization and population history of the
643              Midas cichlid species complex (Amphilophus spp.) in the Nicaraguan crater lakes. *BMC Evol.*
644              *Biol.* 10, 326

645    56     Kusche, H. *et al.* (2015) Sympatric ecological divergence associated with a color polymorphism.
646              *BMC Biol.* 13,

647    57     Pfaender, J. *et al.* (2016) Rugged adaptive landscapes shape a complex, sympatric radiation. *Proc.*
648              *R. Soc. B Biol. Sci.* 283, 11–13

649    58     Hadid, Y. *et al.* (2013) Possible incipient sympatric ecological speciation in blind mole rats
650              (Spalax). *Proc. Natl. Acad. Sci.* 110, 2587–2592

651    59     Elmer, K.R. *et al.* (2010) Rapid evolution and selection inferred from the transcriptomes of
652              sympatric crater lake cichlid fishes. *Mol. Ecol.* 19, 197–211

653    60     Foote, A. *et al.* Genome-culture coevolution promotes rapid divergence of killer whale ecotypes.
654              *nature.com* at <https://www.nature.com/articles/ncomms11693>

655    61     Alcaide, M. *et al.* (2014) Genomic divergence in a ring species complex. *Nature* 511, 83–85

656    62     Igea, J. *et al.* (2015) A comparative analysis of island floras challenges taxonomy-based
657              biogeographical models of speciation. *Evolution (N. Y).* 69, 482–491

658    63     Malinsky, M. *et al.* (2015) Genomic islands of speciation separate Cichlid ectomorphs in an East
659              African crater lake. *Science (80-. ).* 350, 1493–1498

660    64     Martin, C.H. *et al.* (2015) Complex histories of repeated gene flow in Cameroon crater lake
661              cichlids cast doubt on one of the clearest examples of sympatric speciation. *Evolution (N. Y).* 69,
662              1406–1422

663    65     Kautt, A.F. *et al.* (2016) Multispecies Outcomes of Sympatric Speciation after Admixture with the
664              Source Population in Two Radiations of Nicaraguan Crater Lake Cichlids. *PLoS Genet.* 12, 1–33

665    66     Poelstra, J. *et al.* (2018) Speciation in sympatry with ongoing secondary gene flow and an
666              olfactory trigger in a radiation of Cameroon cichlids. *bioRxiv* 229864,

667    67     Richards, E. *et al.* (2018) Don't throw out the sympatric speciation with the crater lake water: fine-
668              scale investigation of introgression provides equivocal support for causal role of secondary gene
669              flow in one of the clearest examples of sympatric speciation. *bioRxiv*

670    68     Machado-Schiaffino, G. *et al.* (2015) Parallel evolution in Ugandan crater lakes: Repeated
671              evolution of limnetic body shapes in haplochromine cichlid fish. *BMC Evol. Biol.* 15,

672    69     Wu, C.-I. and Ting, C.-T. (2004) Genes and speciation. *Nat. Rev. Genet.* 5, 114–122

673    70     Foote, A.D. (2017) Sympatric Speciation in the Genomic Era. *Trends Ecol. Evol.* 33, 85–95

674    71     Martin, C.H. (2016) The cryptic origins of evolutionary novelty : 1000-fold faster trophic
675              diversification rates without increased ecological opportunity or hybrid swarm. DOI:
676              10.1111/evo.13046

72  Dasmahapatra, K.K. *et al.* (2012) Butterfly genome reveals promiscuous exchange of mimicry
adaptations among species. *Nature* 487, 94–98
73  Hedrick, P.W. (2013) Adaptive introgression in animals: Examples and comparison to new
mutation and standing variation as sources of adaptive variation. *Mol. Ecol.* 22, 4606–4618
74  Huerta-Sanchez, E. *et al.* (2014) Altitude adaptation in Tibetans caused by introgression of
Denisovan-like DNA. *Nature* 512, 194–197
75  Stankowski, S. and Streisfeld, M.A. (2015) Introgressive hybridization facilitates adaptive
divergence in a recent radiation of monkeyflowers. *Proc. R. Soc. B Biol. Sci.* 282,
76  Juan, E. *et al.* (2017) Evolution of novel mimicry rings facilitated by adaptive introgression in
tropical butterflies. *Mol. Ecol.* 26, 5160–5172
77  Richards, E.J. and Martin, C.H. (2017) Adaptive introgression from distant Caribbean islands
contributed to the diversification of a microendemic adaptive radiation of trophic specialist
pupfishes. *PLOS Genet.* 13, e1006919
78  Rieseberg, L.H. *et al.* (1999) Transgressive segregation, adaptation and speciation. *Heredity
(Edinb)*. 83, 363–372
79  Kagawa, K. and Takimoto, G. (2017) Hybridization can promote adaptive radiation by means of
transgressive segregation. *Ecol. Lett.* 21, 264–274
80  Schumer, M. *et al.* (2014) How common is homoploid hybrid speciation? *Evolution (N. Y)*. 68,
1553–1560
81  Fishman, L. *et al.* (2013) Chromosomal rearrangements and the genetics of reproductive barriers
in mimulus (monkey flowers). *Evolution (N. Y)*. 67, 2547–2560
82  Guerrero, R.F. and Hahn, M.W. (2017) Speciation as a sieve for ancestral polymorphism. *Mol.
Ecol.* DOI: 10.1111/mec.14290
83  Nelson, T.C. and Cresko, W.A. (2018) Ancient genomic variation underlies repeated ecological
adaptation in young stickleback populations. *Evol. Lett.* 2, 9–21
84  Durand, E.Y. *et al.* (2011) Testing for ancient admixture between closely related populations. *Mol.
Biol. Evol.* 28, 2239–2252
85  Smith, J.M. and Haigh, J. (1974) The hitch-hiking effect of a favourable gene. *Genet. Res.* 23, 23–
35
86  Ravinet, M. *et al.* (2017) Interpreting the genomic landscape of speciation: a road map for finding
barriers to gene flow. *J. Evol. Biol.* 30, 1450–1477
87  Nachman, M.W. and Payseur, B.A. (2012) Recombination rate variation and speciation :
theoretical predictions and empirical results from rabbits and mice. DOI: 10.1098/rstb.2011.0249
88  Cruickshank, T.E. and Hahn, M.W. (2014) Reanalysis suggests that genomic islands of speciation
are due to reduced diversity, not reduced gene flow. *Mol. Ecol.* 23, 3133–3157
89  Bay, R.A. and Ruegg, K. (2017) Genomic islands of divergence or opportunities for introgression?
*Proc. R. Soc. B Biol. Sci.* 284,
90  Clarkson, C.S. *et al.* (2014) Adaptive introgression between Anopheles sibling species eliminates
a major genomic island but not reproductive isolation. *Nat. Commun.* 5, 4248
91  Stölting, K.N. *et al.* (2015) Genome-wide patterns of differentiation and spatially varying selection
between postglacial recolonization lineages of *Populus alba* (Salicaceae), a widespread forest tree.
*New Phytol.* 207, 723–734
92  Felsenstein, J. (1981) Skepticism towards Santa-Rosalia or why are there so few kinds of animals.
*Evolution (N. Y)*. 35, 124–138
93  Ortíz-Barrientos, D. and Noor, M.A.F. (2005) Evidence for a one-allele assortative mating locus.
*Science* 310, 1467
94  Schrider, D.R. and Kern, A.D. (2016) S/HIC: Robust Identification of Soft and Hard Sweeps
Using Machine Learning. *PLOS Genet.* 12, e1005928
95  Kern, A.D. and Schrider, D.R. (2018) diploS/HIC: An Updated Approach to Classifying Selective
Sweeps. *G3 Genes|Genomes|Genetics*
96  Boyle, E.A. *et al.* (2017) An Expanded View of Complex Traits: From Polygenic to Omnigenic.

728   *Cell* 169, 1177–1186

729 97 Siepel, A. *et al.* (2005) Evolutionarily conserved elements in vertebrate, insect, worm, and yeast
730   genomes. *Genome Res.* 15, 1034–1050

731 98 Erickson, P.A. *et al.* (2016) Microinjection for transgenesis and genome editing in threespine
732   sticklebacks. *J. Vis. Exp.* DOI: 10.3791/54055

733 99 Kratochwil, C.F. *et al.* (2017) Tol2 transposon-mediated transgenesis in the Midas cichlid
734   (Amphilophus citrinellus) — towards understanding gene function and regulatory evolution in an
735   ecological model system for rapid phenotypic diversification. *BMC Dev. Biol.* 17, 15

736 100 Hart, J.C. and Miller, C.T. (2017) Sequence-Based Mapping and Genome Editing Reveal
737   Mutations in Stickleback <em>Hps5</em> Cause Oculocutaneous Albinism and the
738   <em>casper</em> Phenotype. *G3 Genes|Genomes|Genetics* 7, 3123 LP-3131

739 101 Cleves, P.A. *et al.* (2018) An intronic enhancer of Bmp6 underlies evolved tooth gain in
740   sticklebacks. *PLOS Genet.* 14, e1007449

741 102 Kronforst, M.R. *et al.* (2013) Hybridization Reveals the Evolving Genomic Architecture of
742   Speciation. *Cell Rep.* 5, 666–677

743 103 Renaut, S. *et al.* (2013) Genomic islands of divergence are not affected by geography of speciation
744   in sunflowers. *Nat. Commun.* 4, 1–8

745 104 Burri, R. *et al.* (2015) Linked selection and recombination rate variation drive the evolution of the
746   genomic landscape of differentiation across the speciation continuum of Ficedula flycatchers.
747   *Genome Res.* 25, 1656–1665

748 105 Irwin, D.E. *et al.* (2016) Recurrent selection explains parallel evolution of genomic regions of high
749   relative but low absolute differentiation in a ring species. *Mol. Ecol.* 25, 4488–4507

750 106 Burri, R. (2017) Interpreting differentiation landscapes in the light of long-term linked selection.
751   *Evol. Lett.* 1, 118–131

752 107 Vijay, N. *et al.* (2017) Genomewide patterns of variation in genetic diversity are shared among
753   populations, species and higher-order taxa. *Mol. Ecol.* 26, 4284–4295

754 108 Van Doren, B.M. *et al.* (2017) Correlated patterns of genetic diversity and differentiation across an
755   avian family. *Mol. Ecol.* 26, 3982–3997

756 109 Ma, T. *et al.* (2018) Ancient polymorphisms and divergence hitchhiking contribute to genomic
757   islands of divergence within a poplar species complex. *Proc. Natl. Acad. Sci.* 115, E236 LP-E243

758 110 Delmore, K.E. *et al.* (2018) Comparative analysis examining patterns of genomic differentiation
759   across multiple episodes of population divergence in birds. *Evol. Lett.* 2, 76–87

760 111 Bush, G.L. (1975) Modes of Animal Speciation. *Annu. Rev. Ecol. Syst.* 6, 339–364

761 112 Sorenson, M.D. *et al.* (2003) Speciation by host switch in brood parasitic indigobirds. 424, 31–32

762 113 Savolainen, V. *et al.* (2006) Sympatric speciation in palms on an oceanic island. *Nature* 441, 210–
763   213

764 114 Kondrashov, A.S. and Kondrashov, F.A. (1999) Interactions among quantitative traits in the
765   course of sympatric speciation. *Nature* 400, 351–354

766 115 Martin, C.H. and Turner, B.J. (2018) Long-distance dispersal over land by fishes: extremely rare
767   ecological events become probable over millennial timescales. *Proc. R. Soc. B Biol. Sci.* 285,

768 116 Martin, C.H. *et al.* (2016) Diabolical survival in Death Valley: recent pupfish colonization, gene
769   flow, and genetic assimilation in the smallest species range on earth. *Proc. R. Soc. B Biol. Sci.* 283,
770   23–34

771 117 A., C.J. and Allen, O.H. (1989) Patterns of speciation in Drosophila. *Evolution (N. Y).* 43, 362–
772   381

773 118 Polechová, J. *et al.* (2005) SPECIATION THROUGH COMPETITION: A CRITICAL REVIEW.
774   *Evolution (N. Y).* 59, 1194–1210

775 119 Norvaišas, P. and Kisdi, E. (2012) Revisiting Santa Rosalia to Unfold a Degeneracy of Classic
776   Models of Speciation. *Am. Nat.* 180, 388–393

777 120 Kopp, M. and Hermisson, J. (2008) Competitive speciation and costs of choosiness. *J. Evol. Biol.*
778   21, 1005–1023

121  Yeaman, S. and Whitlock, M.C. (2011) The genetic architecture of adaptation under migration-
selection balance. *Evolution (N. Y).* 65, 1897–1911

122  Bürger, R. and Akerman, A. (2011) The effects of linkage and gene flow on local adaptation: A
two-locus continent–island model. *Theor. Popul. Biol.* 80, 272–288

123  Akerman, A. and Bürger, R. (2014) The consequences of gene flow for local adaptation and
differentiation: a two-locus two-deme model. *J. Math. Biol.* 68, 1135–1198

124  Yeaman, S. *et al.* (2016) The evolution of genomic islands by increased establishment probability
of linked alleles. *Mol. Ecol.* 25, 2542–2558

125  Cotto, O. and Servedio, M.R. (2017) The Roles of Sexual and Viability Selection in the Evolution
of Incomplete Reproductive Isolation: From Allopatry to Sympatry. *Am. Nat.* 190, 680–693

126  Sachdeva, H. and Barton, N.H. (2017) Divergence and evolution of assortative mating in a
polygenic trait model of speciation with gene flow. *Evolution (N. Y).* 71, 1478–1493

127  Seehausen, O. (2006) African cichlid fish: A model system in adaptive radiation research. *Proc. R.
Soc. B Biol. Sci.* 273, 1987–1998

128  Sparks, J.S. (2004) Molecular phylogeny and biogeography of the Malagasy and South Asian
cichlids (Teleostei: Perciformes: Cichlidae). *Mol. Phylogenet. Evol.* 30, 599–614

129  Martin, C.H. and Wainwright, P.C. (2013) A Remarkable Species Flock of Cyprinodon Pupfishes
Endemic to San Salvador Island, Bahamas. *Bull. Peabody Museum Nat. Hist.* 54, 231–241

130  Ford, A.G.P. *et al.* (2015) High levels of interspecific gene flow in an endemic cichlid fish
adaptive radiation from an extreme lake environment. *Mol. Ecol.* 24, 3421–3440

131  Herder, F. *et al.* (2008) Adaptive sympatric speciation of polychromatic "roundfin" sailfin
silverside fish in Lake Matano (Sulawesi). *Evolution (N. Y).* 62, 2178–2195

132  Sibbing, F.A. *et al.* (1998) Speciation of endemic Lake Tana barbs (Cyprinidae, Ethiopia) driven
by trophic resource partitioning; a molecular and ecomorphological approach. *Aquat. Ecol.* 32,
217–227

133  Elmer, K.R. *et al.* (2010) Rapid sympatric ecological differentiation of crater lake cichlid fishes
within historic times. *BMC Biol.* 8,

134  Kautt, A.F. *et al.* (2018) Lessons from a natural experiment : Allopatric morphological divergence
and sympatric diversification in the Midas cichlid species complex are largely influenced by
ecology in a deterministic way. *Evol. Lett.* DOI: 10.1002/evl3.64

135  Wagner, C.E. *et al.* (2014) Cichlid species-area relationships are shaped by adaptive radiations that
scale with area. *Ecol. Lett.* 17, 583–592

136  Kautt, A.F. *et al.* (2016) Incipient sympatric speciation in Midas cichlid fish from the youngest
and one of the smallest crater lakes in Nicaragua due to differential use of the benthic and limnetic
habitats? *Ecol. Evol.* 6, 5342–5357

137  Kratochwil, C.F. *et al.* (2015) Embryonic and larval development in the Midas cichlid fish species
flock (Amphilophus spp.): A new evo-devo model for the investigation of adaptive novelties and
species differences Evolutionary developmental biology. *BMC Dev. Biol.* 15, 1–15

138  Martin, S.H. *et al.* (2015) Evaluating the use of ABBA-BABA statistics to locate introgressed loci.
*Mol. Biol. Evol.* 32, 244–257

139  Rosenzweig, B.K. *et al.* (2016) Powerful methods for detecting introgressed regions from
population genomic data. *Mol. Ecol.* 25, 2387–2397

140  Schrider, D.R. *et al.* (2018) Supervised machine learning reveals introgressed loci in the genomes
of Drosophila simulans and D. sechellia. *PLOS Genet.* 14, e1007341

141  Hibbins, M.S. and Hahn, M.W. (2018) Population genetic tests for the direction and relative
timing of introgression. *bioRxiv*

142  Elworth, R. *et al.* (2018) DGEN: A Test Statistic for Detection of General Introgression Scenarios.
*bioRxiv*

143  Excoffier, L. *et al.* (2013) Robust Demographic Inference from Genomic and SNP Data. *PLoS
Genet.* 9,

144  Gutenkunst, R.N. *et al.* (2009) Inferring the Joint Demographic History of Multiple Populations

from Multidimensional SNP Frequency Data. *PLOS Genet.* 5, e1000695

145   Pool, J.E. and Nielsen, R. (2009) Inference of Historical Changes in Migration Rate From the
Lengths of Migrant Tracts. *Genetics* 181, 711 LP-719

146   Gravel, S. (2012) Population Genetics Models of Local Ancestry. *Genetics* 191, 607 LP-619

147   Loh, P.-R. *et al.* (2013) Inferring Admixture Histories of Human Populations Using Linkage
Disequilibrium. *Genetics* 193, 1233 LP-1254

148   Vernot, B. and Akey, J.M. (2014) Resurrecting Surviving Neandertal Lineages from Modern
Human Genomes. *Science (80-. ).*

149   Gronau, I. *et al.* (2011) Bayesian inference of ancient human demography from individual genome
sequences. *Nat. Genet.* 43, 1031

150   Patterson, N. *et al.* (2014) The genomic landscape of Neanderthal ancestry in present-day humans.
*Nature* 507, 354–7

151   Prüfer, K. *et al.* (2013) The complete genome sequence of a Neanderthal from the Altai
Mountains. *Nature* 505, 43

152   Seguin-Orlando, A. *et al.* (2014) Genomic structure in Europeans dating back at least 36,200
years. *Science (80-. ).* 346, 1113 LP-1118

153   Thomson, R. *et al.* (2000) Recent common ancestry of human Y chromosomes: Evidence from
DNA sequence data. *Proc. Natl. Acad. Sci.* 97, 7360 LP-7365

154   Tang, H. *et al.* (2002) Frequentist Estimation of Coalescence Times From Nucleotide Sequence
Data Using a Tree-Based Partition. *Genetics* 161, 447 LP-459

155   Hudson, R.R. (2007) The Variance of Coalescent Time Estimates from DNA Sequences. *J. Mol.*
*Evol.* 64, 702–705

156   Coop, G. *et al.* (2008) The Timing of Selection at the Human FOXP2 Gene. *Mol. Biol. Evol.* 25,
1257–1259

157   Peter, B.M. *et al.* (2012) Distinguishing between Selective Sweeps from Standing Variation and
from a De Novo Mutation. *PLOS Genet.* 8, e1003011

158   Beleza, S. *et al.* (2013) The Timing of Pigmentation Lightening in Europeans. *Mol. Biol. Evol.* 30,
24–35

159   Nakagome, S. *et al.* (2016) Estimating the Ages of Selection Signals from Different Epochs in
Human History. *Mol. Biol. Evol.* 33, 657–669

160   Ormond, L. *et al.* (2016) Inferring the age of a fixed beneficial allele. *Mol. Ecol.* 25, 157–169

161   Chen, H. and Slatkin, M. (2013) Inferring Selection Intensity and Allele Age from Multilocus
Haplotype Structure. *G3 Genes,Genomes,Genetics* 3, 1429–1442

162   Chen, H. *et al.* (2015) A hidden Markov model for investigating recent positive selection through
haplotype structure. *Theor. Popul. Biol.* 99, 18–30

163   Smith, J. *et al.* (2018) Estimating time to the common ancestor for a beneficial allele. *Mol. Biol.*
*Evol.* 35, 1003–1017

164   Ekblom, R. and Wolf, J.B.W. (2014) A field guide to whole-genome sequencing, assembly and
annotation. *Evol. Appl.* 7, 1026–1042

165   Matz, M. V (2018) Fantastic Beasts and How To Sequence Them: Ecological Genomics for
Obscure Model Organisms. *Trends Genet.* 34, 121–132

166   Chatterjee, S. and Ahituv, N. (2017) Gene Regulatory Elements, Major Drivers of Human Disease.
*Annu. Rev. Genomics Hum. Genet.* 18, 45–63

167   Witte, J.S. (2010) Genome-Wide Association Studies and Beyond. *Annu. Rev. Public Health* 31,
9–20

168   Price, A.L. *et al.* (2015) Progress and promise in understanding the genetic basis of common
diseases. *Proc. R. Soc. B Biol. Sci.* 282, 20151684

169   Gibson, G. (2018) Population genetics and GWAS: A primer. *PLoS Biol.* 16, 1–6

170   Hart, J.C. and Miller, C.T. (2017) Sequence-based mapping and genome editing reveal mutations
in stickleback Hps5 cause oculocutaneous albinism and the Casper phenotype. *G3*
*Genes|Genomes|Genetics* 7, 3123 LP-3131