Research Article

Screening of candidate substrates and coupling ions of transporters by thermostability

shift assays

Homa Majd[1], Martin S. King[1], Shane M. Palmer[1], Anthony C. Smith[1], Liam D.H. Elbourne[3],

Ian T. Paulsen[3], David Sharples[2], Peter J. F. Henderson[2], Edmund R. S. Kunji[1, *]

[1]Medical Research Council Mitochondrial Biology Unit, University of Cambridge, Wellcome

Trust/MRC Building, Cambridge Biomedical Campus, Hills Road, Cambridge, CB2 0XY, United

Kingdom

[2]Astbury Centre for Structural Molecular Biology and School of BioMedical Sciences,

University of Leeds, Leeds, LS2 9JT, United Kingdom

[3]Department of Molecular Sciences, Macquarie University, Sydney, NSW 2109, Australia

* Author to whom correspondence should be addressed:

Tel.: + 44 1223 252850, Fax.: + 44 1223 252875, e-mail: ek@mrc-mbu.cam.ac.uk

Word count: 3,000 of 5,000 max

Number of figures: 4

Number of tables: 0

Abstract

Substrates of most transport proteins have not been identified, limiting our understanding

of their role in physiology and disease. Traditional identification methods use transport

assays with radioactive compounds, but they are technically challenging and many

compounds are unavailable in radioactive form or are prohibitively expensive, precluding

large-scale trials. Here, we present a high-throughput screening method that can identify

candidate substrates from libraries of unlabeled compounds. The assay is based on the

principle that transport proteins recognize substrates through specific interactions, which

lead to enhanced stabilization of the transporter population in thermostability shift assays.

Representatives of three different transporter (super)families were tested, which differ in

structure as well as transport and ion coupling mechanisms. In each case, the substrates

were identified correctly from a large set of chemically related compounds, including

stereo-isoforms. In some cases, stabilization by substrate binding was enhanced further by

ions, providing testable hypotheses on energy coupling mechanisms.

Introduction

Transport proteins, also called transporters, porters, carriers, translocases or permeases,

encompass a diverse and ubiquitous group of membrane proteins that facilitate the

translocation of ions and molecules across all types of biological membranes, linking

biochemical pathways, maintaining homeostasis and providing building blocks for growth

and maintenance. They comprise 5-15% of the genomic complement in most organisms.

Transport proteins are classified into four major groups; primary active transporters,

secondary active transporters, facilitative transporters, and channels. Human solute

transport proteins have been divided into 65 subfamilies (based on sequence alignments or

experimentally determined substrate specificities), which transport a diverse range of

compounds, including amino acids, sugars, nucleotides, lipids, vitamins, hormones,

inorganic and organic ions, metals, xenobiotics and drugs (http://slc.bioparadigms.org/). Of

the 538 identified solute transport proteins in humans, more than a quarter have no

assigned substrate specificity (http://slc.bioparadigms.org/). Consequently, their role in

human physiology is ill-defined and opportunities for drug intervention are missed (Hediger

et al., 2013). Currently, about half of the transport proteins have been associated with

human disease (Rask-Andersen et al., 2013; Hediger et al., 2013; Lin et al., 2015), but only

12 classes of drugs target them directly, despite their inherent 'druggability' (Hediger et al.,

2013; Lin et al., 2015; Cesar-Razquin et al., 2015; Fauman et al., 2011). The situation is not

better in other eukaryotic organisms and bacterial pathogens, currently on the WHO list

(**Supplementary Table1**) (Elbourne et al., 2017). For most transporters in the sequence

databases the identifications are preliminary because they are based only on sequence

homology without direct experimental evidence for the substrates, even though single

amino acid residue variations in the substrate binding site can alter the specificity

profoundly. Moreover, the substrate specificities of some transporters may have been

incompletely or incorrectly assigned. Finally, there could be membrane proteins with

unassigned function that belong to unidentified transporter families, which are not counted

at all.

One of the major challenges is to find the correct substrates from a large number of

potential candidates. There are approximately ~3,000 metabolites in *E. coli* (Sajed et al.,

2016), ~16,000 in *S. cerevisiae* (Ramirez-Gaona et al., 2017), ~40,000 in *Homo sapiens*

(Wishart et al., 2013), and plants must have even more as they carry out an extensive

secondary metabolism. Some metabolites, such as vitamin B_6, have several interconvertible

species, each of which could be transported. The classical method to identify transport

proteins monitors the accumulation of radiolabeled compounds in whole cells, membrane

vesicles or proteoliposomes. However, these experiments can easily fail when the

expressed transporters are not active due to targeting, insertion or folding issues, when

they are unstable in purification and reconstitution experiments, or when substrate and

coupling ion gradients are not setup correctly. Moreover, some compounds are not

available in radioactive form or are prohibitively expensive, preventing large-scale

identification trials. Given all these technical difficulties, it is often necessary to limit the

number of candidate substrates first by using phylogenetic analysis, by analyzing

phenotypic (patho)physiological data, by complementation studies or by metabolic analysis

of knock-out or mutant strains.

Therefore, there is an unmet demand for the development of new methods to limit

the number of potential substrates for identification of solute carriers. Here, we present a

high-throughput screening method for the identification of substrates of transporters,

which does not require radioactive compounds or prior knowledge. The method uses the

simple concept that transporters recognize their substrates through specific interactions,

enhancing their stabilization in thermostability shift assays. We verify the approach by

defining the substrate specificity of three solute carriers from different bacterial and

eukaryotic protein families and show that these experiments also provide valuable clues

about the ion coupling mechanism.

Results

Principle of the method

Ligands, such as substrates or inhibitors, are recognized by transport proteins through

specific interactions at the exclusion of other molecules. The formation of these additional

bonds leads to an increase in the total number of interactions (Robinson and Kunji, 2006;

Yan, 2017; Yamashita et al., 2005). Consequently, binding leads to an overall increase in the

stability of the ligand-bound species compared to the unliganded species in the population

of protein molecules. We have previously shown that the mitochondrial ADP/ATP carrier

(AAC) from the thermophilic fungus *Thermothelomyces thermophila (*UniProt G2QNH0)

when purified in dodecyl-maltoside is stabilized upon binding of its specific inhibitors

carboxyatractyloside and bongkrekic acid in thermostability shift assays using the thiol-

reactive fluorophore N-[4- (7-diethylamino-4-methyl-3-coumarinyl)-phenyl]-maleimide

(CPM) (Crichton et al., 2015; King et al., 2016). In the assay, the apparent melting

temperature T_m of a population of purified transporters is determined by monitoring the

increase in fluorescence by CPM reacting with thiols that have become exposed due to

thermal denaturation of the proteins (**Fig. 1a**). The apparent melting temperature T_m is the

temperature at which the rate of unfolding for a given population is highest. We tested

whether transported substrates can also cause a shift in thermostability, as their properties

differ quite substantially from those of inhibitors or other tight binders. Transported

substrates bind only transiently and relatively weakly, leading to a conformational change

that in turn causes the release of the substrate on the other side of the membrane, so it

was not obvious that this approach would work. However, the thermostability of the AAC

population was enhanced in the presence of ADP and ATP, but not in the presence of AMP,

which reflects the known substrate specificity of the carrier well (Mifsud et al., 2013) (**Fig.**

1b). This effect is only observed at high concentrations of the substrate, well above the

apparent Km of transport (**Supplementary Fig. 1**). The higher the substrate concentration,

the higher the likelihood that part of the population is prevented from unfolding by binding

of the substrate, leading to the observed shift in thermostability. We reasoned that this

approach could be applied as a screening method to find substrate candidates of

uncharacterized transporters by using compound libraries. To verify the method, we have

tested transporters from three different (super)families, which are distinct in structure and

transport mechanism.

Substrates cause specific thermostability shifts in different transporters

The galactose transporter GalP of *E. coli* (Henderson, 1977) is a prototypical member of the

sugar porter family that belongs to the major facilitator superfamily (MFS) (Pao et al., 1998).

Currently, the MFS contains 24 different transporter families and 320,665 sequence entries

from 5224 species, but the substrates for the vast majority of them have not been formally

identified (Pfam CL0015). In humans, 14 of the 65 solute carrier families belong to the MFS,

and substrates are not known for around 12% of them (http://slc.bioparadigms.org/).

The structure of GalP has not been determined, but those of its mammalian homologs

GLUT1 (Deng et al., 2014), GLUT3 (Deng et al., 2015), and GLUT5 (Nomura et al., 2015) are

available (**Fig. 2a**). GalP contains three cysteine residues, of which only one is readily

accessible to reaction with *N*-ethylmaleimide (McDonald and Henderson, 2001). To

evaluate the strategy, we measured the unfolding curves of purified GalP in dodecyl-

maltoside in the presence of a large number of different sugars. We determined the

temperature shift by subtracting the apparent T_m of unliganded GalP (57.6 ± 0.3 °C) from

the apparent T_m values observed in the presence of compounds (ΔT_m) (**Supplementary Fig.**

2). The GalP population was markedly stabilized by D-glucosamine (ΔT_m; 5.7 ± 0.4 °C), D-

glucose (ΔT_m; 4.2 ± 0.5 °C), D-galactose (ΔT_m; 2.1 ± 0.6 °C), and to a lesser extent by D-talose,

2-deoxy-D-glucose, 6-deoxy-D-glucose. Importantly, the related L-isoforms showed no

significant shift (**Fig. 2b**). These results match the known substrate specificity of GalP well

(Henderson and Maiden, 1990; Walmsley et al., 1994). D-Glucosamine, the most stabilizing

compound, had not been investigated previously, but transport assays showed that this

compound is a new substrate (**Supplementary Fig. 3**), demonstrating that the assay can be

used to discover as well as to extend the substrate specificity of transport proteins.

The transport protein Mhp1 from *Microbacterium liquefaciens* (Uniprot D6R8X8)

transports 5-substituted hydantoins in a sodium-coupled mechanism (Suzuki and

Henderson, 2006; Shimamura et al., 2010) (**Fig 2c**), and is a member of the nucleobase-

cation-symport family of homologous proteins that also transport purines, pyrimidines,

vitamins and related metabolites. The transporter has the LeuT-fold (Shimamura et al.,

2010; Yamashita et al., 2005) and belongs to the amino acid-polyamine-organocation

superfamily (Vastermark and Saier, 2014), which currently contains 20 families and 147,819

sequence entries, the substrates of which have mostly not been identified (Pfam CL0062).

Mhp1 contains three cysteine residues, of which only one is readily accessible to reaction

with *N*-ethylmaleimide (Calabrese et al., 2017). Purified Mhp1 in dodecyl-maltoside had an

apparent T_m of 51.3 °C ± 0.6 °C (**Supplementary Fig. 4**) and to test the binding specificity,

the ΔT_m was determined upon addition of different nucleobases and other compounds (**Fig.**

2d). The only stabilizing compounds were 5-indolylmethyl-L-hydantoin (ΔT_m; 14.1 ± 1.2 °C),

5-benzyl-L-hydantoin (ΔT_m; 13.6 ± 1.2 °C) and 5-bromovinylhydantoin (ΔT_m; 11.9 ± 1.1 °C).

A chemically related inhibitor of Mh1p, 5- (2-naphthylmethyl)-L-hydantoin (L-NMH) also led

to a thermostability shift (ΔT_m; 15.6 ± 1.7 °C). These results match the known substrate and

inhibitor specificity of Mhp1 (Simmons et al., 2014), showing that the assay could identify

substrates from a set of closely related compounds without false positives.

To validate the method further, we also screened the mitochondrial ADP/ATP carrier

from *T. thermophila (*UniProt G2QNH0) (King et al., 2016) (**Fig. 2e**). This transport protein

belongs to the mitochondrial carrier family (MCF), which is involved in the translocation of

chemically diverse compounds across the mitochondrial inner membrane, using uniport,

symport or antiport modes of transport (Kunji, 2012). Currently, there are 89,340 different

sequence entries from 831 different species in the database (Pfam PF00153). In humans,

the MCF is the largest solute carrier family with 53 different members (SLC25), but the

substrate specificity of only half of them has been defined (Kunji, 2012). We screened the

thermostability of purified AAC using a library of mitochondrial compounds

(**Supplementary Fig. 5**). In the presence of ATP, ADP, and dADP the population was

stabilized, showing ΔT_m values of 7.2 ± 0.2, 6.0 ± 0.5 and 1.8 ± 0.2 °C, respectively (**Fig. 1b**

and **2f**). Some other compounds, mostly structurally related nucleotides, also stabilized the

protein, but with significantly smaller shifts (**Fig. 2f**). For each type of nucleotide, the di- and

tri-phosphate species showed larger shifts than the monophosphate forms, similar to the

preference of ATP and ADP over AMP (**Fig. 1d** and **2f**), showing that the assay can also

identify functional groups that are important contributors to substrate binding.

Substrate screening of a mitochondrial carrier from *Tetrahymena thermophila*

To see whether this method can be used to identify candidate substrates for a previously

uncharacterized transporter, we performed a high-throughput screen on a purified

mitochondrial carrier from the thermophilic ciliate *Tetrahymena thermophila (*UniProt

I7M3J0). The carrier is phylogenetically related to the yeast mitochondrial carrier that

transports inorganic phosphate in symport with a proton into the mitochondrial matrix for

ATP synthesis (Runswick et al., 1987), but its substrates have not been identified

experimentally. The population of carriers had an apparent T_m of 56.0 ± 0.8 °C and was

screened against a library of 132 different mitochondrial compounds at pH 6.0 (**Fig. 3**). The

highest shift in the T_m of the population was observed for phosphate (ΔT_m; 4.6 ± 0.4 °C),

followed by glyoxylate (ΔT_m; 3.6 ± 0.8 °C), acetyl-phosphate (ΔT_m; 3.2 ± 0.5 °C) and

phosphoenolpyruvate (ΔT_m; 2.3 ± 0.2 °C) (**Supplementary Fig. 6**), whereas small shifts were

observed for nucleotides.

These compounds either contain phosphate as a functional group or resemble the structure

of phosphate, such as glyoxylate (**Supplementary Fig. 7**). Thus, this assay can be used to

narrow down substantially the number of potential substrate candidates from a library of

unlabeled compounds.

The effect of coupling ions

Secondary active transporters are widespread and often use the electrochemical gradient

of protons or other ions to drive the uptake of substrates against their concentration

gradient. Coupling ions are also recognized by transporters through specific interactions,

often directly associated with the binding of substrates (Kunji and Robinson, 2010;

Krishnamurthy et al., 2009).

5-Benzyl-L-hydantoin is transported by Mhp1 in symport with sodium ions (Suzuki

and Henderson, 2006; Shimamura et al., 2010) and thus we tested whether the

temperature shifts of Mhp1 induced by substrate binding were ion-dependent. When

different ions were tested, only sodium ions induced a large substrate stabilization, whereas

relatively small stabilizations were observed in the presence of potassium, calcium and

magnesium ions (**Fig. 4a**).

Mitochondrial phosphate carriers are proton-driven, coupling phosphate

translocation to proton symport (Kunji and Robinson, 2010). To investigate whether proton-

dependency of substrate binding could be inferred from the assay, the shift in thermal

stability by substrate binding was determined at pH 6.0 and 8.0. The pH can affect the

reaction of maleimide groups of CPM with free thiols and the stability of the protein itself,

but the ΔT_m value shows the effect of pH on substrate binding only. In both cases a

substrate-induced stabilization was observed, but the effect was larger in pH 6.0 than in pH

8.0 (ΔT_m 4.6 ± 0.4 °C and 2.6 ± 0.4 °C, respectively) (**Fig. 4b**).

As a control, we carried out the same experiment with the mitochondrial ADP/ATP

carrier, which is not a proton symporter (**Fig. 4c**). The opposite effect was observed;

substrate binding at pH 8.0 led to a bigger thermostability shift than at pH 6.0. The predicted

substrate binding site of AAC has three positively charged residues that interact with the

negatively charged phosphate groups of ADP and ATP (Kunji and Robinson, 2006; Robinson

and Kunji, 2006). ATP has pKa values of 0.9, 1.5, 2.3 and 7.7, whereas ADP has pKa values

0.9, 2.8 and 6.8. This means that the substrates are more negatively charged at pH 8.0

compared to pH 6.0, which will enhance their binding to the positively charged residues of

the substrate binding site, leading to an increase in thermostability. In the case of the

phosphate carrier the situation is very different. Phosphate has pKa values of 2.2, 7.2 and

12.3, and the predicted binding site contains two positively charged residues to neutralize

the two negative charges on phosphate. However, in addition, the binding site contains a

negatively charged glutamate, which has a pKa of ~4.0. Only when this residue is protonated

can binding of the negatively charged phosphate to the site occur (Kunji and Robinson,

2010), which occurs more readily at pH 6.0 than at pH 8.0, explaining the difference in

thermostability. These results show that substrate binding stabilizes the transporters, but

that this effect is enhanced further by the presence of coupling ions, providing testable

hypotheses on energy coupling mechanisms.

Discussion

We have shown that thermostability assays can be used to study the interaction of

substrates with transporters. This is not intuitive, as substrates only bind transiently and

rather weakly in the transport cycle. Still, for diverse types of transport proteins, which

differ in structure and transport mechanism, a shift in thermostability was observed in the

presence of specific substrates. The shift is observed at concentrations well above the

apparent Km of transport, as under these conditions a larger part of the protein population

is rescued from unfolding by binding of the substrate, increasing the overall number of

interactions.

These studies highlight the different contributions these assays can make to

studying the properties of transport proteins. First, they can be used to limit the number of

potential substrate candidates from libraries of unlabeled compounds, providing important

clues about the substrate specificity (**Fig. 2** and **3**). In the tested cases, the assay correctly

identified the substrates from a set of chemically related compounds, including stereo-

isoforms. Still, these candidates need to be tested by transport or other types of assays, as

the compounds could potentially be inhibitors or regulators. However, no prior knowledge

is required and the number of radioactive compounds that need testing is highly reduced,

meaning that it is easier and cheaper to establish a robust transport assay. Second, the

assays can be used to extend the substrate specificity of known transport proteins, as was

shown for GalP with D-glucosamine (**Fig. 2b**). Third, they can be used to identify functional

groups that are key to substrate binding, as shown for AAC, where nucleotide di- and tri-

phosphates bound more tightly than nucleotide monophosphates (**Fig. 2f**). In the case of

mitochondrial phosphate carriers, the common properties of compounds, such as the

phosphate group, may lead directly to the most probable candidates. Fourth, this assay can

provide important clues on the involvement of coupling ions in substrate binding, as shown

for cases where transport is driven by sodium ion or proton symport (**Fig. 4**).

This method has major advantages over the traditional methods of identification. It

is highly reproducible and can be performed in a high-throughput manner, allowing

screening of about 100 compounds per machine per hour. The assay requires a relatively

small amount of protein per assay (micrograms), depending on the number of buried

cysteine residues. Another advantage is that the unfolding curves are themselves important

quality checks, as they indicate that the protein is folded and therefore competent to bind

substrate.

There are also some limitations, as only proteins that have buried thiols can be used,

although accessible cysteines can be tolerated as they only raise the base line of the assay

without interfering with the unfolding curve. Thiol-containing compounds, such as cysteine

or glutathione, cannot be screened as the probe reacts with them directly. However,

changes in the fluorescence of environmentally-sensitive dyes or endogenous tryptophans

can be used as alternatives to CPM. Furthermore, the assay requires purified and stable

transporter in detergent, although partially purified protein might be sufficient. Advances

in bacterial (King et al., 2015; Ward et al., 1999), yeast (Routledge et al., 2016), insect

(Contreras-Gomez et al., 2014), and mammalian (Goehring et al., 2014; Andrell and Tate,

2013) expression systems have allowed many transporters to be expressed in folded and

functional states. Also, robust purification methods and novel classes of 'stabilizing'

detergents, such as the neopentyl glycol maltoside detergents (Chae et al., 2010), have

allowed even poorly expressed transporters to be purified in the small quantities required

for these studies.

Even though the main purpose of this study was to identify potential substrates of

transporters, the assays may also be used in drug discovery or in the identification of

substrates, inhibitors and regulators of any other soluble or membrane proteins with

occluded cysteines.

Materials and Methods

Materials

Chemicals were obtained from Sigma Aldrich and Thermo Fisher Scientific (USA). Nickel NTA

and sepharose beads were purchased from Qiagen (USA). All enzymes were provided by

New England BioLabs (USA). Lipids were purchased from Avanti Polar Lipids (USA) and

detergents from Anatrace (USA). The hydantoin compounds were a kind gift of Marta Sans,

Maria Kokkinidou and Arwen Pearson (University of Hamburg).

Protein expression

For expression of mitochondrial ADP/ATP carrier of *Thermothelomyces thermophila* and the

putative phosphate carrier of *Tetrahymena thermophila* in yeast mitochondria, gene

constructs were designed to contain an N-terminal tag composed of eight histidine residues

followed by a Thr-Ser-Glu-Asp linker and an Ile-Glu-Gly-Arg Factor Xa protease cleavage site.

The genes were cloned into pYES3/CT vector (Invitrogen, USA) with a constitutively active

promoter (*pMIR* promoter of the *S. cerevisiae* phosphate carrier). The plasmids were

transformed into *S. cerevisiae* WB.12 (MATα ade2-1 trp1-1 ura3-1 can1-100 aac1::LEU2

aac2::HIS3) and W303-1B (MATα leu2-3,112 trp1-1 can1-100 ura3-1 ade2-1 his3-11,15)

strains respectively. Cells were grown in a 50-litre fermenter after which mitochondria were

prepared (Kunji and Harding, 2003).

For expression of GalP in *E. coli* the promoter region and the *galP* gene, which was

modified to encode six histidine residues at the C-terminus of the protein, were cloned into

plasmid pBR322 to form plasmid pGP1, which was transformed into the galactose/glucose

transport-deficient host strain JM1100 (*ptsG ptsM ptsF mgl galP Hfr Δhis gnd thyA galK*)

(Henderson et al., 1977). Cells were grown in basal salts medium supplemented with 30

mM glycerol, 20 µg/ml thymine, 80 µg/ml histidine and 15 µg/ml tetracycline in a fermenter

(30- or 100-liter scale). The gene *hyuP* from *M. liquefaciens*, modified to encode six histidine

residues at the C-terminus, was cloned into plasmid pTTQ18 (Stark, 1987; Suzuki and

Henderson, 2006). The His$_6$-tagged Mhp1 hydantoin transport protein was expressed in *E.*

coli BL21 (DE3) grown in M9 medium supplemented with 20 mM glycerol, 20 mM NH$_4$Cl,

100 µg/ml of carbenicillin, 0.2% w/v casamino acids, 2 mM MgSO$_4$.7H$_2$O, 0.4 mM

CaCl$_2$.2H$_2$O, using induction by IPTG in 100-liter fermenter cultures. In all cases after

harvesting the intact cells preparations were made of inner membranes (Ward et al., 1999),

which were stored at -80 °C in Tris-HCl buffer pH 7.5 until used for purification of each

individual protein.

Protein purification

The mitochondrial ADP/ATP carrier and phosphate carrier were purified using the

detergents dodecyl-maltoside and lauryl maltose neopentyl glycol, respectively, using

established procedures (King et al., 2016).

For the purification of His-tagged GalP, *E. coli* membranes were solubilized in buffer

containing 20 mM Tris-HCl pH 8, 20 mM imidazole, 300 mM NaCl, 20% glycerol and 1%

dodecyl-maltoside for 1 h at 4 °C with gentle mixing to a final protein concentration of 2.5

mg/ml. After centrifugation (108,000 x g, 1 h, 4 °C), the supernatant was mixed with pre-

washed nickel-NTA resin and purified by immobilized nickel affinity chromatography (1 ml

resin per 37 mg of total protein; Superflow, QIAGEN) for batch-binding affinity

chromatography for 1 h at 4 °C. Non-specific proteins were removed with buffer containing

20 mM Tris-HCl pH 8, 20 mM imidazole, 150 mM NaCl, 10% glycerol and 0.02% dodecyl-

maltoside, after which His-tagged GalP was eluted with buffer containing 20 mM Tris-HCl

pH 8, 200 mM imidazole, 100 mM NaCl, 5% glycerol and 0.02% dodecyl-maltoside.

Imidazole was then removed by passing the sample through pre-equilibrated PD-10

desalting column (GE Healthcare). Protein samples were snap-frozen and stored in liquid

nitrogen.

The purification of Mhp1 followed the same procedure as the GalP purification with

the following modifications. Membranes were solubilized for 2 h and incubated with Nickel

NTA resin for 2 h. The wash buffer contained 10 mM Tris-HCl pH 8, 20 mM imidazole, 10%

glycerol and 0.05% dodecyl-maltoside and the elution buffer contained 10 mM Tris-HCl pH

8, 200 mM imidazole, 2.5% glycerol and 0.05% dodecyl-maltoside.

Transport assays

Robotic transport assays were performed using a Hamilton MicroLab Star robot (Hamilton

Robotics Ltd, UK). For GalP transport assays, *E. coli* cells (GalP-expressing strain

pGP1/JM1100 and JM1100 control strains) were diluted to OD_{600} of 20 in MES buffer (5 mM

MES pH 6.6, 150 mM KCl) and incubated at room temperature with 20 mM glycerol for 10

minutes to be energized prior to being loaded into the wells of a MultiScreenHTS+ Hi Flow-

FB (pore size = 1.0 μm, Millipore, USA). Transport assays were initiated by addition of 100

μl of buffer containing 5 μM [^{14}C]-galactose (American Radiolabeled Chemicals, 0.20

GBq/mmol) or 5 μM [^{14}C]-glucosamine (American Radiolabeled Chemicals, 0.204

GBq/mmol). For the competition assay, radiolabeled substrate was mixed with a 4,000-fold

excess of the competitor compound prior to the assay. The transport was stopped after 0,

10, 20, 30, 45, 60, 150 s and, 5, 7.5 and 10 min incubation times with 200 μl ice-cold buffer

and the samples were filtered and washed as above. Levels of radioactivity were measured

by adding 200 μl MicroScint-20 (Perkin Elmer) and measured with a TopCount scintillation

counter (Perkin Elmer).

Preparation of the mitochondrial compound library

To identify mitochondrial metabolites every enzyme in the KEGG metabolic database

(Kanehisa et al., 2017) was evaluated for mitochondrial localization using the MitoMiner

database of mitochondrial localization evidence (Smith and Robinson, 2016). A wide range

of data were considered including large-scale experimental evidence from GFP tagging and

mass-spectrometry of mitochondrial fractions, mitochondrial targeting sequence

predictions, immunofluorescent staining from the Human Protein Atlas (Thul et al., 2017),

and annotation from the Gene Ontology Consortium (The Gene Ontology, 2017). Ensembl

Compara allowed these data to be shared across human, mouse, rat and yeast homologs

(Zerbino et al., 2018). Once a list of enzymes with probable mitochondrial localization was

collated, KEGG was used to produce a corresponding list of potential mitochondrial

compounds. Additional candidates were taken from a computer model of mitochondrial

metabolism that manually partitioned metabolites on either side of the mitochondrial inner

membrane (Smith and Robinson, 2011). Compounds were dissolved in PIPES buffer (10 mM

PIPES pH 7.0, 50 mM NaCl) to a final concentration of 25 mM or in dimethyl sulfoxide to a

final concentration of 100 mM. pH was adjusted if necessary and the stocks were frozen at

-80 °C.

Analysis of the total number of identified transport protein substrates in different species

The TransportDB database contains a large number of organisms for which the transporter

complement has been identified via a bioinformatic pipeline, along with substrate

predictions for the transporters characterized (Elbourne et al., 2017). To acquire the figures

in Supplementary Table 1, a Perl script driving an SQL query of the underlying MySQL

database to TransportDB was developed to quantify those transporters where no prediction

could be made for the substrate for the listed species. The number of transporters with

unassigned specificity represents a minimal number, as the search procedure uses

sequence similarity to characterised transporters as a criterion. However, the substrate

specificity needs to be determined experimentally, as a single mutation in the binding site

can profoundly change substrate recognition.

Thermostability shift assay and library screening

To determine stability, purified protein (typically 1-3 µg) was mixed with 20 µg/ml of the

thiol-reactive fluorophore 7-diethylamino-3- (4'-maleimidylphenyl)-4-methylcoumarin

(CPM) and the fluorescent intensity was measured over a 25-90 °C range of temperature

using a rotatory qPCR multi sample instrument (Rotor-Gene Q, Qiagen, the Netherlands).

Following an initial 18 °C pre-incubation step of 90 s, the temperature was ramped 1 °C

every 15 s, with a 4-s wait between readings, which is equal to a ramp rate of 5.6 °C/min.

The excitation and emission wavelengths were 460 nm and 510 nm, respectively. Five

mg/ml stocks of CPM prepared in dimethyl sulfoxide were diluted to 0.1 mg/ml and

equilibrated in assay buffer for one hour at room temperature in the dark before addition

to the protein sample. The assay buffer was usually the buffer in which the protein was

purified or in a similar buffer with a different pH (MES for pH 6.0, HEPES for pH 8.0) or a

different concentration of other ions. Data analyses and apparent melting temperatures

(T_m, the inflection point of a melting temperature) were determined using software

supplied with the instrument.

For GalP experiments, 500 mM sugar stocks were made in MilliQ water (Merck-

Millipore, USA) as 10 times stocks. For Mhp1 experiments, 100 mM compound stocks were

made in dimethyl sulfoxide as 50 times stocks.

Data analyses and representation

Statistical analyses were performed using Microsoft Excel with the inbuilt function of two-

tailed, two-sample unequal variance Student's *t*-test. The average apparent T_m of 'no ligand'

control samples (three technical repeats within each Rotor-Gene Q run) was subtracted

from the apparent T_m measured for each compound addition in the same run. This assay

was performed with three or five biological repeats using independent batches of purified

protein. The null hypothesis of the *t*-test was that the observed ΔT_m for each compound

was not significantly different from zero.

423
Acknowledgments

This work was supported by grant MC_UU_00015/1 of the Medical Research Council, UK.

H.M. gratefully acknowledges the Cambridge Commonwealth, European and International

Trust for support of her PhD studies. PJFH thanks the Leverhulme Trust for an Emeritus

Research Fellowship (Grant number EM-2014 -045). The fermenters and allied equipment

for protein production in Leeds were funded by the BBSRC (MPSI BBS/B/14418), the

Wellcome Trust (JIF 062164/Z/00/Z) and the University of Leeds. The hydantoin derivatives

were a kind gift from Marta Sans, Maria Kokkinidou and Arwen Pearson (University of

Hamburg).

Author contributions

H.M. and M.S.K. contributed protein purification and thermostability shift assays; H.M.,

M.S.K. contributed transport assays; S.M.P. and D.S. contributed cell culturing; A.C.S.

contributed the mitochondrial compound library; L.D.H.E and I.T.P contributed

bioinformatics analyses of substrate characterization; H.M., M.S.K., P.J.F.H. and E.R.S.K.

contributed the design of the experiments, analyses of the data and writing of the

manuscript.

Competing financial interests

The authors declare no competing interests.

References

Andrell J, Tate CG 2013. Overexpression of membrane proteins in mammalian cells for

structural studies. *Mol. Membr. Biol.* **30:**52-63. 10.3109/09687688.2012.703703

Calabrese AN, Jackson SM, Jones LN, Beckstein O, Heinkel F, Gsponer J, Sharples D, Sans

M, Kokkinidou M, Pearson AR, Radford SE, Ashcroft AE, Henderson PJF 2017.

Topological Dissection of the Membrane Transport Protein Mhp1 Derived from

Cysteine Accessibility and Mass Spectrometry. *Anal. Chem.* **89:**8844-8852.

10.1021/acs.analchem.7b01310

Cesar-Razquin A, Snijder B, Frappier-Brinton T, Isserlin R, Gyimesi G, Bai X, Reithmeier RA,

Hepworth D, Hediger MA, Edwards AM, Superti-Furga G 2015. A Call for Systematic

Research on Solute Carriers. *Cell* **162:**478-87. 10.1016/j.cell.2015.07.022

Chae PS, Rasmussen SG, Rana RR, Gotfryd K, Chandra R, Goren MA, Kruse AC, Nurva S,

Loland CJ, Pierre Y, Drew D, Popot JL, Picot D, Fox BG, Guan L, Gether U, Byrne B,

Kobilka B, Gellman SH 2010. Maltose-neopentyl glycol (MNG) amphiphiles for

solubilization, stabilization and crystallization of membrane proteins. *Nat. Methods*

7:1003-1008. 10.1038/nmeth.1526

Contreras-Gomez A, Sanchez-Miron A, Garcia-Camacho F, Molina-Grima E, Chisti Y 2014.

Protein production using the baculovirus-insect cell expression system. *Biotechnol.*

Prog. **30:**1-18. 10.1002/btpr.1842

Crichton PG, Lee Y, Ruprecht JJ, Cerson E, Thangaratnarajah C, King MS, Kunji ERS 2015.

Trends in thermostability provide information on the nature of substrate, inhibitor,

and lipid interactions with mitochondrial carriers. *J. Biol. Chem.* **290:**8206-8217.

10.1074/jbc.M114.616607

Deng D, Sun P, Yan C, Ke M, Jiang X, Xiong L, Ren W, Hirata K, Yamamoto M, Fan S, Yan N

2015. Molecular basis of ligand recognition and transport by glucose transporters.

Nature **526:**391-6. 10.1038/nature14655

Deng D, Xu C, Sun P, Wu J, Yan C, Hu M, Yan N 2014. Crystal structure of the human

glucose transporter GLUT1. *Nature* **510:**121-5. 10.1038/nature13306

Elbourne LD, Tetu SG, Hassan KA, Paulsen IT 2017. TransportDB 2.0: a database for

exploring membrane transporters in sequenced genomes from all domains of life.

Nucleic Acids Res **45:**D320-D324. 10.1093/nar/gkw1068

Fauman EB, Rai BK, Huang ES 2011. Structure-based druggability assessment--identifying

suitable targets for small molecule therapeutics. *Curr.Opin. Chem. Biol.* **15:**463-

468. 10.1016/j.cbpa.2011.05.020

Goehring A, Lee C-H, Wang KH, Michel JC, Claxton DP, Baconguis I, Althoff T, Fischer S,

Garcia KC, Gouaux E 2014. Screening and large-scale expression of membrane

proteins in mammalian cells for structural studies. *Nat. Protocols* **9:**2574-2585.

10.1038/nprot.2014.173

Hediger MA, Clemencon B, Burrier RE, Bruford EA 2013. The ABCs of membrane

transporters in health and disease (SLC series): introduction. *Mol. Aspects Med.*

34:95-107. 10.1016/j.mam.2012.12.009

Henderson PJ 1977. The multiplicity of components, energization mechanisms and
functions involved in galactose transport into *Escherichia coli*. *Biochem. Soc. Trans.*
5:25-8.

Henderson PJ, Giddens RA, Jones-Mortimer MC 1977. Transport of galactose, glucose and
their molecular analogues by *Escherichia coli* K12. *Biochem. J.* **162:**309-320.

Henderson PJ, Maiden MC 1990. Homologous sugar transport proteins in *Escherichia coli*
and their relatives in both prokaryotes and eukaryotes. *Philos. Trans. R. Soc. Lond.*
B Biol. Sci. **326:**391-410.

Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K 2017. KEGG: new perspectives
on genomes, pathways, diseases and drugs. *Nucleic Acids Res* **45:**D353-D361.
10.1093/nar/gkw1092

King MS, Boes C, Kunji ERS 2015. Membrane Protein Expression in *Lactococcus lactis*.
Methods Enzymol. **556:**77-97. 10.1016/bs.mie.2014.12.009

King MS, Kerr M, Crichton PG, Springett R, Kunji ER 2016. Formation of a cytoplasmic salt
bridge network in the matrix state is a fundamental step in the transport
mechanism of the mitochondrial ADP/ATP carrier. *Biochim. Biophys. Acta* **1857:**14-
22. 10.1016/j.bbabio.2015.09.013

Krishnamurthy H, Piscitelli CL, Gouaux E 2009. Unlocking the molecular secrets of sodium-
coupled transporters. *Nature* **459:**347-55. 10.1038/nature08143

Kunji ERS 2012. Structural and Mechanistic Aspects of Mitochondrial Transport Proteins.
In: FERGUSON, S. (ed.) *Comprehensive Biophysics.* Elsevier.

Kunji ERS, Harding M 2003. Projection structure of the atractyloside-inhibited
mitochondrial ADP/ATP carrier of *Saccharomyces cerevisiae*. *J. Biol. Chem.*
278:36985-36988.

Kunji ERS, Robinson AJ 2006. The conserved substrate binding site of mitochondrial
carriers. *Biochim. Biophys. Acta* **1757:**1237-1248. 10.1016/j.bbabio.2006.03.021

Kunji ERS, Robinson AJ 2010. Coupling of proton and substrate translocation in the
transport cycle of mitochondrial carriers. *Curr. Opin. Struct. Biol.* **20:**440-7.
10.1016/j.sbi.2010.06.004

Lin L, Yee SW, Kim RB, Giacomini KM 2015. SLC transporters as therapeutic targets:

emerging opportunities. *Nat. Rev. Drug Discov.* **14:**543-560. 10.1038/nrd4626

Mcdonald TP, Henderson PJ 2001. Cysteine residues in the D-galactose-H+ symport

protein of *Escherichia coli*: effects of mutagenesis on transport, reaction with N-

ethylmaleimide and antibiotic binding. *Biochem. J.* **353:**709-717.

Mifsud J, Ravaud S, Krammer EM, Chipot C, Kunji ERS, Pebay-Peyroula E, Dehez F 2013.

The substrate specificity of the human ADP/ATP carrier AAC1. *Mol. Membr. Biol.*

30:160-168. Doi 10.3109/09687688.2012.745175

Nomura N, Verdon G, Kang HJ, Shimamura T, Nomura Y, Sonoda Y, Hussien SA, Qureshi

AA, Coincon M, Sato Y, Abe H, Nakada-Nakura Y, Hino T, Arakawa T, Kusano-Arai O,

Iwanari H, Murata T, Kobayashi T, Hamakubo T, Kasahara M, Iwata S, Drew D 2015.

Structure and mechanism of the mammalian fructose transporter GLUT5. *Nature*

526:397-401. 10.1038/nature14909

Pao SS, Paulsen IT, Saier MH, Jr. 1998. Major facilitator superfamily. *Microbiol. Mol. Biol.*

Rev. **62:**1-34.

Ramirez-Gaona M, Marcu A, Pon A, Guo AC, Sajed T, Wishart NA, Karu N, Djoumbou

Feunang Y, Arndt D, Wishart DS 2017. YMDB 2.0: a significantly expanded version

of the yeast metabolome database. *Nucleic Acids Res.* **45:**D440-D445.

10.1093/nar/gkw1058

Rask-Andersen M, Masuram S, Fredriksson R, Schioth HB 2013. Solute carriers as drug

targets: current use, clinical trials and prospective. *Mol. Aspects Med.* **34:**702-10.

10.1016/j.mam.2012.07.015

Robinson AJ, Kunji ERS 2006. Mitochondrial carriers in the cytoplasmic state have a

common substrate binding site. *Proc. Natl. Acad. Sci. U.S.A.* **103:**2617-2622.

Routledge SJ, Mikaliunaite L, Patel A, Clare M, Cartwright SP, Bawa Z, Wilks MDB, Low F,

Hardy D, Rothnie AJ, Bill RM 2016. The synthesis of recombinant membrane

proteins in yeast for structural studies. *Methods* **95:**26-37.

10.1016/j.ymeth.2015.09.027

Runswick MJ, Powell SJ, Nyren P, Walker JE 1987. Sequence of the bovine mitochondrial

544   phosphate carrier protein: structural relationship to ADP/ATP translocase and the

545   brown fat mitochondria uncoupling protein. *EMBO J.* **6:**1367-1373.

546   Sajed T, Marcu A, Ramirez M, Pon A, Guo AC, Knox C, Wilson M, Grant JR, Djoumbou Y,

547   Wishart DS 2016. ECMDB 2.0: A richer resource for understanding the

548   biochemistry of *E. coli*. *Nucleic Acids Res.* **44:**D495-D501. 10.1093/nar/gkv1060

549   Shimamura T, Weyand S, Beckstein O, Rutherford NG, Hadden JM, Sharples D, Sansom

550   MS, Iwata S, Henderson PJ, Cameron AD 2010. Molecular basis of alternating

551   access membrane transport by the sodium-hydantoin transporter Mhp1. *Science*

552   **328:**470-3. 328/5977/470 [pii]

553   10.1126/science.1186303

554   Simmons KJ, Jackson SM, Brueckner F, Patching SG, Beckstein O, Ivanova E, Geng T,

555   Weyand S, Drew D, Lanigan J, Sharples DJ, Sansom MS, Iwata S, Fishwick CW,

556   Johnson AP, Cameron AD, Henderson PJ 2014. Molecular mechanism of ligand

557   recognition by membrane transport protein, Mhp1. *EMBO J.* **33:**1831-1844.

558   10.15252/embj.201387557

559   Smith AC, Robinson AJ 2011. A metabolic model of the mitochondrion and its use in

560   modelling diseases of the tricarboxylic acid cycle. *BMC Syst Biol* **5:**102.

561   10.1186/1752-0509-5-102

562   Smith AC, Robinson AJ 2016. MitoMiner v3.1, an update on the mitochondrial proteomics

563   database. *Nucleic Acids Res* **44:**D1258-61. 10.1093/nar/gkv1001

564   Stark MJ 1987. Multicopy expression vectors carrying the lac repressor gene for regulated

565   high-level expression of genes in *Escherichia coli*. *Gene* **51:**255-267.

566   Suzuki S, Henderson PJ 2006. The hydantoin transport protein from *Microbacterium*

567   *liquefaciens*. *J. Bacteriol.* **188:**3329-3336. 10.1128/JB.188.9.3329-3336.2006

568   The Gene Ontology C 2017. Expansion of the Gene Ontology knowledgebase and

569   resources. *Nucleic Acids Res* **45:**D331-D338. 10.1093/nar/gkw1108

570   Thul PJ, Akesson L, Wiking M, Mahdessian D, Geladaki A, Ait Blal H, Alm T, Asplund A,

571   Bjork L, Breckels LM, Backstrom A, Danielsson F, Fagerberg L, Fall J, Gatto L, Gnann

C, Hober S, Hjelmare M, Johansson F, Lee S, Lindskog C, Mulder J, Mulvey CM,

Nilsson P, Oksvold P, Rockberg J, Schutten R, Schwenk JM, Sivertsson A, Sjostedt E,

Skogs M, Stadler C, Sullivan DP, Tegel H, Winsnes C, Zhang C, Zwahlen M,

Mardinoglu A, Ponten F, Von Feilitzen K, Lilley KS, Uhlen M, Lundberg E 2017. A

subcellular map of the human proteome. *Science* **356**. 10.1126/science.aal3321

Vastermark A, Saier MH, Jr. 2014. Evolutionary relationship between 5+5 and 7+7 inverted

repeat folds within the amino acid-polyamine-organocation superfamily. *Proteins*

82:336-46. 10.1002/prot.24401

Walmsley AR, Martin GE, Henderson PJ 1994. 8-Anilino-1-naphthalenesulfonate is a

fluorescent probe of conformational changes in the D-galactose-H+ symport

protein of *Escherichia coli*. *J. Biol. Chem.* **269:**17009-19.

Ward A, O'reilly J, Rutherford NG, Ferguson SM, Hoyle CK, Palmer SL, Clough JL, Venter H,

Xie H, Litherland GJ, Martin GE, Wood JM, Roberts PE, Groves MA, Liang WJ, Steel

A, Mckeown BJ, Henderson PJ 1999. Expression of prokaryotic membrane

transport proteins in *Escherichia coli*. *Biochem. Soc. Trans.* **27:**893-899.

Wishart DS, Jewison T, Guo AC, Wilson M, Knox C, Liu Y, Djoumbou Y, Mandal R, Aziat F,

Dong E, Bouatra S, Sinelnikov I, Arndt D, Xia J, Liu P, Yallou F, Bjorndahl T, Perez-

Pineiro R, Eisner R, Allen F, Neveu V, Greiner R, Scalbert A 2013. HMDB 3.0--The

Human Metabolome Database in 2013. *Nucleic Acids Res.* **41:**D801-D807.

10.1093/nar/gks1065

Yamashita A, Singh SK, Kawate T, Jin Y, Gouaux E 2005. Crystal structure of a bacterial

homologue of Na (+)/Cl (-)-dependent neurotransmitter transporters. *Nature*.

Yan N 2017. A Glimpse of Membrane Transport through Structures-Advances in the

Structural Biology of the GLUT Glucose Transporters. *J. Mol. Biol.* **429:**2710-2725.

10.1016/j.jmb.2017.07.009

Zerbino DR, Achuthan P, Akanni W, Amode MR, Barrell D, Bhai J, Billis K, Cummins C, Gall

A, Giron CG, Gil L, Gordon L, Haggerty L, Haskell E, Hourlier T, Izuogu OG, Janacek

SH, Juettemann T, To JK, Laird MR, Lavidas I, Liu Z, Loveland JE, Maurel T, Mclaren

W, Moore B, Mudge J, Murphy DN, Newman V, Nuhn M, Ogeh D, Ong CK, Parker A,

Patricio M, Riat HS, Schuilenburg H, Sheppard D, Sparrow H, Taylor K, Thormann A,

Vullo A, Walts B, Zadissa A, Frankish A, Hunt SE, Kostadima M, Langridge N, Martin

FJ, Muffato M, Perry E, Ruffier M, Staines DM, Trevanion SJ, Aken BL, Cunningham

F, Yates A, Flicek P 2018. Ensembl 2018. *Nucleic Acids Res* **46:**D754-D761.

10.1093/nar/gkx1098

Figures

Figure 1. Substrate-induced stabilization of a mitochondrial ADP/ATP carrier. (a) As

protein molecules in a population unfold due to a gradual rise in temperature (25 – 90 °C),

buried cysteine residues become solvent-exposed and accessible to the thiol-specific probe

CPM (blue stick representation) that becomes fluorescent upon reaction. (**b**) Typical

unfolding curves of the mitochondrial ADP/ATP carrier of *Thermothelomyces thermophila*

*(*2 μg) in the absence and presence of AMP, ADP and ATP, shown in red, blue and green,

respectively. (**c**) The apparent melting temperature (T_m) is the peak in the derivative of the

unfolding curve (dF/dT) (below), which is used as an indicator of thermal stability. The

apparent melting temperatures reported in the text are from three independent protein

purifications.

Figure 2. Validation of the method for determining substrate specificity using three unrelated proteins. (a), (**c**) and (**e**) Structural models of GalP (based on GLUT5, PDB 4YBQ), Mhp1 (PDB 2JLN) and AAC (based on Aac2p, PDB 4C9G), respectively. The models are shown in rainbow cartoon and wheat surface representations. Cysteine residues are shown as spheres, except for Cys-19 of GalP, which could not be modeled. (**b**), (**d**) and (**f**) Thermostability screen of GalP (3 μg), Mhp1 (3 μg) and AAC (2 μg) against sugar, nucleobase and nucleotide libraries, respectively. The temperature shift (ΔT_m) is the apparent melting temperature in the presence of compound minus the apparent melting temperature in the absence of compound. The data are represented by the standard deviation of five, three and three independent repeats, respectively. Two-tailed Student's *t*-tests assuming unequal variances were performed for the significance analysis (0.05<*p*-value: not significant; 0.01<*p*-value<0.05: *; 0.001<*p*-value<0.01: **; 0.0001<*p*-value<0.001: ***; *p*-value<0.0001: ****). L-BH, 5-benzyl-L-hydantoin; BVH, 5-bromovinylhydantoin; L-IMH, 5-indolylmethyl-L-hydantoin, L-NMH 5- (2-naphthylmethyl)-L-hydantoin.

Figure 3. Identification of substrates for a mitochondrial phosphate carrier from the thermophilic ciliate *Tetrahymena thermophila*. Purified carrier (1 µg) in lauryl maltose neopentyl glycol was incubated at pH 6.0 with each library compound separately and the ΔT_m was determined, which is the difference between the apparent melting temperatures in the presence and absence of the tested compound. The data are represented by the average and standard deviation of three independent assays. The significance tests were performed as described in the legend to **Fig. 2**.

Figure 4. The effect of coupling ions on the stabilization of transporters by substrate

binding. (a) Thermostability shifts of Mhp1 (3 μg) induced by binding of hydantoins in the

presence of NaCl, KCl, CaCl$_2$ and MgCl$_2$. L-BH: 5-benzyl-L-hydantoin, BVH: 5-

bromovinylhydantoin, L-IMH: 5-indolmethyl-L-hydantoin. (**b**) Thermostability shifts of the

phosphate carrier (1 μg) induced by binding of phosphate-containing compounds at pH 6.0

and 8.0. Acetyl-Pi: acetyl-phosphate, Pi: phosphate. (**c**) Thermostability shifts of the (2 μg)

induced by ADP and ATP binding at pH 6.0 and 8.0, using AMP as control.

Supplementary information

Screening of candidate substrates and coupling ions of

transporters by thermostability shift assays

Homa Majd[1], Martin S. King[1], Shane M. Palmer[1], Anthony C. Smith[1], Liam D.H. Elbourne[3],

Ian T. Paulsen[3], David Sharples[2], Peter J. F. Henderson[2], Edmund R. S. Kunji[1, *]

[1]Medical Research Council Mitochondrial Biology Unit, University of Cambridge, Wellcome

Trust/MRC Building, Cambridge Biomedical Campus, Hills Road, Cambridge, CB2 0XY, United

Kingdom

[2]Astbury Centre for Structural Molecular Biology and School of BioMedical Sciences,

University of Leeds, Leeds, LS2 9JT, United Kingdom

[3]Department of Molecular Sciences, Macquarie University, Sydney, NSW 2109, Australia

 Supplementary Table

 Supplementary Table 1. Status of predictions of transport protein substrate specificity in
 representative archaeal, eubacterial and eukaryotic species (April 2018).

Genus	Species	Transporter Proteins	Proteins with no predicted substrate	Percentage
Acinetobacter	*baumannii*	391	75	19%
Arabidopsis	*thaliana*	1279	311	24%
Caenorhabditis	*elegans*	669	179	27%
Campylobacter	-	197	37	19%
Drosophila	*melanogaster*	662	105	16%
Enterococcus	*faecium*	392	44	11%
Escherichia	*coli*	638	120	19%
Halobacterium	-	208	10	5%
Helicobacter	*pylori*	140	33	24%
Homo	*sapiens*	1472	102	7%
Klebsiella	*pneumoniae*	879	164	19%
Neisseria	*gonorrhoeae*	172	32	19%
Oryza	*sativa*	1285	165	13%
Pseudomonas	*aeruginosa*	703	147	21%
Saccharomyces	*cerevisiae*	343	126	37%
Salmonella	-	605	113	19%
Shigella	-	534	95	18%
Staphylococcus	*aureus*	348	41	12%
Streptococcus	*pneumoniae*	307	50	16%
Sulfolobus	-	205	18	9%
Vibrio	*cholerae*	458	82	18%
	Total	11886	2048	18%

 Supplementary Figures

Supplementary Figure 1. The substrate concentration-dependency of the apparent
melting temperature of purified mitochondrial ADP/ATP carrier. The apparent melting
temperature (T_m) was determined from the peak in the derivative of the unfolding curve for
different concentrations of AMP, ADP, and ATP in 20 mM HEPES pH 8.0, 100 mM NaCl, 0.1
% dodecyl-maltoside, 0.1 mg ml^{-1} tetraoleoyl cardiolipin. The data are represented by the
average and standard deviation of three biological repeats.

Supplementary Figure 2. Substrate-induced stabilization of the galactose transporter
GalP. Typical thermostability profiles of purified GalP (**top**) and their derivatives (dF/dT;
bottom) in the absence and presence of stabilizing substrates D-galactose (red trace), D-
glucose (blue trace) and glucosamine (green trace), which was also causing a significant
shift. The apparent melting temperature (T_m) is the peak in the derivative of the unfolding
curve (dF/dT) and is indicated for each curve. For each reaction, 2.5 µg purified protein was
assayed in the presence of 50 mM compound. The apparent melting temperature reported
in the text is from five independent protein purifications.

696
697

Supplementary Figure 3. Glucosamine is a transported substrate of the galactose
transporter GalP. Transport assays with [^{14}C]-labeled D-glucosamine in whole cells of the
GalP-expressing plasmid pGP1 in the host *E. coli* strain JM1100 (open circles) and control
JM1100 without plasmid (closed circles). Transport in (**a**) the absence or (**b**) presence of
4,000-fold excess of unlabeled D-galactose as described in the "Transport assays" section.
The data are represented by the average and the standard deviation of four technical
repeats.
705

Supplementary Figure 4. Ligand-induced stabilization of Mhp1 in the presence of sodium ions. Typical thermostability profiles of purified Mhp1 (**top**) and their derivatives (dF/dT; **bottom**) in the absence or presence of stabilizing compounds 5-benzyl-L-hydantoin (L-BH, red trace), 5-indolylmethyl-L-hydantoin (L-IMH, blue trace), 5-(2-naphthylmethyl)-L-hydantoin (L-NMH, green trace) and 5-bromovinylhydantoin (BVH, orange trace). The apparent melting temperature (T_m) is the peak in the derivative of the unfolding curve (dF/dT) and is indicated for each curve. For each reaction, 2.5 μg purified protein was assayed in the presence of 140 mM NaCl and 2 mM compound in buffer containing 10 mM Tris-HCl pH 8.0, 2.5% glycerol and 0.05% dodecyl-maltoside. The apparent melting temperature reported in the text is from three independent protein purifications.

Supplementary Figure 5. Substrate screening of the mitochondrial ADP/ATP carrier. Purified ADP/ATP carrier in dodecyl-maltoside was separately mixed with 2.5 mM of library compounds in 20 mM HEPES pH 8.0, 100 mM NaCl, 0.1 % dodecyl-maltoside, 0.1 mg ml^{-1} tetraoleoyl cardiolipin and subjected to a melting regime. The shifts in the apparent T_m (see Online Methods) were recorded and are shown as bars. The data are represented by the average and standard deviation of three independent assays. The significance tests were carried out as in **Figure 2**.

Supplementary Figure 6. Substrate-induced stabilization of the mitochondrial phosphate

carrier. Typical thermostability curves of purified phosphate carrier (**top**) and their

derivatives (dF/dT; **bottom**) in the absence and presence of stabilizing compounds

phosphate (PEP, red trace), glyoxylate (blue trace), acetyl-phosphate (green trace) and

phosphoenolpyruvate (orange trace). The apparent melting temperature (T_m) is the peak in

the derivative of the unfolding curve (dF/dT) and is indicated for each curve. For each

reaction, 2 μg purified protein was assayed in the presence of 2.5 mM compound in 20 mM

MES pH 6.0, 100 mM NaCl, 0.1 % lauryl maltose neopentyl glycol, 0.1 mg ml^{-1} tetraoleoyl

cardiolipin. The apparent melting temperatures reported in the text are from three

independent protein purifications.

phosphate   acetyl phosphate

glyoxylate   phosphoenolpyruvate

739
740
741   **Supplementary Figure 7.** Chemical structures of compounds that showed significant
742   thermostability shifts with the mitochondrial phosphate carrier.
743