Expansion of a core regulon in specialized metabolism by mobile genetic

elements promotes chemical diversity in *Arabidopsis thaliana*

Brenden Barco*, Yoseph Kim**, and Nicole K. Clay*

* Department of Molecular, Cellular & Developmental Biology, Yale University, Kline

Biology Tower 734, 219 Prospect St., New Haven, CT 06511

** Hopkins School, 986 Forest Rd, New Haven, CT 06515

Author Contributions: B.B. and N.K.C performed pathogen assays and ChIP-PCR

experiments. B.B. and Y.K. profiled accessions and species. B.B. performed all other

experiments. B.B. and N.K.C. interpreted the results and wrote the paper.

Abstract

Plant specialized metabolites are ecologically specialized, mostly lineage-specific

molecules whose chemical diversity has been exploited by humans for medical,

agriculture, and industrial applications. The mechanisms that gave rise to these

phenotypic novelties are unclear, particularly those involving the co-option of recently

duplicated genes into functional modules. Here, we show that a LINE retrotransposon

(*EPCOT3*) is responsible for the recruitment of newly duplicated gene *CYP82C2* into

the WRKY33 regulon and the indole-3-carbonylnitrile (ICN) biosynthetic pathway.

WRKY33 is an ancient regulator of plant specialized metabolism, functionally conserved

since the gymnosperm-angiosperm split over 300 million years ago. Preferred WRKY33

binding sites are carried by *EPCOT3*, which inserted upstream of *CYP82C2* and

underwent chromatin remodeling to become an enhancer that coordinately regulates

CYP82C2 gene expression in response to pathogen effectors. The regulatory

neofunctionalization of *CYP82C2* gave rise to pathogen-inducible expression of

species-specific metabolite 4-hydroxy-ICN, which is required for antibacterial defense in

Arabidopsis thaliana. Our results suggest that the transposable element *EPCOT3*

contributed clade/species-specific innovations to a core regulon that functions as an

extended regulon in specialized metabolism and plant innate immunity.

Keywords: specialized/secondary metabolism | regulatory neofunctionalization | LINE

retrotransposon | WRKY33

Summary

Plant secondary or specialized metabolites are essential for plant survival in complex

environments and collectively number in the hundreds of thousands. The genetics and

epigenetics of chemical diversity in plant specialized metabolism remain unclear. Here,

we describe an expansion of the core interactions between an ancient transcription

factor and its target biosynthetic genes by mobile genetic elements that disseminate

transcription factor binding sites in the genome and undergo chromatin remodeling to

become transcriptional enhancers. The extended interactions led to the biosynthesis of

a species-specific antimicrobial metabolite important for plant survival. Our findings

contribute to a growing understanding of chemical innovation, a critically important but

poorly understood process in evolutionary biology.

Plant specialized metabolites exist as adaptations towards co-evolving biotic and

fluctuating abiotic environments. Consequently, plant specialized metabolism is under

constant selective pressure towards chemical innovation (Chae *et al.*, 2014; Weng *et*

al., 2012). The evolutionary process of chemical innovation resulted in the collective

synthesis of hundreds of thousands of chemically diverse and ecologically specialized

metabolites, many of which exhibit narrower taxonomic distributions compared to

primary metabolites (Dixon, 2001; Wink, 2003). Plant specialized metabolic diversity is

thought to evolve through the co-option of pre-existing genes. This is mainly

accomplished via gene duplication of primary or specialized metabolic enzyme-

encoding genes and the neofunctionalization of one or both paralogs to produce

enzymes with new expression patterns and/or protein functions, including an increased

ability to carry out alternate but latent reactions on novel substrates (Ohno, 1970; Force

et al., 1999; Weng *et al.*, 2012).

To respond appropriately to changing environments, plant specialized metabolism must

be highly dynamic and tightly controlled. This is accomplished in large part by

interactions between transcription factors (TFs) and the constituent genes in specialized

metabolic pathways, which are often organized under common TFs into regulons and

thus more co-expressed than those in non-specialized metabolism (Omranian *et al.*,

2015). This organization is thought to fine-tune the timing, amplitude, and tissue-specific

expression of pathway genes and subsequent metabolite accumulation (Grotewold,

2005; Hartmann, 2007; Martin *et al.*, 2010; Tohge & Fernie, 2012). However, very little

is known about how newly duplicated genes enter into these core regulons to promote

specialized metabolic diversity.

Changes in *cis*-regulatory sequences such as enhancers and promoters are a major

driver of phenotypic diversity (Levine and Davidson, 2005; Prud'homme *et al.*, 2007;

Wray, 2007; Wittkopp & Kalay, 2012; Rogers *et al.*, 2013) and likely accelerate the

capture of newly duplicated biosynthetic genes into regulons. Enhancers, which consist

of TF binding sites (TFBSs) that alter the transcription of target genes independent of

orientation and distance (Spitz & Furlong, 2012), are most responsible for *cis*-regulatory

divergence (Wittkopp & Kalay, 2012). Enhancers are derived either through mutation or

transposable element (TE) insertion. TEs can provide TFBSs and other regulatory

innovations upon which selection acts to drive their co-option or exaptation as *cis*-

regulatory elements. The identification of ancient TE co-option or exaptation events by

sequence conservation has led to the hypothesis that TE exaptation events were largely

responsible for a rapid transcriptional rewiring of gene regulatory networks in higher

plants, mammals, and other vertebrates (de Souza *et al.*, 2013; Henaff *et al.*, 2014).

Bacteria elicit two primary immune defense modes in plants, Pattern- and Effector-

triggered immunity (PTI and ETI) (Jones & Dangl, 2006). Pathogenic bacteria

additionally compromise PTI via specific virulence effector proteins (effector-triggered

susceptibility, ETS; Jones & Dangl, 2006). PTI involves the extracellular perception of

conserved molecules known as microbe-associated molecular patterns (MAMPs),

whereas ETI involves the cytosolic perception of effectors. Although ETI results in the

formation of more rapid and robust pathogen-specific response (Jones & Dangl, 2006),

both result in the ability of naïve host cells to generate, through non-self perception and

subsequent transcriptional reprogramming, specialized metabolites necessary for

pathogen defense (Hammerschmidt, 1999; Mansfield, 2000; Clay *et al.*, 2009).

Pathogen-inducible specialized metabolites can be synthesized *de novo* in the plant in

an active state (phytoalexins) or constitutively as precursors that become activated

(phytoanticipins) (VanEtten *et al.*, 1994). Glucosinolates are b-thioglucoside

phytoanticipins produced by plants in the mustard family (Brassicaceae) (Rodman *et al.*,

1998; Mithen *et al.*, 2010). In *A. thaliana,* pathogen inoculation or MAMP treatment

directs the biosynthesis of tryptophan (Trp)-derived indole glucosinolates to 4-

hydroxyindol- and 4-methoxyindol-3-ylmethylglucosionolate (4OH-I3M, 4M-I3M), and

triggers the biosynthesis of phytoalexins 3-thiazol-2'yl-indole (camalexin) and 4-

hydroxyindole-3-carbonylnitrile (4OH-ICN) (Tsuji *et al.*, 1992; Bednarek *et al.*, 2009;

Clay *et al.*, 2009; Rajniak *et al.*, 2015). These molecules are critical for defense

responses against against *Pseudomonas syringae* pv, *tomato* DC3000 (*Pst*) (Clay *et al.*,

2009 Rajniak *et al.*, 2015). Unlike 4M-I3M, whose immune function appears conserved

across Brassicaceae, camalexin biosynthesis is restricted to the Camelineae tribe of

Brassicaceae, thus representing a clade-specific diversification of pathogen-inducible

Trp-derived specialized metabolism (Bednarek *et al.*, 2011). The phylogenetic

conservation of 4OH-ICN biosynthesis has not yet been investigated.

The indole glucosinolate, camalexin and 4OH-ICN biosynthetic pathways share the

conversion of Trp to indole-3-actetaldoxime (IAOx) via the genetically redundant P450

monoxygenases CYP79B2 and CYP79B3 (**Fig. 1a**) (Zhao *et al.*, 2002; Glawischnig *et

al.*, 2004; Rajniak *et al.*, 2015). The camalexin and 4OH-ICN pathways share the

conversion of IAOx to indole-3-cyanohydrin (ICY) by partially redundant P450s

CYP71A12 and CYP71A13 (**Fig. 1a**) (Nafisi *et al.*, 2007; Klein *et al.*, 2013; Rajniak *et

al.*, 2015). CYP71A13 and CYP71B15/PAD3 catalyze further reactions, leading to

camalexin production, whereas the flavin-dependent oxidase FOX1/AtBBE3/Re-Tox1

and P450 CYP82C2 convert ICY to 4OH-ICN (**Fig. 1a**) (Nafisi *et al.*, 2007; Böttcher *et

al.*, 2009; Rajniak *et al.*, 2015).

WRKY transcription factors (TFs) are key regulators of PTI and ETI (Eulgem &

Somssich, 2007), and of plant specialized metabolism (Schluttenhofer & Yuan, 2015).

WRKY33 is the most ancient regulator of plant specialized metabolism; its ortholog in

the green alga *Chlamydomonas reinhardtii* may be ancestral to all higher plant WRKYs

(Rinerson *et al.*, 2015; Schluttenhofer & Yuan, 2015). Predicted WRKY33 orthologs in

higher plants regulate the biosynthesis of lineage-specific alkaloids, terpenes, and

phenylpropanoids (Schluttenhofer & Yuan, 2015). In *A. thaliana*, WRKY33 directly

activates nearly all 4OH-ICN and camalexin pathway genes in response to bacterial

flagellin and fungal pathogen *Botrytis cinerea* (Liu *et al.*, 2015, Birkenbihl *et al.*, 2017)

and may directly activate *CYP71A13* in response to *Pst* and *Pst* carrying the pathogen

effector *avrRpm1* (*Psta*) (Qiu *et al.*, 2008). WRKY TFs bind to the W-box core motif

[TTGAC(T/C)] and require additional motifs to encode binding specificity (Rushton *et al.*,

2010; Liu *et al.*, 2015). WRKY33 has been shown recently to preferentially bind W-

boxes that are within 500nt of the motif [(T/G)TTGAAT] (hereafter referred to as the

WRKY33-specific motif) in response to *B. cinerea* (Liu *et al.*, 2015).

Here, we show that 4OH-ICN is an *A. thaliana*-specific metabolite that is preferentially

metabolized under ETI conditions, displays WRKY33-dependent intraspecific variation,

and is directly correlated with the insertion of a WRKY33-binding LINE retrotransposon

sequence (*EPCOT3*) upstream of *CYP82C2*. Phylogenetic and epigenetic analyses of

related TEs reveal that the preferred WRKY33 binding sequence on *EPCOT3* likely

formed pre-insertion, and that chromatin remodeling occurred post-insertion, leading to

EPCOT3's co-option as a *CYP82C2* enhancer. *EPCOT3* likely contributed to the

expansion of an ancient core regulon in specialized metabolism to accommodate a

clade/species-specific innovation within the conserved framework of pathogen-inducible

Trp-derived metabolism.

Results

4OH-ICN is specific to ETI

To identify the major Trp-derived specialized metabolites induced under ETI in *A.*

thaliana, we compared host transcriptional and metabolic responses to the PTI-eliciting

MAMP flg22, the PTI/ETS-eliciting pathogen *Pst*, and the ETI-eliciting pathogen *Psta*

under similar conditions as those of previous studies (Denoux *et al.*, 2008; Clay *et al.*,

2009). Both flg22 and *Psta* induced genes involved in 4OH-ICN, camalexin and 4M-I3M

biosynthesis, with 4OH-ICN and camalexin biosynthetic genes having a higher level of

induction than those of 4M-I3M in *Psta*-inoculated plants (**SI Appendix, Fig. S1a**;

Denoux *et al.*, 2008). This result is consistent with the largely quantitative differences

observed in transcriptional responses between PTI and ETI (Tao *et al.*, 2003; Navarro

et al., 2004). By contrast, the metabolite responses between PTI and ETI differed

qualitatively. 4OH-I3M and 4M-I3M were present in uninfected plants and accumulated

to modest levels at the expense of parent metabolite I3M in flg22- and *Psta*-inoculated

plants (**SI Appendix, Fig. S1b**) (Clay *et al.*, 2009). By comparison, ICN, 4OH-ICN, and

camalexin were absent in uninfected plants and at trace levels in flg22-inoculated

plants. ICN and camalexin accumulated to high levels in *Pst-* and *Psta*-inoculated

plants, whereas 4OH-ICN solely accumulated to high levels in *Psta*-challenged plants

(**Fig. 1b; SI Appendix, Fig. S1c**). These results suggest that 4OH-I3M, 4M-I3M,

camalexin, and ICN are synthesized in response to multiple PTI elicitors, whereas 4OH-

ICN biosynthesis is specific to ETI.

WRKY33 required and sufficient to activate 4OH-ICN

4OH-ICN biosynthetic genes are highly co-expressed with each other (Rajniak *et al.*,

2015) and with camalexin biosynthetic genes (**SI Appendix, Fig. S1d**), which are in the

WRKY33 regulon (Qiu *et al.*, 2008; Birkenbihl *et al.*, 2012). To determine whether 4OH-

ICN biosynthetic genes are also in the WRKY33 regulon, we compared camalexin, ICN

and 4OH-ICN levels between wild-type and a *wrky33* loss-of-function mutant that

encodes two differently truncated proteins (**Fig. 2a**; Zheng *et al.*, 2006). Consistent with

a previous report (Qiu *et al.*, 2008), *wrky33* was impaired in camalexin biosynthesis in

response to *Psta* and *Pst* carrying the ETI-eliciting pathogen effector *avrRps4* (*Pst*

avrRps4) (**Fig. 2b; SI Appendix, Fig. S2a**). The *wrky33* mutant was similarly impaired

in 4OH-ICN biosynthesis (**Fig. 2b; SI Appendix, Fig. S2a**). These results indicate that

WRKY33 is required for camalexin and 4OH-ICN biosynthesis in response to multiple

ETI elicitors.

To investigate whether *WRKY33* is sufficient to activate the 4OH-ICN pathway, we used

a two-component glucocorticoid-inducible system to generate *wrky33* plants that in the

presence of the glucocorticoid hormone dexamethasone (dex) express a wild-type copy

of WRKY33 with a C-terminal fusion to 1x flag epitope (*wrky33/DEX:WRKY33-flag*; **SI**

Appendix, Figs. S2b-c). Induced expression of *WRKY33-flag* restored camalexin and

4OH-ICN biosynthesis in *Psta*-challenged *wrky33* plants to greater than wild-type levels

(**SI Appendix, Fig. S2d**). These results indicate that *WRKY33* is required and sufficient

to activate camalexin and 4OH-ICN biosynthesis.

Intraspecific variation in *WRKY33* affects 4OH-ICN and immunity

Intraspecific variation in TFs can contribute to gain or loss of phenotypes, such as

branching in maize (Studer *et al.*, 2011) or pelvic loss in three-spined stickleback fish

(Chan *et al.*, 2010). In addition, the wide variation in camalexin biosynthesis reported

among natural accessions of *A. thaliana* (Kagan & Hammerschmidt, 2002) suggests

that a similar variation in 4OH-ICN biosynthesis may exist. To identify additional

transcriptional activators of 4OH-ICN biosynthesis that otherwise might be refractory to

traditional genetic approaches, we compared intraspecific variation in *Psta*-induced

camalexin, ICN and 4OH-ICN among 35 re-sequenced accessions and *wrky33* (Col-0

accession). We found camalexin and 4OH-ICN levels to be positively correlated among

accessions (R^2 = 0.37**; SI Appendix, Fig. S2e**), lending further support to their co-

regulation by WRKY33. Accession Dijon-G (Di-G) was identified to produce less

camalexin and 4OH-ICN and more ICN than its near-isogenic relatives, the Landsberg

accessions L*er*-0 and L*er*-1 (**Fig. 2b; SI Appendix, Figs. S2e-f**). In addition, differences

observed in the metabolite response between Landsberg accessions and Di-G most

closely resembled those between Col-0 and *wrky33* mutant (**Fig. 2b, SI Appendix, Fig.**

S2e). These results led us to hypothesize that genetic variation in a regulatory gene, as

opposed to an immune signaling gene, is responsible for the metabolite phenotypes

observed in Di-G. To test this hypothesis, genetic variation between Di-G and three

sequenced Landsberg accessions (La-0, L*er*-0, and L*er*-1) were used to identify 354

genes that were differentially mutated to high effect in Di-G (**SI Appendix, Fig. S2g**).

Twenty-eight of these mutated Di-G genes were annotated by Gene Ontology to have

roles in defense, including *WRKY33* (**SI Appendix, Table S1**). We confirmed by Sanger

sequencing that Di-G *WRKY33* harbors a nonsense mutation early in the N-terminal

DNA-binding motif (**Fig. 2a**), likely abolishing protein function. Our findings indicate that

camalexin and 4OH-ICN are sensitive to intraspecific variation in *WRKY33*.

Pst infection reduces plant fitness (Kover & Schaal, 2002), and camalexin and 4OH-ICN

promote plant fitness by contributing non-redundantly to disease resistance to *Pst*

(Rajniak *et al.*, 2015). To confirm that disease resistance to *Pst* is also sensitive to

intraspecific variation in *WRKY33*, we measured bacterial growth in adult leaves of

wkry33 and Di-G and their respective (near-)isogenic accessions Col-0 and L*er*-1.

wrky33 and Di-G were more susceptible to *Pst* than their (near)isogenic relatives and

comparable to the 4OH-ICN biosynthetic mutant *cyp82C2* (**Fig. 2c**; Rajniak *et al.*,

2015). In addition, induced expression of *WRKY33-flag* restored wild-type levels of

resistance in *wrky33* (**Fig. 2c**). Together, our results support a specific role of *WRKY33*

in antibacterial defense as an activator of Trp-derived specialized metabolism.

WRKY33 activates 4OH-ICN biosynthesis

To confirm that the 4OH-ICN biosynthetic pathway is in the WRKY33 regulon, we first

compared *WRKY33, CYP71A13, CYP71B15, FOX1* and *CYP82C2* transcript levels

among WT, *wrky33*, and *wrky33/DEX:WRKY33-flag*. Consistent with previous reports

(Qiu *et al.*, 2008), *CYP71A13* and *CYP71B15* expression was down-regulated in *wrky33*

plants in response to *Psta* and upregulated in *wrky33/DEX:WRKY33-flag* (**SI Appendix,**

Fig. S3a). Similarly, *FOX1* and *CYP82C2* expression were unchanged or down-

regulated in *wrky33* plants in response to *Psta*, and significantly upregulated in

wrky33/DEX:WRKY33-flag (**Fig. 3a**). Together with our metabolite analysis, these

findings indicate that WRKY33 mediates camalexin and 4OH-ICN biosynthesis in

response to pathogen effectors.

We then tested for WRKY33 binding to W-box-containing regions upstream of

camalexin and 4OH-ICN biosynthetic genes in dex-treated and *Psta*-infected

wrky33/DEX:WRKY33-flag seedlings by chromatin immunoprecipitation (ChIP)-PCR.

WRKY33 has been shown to bind to a W-box region upstream of *CYP71A12* (Birkenbihl

et al., 2017), a region that also contains three WRKY33-specific motifs and is consistent

with WRKY33's reported binding site preference (Liu *et al.*, 2015). We additionally

observed that *Psta*-induced WRKY33 bound strongly (greater than 5-fold enrichment) to

a single W-box region upstream of *FOX1* and *CYP82C2* (W2 and W4, respectively;

Figs. 3b-c; SI Appendix, Fig. S3b). Both regions also contain one to three WRKY33-

specific motifs. Together with our expression analysis, our findings indicate that

WRKY33 uses preferred WRKY33 binding sites to directly activate 4OH-ICN

biosynthetic genes in response to pathogen effectors.

Interestingly, *Psta*-induced WRKY33 did not bind to the W5 region upstream of

CYP82C2 (**Fig. 3c**), a W-box region that does not contain any WRKY33-specific motifs

and is just upstream of neighboring gene of unknown function *At4g31960* (**Fig. 3b**).

WRKY33 reportedly binds to W5 in response to flg22 and *B. cinerea* (Liu *et al.*, 2015;

Birkenbihl *et al.*, 2017). By contrast, *Psta*-induced WRKY33 bound strongly to W1

region upstream of *CYP71B15* (**SI Appendix, Fig. S3c-d)**, a W-box region that also

does not contain any WRKY33-specific motifs. WRKY33 reportedly binds to a region

encompassing W1 in response to flg22 and *Psta* (Qiu *et al.*, 2008; Birkenbihl *et al.*,

2012). These findings suggest that WRKY33 may use W-box extended motifs or novel

specificity motifs to target camalexin biosynthetic genes in response to pathogen

effectors, or 4OH-ICN biosynthetic genes in response to MAMPs or fungal pathogens.

***CYP82C2* underwent regulatory neofunctionalization**

CYP82C2 catalyzes the last step in 4OH-ICN biosynthesis, hydroxylating ICN to form

4OH-ICN (Rajniak *et al.*, 2015), and likely was the last 4OH-ICN pathway gene to be

recruited to the WRKY33 regulon in *A. thaliana*. To explore the phylogenetic distribution

pattern of 4OH-ICN biosynthesis, we profiled ICN and 4OH-ICN metabolites in close

and distant relatives of *A. thaliana* in response to *Psta*. While ICN biosynthesis was

observed across multiple close relatives, 4OH-ICN was only detected in *A. thaliana*

(**Fig. 4a; SI Appendix, Fig. S4a**). This result suggests that 4OH-ICN manifests a

species-specific diversification of pathogen-inducible Trp-derived metabolism in the

mustard family.

In *A. thaliana*, *CYP82C2* resides in a near-tandem cluster with paralogs *CYP82C3* and

CYP82C4 (**Fig. 4b**). We performed phylogenetic and syntenic analyses to identify

putative *CYP82C2* orthologs in ICN-synthesizing species. All identified homologs are

syntenic to *CYP82C2* or *CYP82C4*, and encode proteins with >88% identity to one

another (**Fig. 4b; SI Appendix, Figs. S4b-c**). *CYP82C3* is present only in *A. thaliana*,

and although more similar to *CYP82C2* than *CYP82C4* in sequence, it is not functionally

redundant with *CYP82C2* (**Fig. 4b; SI Appendix, Fig. S4b**; Rajniak *et al.*, 2015).

CYP82C4 is required for the biosynthesis of sideretin, a widely-conserved

phenylalanine-derived metabolite required for iron acquisition (Rajniak *et al.*, 2018).

CYP82C4 has syntenic orthologs in the mustard family, correlating with the distribution

of sideretin biosynthesis (**Fig. 4b; SI Appendix, Fig. S4b;** Rajniak *et al.*, 2018). By

contrast, *CYP82C2* has syntenic orthologs only within the *Arabidopsis* genus (**Fig. 4b;**

SI Appendix, Fig. S4b). These results suggest that *CYP82C2* duplicated from

CYP82C4 prior to the formation of the *Arabidopsis* genus and then acquired a new

expression pattern and/or catalytic function prior to *A. thaliana* speciation approx. 2

million years later (Hu *et al.*, 2011; Hohmann *et al.*, 2015).

CYP82C2 and CYP82C4 were previously characterized to 5-hydroxylate with equal

efficiency the specialized metabolite 8-methoxypsoralen, a molecule structurally

reminiscent of ICN and sideretin (Kruse *et al.*, 2008). The apparent similarities in

substrate specificity and catalytic function suggest that *CYP82C2* may have diverged

from *CYP82C4* in expression but not protein function. To test this, we first compared the

expression of *CYP82C2* and *CYP82C4* in *A. lyrata* and *A. thaliana* in response to *Psta*.

4OH-ICN biosynthetic genes *CYP79B2, CYP71A12* and *FOX1* were upregulated in both

species, consistent with the common presence of ICN (**Figs. 4a,c**). By contrast,

CYP82C2 levels were respectively upregulated and unchanged in *A. thaliana* and *A.*

lyrata, correlating with the distribution of 4OH-ICN in these species (**Figs. 4a,c**).

CYP82C4 expression was unchanged in both species (**Fig. 4c**). These results indicate

that 4OH-ICN biosynthesis is linked with pathogen-induced expression of *CYP82C2.*

We then compared the aligned upstream sequences of *CYP82C2* and *CYP82C4* in *A.*

lyrata and *A. thaliana* and observed good sequence conservation among orthologs but

poor conservation among paralogs (**SI Appendix, Fig. S4d**), indicating that sequences

upstream of *CYP82C4* and *CYP82C2* were independently derived. We performed

expression analysis in *A. thaliana* to confirm that *CYP82C2* and *CYP82C4* have

different expression patterns. *CYP82C2* expression is upregulated in response to *Psta*

and unchanged under iron deficiency (**Figs. 4c-d; SI Appendix, Fig. S1a;** Rajniak *et*

al., 2015). Conversely, *CYP82C4* is upregulated under iron deficiency and unchanged

in response to *Psta* (**Figs. 4c-d**; Murgia *et al.*, 2011; Rajniak *et al.*, 2018). Finally,

CYP82C4 was unchanged in *Psta*-challenged *wrky33* and *wrky33/DEX:WRKY33-flag*

(**SI Appendix, Fig. S4e**). Our findings suggest that *CYP82C2* diverged from *CYP82C4*

by acquiring WRKY33 regulation for its pathogen-induced expression.

We next assessed dN/dS ratios along branches of the CYP82C phylogenetic tree (**SI**

Appendix, Fig. S4b) and found good support for purifying selection acting on CYP82C

enzymes ($\omega=0.21$), and no support for positive selection acting on CYP82C2/3 enzymes

(**SI Appendix, Table S2**). Lastly, we identified non-conserved amino acid residues

among CYP82C homologs and mapped this information onto a homology model of

CYP82C2. The protein inner core, which encompasses the active site and substrate

channel, is highly conserved among CYP82C homologs (**SI Appendix, Fig. S4g**), and

is consistent with CYP82C2 and CYP82C4's reportedly redundant catalytic functions

(Kruse *et al.*, 2008). Altogether, our findings suggest that *CYP82C2* underwent

regulatory neofunctionalization (Moore & Purugganan, 2005), diverging from *CYP82C4*

in expression but not protein function.

TE *EPCOT3* is an *CYP82C2* enhancer

WRKY33 regulation of *CYP82C2* is mediated by a WRKY33 TFBS in the W4 region

(**Figs. 3, 5a; SI Appendix, Fig. S3c**). Preferential WRKY33 binding at this region

should also be influenced by chromatin features associated with *cis*-regulatory elements

like enhancers and basal promoters (Slattery *et al.*, 2014). To investigate how

CYP82C2 acquired WRKY33 binding for its pathogen-induced expression, we

compared the aligned upstream sequences of *CYP82C* homologs in ICN-synthesizing

species. We observed three large upstream sequences specific to *A. thaliana*

CYP82C2, hereafter named <u>E</u>ighty-two-C2 <u>P</u>romoter <u>C</u>ontained <u>O</u>nly in A. <u>T</u>haliana1-3

(*EPCOT1-3*; **Fig. 5a**). *EPCOT3* in particular is a 240nt region that completely

encompasses W4 (**Fig. 5a**), indicating that the WRKY33's regulation of *CYP82C2* in

response to *Psta* may be species-specific. Further bioinformatics analysis revealed that

EPCOT3 has the epigenetic signature of an active enhancer (Roudier *et al.*, 2011; Liu *et*

al., 2018). Relative to neighboring sequences, *EPCOT3* is enriched with activating

histone mark H3K4me2 and lacks the repressive histone mark H3K27me3 (**Fig. 5b**)

(Heintzman *et al.*, 2007; Hoffman *et al.*, 2010; Roudier *et al.*, 2011; Bonn *et al.*, 2012;

Wang *et al.*, 2014). Our findings suggest that *EPCOT3* functions as an enhancer that

mediates WRKY33 binding and activation of *CYP82C2* in response to pathogen

effectors.

EPCOT3 contains a 3' poly-A tail and is flanked by variable-length target site

duplications (**Fig. 5c; SI Appendix, Fig. S5a**), which are hallmarks of eukaryotic LINE

retrotransposons (Malik *et al.*, 1999). LINE retrotransposition (reverse transcription and

integration) results in frequent 5'-truncation of retrocopies (Luan *et al.*, 1993). We

identified eleven variably truncated retrocopies similar to *EPCOT3* throughout the

genome, including *Ta22*, one of the first LINEs characterized in *A. thaliana* (**Fig. 5c**; **SI**

Appendix, Figs. S5a-b, Table S3; Wright *et al.*, 1996). *EPCOT3*-related LINEs were

sorted into two groups roughly correspondent to their phylogenetic placement:

EPCOT3-LIKE (*EPL*) for those with high identity (>65%) to *EPCOT3* and *Ta22* or *Ta22-*

LIKE (*Ta22L*) for the remainder (**SI Appendix, Fig. S5a; Table S3**). Only *Ta22* and

Ta22L1 are full-length LINEs (**Fig. 5c**), presumably encoding the proteins necessary for

their own transposition and for the transposition of nonautonomous family members like

EPCOT3. We also identified two syntenic species-specific *Ta22Ls*, but no *EPLs*, in *A.*

lyrata (**SI Appendix, Table S3**). Given the 80% overall sequence identity between *A.*

thaliana and *A. lyrata* (Hu *et al.*, 2011), this data indicates that *EPCOT3* and *EPLs*

arose from retrotransposition following the speciation of *A. thaliana.*

Of all the retrocopies, *EPL1* is most similar to *EPCOT3* (85.4% identity), sharing the W-

box and WRKY33-specific motif, whereas *EPL2* is less similar (67%) and lacks the

WRKY33-specific motif (**Fig. 5c; SI Appendix, Table S3, Fig. S5a**). *EPL1* and *EPL2*

are much less truncated than *EPCOT3* (**Fig. 5c**), and lack epigenetic signatures typical

of *cis*-regulatory sequences (**SI Appendix, Fig. S5c**) (Roudier *et al.*, 2011; Liu *et al.*,

2018). To investigate whether the sequence information and chromatin features

associated with *EPL*s are sufficient for WRKY33 binding, we tested for WRKY33 binding

to *EPL* sequences homologous to the W4 region of *EPCOT3* in dex-treated, *Psta*-

infected *wrky33/DEX:WRKY33-flag* plants by ChIP-(q)PCR. Compared to *EPCOT3*

(**Fig. 3c**), WRKY33 respectively bound weakly or not at all to *EPL1* and *EPL2* (**Fig. 5d;**

SI Appendix, Fig. S5d). Our findings suggest the following history: (1) *EPL1* likely

retroduplicated from *EPL2* or its progenitor, which already contained a W-box; (2) *EPL1*

then acquired a WRKY33-specific motif by mutation; (3) *EPCOT3* likely retroduplicated

from *EPL1* and then acquired epigenetic signatures of an enhancer, thereby allowing

selection to act on standing variation rather than *de novo* mutation for *CYP82C2*

recruitment into the 4OH-ICN biosynthetic pathway.

Discussion

TEs were originally conceived to act as "controlling elements" of several loci in the

genome (McClintock, 1956), and are now well understood to provide TFBSs and other

innovations upon which selection acts to drive their exaptation as *cis*-regulatory

elements. The identification of ancient TE-mediated exaptation events by sequence

indicates TEs were largely responsible for the rapid transcriptional rewiring of gene

regulatory networks in many eukaryotes (de Souza *et al.*, 2013). Relatively recent

exaptations of TEs into enhancers of host genes have been described for the beta-like

globin and IFNL1 genes in primates, Cyp6g1 in fruit flies, and tb1 in maize (Pi *et al.*,

2004; Thomson *et al.*, 2009; Schmidt *et al.*, 2010; Studer *et al.*, 2011; de Souza *et al.*,

2013). However, it is unclear whether TE exaptations contribute to physiological

changes that result in interspecific variation/innovation and fitness benefit (de Souza *et*

al., 2013). In this study, we show that *EPCOT3* is a TE-derived enhancer that mediates

WRKY33 binding and activation of *CYP82C2,* leading to 4OH-ICN biosynthesis and

increased disease resistance to a bacterial pathogen. This is to date the first report of a

recent TE exaptation event that resulted in a clade-specific expansion of a core regulon

in plant specialized metabolism.

Although the *EPL1/EPCOT3* progenitor retrotransposed a preferred WRKY33 TFBS in

the form of *EPCOT3* upstream of *CYP82C2*, a further series of epigenetic modifications

were needed to facilitate optimal access of *EPCOT3* by WRKY33. *EPL1* exists in a

silenced heterochromatin state (**SI Appendix, Fig. S5c**), typical for TEs (Slotkin &

Martienssen, 2007), and is bound weakly by WRKY33 (**Fig. 5d**), whereas *EPCOT3* is in

an open chromatin state (**Fig. 5b**; Roudier *et al.*, 2011; Liu *et al.*, 2018) and bound

strongly by WRKY33 (**Fig. 3c**). The more severe 5'-truncation of *EPCOT3* could

account for its release from TE silencing mechanisms, and the initially weak WRKY33

binding could provide a 'seed' for chromatin remodelers to drive the exaptation of newly

retrotransposed *EPCOT3* into a *bona fide* enhancer. Further epigenomic sampling

within *Arabidopsis* is needed to better clarify epigenetic transformations underlying the

EPCOT3 exaptation event.

Accession Di-G is closely related to Landsberg accessions (SI Appendix, Figs. S2e-g;

Hardtke *et al.*, 1996), but synthesizes less camalexin and 4OH-ICN (**Fig. 2b**; Kagan &

Hammerschmidt, 2002), is more susceptible to a range of bacterial and fungal

pathogens (**Fig. 2c**) (Hugouvieux *et al.*, 1998; Kagan & Hammerschmidt, 2002;

Mukherjee *et al.*, 2009), and is more sensitive to ethylene phytohormone (Chatfield *et*

al., 2008). WRKY33 has been implicated in camalexin biosynthesis (Qiu *et al.*, 2008),

antifungal defense (Zheng *et al.*, 2006), and ethylene biosynthesis (Li *et al.*, 2012). We

identified WRKY33 as causal for some if not all of these phenotypes in Di-G. This is the

first report of WRKY33's involvement in antibacterial defense and is consistent with the

contribution of camalexin and 4OH-ICN towards antibacterial defense (Rajniak *et al.*,

2015).

WRKY33 is an ancient transcription factor responsible for many fitness-promoting traits

in plants, thus it is unexpected that an *A. thaliana* accession would have a naturally

occurring *wrky33* mutation (C536T transversion). Di-G is the sole member of 1,135

sequenced accessions to have a high-effect single nucleotide polymorphism (SNP) in

WRKY33 (1001 Genomes Consortium, 2016). Di-G and L*er*-0 have long been models

for studies in mutagenesis (Rédei, 1962, Müller, 1966), and thus a possibility exists that

Di-G may have originated from an ethyl methanesulfonate (EMS) mutagenesis screen

of L*er*-0. Historical EMS mutagenesis experiments generated upwards of tens of

thousands of mutations per cell (Müller 1966; Rédei & Koncz, 1993; Camara *et al.*,

2000), well within the range of ~25,000 SNPs that are not concordant between Di-G and

L*er*-0 (**SI Appendix, Fig. S2f**). These findings open the possibility that Di-G is not a

natural accession but an artificially-derived L*er*-0 *wrky33* mutant.

METHODS

Details of plant materials and growth conditions, plant binary vector construction and

transformation, bacterial infection assays, RNA extraction and qPCR analysis,

extraction and LC-DAD-MS analysis of indole phytoalexins, extraction and LC-DAD-

FLD-MS analysis of glucosinolates, chromatin immunoprecipitation and (q)PCR,

phylogenetic analysis and bioinformatics analysis can be found in **SI Appendix**.

ACKNOWLEDGEMENTS

We thank E.S. Sattely for ICN/ICN-ME, 4OH-ICA/4OH-ICA-ME and camalexin

standards. This work was supported by T32-GM007499 (to B.B.) and

Elsevier/Phytochemistry Young Investigator Award (to N.K.C.).

REFERENCES

1001 Genomes Consortium (2016) 1,135 Genomes Reveal the Global Pattern of

Polymorphism in Arabidopsis thaliana. Cell, 166(2), 481-491.

Birkenbihl RP, Diezel C, Somssich, IE (2012) Arabidopsis WRKY33 is a key

transcriptional regulator of hormonal and metabolic responses toward Botrytis cinerea

infection. Plant Physiology, 159(1), 266-285.

Birkenbihl RP, Kracher B, Somssich, IE (2017) Induced Genome-Wide Binding of Three

Arabidopsis WRKY Transcription Factors during Early MAMP-Triggered Immunity. The

Plant Cell, 29(1), 20-38.

Bonn S, *et al.* (2012) Tissue-specific analysis of chromatin state identifies temporal

signatures of enhancer activity during embryonic development. Nat Genet 44(2): 148-

156.

Böttcher C, *et al.* (2009) The multifunctional enzyme CYP71B15 (PHYTOALEXIN

DEFICIENT3) converts cysteine-indole-3-acetonitrile to camalexin in the indole-3-

acetonitrile metabolic network of Arabidopsis thaliana. Plant Cell 21(6): 1830-1845.

Camara, MD, Ancell CA, Pigliucci M (2000) Induced mutations: a novel tool to study

phenotypic integration and evolutionary constraints in Arabidopsis thaliana. Evolutionary

Ecology Research, 2(8), 1009-1029.

Chae L, Kim T, Nilo-Poyanco R, Rhee SY (2014) Genomic signatures of specialized

metabolism in plants. Science 344: 510-513.

Chatfield SP, Raizada MM (2008) Ethylene and shoot regeneration: hookless1

modulates de novo shoot organogenesis in Arabidopsis thaliana. Plant Cell Reports,

27(4), 655-666.

Chan YF, *et al*. (2010) Adaptive evolution of pelvic reduction in sticklebacks by recurrent

deletion of a Pitx1 enhancer. Science 237: 302-305.

Clay NK, Adio AM, Denoux C, Jander G, Ausubel FM (2009) Glucosinolate metabolites

required for an Arabidopsis innate immune response. Science 323: 95-101.

de Souza, FS. J Franchini, LF Rubinstein M (2013) Exaptation of transposable elements

into novel cis-regulatory elements: is the evidence always strong? Mol Biol Evol 30(6),

1239-1251.

Dixon RA (2001) Natural products and plant disease resistance. Nature 411: 843-847.

Denoux C, *et al*. (2008) Activation of defense response pathways by OGs and Flg22

elicitors in Arabidopsis seedlings. Mol Plant 1: 423-445.

Eulgem T, Somssich IE (2007) Networks of WRKY transcription factors in defense

signaling. Curr Opin Plant Biol 10: 366-371.

Force AM, *et al*. (1999) Preservation of duplicate genes by complementary,

degenerative mutations. Genetics 151: 1531-1545.

Glawischnig E, Hansen BG, Olsen CE, Halkier BA (2004) Camalexin is

synthesized from indole-3-acetaldoxime, a key branching point between primary and

secondary metabolism in Arabidopsis. Proc Natl Acad Sci USA 101: 8245-8250.

Grotewold E (2005) Plant metabolic diversity: a regulatory perspective. Trends Plant Sci

10: 57-62.

Greene EA, *et al*. (2003) Spectrum of chemically induced mutations from a large-scale

reverse-genetic screen in Arabidopsis. Genetics, 164(2), 731-740.

Hammerschmidt R (1999) PHYTOALEXINS: What have we learned after 60 years?

Annu Rev Phytopathol 37: 285-306.

Hardtke CS, Müller J, Berleth T (1996) Genetic similarity among Arabidopsis thaliana

ecotypes estimated by DNA sequence comparison. Plant Molecular Biology, 32(5), 915-

922.

Hartmann T (2007) From waste products to ecochemicals: fifty years research of plant

secondary metabolism. Phytochemistry 68: 2831-2846.

Henaff E, *et al*. (2014) Extensive amplification of the E2F transcription factor binding

sites by transposons during evolution of Brassica species. Plant J 77: 852-862.

Heintzman ND, *et al*. (2007) Distinct and predictive chromatin signatures of

transcriptional promoters and enhancers in the human genome. Nat Genet, 39(3), 311.

Hoffman BG, *et al*. (2010) Locus co-occupancy, nucleosome positioning, and H3K4me1

regulate the functionality of FOXA2-, HNF4A-, and PDX1-bound loci in islets and liver.

Genome Res 20(8): 1037-1061.

Hohmann N, Wolf EM, Lysak MA, Koch MA (2015) A time-calibrated road map of

brassicaceae species radiation and evolutionary history. Plant Cell 27(10): 2770-2784.

Hu TT, *et al*. (2011) The Arabidopsis lyrata genome sequence and the basis of rapid

genome size change. Nat Genet 43(5): 476-481.

Hugouvieux V, Barber, CE, Daniels, MM (1998) Entry of Xanthomonas campestris pv.

campestris into hydathodes of Arabidopsis thaliana leaves: a system for studying early

infection events in bacterial pathogenesis. MPMI, 11(6), 537-543.

Jones JDG, Dangl JL (2006) The plant immune system. Nature 444: 323-329.

Kagan IA, Hammerschmidt R (2002) Arabidopsis ecotype variability in camalexin

production and reaction to infection by Alternaria brassicicola. J Chem Ecol 28(11):

2121-2140.

Klein AP, Anarat-Cappillino G, Sattely ES (2013) Minimum set of cytochromes P450 for

reconstituting the biosynthesis of camalexin, a major Arabidopsis antibiotic.

Angewandte Chemie 52: 13625-13628.

Kover PX, Schaal BA (2002) Genetic variation for disease resistance and tolerance

among Arabidopsis thaliana accessions. Proc Natl Acad Sci USA 99(17): 11270-11274.

Kruse T, *et al*. (2008) In planta biocatalysis screen of P450s identifies 8-

methoxypsoralen as a substrate for the CYP82C subfamily yielding original chemical

structures. Chem Biol 15: 149-156.

Levine M, Davidson EH (2005) Gene regulatory networks for development. Proc Natl

Acad Sci USA 102: 4936-4942.

Li G, *et al.* (2012). Dual-level regulation of ACC synthase activity by MPK3/MPK6

cascade and its downstream WRKY transcription factor during ethylene induction in

Arabidopsis. PLoS genetics, 8(6), e1002767.

Liu S, Kracher B, Ziegler J, Birkenbihl RP, Somssich IE (2015) Negative regulation of

ABA signaling by WRKY33 is critical for Arabidopsis immunity towards Botrytis cinerea

2100. Elife 4: e07295.

Liu Y, *et al.* (2018) PCSD: a plant chromatin state database. Nucleic Acids Research,

46(D1), D1157-D1167.

Luan DD, Korman MH, Jakubczak JL, Eickbush TH (1993) Reverse transcription of

R2Bm RNA is primed by a nick at the chromosomal target site: a mechanism for non-

LTR retrotransposition. Cell 72: 595-605.

Malik HS, Burke WD, Eickbush TH (1999) The age and evolution of non-LTR

retrotransposable elements. Mol Biol Evol 16: 793-805.

Mansfield JW (2000) Antimicrobial compounds and resistance: the role of phytoalexins

and phytoanticipins. In Mechanisms of Resistance to Plant Diseases, A. Slusarenko,

RSS Fraser and LC. van Loon eds (Kluwer Academic Publishers Dordrecht, The

Netherlands), pp. 325-370.

Martin C, Ellis N, Rook F (2010) Do transcription factors play special roles in adaptive

variation? Plant Physiol 154: 506-511.

McClintock B (1956) Controlling elements and the gene. Cold Spring Harb Symp Quant

Biol 21: 197-216.

Mithen R, Bennett R, Marquez J (2010) Glucosinolate biochemical diversity and

innovation in the Brassicales. Phytochemistry 71: 2074-2086.

Moore RC, Purugganan MD (2005) The evolutionary dynamics of plant duplicate genes.

Curr Opin Plant Biol, 8(2), 122-128.

Mukherjee AK, Lev S, Gepstein S, Horwitz, BB (2009) A compatible interaction of

Alternaria brassicicola with Arabidopsis thaliana ecotype DiG: evidence for a specific

transcriptional signature. BMC Plant Biology, 9(1), 31.

Müller, AA (1966) Die Induktion von rezessiven Letalmutationen durch

Äthylmethansulfonat beiArabidopsis. Theoretical and Applied Genetics, 36(5), 201-220.

Murgia I, Tarantino D, Soave C, Morandini P (2011) Arabidopsis CYP82C4 expression

is dependent on Fe availability and circadian rhythm, and correlates with genes involved

in the early Fe deficiency response. J Plant Physiol 168: 894-902.

Nafisi M, *et al.* (2007) Arabidopsis cytochrome P450 monooxygenase 71A13 catalyzes

the conversion of indole-3-acetaldoxime in camalexin synthesis. Plant Cell 19: 2039-

2052.

Navarro L, *et al.* (2004) The transcriptional innate immune response to flg22. Interplay

and overlap with Avr gene-dependent defense responses and bacterial pathogenesis.

Plant Physiol 135: 1113-1128.

Ohno S (1970) Evolution by Gene Duplication. Springer-Verlag: Heidelberg, Germany.

Omranian N, *et al.* (2015) Differential metabolic and coexpression networks of plant

metabolism. Trends Plant Sci 20: 266-268

Pi W, *et al.* (2004) The LTR enhancer of ERV-9 human endogenous retrovirus is active

in oocytes and progenitor cells in transgenic zebrafish and humans. Proc Natl Acad Sci

USA 101: 805-810.

Prud'homme B, Gompel N, Carroll SB (2007) Emerging principles of regulatory

evolution. Proc Natl Acad Sci USA 104 Suppl 1: 8605-8612.

Qiu JL, *et al*. (2008) Arabidopsis MAP kinase 4 regulates gene expression through

transcription factor release in the nucleus. EMBO J 27: 2214-2221.

Rajniak J, Barco B, Clay NK, Sattely ES (2015) A new cyanogenic metabolite in

Arabidopsis required for inducible pathogen defence. Nature 525: 376-379.

Rajniak J, *et al*. (2018) Biosynthesis of redox-active metabolites in response to iron

deficiency in plants. Nat Chem Biol 14: 442-450.

Rédei GP (1962) Single locus heterosis. Zeitschrift für Vererbungslehre, 93(1), 164-170.

Rédei GP, Koncz C (1993) Classical mutagenesis. In *Methods in Arabidopsis research*

(pp. 16-82).

Rinerson CI, Rabara R, Tripathi P, Shen QJ, Rushton PJ (2015) The evolution of WRKY

transcription factors. BMC Plant Biology 15: 66.

Rodman J, Soltis R, Soltis D, Sytsma K, Karol K. (1998) Parallel evolution of

glucosinolate biosynthesis inferred from congruent nuclear and plastid gene

phylogenies. Am J Bot 85: 997.

Rogers WA, *et al.* (2013) Recurrent modification of a conserved cis-regulatory element

underlies fruit fly pigmentation diversity. PLoS Genetics, 9(8), e1003740.

Roudier F, *et al.* (2011) Integrative epigenomic mapping defines four main chromatin

states in Arabidopsis. EMBO J 30(10): 1928-1938.

Rushton PJ, Somssich IE, Ringler P, Shen QJ (2010) WRKY transcription factors.

Trends Plant Sci 15: 247-258.

Schluttenhofer C, Yuan L (2015) Regulation of specialized metabolism by WRKY

transcription factors. Plant Physiol 167: 295-306.

Schmidt JM, *et al.* (2010) Copy number variation and transposable elements feature in

recent, ongoing adaptation at the Cyp6g1 locus. PLoS Genetics 175: 1071-1077.

Slattery M, *et al.* (2014) Absence of a simple code: how transcription factors read the

genome. Trends Biochem Sci 39: 381-399.

Slotkin, R. K, Martienssen R (2007) Transposable elements and the epigenetic

regulation of the genome. Nat Rev Genet, 8(4), 272.

Spitz, F. Furlong EE (2012) Transcription factors: from enhancer binding to

developmental control. Nat Rev Genet 13: 613-626.

Studer A, Zhao Q, Ross-Ibarra J, Doebley J (2011) Identification of a functional

transposon insertion in the maize domestication gene tb1. Nat Genet 43: 1160-1163.

Tam OH, *et al*. (2008) Pseudogene-derived small interfering RNAs regulate gene

expression in mouse oocytes. Nature 453: 534-538.

Tao Y, *et al*. (2003) Quantitative nature of Arabidopsis responses during compatible and

incompatible interactions with the bacterial pathogen Pseudomonas syringae. Plant Cell

15: 317-330.

Thomson SJ, *et al*. (2009) The role of transposable elements in the regulation of IFN-λ1

gene expression. Proc Natl Acad Sci USA 106(28): 11564-11569.

Tsuji J, Jackson EP, Gage DA, Hammerschmidt R, Somerville SC (1992)

Phytoalexin accumulation in Arabidopsis thaliana during the hypersensitive reaction to

Pseudomonas syringae pv syringae. Plant Physiol 98: 1304-1309.

Watanabe T, *et al.* (2008) Endogenous siRNAs from naturally formed dsRNAs regulate

transcripts in mouse oocytes. Nature 453: 539-543.

Tohge T, Fernie AR (2012) Co-expression and co-responses: within and beyond

transcription. Front Plant Sci 3: 248.

VanEtten HD, Mansfield JW, Bailey JA, Farmer EE (1994) Two classes of plant

antibiotics: phytoalexins versus "phytoanticipins." Plant Cell 6: 1191-1192.

Wang Y, Li X, Hu H (2014) H3K4me2 reliably defines transcription factor binding

regions in different cells. Genomics, 103(2), 222-228.

Watanabe T, *et al.* (2008) Endogenous siRNAs from naturally formed dsRNAs regulate

transcripts in mouse oocytes. Nature 453: 539-543.

Weng JK, Philippe RN, Noel JP (2012) The rise of chemodiversity in plants. Science

336: 1667-1670.

Wicker T, *et al.* (2007) A unified classification system for eukaryotic transposable

elements. Nat Rev Genet 8: 973-982.

Wink M (2003) Evolution of secondary metabolites from an ecological and molecular

phylogenetic perspective. Phytochemistry 64: 3-19.

Wittkopp PJ, Kalay G (2012) Cis-regulatory elements: molecular mechanisms and

evolutionary processes underlying divergence. Nat Rev Genet 13: 59-69

Wray GA (2007) The evolutionary significance of cis-regulatory mutations. Nat Rev

Genet 8: 206-216.

Zhao Y, *et al.* (2002) Trp-dependent auxin biosynthesis in Arabidopsis:

involvement of cytochrome P450s CYP79B2 and CYP79B3. Genes Dev 16: 3100-3112.

Zheng Z, Qamar SA, Chen Z, Mengiste T (2006) Arabidopsis WRKY33 transcription

factor is required for resistance to necrotrophic fungal pathogens. Plant J 48: 592-605.

Fig. 1. 4OH-ICN biosynthesis is specific to ETI

A. Schematic of tryptophan (L-Trp)-derived specialized metabolism in *A. thaliana*. White

arrows denote the presence of additional enzymes. ICY, indole cyanohydrin; ANI, *aci-*

nitro indole.

B. LC-DAD-MS analysis of camalexin, ICN, and 4OH-ICN in seedlings elicited with

flg22, *Pst*, or *Psta* for 24 hr. Data represent mean ± SE of four biological replicates.

Different letters denote statistically significant differences ($P < 0.05$, two-tailed *t*-test).

ICA-ME and 4OH-ICA-ME are methanolic degradation products of ICN and 4OH-ICN,

respectively. 4OH-ICA is an aqueous degradation product of 4OH-ICN.

Fig. 2. Intraspecific variation in *WRKY33* affects 4OH-ICN and immunity

A. Schematic of WRKY33 proteins in Col-0, Col-0 *wrky33*, L*er*-1 and Di-G. Black boxes

denote WRKY domain (W) or nuclear localization signal (NLS).

B. LC-DAD-MS analysis of camalexin, ICN, and 4OH-ICN in seedlings inoculated with

Psta for 24 hr. Data represent mean ± SE of four replicates.

757   **C.** Bacterial growth analysis of *Pst* in surface-inoculated leaves pre-treated with 20 μM

758   dex for 6-8 hr. Data represent mean ± SE of 8-12 biological replicates. CFU, colony-

759   forming units. Different letters in **(B-C)** denote statistically significant differences ($P <$

760   0.05, two-tailed *t*-test). Experiments in (**B-C**) were performed at least twice, producing

761   similar results.

Fig. 3. WRKY33 directly activates 4OH-ICN biosynthetic genes.

A. qPCR analysis of 4OH-ICN regulatory and biosynthetic genes in seedlings inoculated

with 20 μM dex and *Psta* for 9 and 12 hr.

B. Schematic of *FOX1* and *CYP82C2* loci, indicating nt positions of W-box-containing

regions (W).

C. ChIP-PCR analysis of W-box-containing regions upstream of *FOX1* and *CYP82C2* in

wrky33/DEX:WRKY33-flag plants co-treated with 20 μM dex (D) or mock solution (M)

772　　and *Psta* for 9 hr. Dashed line represents the 5-fold cutoff between weak and strong TF-

773　　DNA interactions. Data in (**B-C**) represent mean ± SE of four replicates.

Fig. 4. Regulatory neofunctionalization of *CYP82C2*

A. (Right) HPLC-DAD analysis of 4OH-ICN in seedlings inoculated with *Psta* for 30 hr.

(Left) Phylogenetic species tree. Data represent mean ± SE of three independent

experiments (n = 4 biological replicates), each with *A. thaliana* as a positive control.

4OH-ICA and 4OH-ICA-ME are aqueous and methanolic degradations products of

4OH-ICN, respectively. DW, dry weight; n.d., not detected. Experiments were performed

twice, producing similar results.

B. (Right) Synteny map of the *CYP82C* genes. Grey arrows represent non-*CYP82C*

genes. Grey dotted lines represent large (>500nt) sequence gaps. (Left) phylogenetic

species tree.

787   **C-D.** qPCR analysis of 4OH-ICN and sideretin biosynthetic genes in seedlings

788   inoculated with *Psta* (**C**) or grown in iron-deficient medium (**D**). Data represents the

789   mean ± SE of four biological replicates.

Fig. 5. TE *EPCOT3* is a *CYP82C2* enhancer

A. mVISTA plot of *CYP82C2* upstream sequence, indicating nt positions of unique

(*EPCOT1–3;* gray boxes) and conserved regions (≥70% sequence identity; pink) among

homologous sequences. Also indicated are positions of W-boxes (green) and WRKY33-

specific motifs (blue) that are present (solid lines) or absent (dashed lines) in each

homologous sequence, previously known WRKY33 TFBSs (diamonds) and ChIP-tested

regions (W1-5). TSS, transciptional start site; *Al, Arabidopsis lyrata; Ah, Arabidopsis*

halleri; *Cr, Capsella rubella*; *Bs, Boechera stricta; Cg, Capsella grandiflora*.

B. Epigenetic map of *CYP82C2* upstream sequence, indicating nt positions of

significant amounts of H3K4me2 (blue-gray bars), and H3K27me3 (purple bars).

C. (Left) Schematic of *EPCOT3* and related LINE retrotransposons in *A. thaliana* drawn

to scale, indicating nt positions of *CYP82C2* and reverse transcriptase (RT) domain. A

text file of the alignment and a more detailed tree are available as **Datasets S2-3**.

(Right) Phylogenetic maximum likelihood tree. Dashed box represent region containing

W-boxes (green lines) and/or WRKY33-binding motifs (blue lines) within *EPCOT3,*

EPL1 and *EPL2*.

D. ChIP-PCR analysis of W-box-containing regions (W) within *EPL1* and *EPL2* in

wrky33/DEX:WRKY33-flag plants co-treated with 20 μM dex (D) or mock solution (M)

and *Psta* for 9 hr. Data represent mean ± SE of four replicates. Dashed line represents

the 5-fold cutoff between weak and strong TF-DNA interactions.

815
816 **Fig. 6. Model of regulatory neofunctionalization of *CYP82C2*.**