Quantification of differential transcription factor activity and multiomic-based classification into activators and repressors: *diffTF*

Ivan Berest[1,*], Christian Arnold[1,*], Armando Reyes-Palomares[1], Giovanni Palla[1], Kasper Dindler Rasmussen[2,3], Kristian Helin[2,3] & Judith B. Zaugg[1]

[1] Structural and Computational Biology Unit, European Molecular Biology Laboratory, Heidelberg

[2] Biotech Research and Innovation Centre (BRIC), University of Copenhagen, Copenhagen

[3] Novo Nordisk Foundation Center for Stem Cell Biology, Copenhagen

[*] equal contribution

Transcription factor (TF) activity constitutes an important readout of cellular signalling pathways and thus for assessing regulatory differences across conditions. However, current technologies lack the ability to simultaneously assess activity changes for multiple TFs and in particular to determine whether a specific TF acts as repressor or activator. To this end, we introduce a widely applicable genome-wide method *diffTF* to assess differential TF binding activity and classifying TFs as activator or repressor by integrating any type of genome-wide chromatin with RNA-Seq data and in-silico predicted TF binding sites (available at https://git.embl.de/grp-zaugg/diffTF). We apply *diffTF* to a large ATAC-Seq dataset of mutated and unmutated chronic lymphocytic leukemia and identify dozens of TFs that are differentially active. Around 40% of them have a previously described association with CLL while ~60% constitute potentially novel TFs driving the different CLL subtypes. Finally, we validated the method experimentally using the well studied system of hematopoietic differentiation in mouse.

INTRODUCTION

Transcription factors (TFs) are important for orchestrating coordinated and dynamic responses to intra- and extracellular stimuli and regulating a multitude of biological processes. Indeed, since many signaling cascades end in the activation of a particular set of TFs, observing a change in overall TF activity can serve as a good readout of signaling pathways that regulate them[1]. However, even though many methods exist to investigate particular aspects of TF biology, none of them is able to globally measure the "activity" of all TFs. In addition, surprisingly little is known about whether particular TFs act as a transcriptional activator or repressor, despite some efforts that have been made by collecting published studies using text mining techniques[2]. TFs are typically lowly abundant proteins, which makes it difficult to detect them in proteomics experiments[1,3], and even if they can be detected, their abundance and TF activity do not necessarily correspond since TFs are highly regulated at the post-translational level. On the other hand, chromatin immunoprecipitation followed by sequencing (ChIP-Seq), which is the gold-standard technique for measuring the genomic TF binding events, provides information only for one TF at the time and it does not detect global changes in TF activity across conditions or cell types unless specific experimental normalisation methods are used (e.g. spike-ins[4]). Finally, luciferase assays only measure the activity of a specific TF at a specific location and are therefore fairly low throughput.

To correctly interpret the downstream effects of changes in TFs binding, it is crucial to know whether they act as transcriptional activator or repressor for the specific biological environment. Databases like TRRUST[2,5] collect annotations of TF modes' of regulation based on literature mining and provide a great resource for well-studied TF-target interactions. However, for the vast majority of TFs, there is no consensus about their molecular function, and most TFs are annotated as both transcriptional activators and repressors, depending on the target gene, cell type and growth condition. A general framework for classifying TFs based on their molecular mode of action in a cell-type and condition-specific manner is currently lacking.

Towards closing these gaps we have developed an approach called *diffTF* to integrate RNA-Seq and genome-wide chromatin accessibility or histone mark ChIP-Seq data to (i) estimate global changes in TF activities across conditions or cell types, and (ii) to classify TFs based on their molecular role into transcriptional activators and repressors. Applying this approach to a large-scale ATAC-Seq dataset of chronic lymphocytic leukemia (CLL) we first showed that *diffTF* is highly robust with respect to wide range of parameter settings, and that it is capable of

identifying many known TFs involved in CLL. In addition, we report many novel candidate TFs
that have never been shown before to be associated with CLL. Furthermore, our classification
into activators and repressors allowed us to specifically investigate the effects of transcriptional
activators and repressors on open chromatin and derive general principles of regulatory modes.
We have experimentally validated *diffTF* using the well characterized system of hematopoietic
stem cell differentiation tree. Finally, the approach has already been successfully applied to
identify TFs that are specifically associated with TET2, an enzyme involved in DNA
demethylation[6].

RESULTS

Conceptual derivation of using open chromatin as read-out of differential TF activity

We define change in TF "activity" as changes in open or active chromatin as measured by
ATAC-Seq, DNase-Seq or ChIP-Seq for H3K27ac or any other chromatin mark that is known to
be affected by TFs. This definition is based on earlier work where we showed that genetic
variants affecting H3K27ac signal across individuals (hQTLs) can be explained by disruptions of
TF motifs whenever the hQTL-SNP overlaps with a TF binding site (TFBS)[7]. Even though the
exact mechanisms of how changes in TF affinity translate to the chromatin level are unknown,
TF binding likely plays a causal role in mediating the effect of the DNA variant onto chromatin
marks[8]. By reversing this argument, we propose to use the aggregate changes in chromatin
accessibility in the vicinity of putative binding sites of a TF as a readout for its change in
"activity" (**Fig. 1a**).
Based on this concept, we developed *diffTF,* which is a computational approach to globally
assess differential TF activity between two conditions (**Suppl. Fig. 1**). It is based on (i) any data
that measures active/open chromatin, (ii) putative binding sites for TFs of interest, and optionally
(iii) matched RNA-Seq data, which are used to classify TFs into transcriptional activators or
repressors. Briefly, it requires *in silico* TFBS that can be obtained using position weight matrices
(PWMs) from a database (e.g. *HOCOMOCO*[9]) and a PWM scanning algorithm (e.g.
PWMScan[10]) for all TFs, or from a database of ChIP-Seq data, such as *ReMap*[11]. For each
TFBS it then calculates the difference between two conditions and summarizes their change in
accessibility across all binding sites of a given TF. In this step it also normalises for GC content
of the respective TFBS (**Suppl. Fig. 2**). Finally, the significance is assessed using an empirical
approach by permuting the sample labels or in an analytical way if the number of samples is too

low (see Methods and **Suppl. Fig. 3**). As a final result, *diffTF* outputs a differential activity value

along with an empirical FDR for each TF, which allows to identify the set of TFs that show a

significantly higher "activity" in either of the conditions (**Fig. 1b**).

Applying *diffTF* to a large ATAC-Seq and RNA-Seq dataset in chronic lymphocytic

leukemia (CLL)

Chronic lymphocytic leukemia (CLL) is one of the most frequent common type of cancer of the

Western world, particularly among old people. There are two major subtypes of CLL, which are

defined based on the mutation status of the IgHV locus (mutated: M-CLL and unmutated:

U-CLL). Overall, U-CLL associates with worse clinical outcomes such as shorter survival time

and higher frequency of relapse after treatment[12]. In M-CLL, B-cells go through normal affinity

maturation. With the aid of T-helper lymphocites, they undergo multiple rounds of IgHV somatic

hypermutation to produce high affinity B-cell receptors (BCR), which are however clonally

restricted. This process is essential for their development, survival and growth[13]. In contrast,

U-CLL B-cells reach their affinity maturation point in an unregulated manner, and without

involvement of T-helpers[14]. Recent studies have shown that the IgHV mutation status can be

well predicted based on chromatin profiles[15].

Given the vast knowledge about CLL, we thought this is an ideal case to apply *diffTF* to (i)

assess its power to recover relevant biological signal, (ii) gain more specific regulatory insights

into the different subtypes of CLL and (iii) assess its technical robustness. We therefore

obtained publicly available ATAC-Seq data for a cohort of CLL patients characterised by the

mutation status of the IgHV locus[15]. ATAC-Seq was performed in lymphocyte samples derived

from 52 CLL patients (27 mutated vs. 25 unmutated)[15]. For a subset of the patients we also

have RNA-Seq data available[15].

TF activity differences between U-CLL and B-CLL are highly robust within a wide

range of parameter settings

After extensive data processing and quality control (**Suppl. Fig. 4-6**), we applied *diffTF* to

identify the differences in TF activities between U-CLL and M-CLL. In total we identified 67 TFs

that are differentially active (FDR < 10%) between the two subtypes (**Fig. 2a**; **Suppl. Table 1**).

Before going into the biological interpretation of the results we used this dataset to assess the

technical robustness of *diffTF* with respect to a reasonable range of parameter settings.

Predicting putative TF binding sites is inherently noisy and it may result in many false positive predictions when compared to a gold standard such as ChIP-seq experiment for a given TF[16]. To address this point, we assessed the dependency of the *diffTF* results on the binding site predictions. We run *diffTF* only on binding sites that were bound by the TF based on ChIP-Seq data using all TFs comprised in *ReMap* and observed a very strong correlation of the resulting TF activity differences (R=0.81; **Fig. 2b**, **Suppl. Fig. 7**). This indicates that *diffTF* is robust with respect to false positive binding sites. Similarly, we found that neither the p-value cut-offs for motif scanning (**Suppl. Fig. 8**) nor the motif database (*JASPAR* vs. *HOCOMOCO*) had a strong impact on the differential TF activity (R=0.87, R=0.62 and R=0.69, respectively, **Fig. 2c-e**). Varying *diffTF* parameters such as the region size used for extracting the signal did not affect the resulting differential TF activities (see Supplement and **Suppl. Fig. 9**).

Next we sought to assess the power of *diffTF* to detect differential TF activities with respect to the level of accessibility differences across conditions. For this, we run it on a reduced set of peaks from which we removed all regions that were differentially accessible at a 5% FDR (see Methods) and found that the results were very similar for both sets (R=0.89). This demonstrates that *diffTF* can detect differences even in low signal data that have little or no differentially accessible peaks between conditions (**Fig. 2f**).

Finally, we assessed the power of *diffTF* with respect to sample size and sequencing depth. Intriguingly, we found the results highly congruent across a wide range of samples sizes and sequencing depths, with the majority of significant TFs in the full data showing the same direction of change in the subsampled data (**Fig. 2g**, **Suppl. Fig. 10**). Generally, the number of samples appears more important than read depth, and results using the full set were consistent for a coverage as low as 1-5 million processed reads per sample (see Methods). Although these results are dataset-specific and difficult to generalize they provide guidelines for the applicability of *diffTF*.

In summary, these results establish the robustness of *diffTF* in quantifying differences of TF activities, and demonstrate that aggregating signal across all binding sites is a powerful and sensible approach to overcome limitations such as low coverage and little underlying biological signal.

diffTF recovers known CLL TFs and proposes many novel candidates

After establishing the technical robustness of *diffTF* for identifying differentially active TFs, we next focused on the biological interpretation of the 67 TFs that showed significant differential activity at FDR 10% between M-CLL and U-CLL patients (**Fig. 2a**; **Suppl. Table 1**). Intriguingly, about ~41% of the differentially active TFs have previously been associated with CLL and mostly (90%) agree with the reported condition (i.e., mutated or unmutated; **Suppl. Table 2**), thus providing a strong biological validation of the approach. The remaining 50% may represent novel candidate TFs that can advance our understanding of the disease etiology in general and the biological differences of mutated and unmutated patients in particular.

Since *diffTF* is based on aggregating signal across TF binding sites, which in turn are identified by scanning for TF motifs, it is not able to distinguish between TFs with highly similar motifs. Therefore, we used the PWM clustering tool Rsat[17] to group TFs into clusters to allow proper interpretation of the results (clusters are available here https://bit.ly/2J9TaaK). Reassuringly, TF clusters are overall consistent in terms of TF activity changes (**Fig. 3a**). Exceptions include cluster 17 which can be explained by a prominent split into two branches early in the clustering into NFAT and NFκB factors, which show more activity in U-CLL and M-CLL, respectively (**Fig. 3c**).

The most active TF cluster in U-CLL is the IRF family and STAT2 (cluster 40; **Fig. 3d**), which is known to regulate cell-differentiation and induce apoptosis[18,19]. Several CLL susceptibility loci have been described proximal to IRF genes, and IRF expression has been associated with disease onset and progression[20–22]. The next most active cluster is the PAX TFs (cluster 54). One of its members, PAX5, together with PRDM1 (IRF cluster), affects B-cell to plasma cell differentiation, for which a disrupted regulation can lead to decreased cell survival and poor patient prognosis[23]. Another prominent set of regulators are the members of the AP-1 complex (cluster 4), which increase proliferation and have recently been shown to play an important role in driving the invasive nature of U-CLL[24]. Other prominent CLL-associated TFs that were identified with *diffTF* are c-MYC and CEBPB, which are involved in cell-proliferation and differentiation. Specifically, high c-MYC mRNA levels have been detected in U-CLL[25,26]. Several TFs with increased activity in U-CLL have been associated with massive U-CLL linked genomic aberrations such as BCL11A (having a (2;14)(p16;q32) translocation[27]). Overall, for U-CLL, we

observe an increased activity for TFs that induce and increase apoptosis and have been reported to be dysregulated in CLL.

In contrast, for M-CLL we identified TFs that regulate and possibly reduce apoptosis, regulate cell cycle and suggest normal functionality of B-cells through the classical BCR, NF-kB and Wnt signaling pathways. The most active TF family in this set of patients are the POU TFs, also known as Oct factors (cluster 12; **Fig 3b**), which regulate normal B-cell development and immunoglobulin production, therefore promoting survival of the lymphoma cells[28]. The POU cluster is followed by the ROR factors (cluster 36), which together with Wnt5a activates NF-kB-dependent survival signaling in CLL[29], and the GATA family (cluster 16), which is known to prime HSCs towards the lymphoid lineage and increase self-renewal of the stem cells in CLL[30]. Other examples include the EGR family, for which motifs have been found to be enriched in aberrant hypomethylated CpG sites in CLL[31]. Interestingly, we also found PPARD, which has recently been linked to M-CLL through its effect on metabolic pathways in cancer cells[32]. We also found members of the GLI family significantly more active in M-CLL. These are part of the Hedgehog signaling pathway and regulate apoptosis, thereby supporting survival of M-CLL cells[33].

Among the novel factors associated with U-CLL, we found several TFs involved in the circadian clock, such as BMAL1 and CLOCK, which form a complex, as well as their repressor NR1D1, the regulation of which has recently been proposed as another hallmark of cancer[34]. Moreover, we found other members of the basic helix-Loop-Helix family, such as, BHE40, a regulator of mitotic division[35], which has recently been found to be essential for development of B1-a cells[36] and TFAP4, TFE3 and TFEB, for which there are known cases of gene-fusions in renal cell carcinoma[37]. Another set of novel TFs that we observed to be more active in M-CLL were associated with cancer- and B-cell relevant pathways such as escape from apoptosis (ZN784)[38], regulation of cell cycle progression (ZBTB6)[39], selection of B-cells and promotion of fetal B lymphopoesis (ARID3A)[40], and control of the B cell-mediated autoimmunity[41,42]. The GFI1 family (cluster 35) is less active in U-CLL and their expression and activation might influence and decrease rates of apoptosis in B-cells[43].

In summary, these results show that *diffTF* is able to recapitulate much of the known biology of the two subtypes of CLL and additionally identifies numerous additional factors that are likely to be implicated in the disease.

Determination of the molecular function of TFs: transcriptional repressors and activators

The paragraphs above have shown that *diffTF* can identify TFs that alter their activity across different types of CLL patients. However, to gain mechanistic insights into some of the regulatory differences between U-CLL and M-CLL, it is crucial to know whether a TF acts as activator, in which case a higher abundance would generally result in increased transcription of its target genes, or repressor, in which case an increase in abundance would be paralleled by decreased target gene transcription. Surprisingly little is known about whether a certain TF acts mostly as transcriptional repressor or activator, and based on literature text mining most TFs have been annotated multiple times as both activator and repressor[5], indicating that the cell type or other external factors are important in determining a TF's main mode of action (**Suppl. Fig. 11B**). Therefore, we have devised a cell-type specific and data-driven, multiomics approach within the framework of *diffTF* to classify TFs into activators and repressors based on RNA-Seq, ATAC-Seq and putative TFBS.

The rationale behind this classification is that increasing levels of an activator TF should result in more open, active chromatin at its targets sites and increased transcription of its target genes and vice versa for repressors. With an increase in abundance of an activator TF we therefore expect an increase in chromatin accessibility surrounding its binding sites while the chromatin surrounding binding sites of a repressing TF we would expect to close (**Fig. 4a**). To test this rationale, we investigated the ATAC-Seq profile surrounding binding sites of a well-known repressor (REST) and activator (STAT2) that are both expressed in our cell type (**Fig. 4b**). We observed that for REST, the accessibility drops to genome-wide average (within peak areas) within 10 bp from the center of the core motif, while only a small region at the binding site is slightly accessible, likely reflecting the binding of the factor itself. In contrast, for STAT2, we observed increased chromatin accessibility also outside the core binding site, which drops to genome-wide average over a distance of >100 bp from the center of the motif, likely representing the effect of the TF on opening chromatin. This indicates that, while there is a similar pattern of increase in accessibility at the exact binding site of a TF, the surrounding chromatin is highly compact for the repressor while it is open for the activator, thus supporting our rationale that repressors overall decrease chromatin accessibility around their binding sites.

To classify TFs into activators and repressors *diffTF* uses RNA-Seq data as an estimate for TF

abundance. For each TF it then calculates the correlations across individuals between a TF's

expression level and the ATAC-Seq signal in its putative target peak. A TF gets classified as

activator when it shows an overall positive correlation with the ATAC-signal at its putative target

sites, as repressor if it is overall negative, and as undetermined if the distribution of correlations

is not significantly different from the peaks that did not overlap its putative binding sites

(examples in **Fig 4c**, see also Methods and **Suppl. Fig. 11A**).

Applying this to the CLL data, where RNA-Seq data was available for eight individuals, we were

able to classify 50% of the expressed TFs as either activators or repressors (**Fig 4c-f**; n=186).

Among the top activators are the IRFs, which are well known transcriptional activators[44] (see

Fig. 4f for a footprint of IRF2 and **Fig. 4c** for the correlation distribution that led to its

classification). Among the top repressors we found PAX5, which has been shown to repress the

activity of BLIMP-1 and thereby regulates plasma cell differentiation[45] (**Fig. 4f,c**). The circadian

regulators provide an example of why it is important to know the mode of action of a particular

TF: When analysing the differential TF activities, it appears as if BMAL1 is more active in M-CLL

while the other two TFs (CLOCK and NR1D1) are more active in U-CLL (**Fig 4e**). However,

since BMAL1 is an activator, CLOCK a repressor and NR1D1 borderline classified as repressor,

we get a consistent result from this analysis whereby all three circadian factors are more "active"

in M-CLL, albeit with a contrary effect on their target genes.

In addition to the TF-specific molecular insights and better interpretability of the data, this global,

cell-type specific classification of TFs allowed us to explore more details about activators and

repressors in general. Importantly, we found that the aggregate signal across all repressors

produced a footprint similar to that of REST, while the one for activators looks similar to STAT2,

thus again indicating that repressors and activators have very distinct open chromatin footprints

(**Fig. 4g**). When investigating TF families, as defined above with the RSat clusters (**Fig. 3**), we

found that TFs from the same PWM cluster are often classified both as activators and

repressors (**Suppl. Fig. 12**), supporting the hypothesis that the molecular function of a TF is

highly variable. The exceptions are clusters 40, containing mainly members of the IRF family

and cluster 17 that contains both NFAT and NFKB TFs, which are mostly classified as

activators.

As a validation for the repressor/activator classification, we tested two specific hypotheses: first

we expected that activating TFs would show a positive correlation between their differential

expression and differential TF activity of M-CLL vs U-CLL patients, while it would be

anticorrelated for repressors. Indeed, we found both a significant positive correlation for
activators (r=0.19, P = 0.05) and a significant negative one for repressors (r=-0.32, P = 0.0033;
Fig. 5a). Second, we expect that the expression of target genes of a given TF should go in the
same direction as the differential TF activity as assessed by *diffTF*. Again, we found that this
was the case for both activators and repressors (**Fig. 5b,** see Methods). These observations
provide an additional, independent validation that *diffTF* is able to pick up differential TF activity.

It is important to note that TFs can often act as activators and repressors at different genomic
loci e.g. depending on their co-factors, whereas here we predict their main mode of action
based on the mean effect across all their predicted binding sites, and thereby lose any
information about bifunctionality. Furthermore, since the classification is based on correlations, it
is heavily dependent on variation in the RNA-Seq signal across individuals, thus biasing the TFs
that can be classified towards those that are variable across individuals. As a consequence, TFs
whose post-transcriptional regulation is not reflected in their transcript abundance will not get
classified correctly. Another potential misclassification may happen because of the similarity of
PWMs within a cluster, which makes it difficult to distinguish the exact effect of one TF while its
expression level is uniquely defined. As an example we cite PRDM1, which is part of IRF family
(cluster 40) and classified as very strong repressor, its footprint however looks more similar to a
typical activator (data not shown), suggesting that it is not PRDM1 driving the ATAC-Seq signal
in this case, but the IRFs. Thus, for distinguishing the functional roles of TFs from the same
cluster/family further biochemical experiments will be needed. Despite these potential pitfalls,
diffTF provides unique insights into the molecular mechanism of TFs on a global level.

Comparison with similar tools

There are a few tools with a similar or comparable focus[46–48] and we thought to compare them
with *diffTF* (**Suppl. Table 5**). Overall, *diffTF* enables a more flexible and tailored analysis of the
data and the question at hand due to the extensive choice of parameters, diagnostic plots,
TFBS-specific results, visualizations, and pipeline adjustability. It is unique in its ability to directly
integrate RNA-Seq with ATAC-Seq data to classify TFs into activators and repressors. Due to
this flexibility, *diffTF* is computationally expensive, and we provide detailed instructions on
memory and time requirements in the documentation.

To compare *diffTF* with a more traditional TF motif analysis we chose *HOMER*[48], which looks at

motif enrichment in a set of differentially accessible peaks. Strikingly, no enriched motifs were

found in M-CLL with *HOMER*, while the few discovered in U-CLL significantly correlated with

differential TF activity as computed by *diffTF* (**Supp. Fig. 13,** see Methods). This analysis

highlights the power of *diffTF* to capture more significant signal than standard motif

enrichment-approaches.

For comparing *diffTF* with an approach that is also based on TF activities we chose *chromVAR*.

Overall, the results correlate very well, with correlation coefficients between 0.65-0.9 for

Pearson and 0.56-0.81 for Spearman, depending on the set of TFs (i.e., all TFs or only those

deemed significant by *diffTF*, therefore predominantly removing TFs with low signal) and

whether *chromVAR* deviations or deviation scores are compared against (**Supp. Fig. 14**, see

Methods). Differences likely arose from the way of what the scores represent or what they are

compared against (i.e., compared against the mean effect in *diffTF* vs modelled background in

chromVAR).

Overall, the comparable part of the results of *diffTF* are highly correlated with the output of other

tools. However, *diffTF* goes one step beyond the currently available methods by classifying TFs

based on their mode of regulation - activator or repressor, thus providing important additional

insights into their molecular function.

Experimental validation in a well-studied system

Finally, we experimentally validated our approach using the well-studied mouse hematopoietic

system. We generated ATAC-Seq profiles of multipotent progenitor cells (MPP; Lin⁻cKit⁺Sca1⁺

CD150⁻CD48⁺), an early hematopoietic progenitor population capable of supporting multilineage

blood production[49], as well as the more differentiated and myeloid-restricted

granulocyte-monocyte progenitors (GMP; Lin⁻cKit⁺Sca1⁻; CD16/32⁺). The obtained profiles were

processed using an in-house ATAC-Seq pipeline and *diffTF* was subsequently used to identify

TFs that are differentially active between MPP and GMP cells (see Online Methods). Due to the

large number of significant TFs reflecting the high diversity between the two cell types, we used

available RNA-Seq data to filter non-expressed TFs. The small number of samples (two) did not

allow a faithful classification the TFs into activators and repressors since the classification is

based on correlations across samples. In GMPs, members of the well-known class of master

regulators of myelopoiesis, the CEBP family (C/ebpα,-β,-δ,-ε,-γ), showed a strong increase in

activity. In addition, we could detect increased activity of the NFKB (RELB, NFKB2)[50] as well as

the AP-1 (JUN/FOS)[51] complexes, which act as another cue for further granulopoiesis (**Fig. 6**,

Suppl. Fig 15). In addition, we also observed higher activity of the MYC/MYB factors, which are

known to be exclusively active in the GMPs[52] as well as NFIL3, which is involved in the

generation of natural killer cells[53]. Conversely, MPPs show a higher activity for IRF/STAT, ZEB1

and ITF2 (part of the Wnt signaling) as well as TFs from the Homeodomain and Forkhead

families[54], all of which are associated with self-renewal of hematopoietic stem cells.

Overall, these results show that *diffTF* was able to identify the known TFs that drive the

differentiation from MPP to GMP, thus demonstrating that it confidently identifies the biologically

significant signals.

DISCUSSION

We here presented a genome-wide method for quantifying differences in differential TF activity

simultaneously for a large set of TFs, and for classifying them into their molecular mode of

action as transcriptional activators or repressors. The method is available for download at

https://git.embl.de/grp-zaugg/diffTF along with a comprehensive documentation and example

data.

We have shown the robustness of *diffTF* with respect to a large range of parameters, sample

sizes and read depth. In particular, we have shown that *diffTF* is able to overcome the inherent

noisiness of TF binding site predictions by aggregating data across all putative binding sites, as

the results when using putative binding sites and ChIP-Seq validated sites were strongly

correlated. Furthermore, we validated the estimation of differential TF activity by (i) comparing

our results with known biological relationships between TFs and certain biological processes, (ii)

by showing an overall correspondence between changes in transcript abundance and changes

in the activity of a given TF, and finally experimentally (iii) by profiling different cell types along

the well-known hematopoietic differentiation system and identifying the key driving factors.

To demonstrate the power of *diffTF* we have applied it to identify and characterise differences

between M-CLL and U-CLL from a publicly available dataset of ATAC-Seq and RNA-Seq. It is

noteworthy that a TF motif enrichment analysis on the significantly differentially accessible

peaks did not reveal any factor to be significantly enriched in M-CLL, indicating that in this case

(as probably in many patient - control studies) the key driver TFs are not necessarily switching

their target peaks (enhancers and promoters) on and off, but rather mis-regulate many common

regions to a lesser extent. Although signal differences at each individual putative binding site

can be very low and rarely significant, if there is a slight bias in activity of the TF, *diffTF* can still

detect an overall difference in TF binding activity by averaging across all of its putative binding

sites and is therefore more powerful than conventional enrichment analyses.

Unlike other methods that have been published on differential TF activity based on

chromatin[46,47], we have devised an approach within the *diffTF* framework to classify TFs into

activators and repressors based on the correlation of their expression level (RNA-Seq) and the

activity of their putative binding sites (ATAC-Seq). This information is highly relevant when

interpreting the action of TFs since it makes a difference of whether the signal is caused by a

repressor that is less active in condition A or an activator that is more active in condition A.

Notably, this classification could work even for data sets for which no RNA-Seq data are

available, as one can use ATAC-Seq signal summarized across the gene body as proxy for its

expression level. We have observed this correlation to be around 0.63 (Pearson).

Calculating differential TF activity based on aggregating signal across the genome is not a new

idea and has been proposed several times based on aggregating the expression of putative

target genes of a certain TF[55,56]. Using the effect on chromatin instead of expression has several

advantages: first chromatin is a much simpler trait since gene expression is the sum of

transcription and degradation. Second, there are much more peaks than genes, thus allowing

for better statistics and signal to noise ratio. Finally, the effect on chromatin is much more local

than on gene expression - in particular in mammalian genomes, where genes can be regulated

by very distal enhancers. We have compared differential TF activity calculated based on the

average expression change of the target genes to the output of *diffTF* and found that while the

direction of activity between both methods is highly correlated, the signal is much lower when

using the expression of target genes instead of chromatin accessibility at putative binding sites.

While similar methods have been proposed for analysing ATAC-Seq data[46,47], our method has

several advantages when dealing with bulk ATAC-Seq data and can also be used for histone

mark ChIP-Seq data: (i) Unlike other methods that calculate the background theoretically based

on the genome-wide read depth, it is insensitive to sequence or locus dependent biases since

we calculate a fold-change between conditions for each region, thus normalizing for local read

depth biases. This is particularly advantageous for detecting small changes such as between

two heterogeneous cohorts in patient-control studies. (ii) *diffTF* allows integration with matching

RNA-Seq data to classify TFs into activators and repressors in a fully data-driven approach

within the same analysis framework. Such classifications greatly help in interpreting the effects of up/down regulation of a particular factors. And, even if classifying them is not possible, at least non-expressed TFs can be identified. (iii) *diffTF* provides the fold-change value for each TFBS which allows for easy retrospective follow-up analysis, e.g. identifying the most differential regions regulated by a specific set of TFs. (iv) Finally, our method allows to analyse time-course data in an additive manner by calculating the overall change of slope for each TF (see Methods).

Overall, with *diffTF* we present a multiomics data integration strategy of ATAC-Seq and RNA-Seq data that allows to calculate differential TF activity across conditions and classifying TFs based on their molecular mode of action into activators and repressors. With this, *diffTF* is laying the foundation for understanding the regulatory mechanisms that are driving the differences in cell state on a systems-wide scale.

METHODS

Methods, including statements of data availability and any associated accession codes and references, are available in the online version of the paper.

ACKNOWLEDGMENTS

We thank Bernd Klaus for help with the statistical part and EMBL for funding. A.R.P is recipient of a postdoctoral fellowship granted by Fundación Ramón Areces. G.P. was awarded the Otto Bayer Scholarship from Bayer Foundations.

AUTHOR CONTRIBUTIONS

J.Z. and I.B. conceived the study, C.A. and I.B. developed the computational framework and performed the analyses, A.R.P. and G.P. contributed to the development and analysis, K.D.R performed the experiments, J.Z. supervised the study and C.A., I.B, and J.Z wrote the manuscript.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests

Figures and figure captions

Figure 1. Concept and simplified workflow of *diffTF*. (a) Concept of the differential TF activity approach: Based on findings from chromatin QTL studies that have reported changes in chromatin often coincide with SNPs disrupting TF binding sites (TFBS), we propose to detect global differences in TF activity by investigating chromatin accessibility (or modifications) in close proximity to their TFBS. (b) A simplified workflow illustrates the principle *diffTF* is based on: it calculates a fold-change between two conditions for each binding site of a given TF and compares this distribution to a background set of fold-changes obtained from GC-content matched loci that do not contain the putative TFBS. The difference in distribution is assessed in significance and effect size and visualized in a volcano plot where the y-axis indicates statistical significance and the x-axis shows the effect size. (For a detailed workflow see Suppl. Fig. 1 and 2).

A) Conceptual derivation of *diffTF*

SNP disrupts TF binding, which in turn affects chromatin environment

Infer differential TF activity from changes in chromatin marks (*diffTF*)

cis Effect → *infer* → *trans* Effect

Genotype 1

Genotype 2

Low activity of TF

High activity of TF

⬡ Transcription factor (TF)

▬ Putative TF binding sites (TFBS)

☷ Nucleosome

−log10 (fdr)

Δ TF activity [group 1 vs group 2]

Visualize differential TF activity

*less active if TF is a repressor

B) Schematic workflow of *diffTF*

TF1 TF1 TF2 TF1 TF2

ATAC-Seq in group 1 (e.g. U-CLL)

Coverage

TF3 TF3 TF2 TF3 TF2

ATAC-Seq in group 2 (e.g. M-CLL)

Coverage

Per TF: calculate fold changes at each TFBS and compare to background

⎯ TFBS of given TF

···· Back-ground

Density

Log2 fold changes [group 1 / group 2]

Figure 2. *diffTF* results for the CLL dataset, experimental validation and technical robustness of the method. (a) Volcano plot of differential TF activity between U-CLL (n=27) and M-CLL (n=25) patients.. The y-axis denotes statistical significance (-log10 transformed). TFs that pass the significance threshold (5% FDR; dotted line) are labeled and colored according to their novelty status (see Suppl. Table. 2). #TFBS denotes the number of predicted TF binding sites in the peak regions this analysis is based on. (b)-(f): Technical robustness of *diffTF*. Scatterplots of the differential TF activity from all TFs for two different *diffTF* analyses are shown. Each point represents one TF. For (c-f), colors represent significance at 5% FDR (white - not significant in either analysis; light green and light blue - significant for the analysis on the x-axis or y-axis, respectively; purple - significant for both analyses). (b) Comparison of all predicted TFBS and TFBS experimentally validated by ChIP-Seq data from ReMap. See also Suppl. Fig. 7. (c-d) Comparison for different p-value thresholds in PWMScan to predict TFBS: (c) standard (1e-5) vs. stringent (1e-6), (d) standard vs. lenient (5e-5) (n=628 TFs for which binding sites were retrieved for both scanning modes.) (e) Comparison of *diffTF* results based on *HOCOMOCO* v10 vs *JASPAR* 2018 as input (n=412 TFs, for which a motif was available in both databases.) (f) Comparison of the full consensus peak set (*allPeaks*) and only the non-differentially accessible peaks (*noDApeaks*; n=640 TFs from *HOCOMOCO*) (g) Robustness analysis based on sequencing depth and sample size. Each cell in the heatmap shows the fraction of TFs that showed the same direction of change as in the full dataset for varying degrees of down-sampling sequencing depth and number of samples, (5% FDR), averaged over 50 independent repetitions to minimize sampling noise. Only TFs that were deemed significant in the full dataset are considered (see also Suppl. Fig. 8). Sequencing depth is shown as fraction of the original data and median number of reads across samples, while the number of samples is shown in the format unmutated + mutated.

 Figure 3. Functional clustering of TFs based on the similarity of their PWMs. (a) Boxplot of
PWM clusters with at least 3 members as defined by RSAT for the differential TF activity
between U-CLL and M-CLL. In each cluster, the most negative and most positive TF is labeled.
(b) RSAT clustering output and tree for cluster 17. The cluster has two distinct subclusters,
representing the NFAT and NFKB families. (c) and (d) RSAT clustering output and tree for
cluster 12 (POU family) and 40 (IRF family), the most distinct clusters for M-CLL and U-CLL
patients, respectively).

Figure 4. Classification of TFs into activator or repressor based on RNA-Seq and ATAC-Seq data. (a) Schematic description of the classification approach: correlation of TF expression level with the accessibility of its target sites. If the correlation of the RNA-level and the chromatin peak openness is positive, the TF acts more as an activator in the particular biological environment, if negative, it acts more as a repressor. Correlations around 0 are classified as undetermined. (b) Exemplary footprints for a well-known repressor (REST, left) and activator (STAT2, right), respectively. The x-axis depicts the distance in bp from the TFBS center, the y-axis denotes the number of average Tn5 insertions, normalised to the library size and numbers of samples across U-CLL and M-CLL. TFBS predicted by PWMscan and overlapping with open chromatin have been used. Solid black line indicates the average insertion sites within accessible chromatin. (c) Distributions of the Pearson correlations between TF expression and ATAC-Seq signal at all putative TFBS (red line), and background distribution of TFBS not-containing the motif of interest (grey line) for two specific TFs, IRF2 (left) and PAX5 (right). (d) Heatmap of the differences in Pearson correlation of putative targets peaks and background for each TF (i.e. subtracting the black from the red line in (c)). Each TF is one row and is annotated as activator (green), undetermined (grey) or repressor (red) as classified by *diffTF*. (e) Same as Figure 2a, but with their predicted role as activator (green), undetermined (black) or repressor (red). See Figure 2a for details. (f) Footprint analysis for an activator (IRF2; right) and repressor (PAX5; left) as classified by *diffTF*. The footprint based on the normalized number of Tn5 insertions (top; blue: U-CLL, orange: M-CLL) and differences (U-CLL - M-CLL; bottom) are shown. See b) for axis descriptions. (g) Summary footprint for all activators (top, green) and repressors (bottom, red). See b) for axis descriptions.

Figure 5. (a) Pearson correlation of the log2 fold changes from RNA-Seq and differential TF activity for activators (green, left), undetermined TFs (black, middle), and repressors (red, right). Only expressed TFs are shown. (b) Correlation of differential TF activity and median of the differential target gene expression of U-CLL against M-CLL. The x-axis shows the median target gene (TG) log2 fold-change, the y-axis denotes the differential TF activity. Each TF label is colored based on its activator/repressor status (green/red) on a continuous scale (dark to light) based on the the correlation strength (see Figure 4d). (OR=odds ratio, r=Pearson correlation coefficient).

Figure 6. *diffTF* recapitulates known TFs that drive the differentiation from MPP to GMP. Circular plot of differential TF activity between 4 wildtype GMP and 4 wildtype MPP samples are shown. This is an alternative representation equivalent to the Volcano plot shown in Figure 2: values on the x-axis denote the differential TF activity, with higher values indicating a higher differential TF activity between the two conditions. TFs that are similarly active between GMP and MPP samples are close to 0, while TFs more active in either condition are located in the left and right part of the plot. The y-axis (radial position) denotes the statistical significance (adjusted p-values). The significance threshold (5% FDR) is indicated as a red circular line. Due to the high number of significant TFs we labeled only those discussed in the main text. The full list is available in Suppl. Table 4.

REFERENCES

1. Kim, R., Emi, M. & Tanabe, K. Cancer immunoediting from immune surveillance to immune escape. *Immunology* **121,** 1–14 (2007).

2. Han, H. *et al.* TRRUST: a reference database of human transcriptional regulatory interactions. *Sci. Rep.* **5,** 11432 (2015).

3. Teng, M. W. L., Swann, J. B., Koebel, C. M., Schreiber, R. D. & Smyth, M. J. Immune-mediated dormancy: an equilibrium with cancer. *J. Leukoc. Biol.* **84,** 988–993 (2008).

4. Bonhoure, N. *et al.* Quantifying ChIP-seq data: a spiking method providing an internal reference for sample-to-sample normalization. *Genome Res.* **24,** 1157–1168 (2014).

5. Han, H. *et al.* TRRUST v2: an expanded reference database of human and mouse transcriptional regulatory interactions. *Nucleic Acids Res.* **46,** D380–D386 (2018).

6. Rasmussen, K. *et al.* TET2 binding to enhancers facilitates transcription factor recruitment in hematopoietic cells. *BioRxiv* (2018). doi:10.1101/336008

7. Grubert, F. *et al.* Genetic control of chromatin states in humans involves local and distal chromosomal interactions. *Cell* **162,** 1051–1065 (2015).

8. Liu, L., Jin, G. & Zhou, X. Modeling the relationship of epigenetic modifications to transcription factor binding. *Nucleic Acids Res.* **43,** 3873–3885 (2015).

9. Kulakovskiy, I. V. *et al.* HOCOMOCO: a comprehensive collection of human transcription factor binding sites models. *Nucleic Acids Res.* **41,** D195-202 (2013).

10. Ambrosini, G., Groux, R. & Bucher, P. PWMScan: a fast tool for scanning entire genomes with a position-specific weight matrix. *Bioinformatics* (2018). doi:10.1093/bioinformatics/bty127

11. Griffon, A. *et al.* Integrative analysis of public ChIP-seq experiments reveals a complex

multi-cell regulatory landscape. *Nucleic Acids Res.* **43,** e27 (2015).

12. Furman, R. R. *et al.* Idelalisib and rituximab in relapsed chronic lymphocytic leukemia. *N.*

Engl. J. Med. **370,** 997–1007 (2014).

13. Neu, K. E. & Wilson, P. C. Taking the broad view on B cell affinity maturation. *Immunity* **44,**

518–520 (2016).

14. Chiorazzi, N. & Ferrarini, M. Cellular origin(s) of chronic lymphocytic leukemia: cautionary

notes and additional considerations and possibilities. *Blood* **117,** 1781–1791 (2011).

15. Rendeiro, A. F. *et al.* Chromatin accessibility maps of chronic lymphocytic leukaemia

identify subtype-specific epigenome signatures and transcription regulatory networks. *Nat.*

Commun. **7,** 11938 (2016).

16. Landt, S. G. *et al.* ChIP-seq guidelines and practices of the ENCODE and modENCODE

consortia. *Genome Res.* **22,** 1813–1831 (2012).

17. Medina-Rivera, A. *et al.* RSAT 2015: regulatory sequence analysis tools. *Nucleic Acids Res.*

43, W50-6 (2015).

18. Lin, T. S., Mahajan, S. & Frank, D. A. STAT signaling in the pathogenesis and treatment of

leukemias. *Oncogene* **19,** 2496–2504 (2000).

19. Benekli, M., Baumann, H. & Wetzler, M. Targeting signal transducer and activator of

transcription signaling pathway in leukemias. *J. Clin. Oncol.* **27,** 4422–4432 (2009).

20. Arvaniti, E. *et al.* Toll-like receptor signaling pathway in chronic lymphocytic leukemia:

distinct gene expression profiles of potential pathogenic significance in specific subsets of

patients. *Haematologica* **96,** 1644–1652 (2011).

21. Havelange, V. *et al.* IRF4 mutations in chronic lymphocytic leukemia. *Blood* **118,** 2827–2829

(2011).

22. Slager, S. L., Caporaso, N. E., de Sanjose, S. & Goldin, L. R. Genetic susceptibility to

chronic lymphocytic leukemia. *Semin. Hematol.* **50,** 296–302 (2013).

23.   Ghamlouch, H. *et al.* Factors involved in CLL pathogenesis and cell survival are disrupted

by differentiation of CLL B-cells into antibody-secreting cells. *Oncotarget* **6,** 18484–18503

(2015).

24.   Mittal, A. K. *et al.* Role of CTLA4 in the proliferation and survival of chronic lymphocytic

leukemia. *PLoS ONE* **8,** e70352 (2013).

25.   Landau, D. A. *et al.* Mutations driving CLL and their evolution in progression and relapse.

Nature **526,** 525–530 (2015).

26.   Yeomans, A. *et al.* Engagement of the B-cell receptor of chronic lymphocytic leukemia cells

drives global and MYC-specific mRNA translation. *Blood* **127,** 449–457 (2016).

27.   Socola, F., Insuasti-Beltran, G., Henrich Lobo, R., Atrash, S. & Sasapu, A. Chronic

Lymphocytic Leukemia with Translocation (2;14)(p16;q32): A Case Report and Review of

the Literature. *Case Rep. Oncol. Med.* **2016,** 9037436 (2016).

28.   Heckman, C. A., Duan, H., Garcia, P. B. & Boxer, L. M. Oct transcription factors mediate

t(14;18) lymphoma cell survival by directly regulating bcl-2 expression. *Oncogene* **25,**

888–898 (2006).

29.   Minami, Y., Oishi, I., Endo, M. & Nishita, M. Ror-family receptor tyrosine kinases in

noncanonical Wnt signaling: their implications in developmental morphogenesis and human

diseases. *Dev. Dyn.* **239,** 1–15 (2010).

30.   Kikushige, Y. *et al.* Self-renewing hematopoietic stem cell is the primary target in

pathogenesis of human chronic lymphocytic leukemia. *Cancer Cell* **20,** 246–259 (2011).

31.   Oakes, C. C. *et al.* DNA methylation dynamics during B cell maturation underlie a

continuum of disease phenotypes in chronic lymphocytic leukemia. *Nat. Genet.* **48,**

253–264 (2016).

32. Li, Y. J. *et al.* PPAR-delta promotes survival of chronic lymphocytic leukemia cells in

energetically unfavorable conditions. *Leukemia* **31,** 1905–1914 (2017).

33. Kern, D. *et al.* Hedgehog/GLI and PI3K signaling in the initiation and maintenance of

chronic lymphocytic leukemia. *Oncogene* **34,** 5341–5351 (2015).

34. El-Athman, R. & Relógio, A. Escaping circadian regulation: an emerging hallmark of

cancer? *Cell Syst.* **6,** 266–267 (2018).

35. D'Annibale, S. *et al.* Proteasome-dependent degradation of transcription factor activating

enhancer-binding protein 4 (TFAP4) controls mitotic division. *J. Biol. Chem.* **289,**

7730–7737 (2014).

36. Kreslavsky, T. *et al.* Essential role for the transcription factor Bhlhe41 in regulating the

development, self-renewal and BCR repertoire of B-1a cells. *Nat. Immunol.* **18,** 442–455

(2017).

37. Kauffman, E. C. *et al.* Molecular genetics and cellular features of TFE3 and TFEB fusion

kidney cancers. *Nat. Rev. Urol.* **11,** 465–475 (2014).

38. Kasim, V. *et al.* Transcription factor Yin Yang 2 is a novel regulator of the p53/p21 axis.

Oncotarget **8,** 54694–54707 (2017).

39. Chevrier, S. *et al.* The BTB-ZF transcription factor Zbtb20 is driven by Irf4 to promote

plasma cell differentiation and longevity. *J. Exp. Med.* **211,** 827–840 (2014).

40. Zhou, Y. *et al.* Lin28b promotes fetal B lymphopoiesis through the transcription factor

Arid3a. *J. Exp. Med.* **212,** 569–580 (2015).

41. Rathinam, C., Lassmann, H., Mengel, M. & Klein, C. Transcription factor Gfi1 restricts B

cell-mediated autoimmunity. *J. Immunol.* **181,** 6222–6229 (2008).

42. Gómez-Martín, D., Díaz-Zamudio, M., Galindo-Campos, M. & Alcocer-Varela, J. Early

growth response transcription factors and the modulation of immune response: implications

towards autoimmunity. *Autoimmun. Rev.* **9,** 454–458 (2010).

43. Coscia, M. *et al.* IGHV unmutated CLL B cells are more prone to spontaneous apoptosis and subject to environmental prosurvival signals than mutated CLL B cells. *Leukemia* **25,** 828–837 (2011).

44. Yanai, H., Negishi, H. & Taniguchi, T. The IRF family of transcription factors: Inception, impact and implications in oncogenesis. *Oncoimmunology* **1,** 1376–1386 (2012).

45. Yasuda, T. *et al.* B cell receptor-ERK1/2 signal cancels PAX5-dependent repression of BLIMP1 through PAX5 phosphorylation: a mechanism of antigen-triggering plasma cell differentiation. *J. Immunol.* **188,** 6127–6134 (2012).

46. Baek, S., Goldstein, I. & Hager, G. L. Bivariate genomic footprinting detects changes in transcription factor activity. *Cell Rep.* **19,** 1710–1722 (2017).

47. Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR: inferring transcription-factor-associated accessibility from single-cell epigenomic data. *Nat. Methods* **14,** 975–978 (2017).

48. Heinz, S. *et al.* Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol. Cell* **38,** 576–589 (2010).

49. Sun, J. *et al.* Clonal dynamics of native haematopoiesis. *Nature* **514,** 322–327 (2014).

50. Paz-Priel, I., Ghosal, A. K., Kowalski, J. & Friedman, A. D. C/EBPalpha or C/EBPalpha oncoproteins regulate the intrinsic and extrinsic apoptotic pathways by direct interaction with NF-kappaB p50 bound to the bcl-2 and FLIP gene promoters. *Leukemia* **23,** 365–374 (2009).

51. Cai, D. H. *et al.* C/EBP alpha:AP-1 leucine zipper heterodimers bind novel DNA elements, activate the PU.1 promoter and direct monocyte lineage commitment more potently than

C/EBP alpha homodimers or AP-1. *Oncogene* **27,** 2772–2779 (2008).

52. Baker, S. J. *et al.* B-myb is an essential regulator of hematopoietic stem cell and myeloid

progenitor cell development. *Proc Natl Acad Sci USA* **111,** 3122–3127 (2014).

53. Gascoyne, D. M. *et al.* The basic leucine zipper transcription factor E4BP4 is essential for

natural killer cell development. *Nat. Immunol.* **10,** 1118–1124 (2009).

54. Sands, W. A., Copland, M. & Wheadon, H. Targeting self-renewal pathways in myeloid

malignancies. *Cell Commun. Signal.* **11,** 33 (2013).

55. Bussemaker, H. J., Li, H. & Siggia, E. D. Regulatory element detection using correlation

with expression. *Nat. Genet.* **27,** 167–171 (2001).

56. Boorsma, A., Lu, X.-J., Zakrzewska, A., Klis, F. M. & Bussemaker, H. J. Inferring

condition-specific modulation of transcription factor activity in yeast through regulon-based

analysis of genomewide expression. *PLoS ONE* **3,** e3112 (2008).