Transgenerational effect of adult density through paternal effect in *Drosophila melanogaster*

Purbasha Dasgupta[1†], purbasha17@iiserbpr.ac.in

Saubhik Sarkar[1†], sarkar.saubhik.sarkar@gmail.com

Akankshya A. Das[2], ayesa692@gmail.com

Tanya Verma[1], tanya17@iiserbpr.ac.in

Bodhisatta Nandy[1]*, nandy@iiserbpr.ac.in

[†] Equal contribution

* Corresponding author

Present address:

1. Indian Institute of Science Education and Research Berhampur

Berhampur Transit Campus,Govt. ITI Building,

NH 59, Engineering School Junction,Ganjam, Berhampur 760 010, Odisha, India.

Contact number: +91-680-222 7755

2. Department of Zoology, Banaras Hindu University, Varanasi – 221005, Uttar Pradesh, India.

Running title: Paternal density effect on progeny fitness

Word count:

Figure count: 3

Table count: 1

Abstract:

Notwithstanding recent evidences, paternal environment is thought to be a potential but unlikely source of fitness variation that can affect trait evolution. Here we studied transgenerational effect(s) of males' exposure to varying adult density in *Drosophila melanogaster* laboratory populations. We held sires at normal (N), medium (M) and high (H) adult densities for two days, before allowing them to mate with virgin females. We then collected eggs from the dams and measured the egg size. Finally we investigated the effect of the paternal treatment on juvenile and adult (male) fitness components. We found a significant treatment effect on juvenile competitive ability where the progeny sired by the H-males had higher competitive ability. Since we did not find the treatment to affect egg size, this effect is unlikely to be mediated through variation in female provisioning. Male fitness components were also found to have a significant treatment effect: M-sons had lower dry weight at eclosion, higher mating latency and lower competitive mating success. While being the first study to show both adaptive and non-adaptive effect of the paternal density in *Drosophila*, our results highlight the importance of considering paternal environment as important source of fitness variation.

Key words: Sire effect, juvenile competitive fitness, mating latency, male reproductive success, crowding adaptation

Introduction:

Parental environment has the potential to influence offspring traits and fitness through transgenerational effects. While maternal transgenerational effect (henceforth maternal effect) has been investigated in great detail due to its prevalence as well as prominence, examples of similar paternal effect is far fewer. However, there is a growing body of evidence showing the importance of the paternal trangenerational effect on fitness related traits of the offspring [1-4].It is therefore important to assess the extent of such paternal effect under various ecological scenarios and in different organisms.

There are many reports of environment dependent maternal effect mediated through variation in maternal provisioning in egg/offspring [5,6]. Females living under high density may suffer from adverse effects of crowding (such as, malnutrition) and may therefore struggle to allocate resources in maternal provisioning either in the form of stored resources in egg or lactation, which in turn would lead to poor quality progeny [7]. Alternatively, females raised in high density may strategically produce less number of eggs/progeny while investing more resources (e.g., yolk) in each of them – thereby giving the progeny a better start for the impending challenges of high density environment [8-11]. Generally, under fluctuating environment, such parental ability to optimize offspring phenotype has been conjectured to be adaptive [1, 12]. Guppy (*Poeciala reticulata*) females were found to produce larger offspring (a) under food limitation [13] and (b) when they experienced high level of competition – priming the offspring for better competitive ability [14]. The larger eggs produced by *D. melanogaster* females that grew in poor food, survive (egg to adult survivorship) better in poor food and give rise to smaller adults [11]. In contrast, Valtonen et al. (2012) found *D. melanogaster* females grown on poor quality food to produce larger offspring (adult) compared to those grown on normal food [15].

Note that much of the maternal effects discussed above are mediated through variation in resource provisioning by mothers. Not surprisingly, most of the reports of environment dependent paternal transgenerational effect (hereafter, paternal effect) come from animals with paternal provisioning

through nuptial gift transfer to the females during mating [16-20]. However, it is only recently that

studies have started to address if similar paternal effects are also present in species without paternal

provisioning. Males of a solitary Ascidian (*Styela plecata*) produced offspring phenotype

corresponding to the population density experienced by the father [4]. Female Neriid flies

(*Telostylenus angusticollis*) raised on richer diet produced larger eggs and offspring that developed

faster, while males raised on richer diet sired larger offspring with better survival rate, especially

under resource scarcity [3, 21]. Valtonen et al. (2012) reported that fathers fed on poor quality diet

sire larger sons [15]. Our previous study showed that paternal experience of rivals' number adaptively

affected offspring reproductive behaviour in *D. melanogaster* [22]. Islam et al. (1994) showed

paternal social environment to have a significant impact on offspring behavioural traits [23]. Paternal

experience of ambient temperature was also found to affect offspring fecundity in *D. melanogaster*

[24]. Low temperature was found to affect offspring phenotype in two other species of Drosophila –

D. simulans [25] and *D. serrata* [26]. Thus there is a growing body of evidence showing environment

dependent paternal effect, however, the full impact of such effect on adaptation to the relevant

ecological challenge is not yet clear.

Here we investigated the effect of paternal experience of population density on progeny behaviour

and fitness components by using *D. melanogaster* laboratory adapted populations. As discussed

previously, paternal effect has already been reported in these [22] and other populations of *D.*

melanogaster, establishing them as a relevant system to investigate the paternal effect and its

consequences on Darwinian fitness [27]. Further, laboratory adapted populations of *D. melanogaster*

can be investigated for a range of environmental parameters. Among the many components of their

ecology, population density (number of individuals per unit space) is particularly very important.

Fruit flies naturally grow in ephemeral resource patches, such as rotting fruits. Since quality of

resource patch is expected to vary a great deal – both within and across generations, crowding in

transiently available rich patches is expected to be a key component of their natural ecology.

Therefore density of adults in a patch not only determines the extent to which individuals have to

compete for limiting space (e.g., oviposition substrate) but also for other resources, such as perishing

food and suitable mates. Increase in density also leads to an increase in the chance of disease

transmission [28].In essence, density often determines the nature and intensity of selection acting on a

population and is generally addressed within the broader premises of density dependent selection [29-

31]. The laboratory adapted populations of the fruit flies have been extremely successfully used to

investigate various components of the theories of density dependent selection (see [32] for a recent

example and a review of some of the relatively older studies). Population density is known to affect

the expression of a wide range of traits both within and across life-history stages [32-37]. However,

little is known about its transgenerational consequences (but see [8]).

We subjected males to three adult density treatments and then allowed them to produce progeny by

mating the treated males to untreated dams. We then investigated the effect of the paternal density

treatment on progeny fitness components in juvenile (juvenile competitive fitness) and adult stages

(males: mating ability, mating latency, copulation duration, courtship frequency, competitive mating

success). We show that density experienced by the fathers affect both juvenile and adult fitness

component.

Methods:

All the experiments were done using a set of laboratory adapted populations of *D. melanogaster* – BL.

Full laboratory history of these populations can be found in [38]. Briefly, these are a set of five

replicate populations (BL $_{1-5}$) maintained on standard Banana-Jaggery-Yeast food, under 14-day

discrete generation cycle at 25 ^{O}C ambient temperature, 60-80% relative humidity, with population

size ~2800. Larval density is maintained at ~70 per 6-8ml food per vial (25mm×90mm,

diameter×height). Adult density is ~70 per vial for the first couple of days of their adult life and

thereafter ~2800 individuals in a ~6.4 l cage (19cm×14cm×24cm). We also used a derived population,

BL$_{st}$ which was derived from BL$_1$ by introducing an autosomal recessive marker – scarlet eye (st)

[22]. BL$_{st}$ population is maintained under a set of conditions identical to the other BL populations.

<u>Paternal treatment:</u>

Sires and dams were generated from a BL population. The design of the protocol followed to generate

the experimental flies is described in Figure 1. To generate the experimental sires and the dams, eggs

were collected from a BL population and cultured under standard density (i.e., 70 per 6-8ml food per

vial). 100 such vials were set up, of which 65 were used to collect the sires (= sire-vials) and the rest

35 for dams (dam vials). Dams were collected as virgins and held in single sex vials at a density of 25

per vial with ad lib food until the day of the sire-dam mating (see below). In the sire-vials, all the flies

were allowed to eclose. These flies were used to set up three adult density treatments – normal (N: 70

individuals per vial), medium (M: 140 individuals per vial) and high (H: 210 individuals per vial). 10

vials were set up for each of the treatments, using flies that were approximately 1 day old. These vials

were left undisturbed for 2 days, following which males from them were separated and used as sires

in the subsequent step. Here and elsewhere throughout the study, all the fly sorting, including

collection of virgins, were done under light CO_2-anaesthesia, unless mentioned otherwise.

<u>Sire-dam mating:</u>

Following the 2-day long conditioning, 25 males were randomly isolated from each adult density

treatment vials, to be used as sires. They were then combined with dams (see previous section) in

fresh food vials (25 sires + 25 dams in a vial) and allowed a time of one and half hour - sufficient time

for a single round of mating. This method of ensuring single round of mating has been previously

used [39]. In addition, mating was visually observed. 10 mating vials were set up per density

treatment. After mating, the sires were discarded and the already inseminated dams from all 10 vials

of a treatment (i.e., a total of 250 females) were transferred to a 2 litre plastic cage with food smeared

with live yeast. Three such cages were thus set up – one for each density treatment. Two days later

eggs were collected from these cages to set up the remainder of the experiments. To collect the eggs, a

fresh food plate was introduced in the cage. The dams were allowed a short window (2-3 hours) for

oviposition. Using a fine brush, eggs were counted on to a fine Agar-strip, which was then transferred

to the culture vials (see below). These eggs are hereafter referred to as treatment eggs.

<u>Measurement of egg-size:</u>

To test if the sires influenced the size of the eggs laid by the dams [40], a subset of these eggs were

frozen at -20^OC and their size was measured. For this purpose, eggs were mounted on a glass slide on

their dorsal side and photographed using Nikon (SMZ745T) and the area of the two-dimensional

elliptical outline of the eggs were measured in ImageJ, software. This area was taken as a proxy for

the size of each egg. A given egg was measured thrice and the average of these three measurements

was taken as the unit of analysis. 50 eggs per treatment were measured for this purpose.

<u>Experiment 1: Juvenile fitness assay</u>

Egg to adult survivorship was taken as a measure of Juvenile fitness. Survivorship of the treatment

eggs were measured against a back ground of a common competitor (BL_{st}) under two conditions –

crowded (C: 150 larvae per 1.5ml food in each vial) and un-crowded (UC: 70 larvae per 6ml food in

each vial). During the assay, treatmenteggs generated in the previous step were cultured with eggs

from common competitors in the ratio 1:4 (C: 30 target, 120 competitor; UC: 14 target, 56

competitor). These common competitors were collected from an untreated BL_{st} stock. On completion

of development, it was possible to identify the target progeny from the competitor progeny based on

eye colour – all progeny of the competitors were scarlet eyed whereas the target progeny were red

eyed. 10 juvenile competition vials were set up for each of the three treatments (viz., N, M and H) and

two assay conditions (i.e., 10 as C and 10 as UC for each treatment). They were left undisturbed till

12th day, by which time all surviving pupae finished eclosion. The adults were sorted based on eye

colour and counted. Juvenile fitness score (w) was calculated for each vial following the formula:

$$w = \frac{number of red eyed progeny observed}{number of red eyed progeny expected}$$

The number of red eyed progeny expected was 14 and 30 for UC and C assay conditions respectively.

<u>Experiment 2: Assay for behaviour and fitness of the sons</u>

To investigate the effect of the treatment on the male progeny, the treatment eggs were cultured in

food vials in the usual density (i.e., 70 per 6ml food in each vial) and the progeny were allowed to

develop. Upon onset of eclosion, males were collected as virgins (< 6 hours post-eclosion). Four

assays were run with these males. (a)For each treatment, 50 males were immediately frozen at -20 $^{\circ}$C

and were later dried at 60 $^{\circ}$C for 48 hours and weighed in groups of 5. (b) A separate set of males

were similarly collected and held in groups of 5 per vial for further assays. Ten such vials, for each

treatment, were set up and left undisturbed till they were 3days old. These males were then

transferred to fresh food vials (hereafter referred to as mating vials) along with five age-matched,

virgin females. Mating vials were set up without the use of anaesthesia. The females used in this step

came from the same replicate BL population and were generated under their standard maintenance

conditions, collected as virgins and held in groups of five per vial with ample food until the day of the

experiment. 10 mating vials were set up for each of the three treatments. They were observed every

two minutes starting from the time when the females were introduced and total number of mating

pairs (n_x) was noted down at each time point (x = 0, 2, 4, 6…). This was done till all the flies finished

mating. Mean mating latency (ML, time taken by a virgin pair to start mating) and mean copulation

duration (CD, duration for which a pair mated) were calculated following an algorithm mentioned

below.

$$ML = \frac{\sum(n_x - n_{x-2})x}{N}$$

196   For all values of x, until, $n_{x-2} \le n_x$.

$$CD = \frac{\sum(n_{x-2} - n_x)x}{N} - ML$$

For all values of x, until, $n_{x-2} \ge n_x$.

Occasionally, some females/males did not mate within one hour long observation. These flies were

excluded from the analysis. Mating ability (MA) is measured as the proportion of the sons

successfully copulated. MA was calculated for every single vial.

(c) Courtship frequency was quantified for the 3day old (post-eclosion) sons of the three paternal

density treatments by setting up similar mating vials as described in the previous section. Ten vials

were set up for each treatment. Therefore a total of 30 vials were observed. After allowing the first

mating, the courtship observation was initiated after a gap of approximately half an hour. Vials where

all the flies did not mate were removed from the assay. Every 45 minutes, each vial was observed for

30 seconds, during which the total number of courtship bouts (male to female) was noted down. A

total of 8 observations were taken. In *Drosophila,* courtship behaviour includes chasing, tapping,

courtship dance and song, genital licking and attempted mounting [41, 42]. Any of the above

mentioned courtship behaviours, displayed by the five males in each vial was counted as one. The

total number of independent male to female courtship displays was counted within the observation

window [43].The treatment identities were unknown to the observers to avoid observer bias. (d)

Another set of males were similarly collected and held, to be used for quantifying their mating success

under competitive condition (CMS, Competitive mating success). This was done by setting up mating

vials with five 3day old target males, five competitor males (BL_{st}) and five virgin females (BL_{st}). Ten

such mating vials were set up for each of the three treatments. After allowing a single round of mating

for all the females in a mating vial, the females were individually transferred to oviposition test tubes

(12mm diameter × 75mm height) with ample food. The females were allowed to oviposit for 18

hours. Following oviposition, the females were discarded and the tubes were retained to allow the

progeny to develop and eclose. For each female, the identity of their mate (whether target/competitor)

was ascertained by observing the eye colour of the progeny. Progeny sired by target males were red

eyed whereas those sired by competitors were scarlet eyed. For a given vial, average CMS of the five

target males in the vial was calculated as the proportion of the females mated to target males (i.e.,

produced red eyed offspring).

<u>Experimental replications and data analyses:</u>

The entire study was carried out in three randomized blocks, using three different BL populations -

BL_1, BL_3 and BL_5.The blocks were handled in separate days. Number of replications within each

block has been mentioned in the previous sections along with the assay design. Except the egg size

and dry body weight assay, all the experimental replication was done at the level of assay mating vials

or juvenile competition vials. All the assays had 10 replicate vials. Vial means were used as the unit

of analysis. For egg size assay, size of each egg was used as the unit of analysis. For dry body weight,

weight of groups of 5 individuals was used as the unit of analysis. Data were analysed using mixed

model Analysis of Variance (ANOVA). Block was treated as random factor, while paternal density

treatment and assay density (wherever applicable) were treated as fixed factors. Multiple comparisons

were done using Tukey's HSD. All the analyses were done in Statistica, version 10 (Statsoft, Tulsa,

OK, U.S.A.).

Results:

Variation in size of the eggs represents variation in maternal provisioning. The effect of the paternal

density treatment on size of the eggs produced by the dams was not significant (Table 1, mean ±SE,

μm^2, N: 80039.1 ±387.5; M: 79967.4 ±415.3; H: 79611.9 ±411.4). The juvenile competitive fitness

assay quantified overall egg to adult survival of the target juveniles compared to the same of juveniles

from a common background (common competitors). While the data from un-crowded assay condition

reflects the baseline survivorship, those from crowded assay condition represents difference in

juvenile competitive ability across the three paternal density treatments. Paternal density treatment

had a significant effect on Juvenile fitness (Table 1). While there was no significant difference

between N and M-treatments, H-treatment had 8.9% higher juvenile fitness compared to that of the N-

treatment. This relative advantage of the H-treatment was only evident under larval crowding, i.e., C-

assay density (Figure 2), indicating competitive superiority of that H-juveniles. However, the paternal

treatment × assay density interaction was marginally non-significant (Table 1).

We only quantified the effect of the paternal density on male offspring. We found a significant effect

of the treatment on dry body weight, ML and CMS (Table 1, Figure 3). Multiple comparisons using

Tukey's HSD indicated that dry body weight of the M-sons were significantly less than that of the N-

sons, with M-sons having 8.7% lower mean dry body weight. The difference between the dry body

weight of the H and N-sons was not statistically significant. Hence the M-sons were significantly

smaller compared to the other two treatments. In the mating assay, though we found some males to

fail in acquiring mating, there was no effect of the treatment on mating ability of the sons (MA: mean

±SE, N: 0.91 ±0.04; M: 0.91 ±0.04; H: 0.93 ±0.04). The M-treatment sons showed significantly

higher (approximately 35%) ML compared to that showed by the N-treatment sons. While H-

treatment also showed 16% higher ML compared to N-treatment, this difference was not significant.

Therefore M-sons took longer time to start mating with virgin females indicating females' reluctance

to accept them as mate due to either poor performance in courtship or small size. This relative

disadvantage of the M-sons was also evident in terms of their competitive ability in mating

competitions. Multiple comparisons on the CMS results indicated that the M-sons had significantly

lower CMS compared to H and N-treatments. CMS of the M-sons was approximately 34% less than

that of the N-sons. This is however, not due to a reduced courtship performance by the M-sons as we

found the effect of the treatment on CF (mean ±SE, N: 6.7 ±0.6; M: 6.8 ±0.6; H: 7.6 ±0.8) to be non-

significant. We also did not find any effect of the treatment on CD (mean ±SE, minutes, N: 18.6 ±0.4;

M: 17.6 ±0.4; H: 17.9 ±0.4), potentially indicating the lack of the treatment effect on post-copulatory

traits of the sons (Table 1).

Discussion:

Given that very few studies have shown the effect of paternal environment on offspring fitness

components, there were two main objectives of the present study – (a) to assess if paternal exposure to

varying population density affected progeny traits; if yes, then (b) to evaluate the adaptive

significance of such effect. The results clearly showed that at sufficiently high density, males had an

adaptive paternal effect on juvenile competitive fitness. As we did not find any effect of our treatment

on size of the eggs produced by the dams, such paternal effect is unlikely to be mediated by variation

in provisioning by the females. We further show that at intermediate density, males sire smaller sons

which are inferior in acquiring mates. Interestingly, such maladaptive effect of paternal density on

offspring adult fitness was not detected at high density.

In holometabolus insects like fruit flies, juvenile (larva and pupa) survival constitutes one of the most

important components of fitness [31]. In addition, juvenile ecology may also have a major effect on

the life-history and fitness components of the adult stage (Heat shock [44]; cold shock [45, 46];

crowding [47, 32, 36]. The observed paternal effect on juvenile competitive fitness therefore is

extremely consequential. Some relatively recent studies have pointed out that evolving parental ability

to optimize offspring fitness related traits can be adaptation to ecological challenges [48], including

crowding [4]. Given that fruit fly natural ecology regularly involves adult and larval crowding, the

observed paternal effect on juvenile competitive fitness can indicate males' adaptation to crowding.

Interestingly, we observed the paternal effect on juvenile competitive fitness, only at the highest

density, which may indicate a certain threshold density beyond which such paternal effect starts

affecting offspring traits. In addition, when assayed under un-crowded condition the progeny from the

three sire treatments do not show any measurable difference in their egg-to-adult survival. This

suggests that the juvenile competitive ability rather than baseline juvenile viability was affected by the

treatment. Since a number of traits (e.g., feeding rate, waste tolerance, development time etc.) affect

juvenile competitive ability in these flies, it will be interesting to find out the trait responsible for

better competitive ability of the H-sons in our study.

In a wide range of species including *Drosophila melanogaster*, maternal exposure to high density or

poor nutrition has been found to affect offspring fitness components [8, 11, 15]. Such effects can

either be beneficial [10, 11, 14, 49, 50] or detrimental [51] depending on the component of the fitness

under consideration and the prevailing condition. As maternal provisioning and other maternal effects

play vital roles in offspring survival and performance, such maternal density/nutrition effect is not

surprising. However, what is not intuitive is the paternal density to have similar impact on offspring

fitness, as our results suggest, given that *Drosophila* males do not pass on any nutrition to the

offspring. It is well known in the *Drosophila* literature that even the laboratory populations harbour

heritable genetic variation in survival under crowding both as adults and juveniles (see [32] and the

references therein for an updated review). Therefore, one possibility is that genetically superior males,

which are better at surviving under high density, may produce offspring which are better both as

juveniles, explaining at least part of our observations. Though larval competitive ability is known to

respond to experimental evolution, indicating heritable genetic variation [30, 31], such heritable

variation is very unlikely to have led to the observed treatment effect on juvenile competitive fitness.

This is because (a) in our assay, we recorded very little mortality in males during the treatment,

indicating negligible hard selection. In addition, we also ensured that there was no soft selection by

randomly picking the set of males from the treatment vials to use them as sires. (b) Even if there was

selection in the current experimental design, the selection is likely to be weak. Such weak selection is

unlikely to explain the observed differences in some of the traits (viz., 8.9% increase in juvenile

competitive fitness, 35% increase in mating latency), especially within one generation. Alternatively,

males may alter maternal provisioning and thereby indirectly affect offspring fitness components [8,

11]. We, however, did not find any measurable difference in the size of the eggs produced by females

mated to the males belonging to the three treatments, making variation in maternal provisioning an

unlikely explanation. Therefore, although sire-effect on the quality of the eggs produced by the

females cannot be completely ruled out, our results tentatively point at non-genetic paternal effect [1]

as the potential cause behind the observed effect of the treatment.

While we found adaptive paternal effect on juvenile performance, adult performance however, was

found to have a significant maladaptive effect of paternal density. Males that experienced

intermediate density were found to sire sons which (a) are smaller, (b) take longer time to start mating

and (c) have lower mating success. Since we did not find any effect of the treatment on courtship

frequency, reduced mating success and increased mating latency was a likely outcome of females'

reluctance to accept relatively smaller males as their mates, a known fitness consequence of reduced

size in *Drosophila* males [52, 53]. Body size has been reported to be affected by transgenerational

paternal effect in another Dipteran – *Telostylinus angusticollis* [21]. Unlike the maladaptive effect

found in our study, the paternal effect on body size reported by Bonduriansky and Head (2007)[21],

however, was adaptive, especially under certain prevailing conditions. Though prima facie, the

observed body size reduction appears to be maladaptive, it will be interesting to investigate its fitness

consequence under varying adult density. Interestingly, this effect was found only at intermediate

density and not in the high density treatment. At this point, it is, however, difficult to suggest any

reason for such specific expression of the paternal effect at intermediate density.

As variation in population density and crowding related ecological challenges are common in almost

all organisms, including fruit flies, paternal effect of the nature reported here is important to

understand. Though paternal ability to optimize offspring traits is likely to be adaptive, especially

under fluctuating environment, the results reported here show that paternal effect can be both adaptive

and maladaptive. To the best of our knowledge this is the first evidence of the effect of paternal

density on juvenile and adult fitness components in *D. melanogaster*. Importantly, our results

emphasize the importance of considering paternal effect as a source of variation in fitness related

traits. The full impact of such paternal effect in the evolution of life-history traits and the underlying

mechanisms are emerging as an important topic of discussion, which is likely to see an increasing

attention in years to come.

Acknowledgements:

We are immensely thankful to N. G. Prasad for sharing the fly populations, using which the BL

populations were created and also for his critical comments on a previous version of this manuscript.

We are also thankful to Syed Zeeshan Ali and Vanika Gupta for their critical comments on a previous

version of the manuscript. TV and PD thank Indian Institute of Science Education and Research

Berhampur for financial assistance in the form of the Institute Scholarship for Ph.D. programme. The

study was supported by a research grant from Department of Science and Technology (INSPIRE

Faculty award, Grant no. DST/INSPIRE/04/2013/000520).

Author contribution: BN conceived the idea. BN, PD and SS designed the experiments. PD, SS, TV

and AA performed the experiments. Analysis of the data was done by BN and PD. Manuscript was

written by BN with assistance from PD and SS.

Conflict of interest: The authors declare no conflict of interest.

Data intended to be archived in: Dryad

References:

1. Bonduriansky R, Day T. 2009 Nongenetic inheritance and its evolutionary implications. *Annu. Rev. Ecol. Evol. Syst.* **40**, 103–125. (doi: 10.1146/annurev.ecolsys.39.110707.173441)

2. Friberg U, Stewart AD, Rice WR. 2012 X- and Y-chromosome linked paternal effects on a life-history trait. *Biol. Lett.* **8**, 71-73 (doi: 10.1098/rsbl.2011.0608)

3. Adler MI, Bonduriansky R. 2013 Paternal Effects on Offspring Fitness Reflect Father's Social Environment. *Evol. Biol.* **40**, 288-292. (doi: 10.1007/s11692-012-9211-6)

4. Crean AJ, Dwyer JM, Marshall DJ. 2013 Adaptive paternal effects? Experimental evidence that the paternal environment affects offspring performance. *Ecology* **94**, 2575-2582. (doi: 10.1890/13-0184.1)

5. Rossiter MC. 1996 Incidences and consequences of inherited environmental effects. *Annu. Rev. Ecol. Syst.* **27,** 451-476. (doi:10.1146/annurev.ecolsys.27.1.451)

6. Mousseau TA, Fox, CW. 1998 The adaptive significance of maternal effects. *Trends in Ecology and Evolution* **13**, 403-407. (doi: 10.1016/S0169-5347(98)01472-4)

7. Christian JJ, Lemunyan CD. 1958 Adverse effects of crowding on lactation and reproduction of mice and two generations of their progeny. *Endocrinology* **63**, 517-529. (doi: 10.1210/endo-63-5-517)

8. Prasad NG, Shakarad M, Rajamani M, Joshi A. 2003 Interaction between the effects of maternal and larval nutritional levels on pre-adult survival in *Drosophila melanogaster*. *Evol. Ecol. Res.* **5**, 903-911.

9. Holbrook GL, Schal C. 2004 Maternal investment affects offspring phenotypic plasticity in a viviparous cockroach. *Proc. Natl. Acad. Sci. U S A.* **101**, 5595-5597. (doi: 10.1073/pnas.0400209101)

10. Mitchell SE, Read AF. 2005 Poor maternal environment enhances offspring disease resistance in an invertebrate. *Proc. Biol. Sci.***272**, 2601–2607. (doi: 10.1098/rspb.2005.3253)

11. Vijendravarma RK, Narasimha S, Kawecki TJ. 2010 Effects of parental larval diet on egg size and offspring traits in *Drosophila*. *Biol. Lett.* **6**, 238-241. (doi: 10.1098/rsbl.2009.0754)

12. Kuijper B, Hoyle RB. 2015 When to rely on maternal effects and when on phenotypic

plasticity? *Evolution* **69**, 950-968. (doi:10.1111/evo.12635)

13. Reznick DN, Reznick YA. 1993 The influence of fluctuating resources on life-history:

Patterns of allocation and plasticity in female Guppies. *Ecology* **74**, 2011-2019. (doi:

10.2307/1940844)

14. Bashey F. 2006 Cross-generational environmental effects and the evolution of offspring size

in the Trinidadian Guppy *Poecilia reticulate*. *Evolution* **60**: 348-361. (doi:10.1111/j.0014-

3820.2006.tb01111.x)

15. Valtonen TM, Kangassal K, Pölkki M, Rantala MJ. 2012 Transgenerational Effects of

Parental Larval Diet on Offspring Development Time, Adult Body Size and Pathogen

Resistance in *Drosophila melanogaster. PLoS ONE* **7**(**2**), e31611

.(doi:10.1371/journal.pone.0031611)

16. Zeh DW, Smith RL. 1995 Paternal investment by terrestrial arthropods. *Am. Zool.* **25**, 785-

805. (doi: 10.1093/icb/25.3.785)

17. Dussourd DE, Ubik K, Harvis C, Resch J, Meinwald J. and Eisner T. 1988 Biparental

defensive endowment of eggs with acquired plant alkaloid in the moth *Utetheisaornatrix*.

Proc. Natl. Acad. Sci. U.S.A. **85**, 5992-5996.

18. Gwynne DT. 1988 Courtship feeding in katydids benefits the mating male's offspring. *Behav.*

Ecol. Sociobiol. **23**, 373-377. (doi: 10.1007/BF00303711)

19. Smedly SR, Eisner T. 1996 Sodium: A male moth's gift to its offspring. *Proc. Natl. Acad. Sci.*

U.S.A. **93**, 809-813. (doi: 10.1073/pnas.93.2.809)

20. Vahed K. 1998 The function of nuptial feeding in insects: a review of empirical studies. *Biol.*

Rev. **73,** 43-78. (doi: 10.1111/j.1469-185X.1997.tb00025.x)

21. Bondurianslky R, Head M. 2007 Maternal and paternal condition effects on offspring

phenotype in *Telostylinus angusticollis* (Diptera: Neriidae). *Int. J. Evol. Biol.* **20**, 2379-2388.

(doi:10.1111/j.1420-9101.2007.01419.x)

22. Dasgupta P, Halder S, Nandy B. 2016 Paternal social experience affects male reproductive

behaviour in *Drosophila melanogaster*. *J. Genet.* **95**, 725-727. (doi: 10.1007/s12041-016-

0670-x)

23. Islam MS, Roessingh P, Simpson SJ, McCaffery AR. 1994 Parental effects on the behaviour

and colouration of nymphs of the desert locust *Schistocercagregaria*. *J. Insect Physiol.* **40,**

173-181. (doi: 10.1016/0022-1910(94)90089-2)

24. Huey RB, Wakefield T, Crill WD, Gilchrist G. 1995 Within- and between-generation effects

of temperature on early fecundity of Drosophila melanogaster. *Heredity* **74**, 216-223. (doi:

10.2307/2410661)

25. Watson MJO, Hoffmann AA. 1995 Cross-Generation Effects for Cold Resistance in Tropical

Populations of *Drosophila melanogaster* and *Drosophila simulans. Aust. J. Zool.* **43,** 51 – 58.

26. Magiafoglou A, Hoffmann AA. 2003 Cross-generation effects due to cold exposure in

Drosophila serrata. Funct. Ecol. **17***,* 664–672. (doi: 10.1046/j.1365-2435.2003.00774.x)

27. William RR, Stewart AD, Morrow EH, Linder JE, Orteiza N, Byrne PG. 2006 Assessing

sexual conflict in the *Drosophila melanogaster* laboratory model system. *Philos. Trans. R.*

Soc. Lond. B. Biol. Sci. **361**, 287-299(doi:10.1098/rstb.2005.1787)

28. Barnes AI, Siva MT. 2000 Density-dependent prophylaxis in the mealworm beetle *Tenebrio*

molitor L. (Coleoptera: Tenebrionidae): cuticular melanization is an indicator of investment

in immunity. *Proc. R. Soc. Lond. B* **267**, 177-182. (doi: 10.1098/rspb.2000.0984)

29. MacArthur RH, Wilson EO. 1967 *The theory of island biogeography.* Princeton Univ. Press,

Princeton, NJ.

30. Mueller LD. 1997 Theoretical and Empirical Examination of Density-Dependent Selection.

Annu. Rev. Ecol. Syst. **28,** 269-288. (doi: 10.1146/annurev.ecolsys.28.1.269)

31. Prasad NG, Joshi A. 2003 What have two decades of laboratory life-history evolution studies

on *Drosophila melanogaster* taught us?*J. Genet.* **82**, 45–76. (doi:10.1007/BF02715881)

32. Sarangi M, Nagarajan A, Dey S, Bose, J, Joshi A. 2016 Evolution of increased larval

competitive ability in *Drosophila melanogaster* without increased larval feeding rate. *J.*

Genet. **95**, 491-503. (doi: 10.1007/s12041-016-0656-8)

33. Mueller LD, Sweet VF. 1986 Density-dependent natural selection in *Drosophila*: Evolution of

pupation height. *Evolution* **40**, 1354-1356. (doi: 10.1111/j.1558-5646.1986.tb05761.x)

34. Mueller LD, Guo PZ, Ayala FJ. 1991 Density-dependent natural selection produces trade-offs

in life history traits. *Science* **253**, 433-435. (doi: 10.1126/science.1907401)

35. Nagarajan A, Natarajan SB, Jayaram M, Joshi A. 2016 Adaptation to larval crowding in

Drosophila ananassae and *Drosophila nasutanasuta*: increased larval competitive ability

without increased larval feeding rate. *J. Genet.* **95,** 411-425 (doi: 10.1007/s12041-016-0655-

9)

36. Shenoi VN, Banerjee SM, Guruswamy B, Sen S, Ali ZS, Prasad NG. 2016 *Drosophila*

melanogaster males evolve increased courtship as a correlated response to larval

crowding. *Anim. Behav.* **120**, 183 -193. (doi: 10.1016/j.anbehav.2016.08.004)

37. Shenoi VN, Prasad NG. 2016 Local adaptation to developmental density does not lead to

higher mating success in *Drosophila melanogaster*. *Int.J. Evol. Biol*, **29**, 407-417. (doi:

10.1111/jeb.12927)

38. Nandy B, Dasgupta P, Halder S, Verma T. 2016 Plasticity in aggression and the correlated

changes in the cost of reproduction in male *Drosophila melanogaster*. *Anim. Behav,* **114**, 3-9.

(doi:10.1016/j.anbehav.2016.01.019)

39. Nandy B, Joshi A, Ali SZ, Sen S, Prasad NG. 2012 Degree of adaptive male mate choice is

positively correlated with female quality variance. *Sci. Rep.* **2**: 447.

40. Pischedda A, Stewart AD, Little MK, Rice WR. 2010 Male genotype influences female

reproductive investment in *Drosophila melanogaster*. *Proc R Soc B* **278**,2165–2172(doi:

10.1098/rspb.2010.2272)

41. Bastock M, Manning A. 1955 The Courtship of *Drosophila melanogaster*. *Behaviour* **8**, 85-

111. (doi:10.1163/156853955X00184)

42. Sokolowski MB. 2010 Social interactions in "simple" model systems. *Neuron*, **65**,780-794.

(doi:10.1016/j.neuron.2010.03.007)

43. Nandy B, Gupta V, Sen S, Udaykumar N, Samant MA, Ali SZ , Prasad NG. 2013 Evolution

of mate-harm, longevity and behaviour in male fruit flies subjected to different levels of

interlocus conflict. *BMC EvolBiol* **13**, 212. (doi: 10.1186/1471-2148-13-212)

44. Khazaeli AA, Tatar M, Pletcher SD, Curtsinger JW. 1997 Heat-induced longevity extension

in Drosophila. I. Heat treatment, mortality, and thermotolerance. *J. Gerontol.* **52A**, B48–B52.

45. Singh K, Kochar E, Prasad NG. 2015 Egg viability, mating frequency and male mating ability

evolve in populations of *Drosophila melanogaster* selected for resistance to cold shock. *PLoS*

ONE **10**, e0129992. (doi:10.5061/dryad.vq080)

46. Singh K, Prasad NG. 2016 Evolution of pre- and post-copulatory traits in female *Drosophila*

melanogaster as a correlated response to selection for resistance to cold stress. *J. Insect*

Physiol. **91-92**, 26-33. (doi: 10.1016/j.jinsphys.2016.06.005)

47. Joshi A, Mueller LD. 1988 Evolution of higher feeding rate in Drosophila due to density-

dependent natural selection. *Evolution*, **42**, 1090-1093. (doi: 10.1111/j.1558-

5646.1997.tb02429.x)

48. Galloway LF, Etterson JR. 2007 Transgenerational plasticity is adaptive in the wild. *Science*

318, 1134-1136. (doi:10.1126/science.1148766)

49. Allen RM, Buckley YM, Marshall DJ. 2008 Offspring Size Plasticity in Response to

Intraspecific Competition: An Adaptive Maternal Effect across Life-History Stages. *Am.*

*Nat.***171**, 225-37. (doi: 10.1086/524952)

50. Gorbi G, Moroni F, Sei S, Rossi V. 2011 Anticipatory maternal effects in two different clones

of *Daphnia magna* in response to food shortage. *J. Limnol.* **70**, 222-230. (doi: 10.3274/JL11-

70-2-05)

51. Meylan S, Clobert J, Sinervo B. 2007 Adaptive significance of maternal induction of

density-dependent phenotypes. *Oikos* **116**, 650-661. (doi: 10.1111/j.0030-1299.2007.15432.x)

52. Partridge L, Ewing A, Chandler A. 1987 Male size and mating success in *Drosophila*

melanogaster: the roles of male and female behaviour. *Anim. Behav.***35**, 555-562. (doi:

10.1016/S0003-3472(87)80281-6)

506 53. Jagadeeshan S, Shah U, Chakrabarti D, Singh RS. 2015 Female Choice or Male Sex Drive?

507 The Advantages of Male Body Size during Mating in *Drosophila Melanogaster. PLoS ONE*

508 **10**, e0144672. (doi: 10.1371/journal.pone.0144672)

Trait	Effect	SS	DF	MS	DF Den	MS Den	F	p
Egg size	Paternal density (PD)	1.75×10^7	2	8.75×10^6	4.02	3.53×10^6	2.48	0.20
	Block	9.84×10^8	2	4.92×10^8	4.03	3.54×10^6	139.11	**<0.01**
	PD×Block	1.41×10^7	4	3.53×10^6	438.00	2.25×10^7	0.16	0.96
Juvenile fitness	Paternal density (PD)	0.21	2	0.11	4.03	0.01	9.93	**0.03**
	Assay density (AD)	0.97	1	0.97	2.00	0.10	9.31	0.09
	Block	0.40	2	0.20	1.58	0.09	2.15	0.35
	PD×AD	0.23	2	0.11	4.01	0.02	5.42	0.07
	PD×Block	0.04	4	0.01	4.00	0.02	0.51	0.74
	AD×Block	0.21	2	0.10	4.00	0.02	4.96	0.08
	PD × AD × Block	0.08	4	0.02	148.00	0.02	1.24	0.30
Dry body weight	Paternal density (PD)	2.67×10^{-7}	2	1.34×10^{-7}	4.00	6.22×10^{-9}	21.49	**< 0.01**
	Block	7.78×10^{-7}	2	3.89×10^{-7}	4.00	6.22×10^{-9}	62.50	**< 0.01**
	PD×Block	2.49×10^{-8}	4	6.22×10^{-9}	80	1.27×10^{-8}	0.49	0.74
Mating latency	Paternal density (PD)	41.810	2	20.90	4	1.12	18.59	**0.01**
	Block	51.158	2	25.58	4	1.12	22.75	**0.01**
	PD×Block	4.477	4	1.12	77	4.59	0.24	0.91
Copulation duration	Paternal density (PD)	14.43	2	7.21	4	1.81	3.98	0.11
	Block	80.87	2	40.44	4	1.81	22.32	**0.01**
	PD×Block	7.22	4	1.81	77	5.51	0.33	0.86
Mating ability	Paternal density (PD)	0.009259	2	0.00	4	0.04	0.10	0.90
	Block	0.046678	2	0.02	4	0.04	0.52	0.63
	PD×Block	0.179963	4	0.04	77	0.02	2.64	**0.04**
Courtship frequency	Paternal density (PD)	13.28	2	6.64	4.03	10.79	0.62	0.58
	Block	500.78	2	250.39	4.00	10.81	23.16	**0.01**
	PD×Block	43.24	4	10.81	73.00	7.36	1.47	0.22
Competitive mating success	Paternal density (PD)	0.64	2	0.32	4.03	0.02	19.47	**0.01**
	Block	0.09	2	0.05	4.06	0.02	2.86	0.17
	PD×Block	0.07	4	0.02	75.00	0.04	0.41	0.80

Table 1: Summary of results of mixed model ANOVA on the various traits under investigation.

Paternal density and assay density (wherever applicable) were considered as fixed factor, while block

as random factor. All tests were done considering α=0.05 and significant p-values are mentioned in

bold fonts.

Figure 1: The design of the assay. The schematic diagram shows the design of the entire study, which

spanned two generations. Treatment was given in the paternal generation. Untreated dams were mated

to the treated sires, followed by the collection of eggs from the dams. Assays were done with the eggs

and the offspring emerging out of the eggs. Some eggs were subjected to mixed culture (along with

competitor eggs) and juvenile competitive fitness vials were set up. Some eggs were cultured as

monocultures (without any competitor eggs) – male progeny emerging from these vials were used for

further assays, such as, mating ability, mating latency, copulation duration, competitive mating

success and courtship frequency.

Figure 2: Effect of the paternal density treatment on juvenile competitive fitness, under crowded and

un-crowded assay conditions. Target eggs (eggs produced by dams mated to treatment sires) were

cultured with competitor eggs (eggs produced by untreated females and males) in juvenile

competition vials – under un-crowded and crowded conditions. Proportion of the target progeny

successfully emerging as adults is considered as the measure of juvenile competitive fitness. Black,

blue and red colour coding represent the progeny of N, M and H males respectively. Paternal

treatment was found to have a significant effect only when the assay was carried out under crowded

condition (represented by *asterisk). The entire experiment was done following a randomized block

design and the data were analysed using three factor mixed model ANOVA with paternal treatment

and assay condition as fixed factors and block as random factor.

Figure 3: Effect of the paternal density treatment on male offspring. (a) Dry body weight at eclosion

of the M-sons was found to be significantly less than that of the other treatments; (b) Mean mating

latency (time taken by a virgin male-female pair to start copulation) of the was significantly higher in

M-sons; (c) When subjected to competition with (untreated) rival males, the competitive mating

success of the M-sons was found to be significantly less than those of the other treatment sons. The

blue broken line indicates the expected value of CMS if there is no mating bias. Black, blue and red

colour coding represent data from the progeny of N, M and H males respectively. Treatments not

sharing common alphabet were found to be significantly different from each other. The entire

experiment was done following a randomized block design and the data were analysed using three

factor mixed model ANOVA with paternal treatment and assay condition as fixed factors and block as

random factor.