Recurrent pregnancy loss is associated with a pro-senescent decidual response during

the peri-implantation window

Emma S Lucas[1,2†], Pavle Vrljicak[1,2†], Joanne Muter[1,2], Maria M Diniz-da-Costa[1,2], Paul J

Brighton[2], Chow-Seng Kong[2], Julia Lipecki[3], Katherine Fishwick[2], Joshua Odendaal[2], Lauren

J. Ewington[2], Siobhan Quenby[1,2], Sascha Ott[1,4], and Jan J Brosens[1,2,*]

[1]Tommy's National Centre for Miscarriage Research, University Hospitals Coventry &

Warwickshire, Coventry, CV2 2DX, United Kingdom.

[2]Division of Biomedical Sciences, Clinical Sciences Research Laboratories, Warwick Medical

School, University of Warwick, Coventry CV2 2DX, United Kingdom.

[3]School of Life Sciences, Gibbet Hill Campus, University of Warwick, Coventry CV4 7AL,

United Kingdom.

[4]Department of Computer Science, University of Warwick, Coventry, CV4 7AL, United

Kingdom.

[*]Corresponding Author and Lead Contact: Jan Brosens M.D., Ph.D. Clinical Sciences

Research Laboratories, Warwick Medical School, University of Warwick, Coventry CV2 2DX,

United Kingdom. Tel: +44 2476968704; FAX: +44 2476968653; Email:

J.J.Brosens@warwick.ac.uk

[†]These authors contributed equally

Abstract

Breakdown of the feto-maternal interface in early pregnancy causes miscarriage. The cycling endometrium becomes poised to transition to a pregnant state during the midluteal implantation window, coinciding with differentiation of stromal cells into decidual cells (DC) and emergence of senescent decidual cells (snDC). Emerging evidence suggests that DC engage uterine natural killer cells to eliminate their senescent counterparts, thus enabling formation of a robust decidual matrix in pregnancy. To examine if failure to constrain snDC during the peri-implantation window increases the risk of miscarriage, we reconstructed the decidual pathway at single-cell level *in vitro* and demonstrated that, without immune surveillance, secondary senescence rapidly transforms DC into progesterone-resistant cells that abundantly express extracellular matrix remodelling factors. Additional single-cell analysis of midluteal endometrium identified *DIO2* and *SCARA5* as marker genes of a diverging decidual response *in vivo*. Finally, we report a conspicuous link between a pro-senescent decidual response in luteal phase endometrium and recurrent pregnancy loss, suggesting that pre-pregnancy screening and intervention may reduce the burden of miscarriage.

Introduction

Approximately 15% of clinical pregnancies result in miscarriage [1], most often during the first trimester. Fetal chromosomal abnormalities account for 50-60% of sporadic miscarriages [2], although the incidence is lower in recurrent pregnancy loss (RPL) [3,4], defined as two or more losses [5,6]. Further, with each additional miscarriage, the frequency of euploid loss increases whereas the likelihood of a successful pregnancy decreases [7], indicating that uterine factors drive higher-order miscarriages. Unfortunately, few interventions improve live birth rates in RPL [5], reflecting that in most cases the underlying mechanisms are incompletely understood.

Following the postovulatory rise in circulating progesterone levels, the endometrium becomes transiently receptive to embryo implantation during the midluteal phase of the cycle. This implantation window also heralds the start of intense tissue remodelling [8], driven in the stroma by differentiation of endometrial stromal cells (EnSC) into specialized decidual cells (DC) and accumulation of uterine natural killer (uNK) cells [9]. Upon embryo implantation, DC rapidly encapsulate the conceptus [10], engage in embryo biosensing [11], and then form a decidual matrix that controls trophoblast invasion [12]. At a molecular level, decidual transformation of EnSC encompasses genome-wide remodelling of the chromatin landscape [13], reprogramming of multiple signalling pathways [14-16], and activation of decidual gene networks [17,18]. This multistep differentiation process starts with an evolutionarily conserved acute cellular stress response [19], marked by a burst of reactive oxygen species (ROS) and release of proinflammatory cytokines [9,20,21]. After a lag period of several days, EnSC lose their fibroblastic appearance and emerge as secretory DC with abundant cytoplasm and prominent endoplasmic reticulum [8]. A hallmark of DC is resistance to multiple stress signals. Several mechanisms underpin decidual stress resistance, including silencing of the c-Jun N-terminal kinase (JNK) pathway and upregulation of various stress defence proteins and ROS scavengers [8,15,22]. In addition, DC highly express 11β-hydroxysteroid dehydrogenase type 1 [23], which converts inactive cortisone into cortisol, a potent anti-inflammatory glucocorticoid. Thus, compared to EnSC, DC are exquisitely adapted to withstand the hyperinflammation and stress associated with deep haemochorial placentation.

Recently, we demonstrated that decidualization also results in the emergence of senescent

decidual cells (snDC), both *in vitro* and *in vivo* [9,24]. In the stroma, the abundance of cells

expressing p16^{INK4}, a tumour suppressor and canonical senescence marker, peaks transiently

during the midluteal phase before rising again prior to menstruation [9]. Cellular senescence is

defined by a state of permanent cell-cycle arrest and prominent secretion of various bioactive

molecules, including ROS, extracellular matrix (ECM) remodelling proteins, proinflammatory

cytokines, chemokines and growth factors, referred to as senescence-associated secretory

phenotype (SASP) [25-27]. Different types of senescent cells underpin pathological and

physiological processes. Chronic senescent cells accumulate progressively in response to

various stressors and cause gradual loss of organ function during ageing and in age-related

diseases mediated by the deleterious effects of the SASP on tissue homeostasis [25,27]. By

contrast, acute senescent cells are linked to biological processes that involve programmed

tissue remodelling, including embryogenesis and wound healing [27-29]. They are induced in

response to specific signals, produce a transient SASP with defined paracrine functions, and

are promptly cleared by immune cells [25]. Recently we demonstrated that snDC exhibit

hallmarks of acute senescent cells [9]. First, DC and snDC both emerge in response to FOXO1

activation, a pivotal decidual transcription factor downstream of the protein kinase A (PKA)

and progesterone signalling pathways. Second, their associated SASP critically amplifies the

initial decidual inflammatory response, which not only drives differentiation of EnSC but is also

linked to induction of key receptivity genes [21]. Further, we have provided evidence that DC

recruit and activate uNK cells, which in turn may eliminate snDC through perforin- and

granzyme-containing granule exocytosis [9,24].

Although snDC constitute a relatively minor and variable stromal population in midluteal

endometrium, they have the potential to impact profoundly on the unfolding decidual response

in a manner that either promotes or precludes pregnancy progression. For example, a

transient SASP associated with acute senescent cells has been shown to promote tissue

plasticity by expanding resident progenitor populations [9,30]. By contrast, senescent cells that

persist (i.e. chronic senescent cells) can induce senescence in neighbouring cells though

juxtracrine signalling (termed 'secondary' or 'bystander' senescence), leading to

spatiotemporal propagation of the phenotype and loss of tissue function [25,31,32].

Recently we reported loss of clonal mesenchymal stem-like cells (MSC) in midluteal

endometrium of RPL patients but how this is linked to subsequent breakdown of the decidual-

placental interface in pregnancy is unclear [33]. We hypothesized that MSC deficiency may drive

a pro-senescent decidual response during the peri-implantation window, culminating in

chronic inflammation in early pregnancy, proteolysis of the decidual-placental interface, and

miscarriage. To test this conjecture, we first performed single-cell transcriptomic analysis on

decidualizing primary EnSC cultures, mapped the emergence of DC and snDC, and identified

co-regulatory gene networks underpinning the multistep decidual pathway. We then extended

the single-cell analysis to peri-implantation endometrial biopsies and screened for stroma-

specific marker genes that signal divergence of the decidual response towards a pro-

senescent decidual state. The expression of two marker genes, *DIO2* and *SCARA5*, and the

abundance of uNK cells were then analysed in peri-implantation endometrial biopsies from 89

RPL patients and 90 control subjects.

Results

Single-cell analysis of the decidual pathway *in vitro*

To identify putative marker genes of DC and snDC, we first reconstructed the decidual

pathway *in vitro* using single-cell transcriptomics. As depicted schematically in Figure 1a,

primary EnSC were decidualized with a progestin (medroxyprogesterone acetate, MPA) and a

cyclic adenosine monophosphate analogue (8-bromo-cAMP, cAMP) for 8 days. The

differentiation signal was then withdrawn for 2 days to assess progesterone/cAMP-

dependency of decidual subsets. Cells were recovered every 48 h and subjected to single-cell

analysis using nanoliter droplet barcoding and high-throughput RNA-sequencing [34].

Approximately 800 cells were sequenced per timepoint, yielding on average 1,282 genes per

cell. After computational quality control (Supplementary Figure S1), 4,580 cells were assigned

to 7 transcriptional cell states using Shared Nearest Neighbour (SNN) and *t*-Distributed

Stochastic Neighbour Embedding (t-SNE) methods. Figure 1b shows cells colour-coded by

day of treatment and Figure 1c by transcriptional state (S1-7). The top 10 differentially

expressed genes (DEG) between the 7 cell states are presented as a heatmap

(Supplementary Figure S2 & Table S1). Apart from a discrete population of proliferative cells

(S1), the bulk of undifferentiated EnSC were in S2. By day 2 of decidualization, most cells had

transitioned to S3, which differed from S2 by 898 DEG (Supplementary Table S2). This

precipitous transcriptomic response is in keeping with an acute cellular stress response and

coincided with a transient rise in IL-6 and IL-8 secretion independent of transcription

(Supplementary Figure S3). Decidualizing cells then progressed in synchrony to S4 by day 4

after which they segregated into two transcriptionally distinct subsets, S5 and S6 (Fig. 1c).

Analysis of independent EnSC cultures confirmed that decidualization polarizes cells into two

subsets (Supplementary Figure S4). Known decidual stress defence genes (e.g. *CRYAB*,

HSD11B1, and *GLRX*) [35-37] were enriched in S5 (designated DC) whereas genes involved in

oxidative stress signalling and cellular senescence, including *KIAA1199*, *CLU* and *ABI3BP* [38-

41], prevailed in S6 (designated snDC) (Fig. 1d and Supplementary Figure S5). Notably, on day

6, the proportion of DC and snDC was 78% and 13%, respectively (Fig. 1e and

Supplementary Table S3). By day 8, snDC were almost as abundant as DC (42% and 45%,

respectively), suggesting rampant secondary senescence. Withdrawal of the decidualization

signal on day 8 elicited a further transcriptional change (S7), which was accounted for by

partial de-differentiation of >95% of DC (Supplementary Table S2 and S3). By contrast, snDC

were much more refractory to withdrawal, suggesting loss of progesterone/cAMP-

dependency.

To explore the 'switching' of DC into snDC further, we ordered decidualizing cells (D2-8) in

pseudotime and colour-coded them by transcriptional state. As shown in Figure 1f, this

analysis revealed that decidualizing EnSC progress along a continuous trajectory towards

senescence. This trajectory was interrupted by a single branchpoint, marking the divergence

of DC and a subset of DC already transitioning towards a senescent phenotype. The top 50

genes underpinning this branchpoint are depicted in a modified heatmap and clustered

hierarchically to visualize modules of genes with similar expression patterns (Fig. 1g). *DIO2*,

coding iodothyronine deiodinase 2, was identified as a major branch gene in the senescent

pathway. This enzyme catalyses the conversion of prohormone thyroxine (T4) into bioactive

triiodothyronine (T3) [42], suggesting increased energy metabolism in snDC. Multiple genes

coding for senescence-associated ECM remodelling factors were co-regulated with *DIO2*,

including *LUM* (lumican) [43], *CLU* (clusterin) [41], *ADAMTS5* (ADAM metallopeptidase with

thrombospondin type 1 motif 5) [44], *KIAA1199* (also known as cell migration inducing

hyaluronidase 1, CEMIP) [39], and *ABI3BP* (ABI family member 3 binding protein) [38]. Notably,

IGFBP1, a widely used decidual marker gene [8], was also part of this module. *FTL* and

SCARA5, along with known decidual genes such as *GLRX* and *IL1RL1*, were part of a

prominent branching module in the non-senescent decidual pathway (Fig. 1g). *FTL* encodes

ferritin light chain (L-ferritin) and *SCARA5* (scavenger receptor class A member 5) the L-

ferritin receptor, suggesting increased iron storage and detoxification capacity in DC.

Taken together, the single-cell analysis confirmed that decidualization is a multistep

process that starts with an acute auto-inflammatory response, which in turn synchronizes

transition of cells through intermediate transcriptional states before emerging mainly as DC

and some snDC. We also demonstrated that snDC rapidly perpetuate the senescent

phenotype across the culture, resulting in chronic senescence and increased expression of

ECM constituents and proteases and other SASP components.

Co-regulated decidual gene networks

We used k-means cluster analysis to identify networks of co-regulated genes across the

decidual pathway (Supplementary Table S4). Analysis of 1748 DEG yielded 7 networks of

uniquely co-regulated genes. Figure 2 depicts individual networks annotated for selected

transcription factor (TF) genes with core roles in decidualization. Network A1 genes are

rapidly downregulated within the first 48 hours of the decidual process after which expression

remains largely stable. The most notable TF to be 'reset' in this manner upon decidualization

is the progesterone receptor (PGR). This observation is in keeping with a previous study

purporting that overexpressed PGR blocks the formation of multimeric transcriptional

complexes upon decidualization by squelching key co-regulators [45]. Network A2 genes,

including *HOXA10*, are progressively down-regulated upon decidualization. Networks B1 and

B2 comprise of biphasic genes that peak prior to the emergence of DC and snDC. They

include several genes coding for pivotal decidual TFs, such as *STAT3*, *MEIS1*, *KLF9*, *WT1*

and *FOXO1* [8,19]. Genes in network C1 are gradually induced upon decidualization and then

peak in DC. Prominent TFs in this network are heart and neural crest derivatives expressed 2

(HAND2) and FOS like 2 (FOSL2), a potent PGR co-regulator [46]. By contrast, two distinct

networks underpinned the emergence of snDC. Network C2 genes rise gradually during the

initial decidual phase but expression is then rapidly accelerated, especially in snDC.

Interestingly, this network is enriched in gene ontology terms 'secreted' (Benjamini adjusted *P*

$= 3.7 \times 10^{-5}$, modified Fisher Exact test) and 'type I interferon signalling pathway' (Benjamini

adjusted *P* $= 1.4 \times 10^{-6}$, modified Fisher Exact test) (Supplementary Table S5), both canonical

features of cellular senescence [24]. A prominent TF in this senescence-associated network is

signal transducer and activator of transcription 1 (STAT1), which is not only activated by

interferon signalling but is also a potent inhibitor of PGR signalling in endometrial cells [47]. A

second biphasic network, designated network D, is also associated with snDC. Genes in this

network are initially repressed upon decidualization and then re-expressed predominantly in

snDC. Intriguingly, this network is enriched in genes known to be repressed by PGR, including

the TF genes *SOX4* and *FOXP1* [14]. Taken together, the data show that complex networks of

co-regulated genes underpin progression of cells along the decidual pathway and suggest that

altered expression levels of PGR co-regulators, such as FOSL2 and STAT1, may account for

progesterone-dependency of DC and progesterone-resistance of snDC.

Coordinated activation of immune surveillance genes

We mined the networks further for genes encoding factors implicated in immune surveillance

of senescent cells. CXCL14 and IL-15 drive uNK cell chemotaxis and proliferation and

activation, respectively [9,48]. TIMP-3 (TIMP metallopeptidase inhibitor 3) plays a critical role in

immune recognition of stressed or senescent cells by inhibiting proteolytic cleavage of surface

stress ligands needed for NK cell recognition [49]. Intriguingly, *CXCL14*, *IL15* and *TIMP3* belong

to the same biphasic gene network (B2), characterized by peak expression immediately prior

to the emergence of DC and snDC (Fig. 3a). Subsequently, expression of these genes drops

markedly in snDC but much less so in DC. To explore this concept of 'programmed' immune

surveillance further, we monitored the secreted levels of CXCL14, IL-15, and TIMP-3 every 48

hours over an 8-day time-course in 4 independent decidualizing cultures. As shown in Figure

3b, secreted levels of all 3 factors rise quickly during the initial decidual phase. While the

levels of CXCL14 and TIMP-3 then appear to plateau, IL-15 continues to accumulate in the

supernatant.

Clusterin (CLU) and the soluble IL-33 decoy receptor sST2 (encoded by *IL1RL1*) are

putative secreted markers of snDC and DC, respectively (Fig. 1g). As shown in Figure 3c,

secreted levels of both markers increase markedly upon decidualization, although the rise in

sST2 levels generally preceded secretion of CLU. We speculated that co-culture with uNK

cells would selectively attenuate the secretion of snDC markers. To test this hypothesis, uNK

cells were isolated from the supernatants of freshly cultured EnSC using magnetic-activated

cell sorting (MACS) and the purity and viability of cells confirmed by flow cytometry

(Supplementary Figure S6). The experimental design is depicted in Figure 3d.To monitor

uNK-cell killing of senescent cells, we measured senescence-associated β-galactosidase

(SAβG) activity, a widely used senescence marker [26], in undifferentiated and decidualizing

cultures with or without added primary uNK cells. As reported previously [9], the increase in

SAβG activity upon decidualization was entirely abrogated upon co-culturing with uNK cells

(Fig. 3e, right panel). Conversely, uNK cells had no impact on basal SAβG activity in

undifferentiated cultures. Co-culturing of uNK cells with decidualizing cells also reduced the

secreted levels of CLU, a stress-induced molecular chaperone molecule, [41] by ~ 50% (Fig. 3e,

middle panel). By contrast, uNK cells had little or no impact on secreted sST2 levels (Fig. 3e,

right panel), which is secreted by DC (Fig. 1g). To confirm that uNK cells target snDC, six

independent primary EnSC cultures were decidualized for 8 days in the presence or absence

of dasatinib, a potent senolytic drug [9]. As shown in Supplementary Figure S7, dasatinib also

markedly reduced SAβG activity and selectively inhibited CLU whereas the effect on sST2

secretion was minimal, thus recapitulating the actions of uNK cells. Taken together, the data

show that the expression and secretion of decidual factors involved in uNK cell recruitment

and activation is hardwired to coincide with the emergence of snDC. If recapitulated *in vivo*,

these observations further suggest an important role for uNK cells in limiting the detrimental

impact of snDC in early pregnancy

Single-cell analysis of peri-implantation endometrium

Decidualization involves activation of numerous genes that are either constitutively expressed

or induced in glandular epithelial cells following ovulation [8]. To identify stroma-specific marker

genes of a divergent peri-implantation decidual response, we first subjected freshly isolated

endometrial cells to scRNA-seq. Biopsies were timed relative to the pre-ovulatory luteinizing

hormone (LH) surge to coincide with the midluteal implantation window (LH+8; n=3) or the

start of the late-luteal phase (LH+10; n=3). Following quality control (Supplementary Figure

S8), t-SNE analysis assigned 2,847 cells to 5 clusters colour-coded to indicate the day of

biopsy (Fig. 4a, left panel). Additional dimensionality reduction analysis was performed on

immune cells (Fig. 4a, right panel). Clusters were designated based on canonical marker

genes as endothelial cells (EC; n=141), epithelial cells (EpC; n=395), immune cells (IC;

n=352), and EnSC (n=1,943) (Fig. 4c and Supplementary Table S6). In addition, a discrete

but as yet uncharacterized cluster of proliferating cells (PC; n=16) was identified (Fig. 4a).

EpC segregated in 4 clusters with the most abundant population (EpC1) expressing prototypic

receptivity genes (e.g. *GPX3*, *PAEP*, and *DPP4*) (Fig. 4b) [50]. EpC2 are ciliated epithelial cells

found interspersed throughout endometrial glands (Supplementary Figure S9). EpC3 were

derived predominantly, but not exclusively, from a single biopsy whereas EpC4 represented

an ambiguous population expressing both epithelial and stromal marker genes (Fig. 4b). uNK

cells, representing 89% of all IC, clustered into 3 subpopulations (NK1-3; Fig. 4a, right panel).

Notably, different NK populations have also recently been described in pregnant decidua [51].

Based on CIBERSORT analysis and cross-referencing of cluster-defining transcripts with

canonical markers curated from the literature, the remaining immune populations were

identified as naive B-cells (IC1), monocytes (IC2), and macrophage/dendritic cells (IC3) (Fig.

4c and Supplementary Table S7). The abundance of various cell types and subsets in each

biopsy is tabulated in Supplementary Table S3.

Marker genes of diverging decidual states

Next, we focused on the EnSC, which clustered prominently by day of biopsy in the *t*-SNE plot

(Fig. 4a). Progression from LH+8 to LH+10 was associated with altered expression of 518

genes in EnSC (Supplementary Table S8), 49% of which are also part of the 7 co-regulated

gene networks *in vitro* ($P<10^{-143}$, hypergeometric test). Although our single-cell analysis was

limited to six biopsies obtained on only two time-points, we surmised that the relationship

between genes that drive the divergence of DC and snDC *in vitro* would, at least partly, be

conserved in stromal cells *in vivo*. To test this hypothesis, we selected the top 50 genes of the

branchpoint in the decidual time-course and determined the Pearson correlations for each

gene pair in EnSC *in vitro* and *in vivo* on LH+8 and LH+10, respectively. We then calculated

the sum of coefficients, ranging from +2 to -2. Congruency was defined as the sum of

correlation coefficients of >1 or <-1 for positively and negatively co-regulated genes,

respectively (Supplementary Figure S10). Using this criterion, 27% of gene pairs were

congruent on both LH+8 and on LH+10, which is significantly more than expected by chance

alone ($P<10^{-30}$ and $P<10^{-58}$, respectively; hypergeometric test).

We reasoned that informative marker genes of a divergent decidual response *in vivo*

should be highly enriched in stromal cells, not regulated in glandular epithelium, and have a

temporal profile across the luteal phase commensurate with the expected switch of DC to

snDC prior to menstruation. Out of 50 branch genes, 5 genes (*TIMP3*, *IGF2*, *DIO2*, *SCARA5*

and *ABI3BP*) were highly enriched in EnSC compared to EpC, EC or IC. When cross-

referenced against two publicly available datasets [Gene Expression Omnibus (GEO) ID:

GSE84169 and GEO ID: GDS2052), only *SCARA5* and *DIO2* met all criteria. Notably,

SCARA5 belongs to the C1 network of genes whose expression peaks in DC whereas *DIO2*,

a gene repressed by the activated PGR in decidualizing EnSC [14], belongs to the D network of

senescence-associated genes (Fig. 5a). Figure 5b shows the relative expression of both

genes in different endometrial cell types as well their temporal expression across the cycle.

Next, we measured *SCARA5* and *DIO2* transcript levels by RT-qPCR in 250 samples

obtained across the implantation window (LH+6-11) to generate percentile graphs based on

the statistical distribution in gene expression for each day (Fig. 5c). To determine if *SCARA5*

and *DIO2* transcripts are co-expressed or mark different decidual cells, we performed

multiplexed single-molecule in situ hybridization (smISH) on endometrial biopsies obtained on

the same cycle day but deemed SCARA5^{HIGH} / DIO2^{LOW}, SCARA5^{LOW} / DIO2^{HIGH} or

SCARA5^{AVERAGE} / DIO2^{AVERAGE} based on the corresponding percentile graphs (Fig. 5d). Most

EnSC were *SCARA5*$^{+}$ but *DIO2*$^{-}$ in SCARA5^{HIGH} / DIO2^{LOW} biopsies whereas the opposite

pattern was observed in SCARA5^{LOW} / DIO2^{HIGH} samples. However, SCARA5^{AVERAGE} /

DIO2^{AVERAGE} samples consisted of mixture of *SCARA5*$^{+}$ or *DIO2*$^{+}$ cells as well as intermediate

cells expressing both transcripts. Finally, we used a computational approach to isolate EnSC

relatively enriched in *SCARA5* transcripts but reduced *DIO2* expression and *vice versa*. As

shown in Supplementary Figure S11, *SCARA5*^{enriched} / *DIO2*^{reduced} cells are characterized by the

expression multiple decidual TF and stress-resistance genes whereas a stress gene signature

was prominent *SCARA5*^{reduced} / *DIO2*^{enriched} cells (Supplementary Table S9).

RPL is associated with an aberrant decidual response

To examine if an aberrant decidual response in cycling endometrium is linked to increased

risk of pregnancy loss, we analysed LH-timed endometrial biopsies from 90 control subjects

and 89 RPL patients. Demographic and clinical characteristics are presented in Table 1. Each

biopsy was divided and processed for RT-qPCR analysis and immuno-histochemistry.

SCARA5 and *DIO2* transcript levels were measured and normalized for the day of the cycle

based on their respective percentile graphs (Fig. 5c). Immunohistochemistry was used to

quantify uNK cells using a standardized and validated approach that involves measuring the number of CD56[+] uNK cells per 100 stromal cells in proximity of the luminal epithelium [52]. The uNK cell data were then normalized for the day of the biopsy in the cycle as reported previously [9]. As shown in Figure 6a, lower *SCARA5* but higher *DIO2* percentiles in RPL samples indicated a diverging pro-senescent decidual response. Further, uNK cells were significantly less abundant in RPL compared to control subjects ($P<0.0001$, Welch two-sided *t*-test). We reasoned that *SCARA5*[+] DC and uNK cells drive successful transformation of the stroma into the decidua of pregnancy. Hence, we created 4 bins based on the sum of *SCARA5* and uNK percentiles and determined the number of RPL and control subjects in each bin. As shown in Figure 6b, 81% (25/31) of subjects assigned to the lowest bin were RPL patients whereas 79% (15/19) in the highest bin were control subjects ($P<0.0001$; Fisher's exact Test). Our analysis also enabled a definition of putative defects across the pathway, including 'decidualization failure' (*SCARA5*: ≤30[th] percentile, *DIO2*: ≤30[th] percentile), 'excessive decidual senescence' (*SCARA5*: ≤30th percentile, *DIO2*: ≥70[th] percentile), and 'uNK cell deficiency' (uNK cells: ≤30[th] percentile, *SCARA5* and *DIO2*: >30[th]-<70[th] percentile) (Fig. 6c). Excessive decidual senescence and uNK cell deficiency were significantly more common in RPL patients compared to control subjects ($P=0.0016$ and $P=0.02$, respectively; χ^2 test). Taken together, the incidence of an aberrant decidual response (i.e. encompassing all defects) was ~3-times higher in RPL compared to control subjects (44% versus 14%, respectively, $P=0.00013$; Fisher's Exact test).

Discussion

Decidualization of the endometrium occurs in all eutherian (placental) mammals where placentation involves trophoblast invasion of the endometrial stroma [53]. In menstruating species, including humans, decidualization is not under the control of an implanting embryo but initiated during the midluteal phase of each cycle [54]. Once triggered, the decidualizing endometrium becomes inextricably dependent on sustained progesterone signalling. In the absence of implantation of a competent embryo, falling progesterone levels elicit an

inflammatory decidual response which, upon recruitment and activation of leukocytes, leads to

tissue breakdown, focal bleeding and menstrual shedding [55,56]. Thus, a critical challenge at

implantation is to simultaneously avoid imminent endometrial breakdown while transforming

the cycling endometrium into a semi-permanent tissue, the decidua, maintained throughout

pregnancy. We recently argued that successful transformation of the endometrium in early

pregnancy requires both successful differentiation of EnSC into DC and prompt elimination of

snDC by activated uNK cells; and conversely, that defects in this process predisposes for

early pregnancy loss [9]. To test our hypothesis, we set out to identify putative marker genes of

an aberrant decidual response using single-cell RNA-seq.

We first generated a detailed transcriptional map of primary EnSC decidualized in culture

over 8 days followed by withdrawal of the differentiation signal for 2 days. This analysis

revealed a multi-step process that starts with a precipitous transcriptional response in

differentiating EnSC, which is followed by synchronous transition of cells through intermediate

states and then the emergence of DC and some snDC. A recent study asserted that DC

emerged in evolution from rewiring of an ancestral cellular stress response [19]. This conclusion

was based on the observation that endometrial fibroblasts isolated from marsupials, which

diverted from eutherians 60 to 80 million years ago [57], activate similar core regulatory genes

as human EnSC in response to a decidualizing stimulus and then mount an acute

inflammatory stress response akin to the initial decidual phase [19]. The key difference is that

most EnSC emerge from this process of inflammatory reprogramming as DC expressing

multiple anti-inflammatory and stress-resistance genes. However, some EnSC fail to activate

essential decidual effector genes and emerge as snDC, thus recapitulating the ancestral

response. We also demonstrated that snDC rapidly convert DC into secondary senescent

cells, a process abrogated upon co-culturing of uNK cells. Importantly, secondary senescence

had a dramatic impact on decidual gene expression, characterized by marked upregulation of

genes encoding for ECM proteins and proteases and other SASP components. The

mechanisms driving secondary senescence of DC in culture warrant further exploration,

including the possibility that EnSC become sensitized to juxtracrine senescence upon

differentiation into DC. A hallmark of snDC is progesterone-resistance, exemplified by

expression of PGR-repressed genes, such as *SOX4* and *FOXP1*, [14] and lack of

responsiveness to progesterone and cAMP withdrawal. Analysis of co-regulated gene

networks indicated that a switch in PGR co-regulators from FOSL2 to STAT1 may account for

progesterone-resistance in snDC, although this conjecture will need to be tested

experimentally. Apart from generating a temporal map of TF and effector genes underpinning

progression of cells along the decidual pathway, the network analysis also revealed that

multiple genes involved in immune surveillance of senescent cells, including *CXCL14*, *IL15*

and *TIMP3*, are hardwired to be activated prior to the emergence of snDC. Interestingly,

human chorionic gonadotrophin has been shown to stimulate uNK cell proliferation directly [58],

which suggests potential cooperation between the implanting embryo and DC in eliminating

snDC.

Our *in vitro* analysis suggested that the default trajectory of decidualizing EnSC is towards

cellular senescence, which can arguably only be avoided by timely clearance of snDC by uNK

cells. Failure to do so leads to chronic senescence, at least in culture. There are multiple

mechanisms by which chronic senescent cells promote tissue dysfunction, including

perturbation of stem cells, disruption of ECM, induction of tissue inflammation, and

propagation of senescence in neighbouring cells [25,27,32]. When extrapolated to the *in vivo*

situation, this means that failure of decidualizing EnSC to 'escape' the default pathway during

the peri-implantation window would inevitably lead to a pro-senescent decidual response and

the formation of an intrinsically unstable decidual-placental interface in pregnancy. To test this

hypothesis, we first set out to identify stroma-specific marker genes that discriminate between

a physiological and pathological decidual response in cycling endometrium. Single-cell

transcriptomic analysis of midluteal endometrium yielded a number of notable results,

including the discovery of a discrete but as yet uncharacterized population of proliferative

mesenchymal cells, the characterization of different epithelial cell subsets, and the

identification of 3 uNK cell states, which broadly corresponded to the different decidual NK cell

subsets recently identified by single-cell analysis in early pregnancy [51]. Different NK cell states

were in part defined by the relative abundance of cell cycle genes. For example, the NK1

population, representing the most proliferating uNK cells, abundantly express genes involved

in granule exocytosis (e.g. *PRF1*, *GNLY*, *GZMA* and *GZMB*). By contrast, NK3 cells express

low levels of cell cycle genes but are defined by *CCL5* and *CXCR4* expression. Notably,

CXCR4[+] uNK cells have previously been implicated in vascular remodelling in pregnancy [59].

   As the *in vivo* single-cell analysis was restricted to few samples across two time-points, we

restricted our search for putative marker genes to the 50 top *in vitro* branching genes that

marked divergence of DC and DC already transitioning towards senescence. Out of 50 genes,

SCARA5 and *DIO2* were deemed the best markers genes of DC and snDC, respectively,

based on their temporal regulation across the cycle in whole endometrial biopsies, lack of

regulation in glandular epithelium, and level of enrichment in stromal cells. Further,

multiplexed smISH showed that SCARA5[+] cells are distinct from DIO2[+] cells, which was

further corroborated by the different transcriptomic profiles of SCARA5[enriched] / mDIO2[reduced]

and SCARA5[reduced] / DIO2[enriched] EnSC in the *in vivo* single-cell data set. To test our hypothesis

that an aberrant decidual trajectory predisposes to pregnancy loss, we first generated

percentile graphs for *SCARA5* and *DIO2* expression using 250 endometrial midluteal biopsies

(LH+6-11). This approach enables comparison of the relative level of gene expression in

biopsies obtained at different days of the cycle. The relative abundance of uNK cells was also

determined using a previously established percentile graph based on analysis of 1,997

biopsies [9]. Next, we quantified the abundance of uNK cells and the expression of *SCARA5*

and *DIO2* in timed endometrial biopsies from RPL and control subjects. Sample selection was

based solely on reproductive history; all RPL patients had 3 or more consecutive miscarriages

whereas control subjects were women with male-factor, tubal or unexplained infertility

awaiting IVF treatment. Overall, our analysis indicated that pre-pregnancy endometrium in

RPL is characterized by uNK cell deficiency in parallel with a shift from a preponderance of

DC to snDC. Next, we examined the frequency of different putative defects along the decidual

pathway which may compromise the placental-decidual interface in early pregnancy. Based

on pre-specified but arbitrary criteria, we found that RPL is associated with 'excessive

decidual senescence' and 'uNK cell deficiency' but not 'decidualization failure'. While

promising, prospective studies are needed to validate our observations and to define

prognostic criteria that could be exploited for pre-pregnancy screening of women at risk of

RPL.

Decidualization is an iterative process in cycling endometrium. At present, it is not

clear if an aberrant decidual response persists from cycle-to-cycle or whether it represents an

intermittent defect that affects some but not all cycles. The relative high cumulative live-birth

rate in RPL favours the latter scenario [60]. Two distinct regulatory mechanisms control cellular

homeostasis in cycling endometrium. Aside from resident progenitor cells in the basal layer,

the endometrium actively recruits bone marrow-derived cells (BDMC) capable of

differentiating into stromal, epithelial and endothelial cells [61]. Experimental studies have

shown that BDMC enhance the regenerative capacity of non-pregnant endometrium [61], and

plausibly contribute to rapid decidual expansion in pregnancy. On the other hand, rapid

accumulation of uNK cells during the midluteal phase engenders selective elimination of snDC

through granule exocytosis, *de facto* rejuvenating the endometrium at the time of embryo

implantation [9]. Thus, by balancing recruitment of BMDC and uNK cells, the endometrium is

intrinsically equipped to fine-tune the decidual response to ensure implantation competence

from one cycle to the next, and perhaps to adapt following pregnancy failure. However,

pathological cues that disrupt these regulatory processes are predicted to increase the

frequency of a pro-senescent decidual response and, by extension, the likelihood of

miscarriage. For example, obesity is associated with loss of clonogenic progenitor cells in

cycling endometrium [33,62] as well as uNK cell deficiency [63,64]. Obesity is also strongly

associated with miscarriage, especially euploid pregnancy loss [65]. Thus, the prognostic value

of a screening test based on endometrial analysis may be enhanced by incorporating clinical

variables, such as maternal BMI, age and number of previous losses.

In summary, based on single-cell transcriptomic analysis, we propose that

decidualization is a multi-step process that ultimately leads to chronic senescence, a cellular

state incompatible with the formation of a functional decidual-placental interface. Our data

suggest that DC must engage uNK cells to eliminate snDC to escape this default pathway,

although additional *in vivo* studies are required to substantiate this conjecture. Nevertheless,

we identified *SCARA5* and *DIO2* as selective marker genes for DC and progesterone-resistant

snDC, respectively. When combined with uNK cells, these marker genes can be used to map

putative defects along the decidual pathway in cycling endometrium. Our findings raise the

possibility of a simple screening test to identify women at risk of miscarriage and to monitor

the effectiveness of pre-pregnancy interventions.

Materials and Methods

Ethical approval and sample collection. The study was approved by the NHS National Research Ethics-Hammersmith and Queen Charlotte's & Chelsea Research Ethics Committee (1997/5065). Endometrial biopsies were obtained from women attending the Implantation Clinic, a dedicated research clinic at University Hospitals Coventry and Warwickshire (UHCW) National Health Service Trust. Timed endometrial biopsies, relative to the preovulatory LH surge, were obtained using a Wallach Endocell™ endometrial sampler with written informed consent and in accordance with The Declaration of Helsinki (2000) guidelines.

Primary endometrial stromal cell (EnSC) culture. Endometrial biopsies were collected in DMEM-F12 media supplemented with 10 % dextran coated charcoal-stripped FBS (DCC) and processed for primary EnSC culture as described [9]. For decidualization studies, confluent monolayers of human endometrial stromal cells (EnSC) were incubated overnight at 37 °C with 5 % CO_2 in phenol red-free DMEM/F-12 containing 2 % DCC, containing antibiotic/antimycotic and L-glutamine (2 % media). To induce differentiation, cells were treated with 0.5 mM 8-bromo-cAMP (Sigma-Aldrich, Poole, UK) and 1 µM medroxyprogesterone acetate (MPA; Sigma-Aldrich) for the indicated time-points. For inhibition of senescence, cells were treated with 250 nM dasatinib (CST, Leiden, The Netherlands) throughout the differentiation time-course.

Droplet generation and single cell sequencing (Drop-Seq). Single-cell transcriptomes were captured in aqueous droplets containing barcoded beads using a microfluidic system (scRNAseq: Dolomite Bio, Royston, UK) according to the manufacturer's protocol and based on the Drop-Seq method described by Macosko and colleagues [34]. Briefly, cells in suspension were placed into the remote chamber of the scRNAseq system. Barcoded beads (Barcoded Bead SeqB; Chemgenes Corp., USA) in lysis buffer at a concentration of 280 beads/µl were loaded into the sample loop. Cell and bead solutions were run at a flow rate of 30 µl/min into a fluorophilic glass microfluidic chip with 100 µm etch depth (Single Cell RNA-seq Chip, Dolomite Bio) with droplet generation oil (Bio-Rad Laboratories, UK) at a flow rate of 200 µl/

min for 15-18 minutes. Droplets were collected into a 50 ml Falcon tube, quality checked using

a C-Chip Fuchs-Rosenthal Haemocytometer (Labtech, Heathfield, UK), and bead doublets

counted. Droplet breakage, bead isolation and reverse transcription were performed exactly

as described by Macosko and Goldman (Drop-Seq Laboratory Protocol version 3.1) on 8000

beads per reaction, with two reactions per sample, to give ~800 single-cell transcriptomes

attached to microparticles (STAMPS) per timepoint. PCR conditions were as per Macosko and

Goldman. Clean-up with Agencourt AMPure XP beads (Beckman Coulter, High Wycombe,

UK) was performed according to standard Illumina RNAseq protocols with a 0.6 × beads to

sample ratio. cDNA was eluted in 12 µl and quality and size assessed using an Agilent

Bioanalyzer High Sensitivity DNA chip (Agilent, Stockport, UK). For tagmentation 600 pg

cDNA, as determined by Qubit High Sensitivity DNA assay (ThermoFisher, Paisley, UK), was

processed according to Macosko and Goldman using Illumina Nextera XT DNA Sample Kit

and Indexing Kit (Illumina, Cambridge, UK). Tagmented libraries were cleaned up using

AMPure XP beads as before, with a 0.6 × beads ratio followed by a repeat clean-up using 1×

beads. Eluted libraries were analysed using Agilent Bioanalyzer High Sensitivity DNA chip to

assess quality and determine library size, and concentration was determined by Qubit High

Sensitivity DNA assay. Library dilution and denaturation was performed as per standard

Illumina protocols and sequenced using NextSeq High Output 75 cycle V2 kit.

Drop-Seq data alignment and quantification. Initial Drop-Seq data processing was

performed using Drop-Seq_tools-1.0.1 following the protocol described by Nemesh

(seqAlignmentCookbook_v1.2Jan2016.pdf, http://mccarrolllab.com/dropseq). Briefly, reads

with low-quality bases in either cell or molecular barcode were filtered and trimmed for

contaminating primer or poly-A sequence. Sequencing errors in barcodes were inferred and

corrected, as implemented by Drop-Seq_tools-1.0.1. Reads were aligned to the hg19

(Human) reference genome concatenated with ERCC annotations using STAR-2.5.3a [66], with

the Gencode21 (Human) as reference transcriptome. Uniquely mapped reads, with ≤ 1

insertion or deletion, were used in quantification. Finally, the DigitalExpression tool [34] was

used to obtain the digital gene expression matrix for each sample. Cell numbers were

selected computationally from the inflection point in a cumulative distribution of reads plotted

against the cell barcodes ordered by descending number of reads [67]. Cell barcodes beyond

the inflection point are believed to represent 'ambient RNA' (e.g. contaminating RNA from

damaged cells), not cellular transcriptomes, and therefore excluded from further analysis. This

resulted in ~800 cells per time-point, matching the number anticipated from processed bead

counts.

Cell aggregation analysis. Analysis of DGE data was performed with Seurat v2 [68]. To select

high-quality data for analysis, cells were included when at least 200 genes were detected,

while genes were included if they were detected in at least 3 cells. Cells which had more than

4500 genes were excluded from the analysis as were cells with more than 5% mitochondrial

gene transcripts to minimize doublets and low-quality (broken or damaged) cells, respectively.

After scaling and normalization of the raw counts in the DGE matrix, cell-cycle regression was

applied. For cell aggregation, a set of highly variable genes was first identified, with an

average expression mean between 0.0125 and 3 and a Log Variant to Mean Ratio of at least

0.5, which were used to perform principal component (PC) analysis. Based on statistical

significance and the robustness of the results, the first 10 PCs were subsequently used as

inputs for clustering via shared nearest neighbour (SNN) and subsequent t-distributed

stochastic neighbour embedding (t-SNE) representation. The Seurat function 'FindAllMarkers'

employing the Wilcoxon test was used to identify marker genes for each cell state cluster in

the t-SNE representation. To obtain independent estimation of the number of unique cell-types

we used SC3 v1.3.18 (5), applying a consensus strategy and Tracy-Widom theory on random

matrices to estimate the optimal number of clusters (k).

Co-regulated gene networks. K-means cluster analysis was performed in MeV v4.8 (6) to

group marker genes based on co-expression across cell state clusters. Figure of Merit (FoM)

was run first to determine the number of expression patterns (k). The predictive power of the

k-means algorithm was estimated using a Fig. of merit (FOM) values for k from 1 to 20. K-

means clustering was run using Pearson correlation metric for a maximum of 50 iterations.

Gene ontology analysis was performed on the clustered genes using DAVID [69].

Trajectory analysis. Trajectory analysis was performed with Monocle v2.6.1 [70]. Branch-point

genes were identified with Branched Expression Analysis Modeling (BEAM) function.

Gene expression correlation. To test gene expression correlation between pairs of genes,

expression was imputed for every cell using Markov Affinity-based Graph Imputation of Cells

(MAGIC) [71]. Pearson correlation coefficients were calculated for top 50 genes determining the

different trajectories at informative branch-points. To assess congruency between the time-

course and biopsy datasets, the correlation coefficients were added resulting in sum of

coefficients between -2 and +2. To infer gene expression associated with a pro-senescent

decidual response, $SCARA5^{enriched}/DIO2^{reduced}$ and $SCARA5^{reduced}/DIO2^{enriched}$ EnSCs were

selected after MAGIC imputation using the midpoint of the expression levels as thresholds.

Differential gene expression on the selected cells was determined using the Seurat function

'FindMarkers' employing the Wilcoxon test, with GO analysis performed using DAVID [69].

Drop-Seq analysis of timed endometrial biopsies. Six LH-timed endometrial biopsies were

processed as described in detail elsewhere [9]. Single-cell fractions were then subjected to

Drop-Seq analysis. The time from biopsy in the clinic to cDNA synthesis was less than 4 hours

for all samples. Anonymized endometrial biopsies were obtained from women aged between

31 and 42 years with regular cycles, body mass index between 23 and 32 kg/m2, and

absence of uterine pathology on transvaginal ultrasound examination.

Reverse transcription quantitative PCR (RT-qPCR). RNA was extracted from endometrial

biopsies which had been snap frozen in clinic (<1 min after collection), using STAT-60 (AMS

Biotechnology, Oxford, UK) according to the manufacturer's instructions. Reverse

transcription was performed from 1 μg RNA using the Quantitect Reverse Transcription Kit

(QIAGEN, Manchester, UK) and cDNA was diluted to 10 ng/μl equivalent before use in qPCR.

Amplification was performed on a 7500 Real-Time PCR system (Applied Biosystems, Paisley,

UK) in 20 μl reactions using 2 × PrecisionPlus Mastermix with SYBR Green and low ROX

(PrimerDesign, Southampton, UK), with 300 nM each of forward and reverse primers. *L19*

was used as a reference control. Primer sequences were as follows: *SCARA5* forward: 5'-

CAT GCG TGG GTT CAA AGG TG-3', *SCARA5* reverse: 5'-CCA TTC ACC AGG CGG ATC

AT-3'; *DIO2* forward: 5'-ACT CGG TCA TTC TGC TCA A-3', *DIO2* reverse: 5'-TTC CAG ACG

CAG CGC AGT-3', *L19* forward: 5'-GCG GAA GGG TAC AGC CAA T-3', L19 reverse: 5'-GCA

GCC GGC GCA AA-3'. Centile calculations were performed on dCt values using R software.

Multiplexed single-molecule in situ hybridization. Formalin-fixed paraffin-embedded

(FFPE) samples were cut to 5 μm sections. RNA in situ hybridization was carried with

RNAscope® 2.5 HD Duplex Reagent Kit (ACD, California, USA) with probes for SCARA5

(574781-C1) and DIO2 (562211-C2) according to manufacturer's guidelines. Following

hybridization and amplification, slides were counterstained with 50 % haematoxylin. Images

were obtained using an EVOS AUTO microscope (ThermoFisher Scientific) with a 40×

objective lens.

Isolation and culture of uNK cells. Primary uNK cells were isolated from luteal phase

endometrial biopsies as described previously[9]. Briefly, supernatant from freshly digested

EnSC cultures was collected uNK cells were isolated by magnetic activated cell separation

(MACS; Miltenyi Biotec, Bergisch Gladbach, Germany) using phycoerythrin (PE)-conjugated

anti-CD56 antibody (Bio-Legend, San Diego, CA, USA), as per manufacturer's instructions.

The CD56[+] positive fraction was collected by centrifugation and cultured in suspension for up

to 5 days in RPMI media (Sigma-Aldrich) supplemented with 10% DCC-FBS, 1× Antibiotic-

Antimycotic, and 2 ng/ml IL-15 (Sigma-Aldrich) to aid uNK cell maturation. To increase yield,

uNK cells from 3-5 subjects were pooled. For co-culture experiments, uNK cells were pelleted

and re-suspended in DMEM/F-12 containing 2 % DCC without IL-15. A total of 5,000 uNK

cells were added to 50,000 EnSC, decidualized or not, per well of a 96-well plate. MACS

isolated uNK cells were analysed by flow cytometry after labelling with BD Horizon™ Fixable

Viability Stain (FVS) 660 (BD Biosciences) in 1×PBS (1:1000) and PE-conjugated CD56 (BD

Biosciences; clone: B159; catalogue no: 555516; 1:5) according to manufacturer's

instructions. Red blood cells (RBC) were removed from the negative fraction using 1×RBC

lysis solution (BD Biosciences, catalogue number: 349202) for 10 minutes at room

temperature. Samples were analysed by BD FACSMelody™ Cell Sorter (BD Biosciences) and

data analysis was performed using FlowJo v10.

Quantitative analysis of uNK cells in endometrial biopsies. FFPE tissue sections were

stained for CD56 (a uNK cell-specific cell surface antigen) using a 1:200 dilution of

concentrated CD56 antibody (NCL-L-CD56-504, Novocastra, Leica BioSystems). Stained

slides were de-hydrated, cleared and cover-slipped in a Tissue-Tek® Prisma® Automated Slide

Stainer, model 6134 (Sakura Flinetek Inc. CA, USA) using DPX coverslip mountant. Bright-

field images were obtained on a Mirax Midi slide scanner using a 20 × objective lens and

opened in Panoramic Viewer v1.15.4 (3DHISTECH Ltd, Budapest, Hungary) for analysis. To

avoid inconsistencies that reflect reduced uNK cell densities at greater endometrial depths,

$CD56^+$ cells were quantified in compartments directly underlying the luminal epithelium. Here,

three randomly selected areas of interest for each biopsy were identified and captured within

Panoramic Viewer before analysis in ImageJ image analysis software. Both luminal and

glandular epithelial cells were removed manually before colour deconvolution into constituent

brown (CD56+ staining) and blue (hematoxylin staining – stromal cells). The area of positive

staining above a manually determined background threshold was used to quantify staining

intensity. The uNK cell percentage was calculated as the number of $CD56^+$ cells per 100

stromal cells and averaged from 3 images [9].

Enzyme-linked immunosorbent assay (ELISA). Culture supernatants were collected every

2 days and centrifuged to clear cell debris prior to storage at -20°C. Analytes in collected

supernatant were measured by ELISA as per manufacturer's instructions (DuoSet ELISA kits,

Bio-Techne, Abingdon, UK). Standard curves were fitted to a 4-parameter logistic fit curve in

GraphPad Prism software and sample concentrations interpolated from these graphs.

Quantitation of SAβG activity. SAβG activity in cultured cells was quantified using the 96-

Well Cellular Senescence Activity Assay kit (CBA-231, Cell Biolabs Inc; CA, USA) as

described previously [9].

Statistical analysis. Statistical analysis of ELISA and SaβG data was performed using

GraphPad Prism 8.0.1 and data are presented as mean ± SEM. Chi-square and Fisher's

exact test on the decidual pathway defects were performed using online calculators at

https://www.socscistatistics.com/. All other analyses were implemented in R.

Data availability. The Drop-Seq data were deposited in the GEO repository (accession

number: GSE127918).

Author Contributions

Conceptualization, J.J.B.; Methodology, P.V., E.S.L, J.L. and S.O; Investigation, E.S.L., P.V.,

J.M., M.D.C., P.J.B., C-S.K., K.F.; Writing – Original Draft, J.J.B., P.V. and E.S.L.; Funding

Acquisition, S.Q., S.O. and J.J.B.; Resources, J.O., L.J.E., S.Q. and J.J.B.; Supervision, S.O.

and J.J.B.

Acknowledgement

We are grateful to all the women who participated in this research. This work was supported

by funds from the Tommy's National Miscarriage Research Centre and Wellcome Trust

Investigator Award to J.J.B and S.O. (212233/Z/18/Z).

References

1   Rai, R. & Regan, L. Recurrent miscarriage. *Lancet* **368**, 601-611, (2006).

2   Hardy, K., Hardy, P. J., Jacobs, P. A., Lewallen, K. & Hassold, T. J. Temporal changes in
chromosome abnormalities in human spontaneous abortions: Results of 40 years of
analysis. *Am J Med Genet A* **170**, 2671-2680, (2016).

3   Carp, H. *et al.* Karyotype of the abortus in recurrent miscarriage. *Fertil Steril* **75**, 678-
682, (2001).

4   Stephenson, M. D., Awartani, K. A. & Robinson, W. P. Cytogenetic analysis of
miscarriages from couples with recurrent miscarriage: a case-control study. *Hum
Reprod* **17**, 446-451, (2002).

5   ESHRE. Recurrent Pregnancy Loss: A Guideline of the European Society of Human
Reproduction and Embryology. (2017).

6   Practice Committee of the American Society for Reproductive, M. Evaluation and
treatment of recurrent pregnancy loss. *Fertility and Sterility* **5**, 1103-1111, (2012).

7   Ogasawara, M., Aoki, K., Okada, S. & Suzumori, K. Embryonic karyotype of abortuses in
relation to the number of previous miscarriages. *Fertil Steril* **73**, 300-304, (2000).

8   Gellersen, B. & Brosens, J. J. Cyclic decidualization of the human endometrium in
reproductive health and failure. *Endocr Rev* **35**, 851-905, (2014).

9   Brighton, P. J. *et al.* Clearance of senescent decidual cells by uterine natural killer cells
in cycling human endometrium. *eLife* **6**, (2017).

10   Weimar, C. H. *et al.* Endometrial stromal cells of women with recurrent miscarriage
fail to discriminate between high- and low-quality human embryos. *PLoS One* **7**,
e41424, (2012).

11   Brosens, J. J. *et al.* Uterine selection of human embryos at implantation. *Sci Rep* **4**,
3894, (2014).

12   Brosens, J. J., Pijnenborg, R. & Brosens, I. A. The myometrial junctional zone spiral
arteries in normal and abnormal pregnancies: a review of the literature. *Am J Obstet
Gynecol* **187**, 1416-1423, (2002).

13   Vrljicak, P. *et al.* Analysis of chromatin accessibility in decidualizing human
endometrial stromal cells. *FASEB J* **32**, 2467-2477, (2018).

14   Cloke, B. *et al.* The androgen and progesterone receptors regulate distinct gene
networks and cellular functions in decidualizing endometrium. *Endocrinology* **149**,
4462-4474, (2008).

15   Leitao, B. *et al.* Silencing of the JNK pathway maintains progesterone receptor activity
in decidualizing human endometrial stromal cells exposed to oxidative stress signals.
FASEB J **24**, 1541-1551, (2010).

16   Muter, J. *et al.* Progesterone-Dependent Induction of Phospholipase C-Related
Catalytically Inactive Protein 1 (PRIP-1) in Decidualizing Human Endometrial Stromal
Cells. *Endocrinology* **157**, 2883-2893, (2016).

17   Lynch, V. J., Leclerc, R. D., May, G. & Wagner, G. P. Transposon-mediated rewiring of
gene regulatory networks contributed to the evolution of pregnancy in mammals. *Nat
Genet* **43**, 1154-1159, (2011).

18   Lynch, V. J. *et al.* Ancient transposable elements transformed the uterine regulatory
landscape and transcriptome during the evolution of mammalian pregnancy. *Cell Rep*
10, 551-561, (2015).

19   Erkenbrack, E. M. *et al.* The mammalian decidual cell evolved from a cellular stress
response. *PLoS Biol* **16**, e2005594, (2018).

20 Al-Sabbagh, M. *et al.* NADPH oxidase-derived reactive oxygen species mediate
decidualization of human endometrial stromal cells in response to cyclic AMP
signaling. *Endocrinology* **152**, 730-740, (2011).
21 Salker, M. S. *et al.* Disordered IL-33/ST2 activation in decidualizing stromal cells
prolongs uterine receptivity in women with recurrent pregnancy loss. *PLoS One* **7**,
e52252, (2012).
22 Kajihara, T. *et al.* Differential expression of FOXO1 and FOXO3a confers resistance to
oxidative cell death upon endometrial decidualization. *Mol Endocrinol* **20**, 2444-2455,
(2006).
23 Kuroda, K. *et al.* Induction of 11beta-HSD 1 and activation of distinct mineralocorticoid
receptor- and glucocorticoid receptor-dependent gene networks in decidualizing
human endometrial stromal cells. *Mol Endocrinol* **27**, 192-202, (2013).
24 Antonangeli, F., Zingoni, A., Soriani, A. & Santoni, A. Senescent cells: Living or dying is
a matter of NK cells. *J Leukoc Biol* **105**, 1275-1283, (2019).
25 Childs, B. G. *et al.* Senescent cells: an emerging target for diseases of ageing. *Nat Rev*
Drug Discov, (2017).
26 Hernandez-Segura, A., Nehme, J. & Demaria, M. Hallmarks of Cellular Senescence.
Trends Cell Biol **28**, 436-453, (2018).
27 van Deursen, J. M. The role of senescent cells in ageing. *Nature* **509**, 439-446, (2014).
28 Jun, J. I. & Lau, L. F. The matricellular protein CCN1 induces fibroblast senescence and
restricts fibrosis in cutaneous wound healing. *Nat Cell Biol* **12**, 676-685, (2010).
29 Storer, M. *et al.* Senescence is a developmental mechanism that contributes to
embryonic growth and patterning. *Cell* **155**, 1119-1130, (2013).
30 Ritschka, B. *et al.* The senescence-associated secretory phenotype induces cellular
plasticity and tissue regeneration. *Genes Dev* **31**, 172-183, (2017).
31 Acosta, J. C. *et al.* Chemokine signaling via the CXCR2 receptor reinforces senescence.
Cell **133**, 1006-1018, (2008).
32 Ito, Y., Hoare, M. & Narita, M. Spatial and Temporal Control of Senescence. *Trends Cell*
Biol **27**, 820-832, (2017).
33 Lucas, E. S. *et al.* Loss of Endometrial Plasticity in Recurrent Pregnancy Loss. *Stem Cells*
34, 346-356, (2016).
34 Macosko, E. Z. *et al.* Highly Parallel Genome-wide Expression Profiling of Individual
Cells Using Nanoliter Droplets. *Cell* **161**, 1202-1214, (2015).
35 Kuroda, K. *et al.* Elevated periimplantation uterine natural killer cell density in human
endometrium is associated with impaired corticosteroid signaling in decidualizing
stromal cells. *The Journal of clinical endocrinology and metabolism* **98**, 4429-4437,
(2013).
36 Song, J. J. *et al.* Role of glutaredoxin in metabolic oxidative stress. Glutaredoxin as a
sensor of oxidative stress mediated by H2O2. *J Biol Chem* **277**, 46566-46575, (2002).
37 Zuo, R. J. *et al.* Crystallin alphaB acts as a molecular guard in mouse decidualization:
regulation and function during early pregnancy. *FEBS Lett* **588**, 2944-2951, (2014).
38 Latini, F. R. *et al.* ABI3 ectopic expression reduces in vitro and in vivo cell growth
properties while inducing senescence. *BMC Cancer* **11**, 11, (2011).
39 Michishita, E., Garces, G., Barrett, J. C. & Horikawa, I. Upregulation of the KIAA1199
gene is associated with cellular mortality. *Cancer Lett* **239**, 71-77, (2006).
40 Petropoulou, C., Trougakos, I. P., Kolettas, E., Toussaint, O. & Gonos, E. S.
Clusterin/apolipoprotein J is a novel biomarker of cellular senescence that does not

738		affect the proliferative capacity of human diploid fibroblasts. *FEBS Lett* **509**, 287-297,
739		(2001).
740	41	Trougakos, I. P. The molecular chaperone apolipoprotein J/clusterin as a sensor of
741		oxidative stress: implications in therapeutic approaches - a mini-review. *Gerontology*
742		**59**, 514-523, (2013).
743	42	Bianco, A. C. & Kim, B. W. Deiodinases: implications of the local control of thyroid
744		hormone action. *J Clin Invest* **116**, 2571-2579, (2006).
745	43	Severino, V. *et al.* Insulin-like growth factor binding proteins 4 and 7 released by
746		senescent cells promote premature senescence in mesenchymal stem cells. *Cell Death*
747		*Dis* **4**, e911, (2013).
748	44	Le Maitre, C. L., Freemont, A. J. & Hoyland, J. A. Accelerated cellular senescence in
749		degenerate intervertebral discs: a possible role in the pathogenesis of intervertebral
750		disc degeneration. *Arthritis Res Ther* **9**, R45, (2007).
751	45	Brosens, J. J., Hayashi, N. & White, J. O. Progesterone receptor regulates decidual
752		prolactin expression in differentiating human endometrial stromal cells. *Endocrinology*
753		**140**, 4809-4820, (1999).
754	46	Mazur, E. C. *et al.* Progesterone receptor transcriptome and cistrome in decidualized
755		human endometrial stromal cells. *Endocrinology* **156**, 2239-2253, (2015).
756	47	Christian, M. *et al.* Interferon-gamma modulates prolactin and tissue factor expression
757		in differentiating human endometrial stromal cells. *Endocrinology* **142**, 3142-3151,
758		(2001).
759	48	Mokhtar, N. M. *et al.* Progestin regulates chemokine (C-X-C motif) ligand 14 transcript
760		level in human endometrium. *Mol Hum Reprod* **16**, 170-177, (2010).
761	49	Raneros, A. B. *et al.* Increasing TIMP3 expression by hypomethylating agents
762		diminishes soluble MICA, MICB and ULBP2 shedding in acute myeloid leukemia,
763		facilitating NK cell-mediated immune recognition. *Oncotarget* **8**, 31959-31976, (2017).
764	50	Altmae, S. *et al.* Meta-signature of human endometrial receptivity: a meta-analysis
765		and validation study of transcriptomic biomarkers. *Sci Rep* **7**, 10077, (2017).
766	51	Vento-Tormo, R. *et al.* Single-cell reconstruction of the early maternal-fetal interface
767		in humans. *Nature* **563**, 347-353, (2018).
768	52	Lash, G. E. *et al.* Standardisation of uterine natural killer (uNK) cell measurements in
769		the endometrium of women with recurrent reproductive failure. *J Reprod Immunol*
770		**116**, 50-59, (2016).
771	53	Ramsey, E. M., Houston, M. L. & Harris, J. W. Interactions of the trophoblast and
772		maternal tissues in three closely related primate species. *Am J Obstet Gynecol* **124**,
773		647-652, (1976).
774	54	Emera, D., Romero, R. & Wagner, G. The evolution of menstruation: a new model for
775		genetic assimilation: explaining molecular origins of maternal responses to fetal
776		invasiveness. *Bioessays* **34**, 26-35, (2012).
777	55	Evans, J. & Salamonsen, L. A. Inflammation, leukocytes and menstruation. *Rev Endocr*
778		*Metab Disord* **13**, 277-288, (2012).
779	56	Evans, J. & Salamonsen, L. A. Decidualized human endometrial stromal cells are
780		sensors of hormone withdrawal in the menstrual inflammatory cascade. *Biol Reprod*
781		**90**, 14, (2014).
782	57	O'Leary, M. A. *et al.* The placental mammal ancestor and the post-K-Pg radiation of
783		placentals. *Science* **339**, 662-667, (2013).

58   Kane, N., Kelly, R., Saunders, P. T. & Critchley, H. O. Proliferation of uterine natural
killer cells is induced by human chorionic gonadotropin and mediated via the mannose
receptor. *Endocrinology* **150**, 2882-2888, (2009).
59   Gibson, D. A., Greaves, E., Critchley, H. O. & Saunders, P. T. Estrogen-dependent
regulation of human uterine natural killer cells promotes vascular remodelling via
secretion of CCL2. *Hum Reprod* **30**, 1290-1301, (2015).
60   Ewington, L. J., Tewary, S. & Brosens, J. J. New insights into the mechanisms
underlying recurrent pregnancy loss. *J Obstet Gynaecol Res* **45**, 258-265, (2019).
61   Santamaria, X., Mas, A., Cervello, I., Taylor, H. & Simon, C. Uterine stem cells: from
basic research to advanced cell therapies. *Hum Reprod Update* **24**, 673-693, (2018).
62   Murakami, K. *et al.* Deficiency in clonogenic endometrial mesenchymal stem cells in
obese women with reproductive failure--a pilot study. *PloS one* **8**, e82582, (2013).
63   Castellana, B. *et al.* Maternal obesity alters uterine NK activity through a functional
KIR2DL1/S1 imbalance. *Immunol Cell Biol* **96**, 805-819, (2018).
64   Perdu, S. *et al.* Maternal obesity drives functional alterations in uterine NK cells. *JCI
Insight* **1**, e85560, (2016).
65   Boots, C. E., Bernardi, L. A. & Stephenson, M. D. Frequency of euploid miscarriage is
increased in obese women with recurrent early pregnancy loss. *Fertil Steril* **102**, 455-
459, (2014).
66   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21,
(2013).
67   Lun, A. T. L. *et al.* EmptyDrops: distinguishing cells from empty droplets in droplet-
based single-cell RNA sequencing data. *Genome Biol* **20**, 63, (2019).
68   Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial reconstruction of
single-cell gene expression data. *Nat Biotechnol* **33**, 495-502, (2015).
69   Huang da, W., Sherman, B. T. & Lempicki, R. A. Systematic and integrative analysis of
large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57, (2009).
70   Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by
pseudotemporal ordering of single cells. *Nat Biotechnol* **32**, 381-386, (2014).
71   van Dijk, D. *et al.* Recovering Gene Interactions from Single-Cell Data Using Data
Diffusion. *Cell* **174**, 716-729 e727, (2018).
815
816

     Figure legends

Fig. 1. Single-cell reconstruction of the decidual pathway in culture. **a** Schematic

representation of *in vitro* time-course experiment. **b** *t*-SNE projection of 4,580 EnSC, colour-

coded according to days of decidualization (D0-8) and upon withdrawal (WD) of the

differentiation signal for 48 hours. D0 represents undifferentiated EnSC. **c** The same *t*-SNE

plot now color-coded according to transcriptional state (S1-7). **d** Violin plots showing log-

transformed, normalized expression levels for indicated genes in decidual cells (DC, state S5)

and senescent decidual cells (snDC, state S6). **e** Relative proportion of total cells assigned to

states S5 (DC), S6 (snDC) and S7 at each experimental timepoint. **f** Decidualizing EnSC were

placed in pseudotime to reconstruct the trajectory of differentiation, revealing a continuous

trajectory towards senescence with a single branchpoint marking the divergence of DC. **g**

Heat map showing gene dynamics during cell state transition at the branchpoint shown in

panel f. Columns are points in pseudotime while rows represent the 50 most dynamic genes

at the branch point. The beginning of pseudotime is in the middle of the heat map and the

trajectory towards DC and snDC are indicated by the arrows. Hierarchical clustering visualizes

modules of genes with similar lineage-dependent expression patterns.

Fig. 2. K-means cluster analysis identifies co-regulated decidual gene networks

Analysis of 1,748 DEG across the decidual pathway (D0-D8) yielded seven networks of

uniquely co-regulated genes. Networks are annotated for selected transcription factor genes

with core roles in decidualization.

Fig. 3. Co-ordinated expression of decidual immune surveillance genes. a Decidual gene

network B2 annotated to highlight genes implicated in uNK cell activation and immune

surveillance of senescence cells. **b** Primary EnSC cultures were decidualized with 8-bromo-

cAMP and MPA (C+M) for the indicated days. ELISAs were performed on spent medium

collected at 48 hours intervals to examine secreted levels of CXCL14, IL-15 and TIMP-3.

Grey dotted lines indicate show secreted levels in individual cultures (n=4). Black solid lines

indicate the median level of secretion. **c** Four independent primary EnSC cultures were

decidualized with 8-bromo-cAMP and MPA (C+M) for the indicated days. ELISAs were

performed on spent medium collected at 48 h intervals to examine secreted levels of clusterin

(CLU) and sST2. Cultures established from the same biopsy are colour matched between

plots (dotted lines). Black solid lines indicate the median level of secretion. **d** Schematic

representation of uNK cell co-culture experiments. A total of 5,000 primary uNK cells were co-

cultured for 48 hours from the indicated timepoint with 50,000 decidualized cells seeded in 96-

well plates. Four independent EnSC cultures were used. **e** SAβG activity in undifferentiated

and decidualized (C+M) cells co-cultured with or without uNK cells. Four independent EnSC

cultures were used. **f** Secretion of CLU and sST2 in decidualized (C+M) cells co-cultured with

uNK cells relative to decidualized cells cultured without uNK cells. Four independent EnSC

cultures were used, lines connect paired samples.

Fig. 4. Identification of endometrial cell types and subsets during the implantation

window. **a** Left panel: t-SNE plot of 2,847 cells isolated from 6 LH-timed biopsies, coloured to

indicate reported cycle day (LH+8 or LH+10), captures all major endometrial cell types,

including epithelial cells (EpC), immune cells (IC), endothelial cells (EC), stromal cells (EnSC)

and a discrete but transcriptionally distinct proliferative (PC) stromal subpopulation. EpC

segregated in 4 subpopulations (EpC1-4). * indicates EpC3 contributed predominantly by a

single sample. Right panel: additional dimensionality reduction separated immune cells into 3

uNK cell subsets (NK1-3), naive B-cells (IC1), monocytes (IC2), and macrophage/dendritic

cells (IC3) **b** Heatmap showing relative expression (z-score) of markers defining cell-types

and EpC subpopulations. *MYO6, RASD1* and *ALCAM* are included as pan-epithelial genes. **c**

Heatmap showing relative expression of markers defining endometrial IC populations during

the implantation window, including three uNK cell subsets.

Fig. 5. *SCARA5* and *DIO2* as marker genes for divergent decidual states. a *SCARA5* and

DIO2 belong to two distinct decidual gene networks with peak expression in DC and snDC,

respectively. **b** Spatial and temporal expression of *SCARA5* and *DIO2* in cycling human

endometrium. Left panels, violin plots showing expression of *SCARA5* and *DIO2 in vivo* in

EnSC, EC (endothelial cells), EpC (epithelial cells), and IC (immune cells) (Wilcoxon rank sum

test with Bonferroni correction). Right panels, expression of *SCARA5* and *DIO2* in proliferative

and early-, mid-, late-luteal phase endometrium. Each bar represents an individual biopsy.

The data were retrieved from microarray data deposited in the Gene Expression Omnibus

(GEO Profiles, GDS2052). **c** *SCARA5* and *DIO2* transcript levels quantified by RT-qPCR

analysis in 250 endometrial biopsies obtained between LH+6 to LH+11. Centile calculations

were performed on dCt values using R software and graphs were generated based on the

distribution of expression on each day. The median number of samples for each day was 43

(range: 30 to 46). **d** Multiplexed single molecule *in situ* hybridization (smISH) on 3 endometrial

biopsies (LH+10) deemed *SCARA5*^{HIGH}/*DIO2*^{LOW} (95th and 20th percentile, respectively),

SCARA5^{AVERAGE}/*DIO2*^{AVERAGE} (60th and 52th percentile, respectively), and *SCARA5*^{LOW}

/*DIO2*^{HIGH} (4th and 91st percentile, respectively). Insert in the left panel shows hybridization with

a negative control probe. Insert in middle panel shows *SCARA5*$^{+}$ cells (blue, closed arrows)

and *DIO2*$^{+}$ cells (pink, open arrows) in close proximity. Scale bar: 100 μM. Original

magnification: ×40.

Fig. 6. Impaired fate divergence of decidual cells in RPL. a Distribution of uNK cells and

SCARA5 and *DIO2* percentiles in timed endometrial biopsies of control subjects (n=90) and

RPL patients (n=89) (Welch two-sided *t*-test). **b** Number of RPL and control subjects across 4

bins defined by the sum of SCARA5 and uNK cell percentiles. **c** Diagram illustrating the

decidual pathway. Fate divergence of EnSC upon decidualization relates to the level of

replicative stress (indicating by nuclear shading) incurred by individual cells during the

proliferative phase. Stress-resistant decidual cells recruit and activate uNK cells to eliminate

stressed/senescent decidual cells through granule exocytosis. Different defects along the

decidual pathway can be identified, including 'decidualization failure', 'excessive decidual

senescence', and 'uNK cell deficiency'. The frequency of each defect in RPL and control

subjects is shown (χ^2 test).

Table 1: Subject Demographics

	Control* (n=90)	RPL** (n=89)	*P*-value†
Age (years) (Median ± IQR)	36(33-37)	36 (33-38)	0.253
BMI (Median ± IQR)	22 (21-25)	26 (22-30)	<0.0001
LH+day (Median ± IQR)	9 (7-10)	9 (7-10)	0.968
First trimester loss [Median (Range)]	0 (0-2)	5 (3-18)	<0.0001
Live births [Mean (Range)]	0 (0-2)	0.35 (0-2)	<0.0001

* Control subjects were women awaiting IVF treatment for a variety of reasons, including male-factor,

unexplained, and tubo-ovarian infertility. All subjects had regular cycles and considered to have good

prognosis; recurrent IVF failure (RIF: > 3 consecutive IVF failures with good quality embryos) patients

were excluded.

** All Recurrent pregnancy loss (RPL) patients in this cohort had 3 or more consecutive miscarriages.

† Data were tested for normality using Shapiro-Wilk test. *P*-value was calculated by two-tailed unpaired

student's t-test for normally distributed data (age) or two-tailed Mann Whitney test for non-normally

distributed data (BMI, LH+, First trimester loss and Live births).

Fig. 1. Single-cell reconstruction of the decidual pathway in culture.

Fig. 2. K-means cluster analysis identifies co-regulated decidual gene networks.

Fig. 3. Co-ordinated expression of decidual immune surveillance genes.

Fig. 4. Identification of endometrial cell types and subsets during the implantation window.

Fig. 5. *SCARA5* and *DIO2* as marker genes for divergent decidual states.

Fig. 6. Impaired fate divergence of decidual cells in RPL.

Supplementary Information for

Recurrent pregnancy loss is associated with a pro-senescent decidual response during the peri-implantation window

Emma S Lucas, Pavle Vrljicak, Joanne Muter , Maria M Diniz-da-Costa, Paul J Brighton, Chow-Seng Kong, Julia Lipecki, Katherine J Fishwick, Joshua Odendaal, Lauren J Ewington, Siobhan Quenby, Sascha Ott, and Jan J Brosens

Corresponding author: Professor Jan Brosens, Division of Biomedical Sciences, Clinical Sciences Research Laboratories, Warwick Medical School, University of Warwick, Coventry CV2 2DX, United Kingdom. Email: J.J.Brosens@warwick.ac.uk

This PDF file includes:

Figs S1-S11

Captions for Tables S1-S9

Supplementary references

Other supplementary information for this manuscript includes the following:

Tables S1 to S9

Fig. S1. Quality control of timecouse scRNA-seq libraries. Cumulative read distribution for main (**a**) and additional (**b**) timecourse samples. Knee and inflection points calculated with the DropUtils R package are presented as blue and green dashed lines, respectively. **c**, Violin plots showing distribution of number of genes (nGene), number of transcripts (nUMI) and percent of mitochondrial reads (percent.mito) per timecourse samples**.**

Fig. S2. Heatmap of the top 10 marker genes for each of the 7 cell states. Each column represents one of 4,580 EnSC. Expression for each gene is centered to the average expression across cells and scaled by their standard deviation. Yellow, black and purple represents high, medium and low expression, respectively.

Fig. S3. Transient IL-6 and IL-8 secretion during the initial decidual phase. Temporal profiles of *IL6* and *IL8* transcript levels (blue line) and corresponding secreted levels across the decidual pathway (D0-D8) and upon withdrawal (WD) of differentiation signals. Note that the initial rise in IL-6 or IL-8 secretion does not coincide with a corresponding increase in mRNA levels. A second rise in IL-8 secretion coincides with the emergence of snDC.

Fig. S4. Single-cell analysis of three independent primary cultures shows cells aggregating by transcriptional state. Three independent primary EnSC cultures (designated a, b, and c) were decidualized for different time-points and then subjected to Drop-seq analysis. Culture 'a' represents the full time-course [D0-D8 plus withdrawal (WD)]; culture 'b' was decidualized for 2 and 8 days whereas culture 'c' was decidualized for 2 days. **a** *t*-SNE plot with cells colour-coded by culture and day of decidualization. **b** *t*-SNE plot with cells colour-coded by cell state. The plot was further annotated to indicate day of decidualization and withdrawal (WD) of differentiation signals.

Fig. S5. Stress-defence mechanisms in DC. **a** Multiple mechanisms underpin stress-resistance of DC, including progesterone-dependent induction of dual specificity phosphatase 1 (DUSP1, also known as mitogen-activated protein (MAP) kinase phosphatase-, MKP1), which silences the c-Jun NH-terminal kinase (JNK) stress signaling pathway and blocks stress-dependent sumoylation of numerous targets, including the liganded progesterone receptor (PGR)[1,2]. Progesterone also regulates the expression the serum- and glucocorticoid-inducible kinase SGK1[3], a kinase that targets and inactivates FOXO3, a key transcription factor involved in oxidative cell death responses in endometrial cells[4]. PLCL1, coding phospholipase C like 1 (inactive), is a progesterone-inducible scaffold protein that uncouples phospholipase C activation downstream of Gq-protein-coupled receptors from intracellular Ca²⁺ release by attenuating inositol trisphosphate (IP3) signaling[5]. Progesterone further upregulates 11β-hydroxysteroid dehydrogenase type 1 (encoded by HSD11B1)[6,7], the enzyme that converts inert cortisone into active cortisol, a powerful anti-inflammatory hormone. A highly-induced decidual gene is

_IL1RL1_, which encodes the IL-33 transmembrane receptor ST2L as well as the secreted decoy receptor sST2[8], a potent anti-inflammatory mediator that binds and inactivates IL-33. The main non-mitochondrial source of reactive oxygen species in endometrial stromal cells is NADPH oxidase NOX4[9]. Although initiation of the decidual process requires NOX4 activation[9], it is also a mediator of cellular senescence[10]. **b** Log-transformed, normalized expression levels for indicated genes in stress-resistant DC (S5) and snDC (S6).

Fig. S6. Flow cytometric analysis of CD56+ cells following MACS separation. uNK cells isolated by MACS from the supernatants of 4 independent freshly established EnSC cultures were subjected to flow cytometry to confirm enrichment of CD56+ cells (CD56-PE) and cell viability (FVS660) in the positive fraction and the absence of CD56+ cells in the negative fraction. In the positive fraction, approximately 86% of cells were confirmed to be viable uNK cells.

Fig S7. Treatment of decidualizing cells with dasatinib mimics uNK co-cultures Six independent EnSC cultures were decidualized with cAMP and MPA (C+M) for eight days in the presence or absence of the senolytic agent, dasatinib (250 nM). Left panel: Schematic representation of dasatinib experiments. Right panel: fold-change in SAβG activity and secretion of clusterin (CLU) and sST2 (encoded by *IL1RL1*) in decidualized cells treated with dasatinib when compared to decidualized cells cultured without dasatinib (dashed line). Individual cultures are indicated by different colours.

Fig. S8. Quality control of biopsy scRNA-seq libraries. a Cumulative read distribution for biopsy samples. Knee and inflection points calculated with the DropUtils R package are presented as blue and green dashed lines, respectively. **b** Violin plots showing distribution of number of genes (nGene), number of transcripts (nUMI) and percent of mitochondrial reads (percent.mito) per biopsy sample.

Fig. S9. Distribution of ciliated cells in endometrial glandular epithelium. Arrows indicate the expression of ciliated epithelial markers, dynein axonemal assembly factor 1 (DNAAF1), radial spoke head component 1 (RSPH1), and cilia and flagella associated protein 157 (CFAP157), within the glandular compartment of human endometrium. Images were retrieved from the Human Protein Atlas v18.1 (https://www.proteinatlas.org/)[11].

Fig. S10. Analysis of *in vitro* branchpoint genes in luteal phase EnSC *in vivo*. Heatmaps depicting the sum of correlation coefficients of the top 50 gene-gene interactions involved in lineage divergence of decidualizing EnSC *in vitro* compared with *in vivo* expression at LH+8 (**a**) and LH+10 (**b**). The colour key indicates the level of congruency, defined as the sum of correlation coefficients of >1 or <-1 for positively (red) and negatively (blue) co-regulated genes, respectively.

Fig. S11. Differential gene expression in *SCARA5*enriched/*DIO2*reduced EnSC versus *SCARA5*reduced/*DIO2*enriched EnSC *in vivo*. a Heatmap showing relative expression (z-score) of 104 DEG, based Wilcoxon rank sum test with Bonferroni corrected *P*-value < 0.05, between *SCARA5*reduced/*DIO2*enriched EnSC (n=368) and *SCARA5*enriched/*DIO2*reduced EnSC (n=271). Columns represent individual EnSC grouped according to their *SCARA5* and *DIO2* transcript levels. *SCARA5*enriched/*DIO2*reduced EnSC were enriched in genes encoding several key decidual TF, including *CEBPB* [12,13], *NR4A1* (also known as *NUR77*) [14] and members of the AP-1 gene family (*FOS*, *FOSB*, and *JUN*) [14,15]. *SCARA5*enriched/*DIO2*reduced EnSC were also enriched in genes involved in stress defences, including *SOD2* (coding superoxide dismutase 2) [16], *DUSP1* (coding MKP1) [1], *TNFAIP3* (TNF alpha induced protein 3), and *NFKBIA* (NFKB inhibitor alpha, IκBα) TNFAIP3 is a ubiquitin-modifying enzyme and potent inhibitor of NF-kappa B activation as well as TNF-mediated apoptosis. It plays a critical role in preventing inflammation *in vivo* [17]. Likewise, IκBα potently suppresses inflammation by binding to NF-kappa B. By contrast, enhanced expression of genes coding various heat shock proteins (HSPs) in *SCARA5*reduced/*DIO2*enriched cells is indicative of a proteotoxic stress response [18]. **b** Gene ontology analysis revealed significantly enriched biological themes associated with *SCARA5*reduced *DIO2*enriched versus *SCARA5*enriched/*DIO2*reduced EnSC. The X-axis shows the proportion of

differentially expressed genes found in each biological category. Benjamini-adjusted *P*-values (Fisher's exact test) are shown.

Supplementary Tables (separate file):

Table S1. Marker genes for each cell state

Table S2. Differentially expressed genes (DEG) *in vitro* (S1-S7)

Table S3. Contribution of individual samples to the endometrial cell populations

Table S4. k-means cluster analysis (k=7)

Table S5. GO analysis of co-expressed categories

Table S6. Marker genes of endometrial cell types and subsets

Table S7. Marker genes of immune cell clusters and sub-types

Table S8. DEG in LH8 vs LH10 EnSCs

Table S9. DEG in *SCARA5*[enriched]/*DIO2*[reduced] EnSC vs *SCARA5*[reduced]/*DIO2*[enriched] EnSC *in vivo*.

Supplementary References:

Leitao, B. *et al.* Silencing of the JNK pathway maintains progesterone receptor activity in decidualizing human endometrial stromal cells exposed to oxidative stress signals. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology* **24**, 1541-1551 (2010).

Leitao, B. B., Jones, M. C. & Brosens, J. J. The SUMO E3-ligase PIAS1 couples reactive oxygen species-dependent JNK activation to oxidative cell death. *FASEB J* **25**, 3416-3425, doi:10.1096/fj.11-186346 (2011).

Salker, M. S. *et al.* Deregulation of the serum- and glucocorticoid-inducible kinase SGK1 in the endometrium causes reproductive failure. *Nature medicine* **17**, 1509-1513, doi:10.1038/nm.2498 (2011).

Kajihara, T. *et al.* Differential expression of FOXO1 and FOXO3a confers resistance to oxidative cell death upon endometrial decidualization. *Mol Endocrinol* **20**, 2444-2455, doi:10.1210/me.2006-0118 (2006).

Muter, J. *et al.* Progesterone-Dependent Induction of Phospholipase C-Related Catalytically Inactive Protein 1 (PRIP-1) in Decidualizing Human Endometrial Stromal Cells. *Endocrinology* **157**, 2883-2893, doi:10.1210/en.2015-1914 (2016).

Kuroda, K. *et al.* Elevated periimplantation uterine natural killer cell density in human endometrium is associated with impaired corticosteroid signaling in decidualizing stromal cells. *The Journal of clinical endocrinology and metabolism* **98**, 4429-4437, doi:10.1210/jc.2013-1977 (2013).

Kuroda, K. *et al.* Induction of 11beta-HSD 1 and activation of distinct mineralocorticoid receptor- and glucocorticoid receptor-dependent gene networks in decidualizing human endometrial stromal cells. *Mol Endocrinol* **27**, 192-202, doi:10.1210/me.2012-1247 (2013).

Salker, M. S. *et al.* Disordered IL-33/ST2 activation in decidualizing stromal cells prolongs uterine receptivity in women with recurrent pregnancy loss. *PLoS One* **7**, e52252, doi:10.1371/journal.pone.0052252 (2012).

Al-Sabbagh, M. *et al.* NADPH oxidase-derived reactive oxygen species mediate decidualization of human endometrial stromal cells in response to cyclic AMP signaling. *Endocrinology* **152**, 730-740, doi:10.1210/en.2010-0899 (2011).

Weyemi, U. *et al.* ROS-generating NADPH oxidase NOX4 is a critical mediator in oncogenic H-Ras-induced DNA damage and subsequent senescence. *Oncogene* **31**, 1117-1129, doi:10.1038/onc.2011.327 (2012).

Uhlen, M. *et al.* Proteomics. Tissue-based map of the human proteome. *Science* **347**, 1260419, doi:10.1126/science.1260419 (2015).

Christian, M., Pohnke, Y., Kempf, R., Gellersen, B. & Brosens, J. J. Functional association of PR and CCAAT/enhancer-binding protein beta isoforms: promoter-dependent cooperation between PR-B and liver-enriched inhibitory protein, or liver-enriched activatory protein and PR-A in human endometrial stromal cells. *Mol Endocrinol* **16**, 141-154, doi:10.1210/mend.16.1.0763 (2002).

Christian, M. *et al.* Cyclic AMP-induced forkhead transcription factor, FKHR, cooperates with CCAAT/enhancer-binding protein beta in differentiating human endometrial stromal cells. *J Biol Chem* **277**, 20825-20832, doi:10.1074/jbc.M201018200 (2002).

Jiang, Y. *et al.* The orphan nuclear receptor Nur77 regulates decidual prolactin expression in human endometrial stromal cells. *Biochem Biophys Res Commun* **404**, 628-633, doi:10.1016/j.bbrc.2010.12.027 (2011).

Mazur, E. C. *et al.* Progesterone receptor transcriptome and cistrome in decidualized human endometrial stromal cells. *Endocrinology* **156**, 2239-2253, doi:10.1210/en.2014-1566 (2015).

Kajihara, T. *et al.* Human chorionic gonadotropin confers resistance to oxidative stress-induced apoptosis in decidualizing human endometrial stromal cells. *Fertil Steril* **95**, 1302-1307, doi:10.1016/j.fertnstert.2010.05.048 (2011).

Lee, E. G. *et al.* Failure to regulate TNF-induced NF-kappaB and cell death responses in A20-deficient mice. *Science* **289**, 2350-2354, doi:10.1126/science.289.5488.2350 (2000).

Nollen, E. A. & Morimoto, R. I. Chaperoning signaling pathways: molecular chaperones as stress-sensing 'heat shock' proteins. *J Cell Sci* **115**, 2809-2816 (2002).