Tip rates, phylogenies, and diversification: what are we estimating, and how good are the

estimates?

Pascal O. Title[1,2,3], Daniel L. Rabosky[1]

[1]Department of Ecology and Evolutionary Biology and Museum of Zoology, University of

Michigan, Ann Arbor, MI 48109, USA

[2]Environmental Resilience Institute, Indiana University, Bloomington, IN 47408, USA

[3]ptitle@iu.edu

<u>Abstract</u>

1. Species-specific diversification rates, or "tip rates", can be computed quickly from

phylogenies and are widely used to study diversification rate variation in relation to

geography, ecology, and phenotypes. These tip rates provide a number of theoretical and

practical advantages, such as the relaxation of assumptions of rate homogeneity in trait-

dependent diversification studies. However, there is substantial confusion in the literature

regarding whether these metrics estimate speciation or net diversification rates.

Additionally, no study has yet compared the relative performance and accuracy of tip rate

metrics across simulated diversification scenarios.

2. We compared the statistical performance of three model-free rate metrics (inverse

terminal branch lengths; node density metric; DR statistic) and a model-based approach

(BAMM). We applied each method to a large set of simulated phylogenies that had been

generated under different diversification processes; scenarios included multi-regime time-

constant and diversity-dependent trees, as well as trees where the rate of speciation

evolves under a diffusion process. We summarized performance in relation to the type of

rate variation, the magnitude of rate heterogeneity and rate regime size. We also

compared the ability of the metrics to estimate both speciation and net diversification

rates.

3.   We show decisively that model-free tip rate metrics provide a better estimate of the rate

of speciation than of net diversification. Error in net diversification rate estimates

increases as a function of the relative extinction rate. In contrast, error in speciation rate

estimates is low and relatively insensitive to extinction. Overall, and in particular when

relative extinction was high, BAMM inferred the most accurate tip rates and exhibited

lower error than non-model-based approaches. DR was highly correlated with true

speciation rates but exhibited high error variance, and was the best metric for very small

rate regimes.

4.   We found that, of the metrics tested, DR and BAMM are the most useful metrics for

studying speciation rate dynamics and trait-dependent diversification. Although BAMM

was more accurate than DR overall, the two approaches have complementary strengths.

Because tip rate metrics are more reliable estimators of speciation rate, we recommend

that empirical studies using these metrics exercise caution when drawing biological

interpretations in any situation where the distinction between speciation and net

diversification is important.

Keywords: tip rates, diversification, terminal branch length, node density, DR, BAMM, trait-

dependent diversification

<u>Introduction</u>

Rates of speciation and extinction vary through time and among lineages (Nee, Mooers &

Harvey 1992; Sanderson & Donoghue 1996; Etienne & Haegeman 2012; Jetz et al. 2012; Moen

& Morlon 2014; Alfaro et al. 2018), contributing to dramatic heterogeneity in species richness

across the tree of life (Alfaro et al. 2009; Jetz et al. 2012; Barker et al. 2013). By characterizing

variation in rates of speciation and extinction, we can better understand the dynamics of

biological diversity through time, across geographic and environmental gradients (Zink, Klicka

& Barber 2004; Ricklefs 2006; Mittelbach et al. 2007; Silvestro, Schnitzler & Zizka 2011;

Rabosky, Title & Huang 2015), and in relation to traits and key innovations (FitzJohn, Maddison

& Otto 2009; Near et al. 2012; Beaulieu & O'Meara 2016). Consequently, there has been great

interest in statistical methods for inferring rates of speciation and extinction from molecular

phylogenies.

Although rates of diversification have traditionally been quantified for clades, there has

been a growing interest in estimating species-specific rates of diversification, which we refer to

here as "tip rates". Tip rates are increasingly used to describe patterns of geographic and trait-

associated variation in diversification (Freckleton, Phillimore & Pagel 2008; Jetz et al. 2012;

Kennedy et al. 2016; Harvey & Rabosky 2017; Quintero & Jetz 2018; Rabosky et al. 2018). It

may seem strange to view evolutionary rates as a property of individual lineages, but such rates

emerge naturally from the birth-death model we typically use to conceptualize the diversification

process (Nee, Mooers & Harvey 1992; Nee, May & Harvey 1994). Under the birth-death

process, individuals (species) are characterized by per-lineage rates of species origination

(speciation, λ) and extinction (μ). For the purposes of inference, these rates are typically

assumed to be constant among contemporaneous members of a focal clade. However, tip rates

can be viewed as our best estimate of the present-day rate of speciation or extinction for an

individual lineage, conditional on past (usually recent) evolutionary history. As such, they

provide information about the expected amount of time that will elapse before a lineage splits or

becomes extinct.

A number of approaches have been used to estimate tip rates, including both model-based

and non-model-based approaches (i.e., models that are parameterized with speciation and

extinction rates, vs metrics that simply rely on branch lengths and splitting events). These

approaches vary in terms of how much information they derive from a focal species (i.e., a

terminal branch) relative to the amount of information they incorporate from other regions of the

phylogeny. On one end of the spectrum, tree-wide estimates (i.e., one rate for the entire

phylogeny) of speciation and extinction rates under a constant-rate birth-death (CRBD) model

provide tip rates that are maximally auto-correlated (identical) across species in the clade; such

rates for any given species are not independent of rates for any other species in the group of

interest. On the other end of the spectrum, terminal branch lengths can be used to derive a

censored estimate of the rate of speciation that is minimally autocorrelated with rates for other

species in the focal clade. Terminal branch lengths are largely unique to each species (rates

might be identical only for sister taxa), but provide a noisy measure of speciation, due to the

stochasticity inherent in the diversification process (Nee, May & Harvey 1994), and they have

been employed as a summary statistic in assessing model adequacy (Bromham, Hua & Cardillo

2016; Gomes, Sorenson & Cardoso 2016). In contrast to single (terminal) branch estimates, tree-

wide estimates should be less susceptible to stochastic noise, because they incorporate

information from the entirety of the tree (e.g., multiple branches are used in the estimates). Of

course, the tree-wide estimate necessarily assumes that all tips share a common underlying

diversification process. Other tip rate metrics fall somewhere between these two extremes,

incorporating some tree-wide information but relaxing the assumption of homogeneous rates

across all lineages (node density metric: Freckleton, Phillimore & Pagel 2008; DR: Jetz et al.

2012). The estimation of tip-specific rates thus entails a tradeoff between the precision of

individual estimates and the stochastic error associated with those estimates.

BAMM (Bayesian Analysis of Macroevolutionary Mixtures, Rabosky 2014) is a model-

based approach that can accommodate heterogeneity in the rate of diversification through time

and among lineages. BAMM simulates a posterior distribution of macroevolutionary rate shift

configurations given a phylogeny of interest; marginal rates of speciation and extinction for

individual taxa can then be extracted from this distribution. In this framework, the correlation in

rates between any pair of species is a function of the posterior probability that they share a

common macroevolutionary rate regime (Rabosky et al. 2014). If the tree-wide posterior

probability of rate variation is low, the marginal rates estimates for individual species will be

similar across the entire tree, as under a CRBD model. Likewise, any pair of taxa that are

consistently assigned to the same macroevolutionary rate regime will necessarily have identical

tip rates.

Tip rates are best suited to a host of questions and hypotheses where the diversification

dynamics over the evolutionary history of a group are either less relevant, or no more relevant,

than the rates of diversification closer to the present day. For example, many hypotheses

involving trait-dependent diversification implicitly assume a time-homogeneous, or constant

through time, effect of the trait on diversification rate (Coyne & Orr 2004; Kay et al. 2006;

Jablonski 2008; FitzJohn 2010; Claramunt et al. 2011). Harvey & Rabosky (2017) found that the

use of tip rates for assessing correlations between continuous traits and diversification has good

performance across a range of diversification scenarios. Furthermore, hypotheses pertaining to

non-historical geographic patterns of diversity are also better addressed with recent rates of

diversification. For example, many hypotheses for the latitudinal diversity gradient propose time-

homogeneous effects of particular environmental factors (temperature, energy, geographic area)

on rates of diversification (Mittelbach et al. 2007; Kennedy et al. 2014; Rabosky, Title & Huang

2015; Schluter 2016; Rabosky et al. 2018). Put simply, if such time-homogeneous processes

have shaped the latitudinal diversity gradient (e.g., correlation between speciation and

temperature: Rohde 1992), then the effect should be manifest in the distribution of present-day

evolutionary rates.

At present, there is substantial confusion in the literature over what quantity various tip

rate metrics actually measure. The DR statistic (Jetz et al. 2012) was originally described as a

measure of the "species-level lineage diversification rate". While supplemental analyses and

subsequent work suggested that DR was a better measure of speciation rate than net

diversification (Jetz et al. 2012, Belmaker & Jetz 2015, Quintero & Jetz 2018), many studies

have nonetheless continued to describe DR as an estimate of the lineage-level net diversification

rate (Marin & Hedges 2016; Oliveira et al. 2016; Cai et al. 2017; and many others).The node

density metric of Freckleton, Phillimore & Pagel (2008) has also been described as a measure of

net diversification. Whether these metrics more accurately measure speciation or net

diversification is critically important for interpreting biodiversity patterns (e.g., two regions

might differ dramatically in speciation rate, but net diversification rates in each might

nonetheless be zero). An objective of our study is thus to compare the ability of DR, node

density, and other metrics to estimate speciation and net diversification rates.

Despite the potential utility of tip rates in geographic and trait-based analyses of

speciation rate heterogeneity (Jetz et al. 2012; Belmaker & Jetz 2015; Oliveira et al. 2016;

Quintero & Jetz 2018), there has yet been no comprehensive comparative assessment of the

accuracy and precision of the estimates, save for supplemental analyses in Jetz et al. (2012) and

Quintero & Jetz (2018). BAMM has low power to infer small rate regimes (Rabosky, Mitchell &

Chang 2017; Meyer & Wiens 2017), leading to the possibility that other approaches might

perform better for smaller phylogenies or when the variation in rates among clades is subtle.

However, DR and related methods will always identify variation in tip rates, even when none

exists, provided there is stochastic variation in branch lengths. A goal of this study is therefore to

evaluate the trade-off between the stochastic noise inherent in non-model-based approaches, and

the conservative but less noisy estimates from model-based metrics. We compare the

performance of these metrics across a range of simulation scenarios, which include both discrete

and continuous variation in rates.

Methods

Tip rate metrics

We assessed the accuracy of four tip rate metrics in this study at quantifying rates of

speciation. As we demonstrate below (see also Supplementary Figure 5 in Jetz et al. 2012;

extended Figure 5 in Quintero & Jetz 2018; Belmaker & Jetz 2015), these metrics are estimators

of speciation rate and not net diversification rate, and we refer to them as such throughout. The

first metric is the inverse of the equal splits measure (Redding and Mooers 2006), also called the

DR statistic (Jetz et al. 2012), *DivRate* (Belmaker & Jetz 2015; Oliveira et al. 2016), *ES* (Harvey

& Rabosky 2017) or *tip DR* (Quintero & Jetz 2018), which we denote in this study as λ_{DR}. This

species-specific measure incorporates the number of splitting events and the internode distances

along the root-to-tip path of a phylogeny, while giving greater weight to branches closer to the

present (Redding & Mooers 2006; Jetz et al. 2012). λ_{DR} is computed as:

166
$$\lambda_{DR_i} = \sum_{j=1}^{N_i} b_j \frac{1}{2^{j-1}}$$

where λ_{DR_i} is the tip rate for species i, N_i is the number of branches between species i and the

root, b_j is the length of branch j, starting at the terminal branch ($j = 1$) and ending with the root.

Jetz et al. (2012) demonstrated that, for trees deriving from a Yule process, and with mild

extinction, the mean λ_{DR} across tips converges on the true speciation rate.

We also considered a simpler metric, node density (Freckleton, Phillimore & Pagel 2008;

denoted by λ_{ND}). This is simply the number of splitting events along the path between the root

and tip of a phylogeny, divided by the age of the phylogeny. While λ_{DR} down-weights the

contribution of branch lengths that are closer to the root, λ_{ND} equally weights the contributions of

all branches along a particular root-to-tip path, regardless of where they occur in time. Under a

pure-birth model ($\mu = 0$), both λ_{DR} and λ_{ND} should yield unbiased estimates of the rate of

speciation.

The third measure we considered is the inverse of the terminal branch lengths (λ_{TB}).

Rapid speciation rates near the present should be associated with proportionately shorter terminal

branches; smaller values of λ_{TB} should thus characterize species with faster rates of speciation.

This measure has recently been used as a summary statistic to assess model adequacy in trait-

dependent diversification studies (Bromham, Hua & Cardillo 2016; Gomes, Sorenson & Cardoso

2016; Harvey & Rabosky 2017). Following Steel & Mooers (2010), we note that the terminal

branch lengths can be used to derive an estimate of the speciation rate; this follows from the fact

that interior and terminal branches have the same expected value under the Yule process (Steel &

Mooers 2010). The corresponding estimator for the i'th tip, λ_{TB} is approximately $1 / 2b$ where b is

the length of a given terminal branch (Steel & Mooers 2010). To our knowledge, λ_{TB} has not

been used to explicitly estimate tip rates as we do here, but given its utility as a summary statistic

and general theoretical properties (Steel & Mooers 2010), we see value in comparing the

performance of this metric to others currently in use.

Finally, we considered a Bayesian, model-based approach to estimating tip rates. BAMM

(Rabosky 2014) assumes that phylogenies are generated by set of discrete diversification

regimes. Using MCMC, the program simulates a posterior distribution of rate shift regimes, from

which marginal posterior rate distributions can be extracted for each tip in the phylogeny. Priors

for BAMM analyses were set using default settings from the setBAMMpriors function from

BAMMtools (Rabosky et al. 2014). The prior parameterizations specified by this function ensure

that the prior density on relative rate changes across the tree is invariant to the scale of the tree

(e.g., multiplying branch lengths by 10^6 will not change inferences about relative rates across the

tree). We denote BAMM tip speciation rates (mean of the marginal posterior) as λ_{BAMM}. As

BAMM also estimates extinction rates for each regime, we also calculated tip-specific net

diversification rate as λ_{BAMM} - μ_{BAMM}, denoted as r_{BAMM}.

<u>Tip rate metrics estimate speciation, not net diversification</u>

As suggested previously (Belmaker & Jetz 2015; supplemental analyses in Jetz et al.

2012), DR and presumably other tip-based measurements, more accurately estimate the rate of

speciation than the rate of net diversification. However, numerous studies continue to refer to DR

as a measure of net diversification (Marin & Hedges 2016; Oliveira et al. 2016; Cai et al. 2017;

Quintero & Jetz 2018; and many others). This is incorrect and it is straightforward to

demonstrate that $\lambda_{TB,}$ λ_{ND} and λ_{DR} are more reliable measures of speciation rates and not net

diversification rates, at least when extinction is moderate to high.

To illustrate this property of the metrics, we applied all approaches to constant-rate birth-

death phylogenies simulated across a range of extinction fractions ($\varepsilon = \mu / \lambda$) , including pure-

birth trees ($\varepsilon = 0$) as well as trees exhibiting very high turnover ($\varepsilon = 1$). To evaluate accuracy of

speciation estimates as a function of ε, we generated 1000 phylogenies with 100 tips each, where

λ and ε were drawn from uniform distributions (λ: [0.05, 0.3]; ε: [0, 1]). Importantly, when λ is

sampled uniformly with respect to ε, the distribution of r is not uniform: the mean, range and

variance in r decrease dramatically as ε increases. To evaluate the accuracy of r as a function of

ε, we thus generated a second set of trees by sampling r and ε from uniform distributions (r:

[0.05, 0.3], ε [0, 1]). As a result, λ has constant mean and variance with respect to ε in the first

set of simulations, and the same is true for r in the second set of simulations (Figure S1). All

phylogeny simulations were conducted with the TreeSim package in R (Stadler 2011).

222   We compared tip rate metrics to true speciation rates λ_{TRUE} (with the first simulation set)

223   and to true net diversification rates r_{TRUE} (with the second simulation set). We evaluated mean

224   per-tip accuracy of the tip rate metrics with three measures of error:

$$\text{mean absolute error} = \sum_{i=1}^{N_i} |\lambda_i - \lambda_{TRUE_i}|/N$$

$$\text{RMSE} = \sqrt{\sum_{i=1}^{N_i} (\lambda_i - \lambda_{TRUE_i})^2/N}$$

$$\text{mean proportional error} = \sum_{i=1}^{N_i} \frac{\lambda_i - \lambda_{TRUE_i}}{\lambda_{TRUE_i}}/N$$

where λ_i is the estimated tip rate for species i out of N total species, λ_{TRUE} is the true tip rate.

Mean absolute error and root mean square error capture the magnitude in error in tip rates, and

mean proportional error quantifies the bias in tip rates, as a function of the true tip rates

(Rabosky et al. 2014). In analyses below, all error summaries yield generally congruent results;

results for mean absolute error are presented in the main text, and others in supplement.

<u>Assessment of tip rate metrics</u>

We tested the performance of the metrics by compiling publicly-available datasets from a

number of simulation-based studies (Table 1). By focusing on simulations from previously-

published work, we thus ensured that the simulation process itself was effectively blinded to the

objectives of the present study. We further note that our trial datasets included several studies

that were critical of BAMM (Moore et al. 2016, Meyer & Wiens 2017). These simulated trees

include rate heterogeneity in time and across lineages. Together, these phylogenies present a

wide range of tree sizes and diversification rate shifts, providing an ideal comparative dataset for

our purposes. To more easily distinguish between these tree types in the text, we refer to the

BAMM-type, multi-regime time-constant phylogenies simply as "multi-regime", and the multi-

regime diversity-dependent phylogenies simply as "diversity-dependent", even though discrete

rate shifts are present in both types of trees. In addition to discrete-shift scenarios (e.g., BAMM-

type process), we simulated phylogenies under an "evolving rates" model of diversification

(Rabosky 2010; as corrected in Beaulieu & O'Meara 2015) to explore performance of tip rate

metrics when diversification rates change continuously and independently along branches, as

might occur if diversification rates are correlated with an underlying continuous trait (FitzJohn

2010). In these simulations, we allowed the logarithm of λ to evolve across the tree under a

Brownian motion process, while holding ε constant. The magnitude of rate heterogeneity among

branches is controlled by the diffusion parameter σ, where greater values lead to greater

heterogeneity in speciation rates. Although published phylogenies with rate data were

unavailable for this simulation scenario, we used simulation code and parameters taken directly

from Beaulieu & O'Meara (2015) to generate trees with similar statistical properties to those in

their study. Simulations were performed with the following parameters: $\lambda = 0.078, 0.103, 0.145,$

0.249 and $\varepsilon = 0.0, 0.25, 0.50, 0.75$. We simulated 100 phylogenies for each (λ, ε) pair, and for

three values of σ ($\sigma = 0.03, 0.06, 0.12$). We evaluated tip rate accuracy by comparing estimated

to true tip rates, using the absolute and proportional error metrics described above. We also

examined the correlation between true and estimated tip rates, combining tip rates from all

phylogenies generated under the same class of diversification process, and visualizing these data

as density scatterplots, generated with the LSD package in R (Schwalb et al. 2018), where colors

indicate the density of points.

Size of diversification rate regimes might be an important factor in a tip rate metric's

ability to accurately estimate rates. For example, BAMM's statistical power in detecting a shift

to a new rate regime is a function of the number of taxa in that rate regime, and tip rates for taxa

from small regimes will more likely be parameterized according to the larger parent regime or

the tree-wide average rate (Rabosky, Mitchell & Chang 2017); this is the expected behavior

when BAMM fails to identify a rate shift. However, non-model-based approaches such as those

examined in this study might be more accurate for small regimes. To explore how rate regime

size influences the accuracy of tip rate metrics, we calculated the mean tip rate for each true rate

regime from all multi-regime phylogenies (simulation datasets from Moore et al. 2016; Rabosky,

Mitchell & Chang 2017; Meyer & Wiens 2017; Mitchell, Etienne & Rabosky 2018). We then

calculated the Pearson correlation coefficient and the slope of a linear model between true and

estimated mean regime rates. We explored the performance of all metrics with respect to regime

sample size, as in Rabosky, Mitchell & Chang (2017: Figure 13). For comparison, we repeated

all performance summaries on tip rates estimated by applying a simple constant-rate birth-death

(CRBD) process to each simulated phylogeny. This exercise is an important control, because it

indicates how much error we would expect for each simulated phylogeny under the simplifying

(incorrect) assumption that rates are constant among lineages and through time for each dataset.

Results

<u>Speciation or net diversification?</u>

As expected, the tip rate metrics examined in this study are more accurate estimators of

the rate of speciation (λ) and not the net rate of species diversification (r). Mean absolute error

increased exponentially with respect to the extinction fraction ε (Figure 1). However, mean

absolute error in speciation rate was largely invariant with respect to ε (0.95 quantile of r-based

and λ-based mean absolute error for λ_{DR}: 2.28 and 0.17, respectively). Nearly identical patterns

were found with RMSE (Figure S2). Note that r and λ for these simulations were drawn from

identical uniform distributions, and absolute error in the rates is thus comparable. Proportional

error generally exhibited the same pattern, and in terms of λ versus r, differences in speciation-

based error varied across ε (Figure S3). There was a weak but significant trend towards

progressively greater underestimation of speciation rates with increasing values of relative

extinction (linear model slopes: -0.08, -0.014, -0.011 for λ_{ND}, λ_{DR} and λ_{BAMM}, respectively).

Overall, error was highest for λ_{TB} by an order of magnitude (Figure S4), and decreased

progressively with λ_{ND} and λ_{DR}, with the lowest overall error in λ_{BAMM}. BAMM estimates of net

diversification rate were relatively accurate, except at the highest values of ε (Figures 1, S2, S3).

<u>Tip rate accuracy across rate-variable phylogenies</u>

Tip rates estimated with BAMM were consistently more accurate than those obtained

using the other methods across all diversification scenarios considered, including multi-regime,

diversity-dependent and evolving rates trees (Figure 2). λ_{DR} was the second-most accurate

metric, although its relationship with true rates was substantially weaker than λ_{BAMM}. λ_{ND} and

λ_{TB} were correlated with true rates but performed relatively poorly overall. However, λ_{TB}

performed better than λ_{ND}, and just as well as λ_{DR} at estimating speciation rates for diversity-

dependent trees (Figure 2, S5). All metrics performed best for multi-regime trees, followed by

evolving rates and diversity-dependent trees, respectively. For diversity-dependent trees, λ_{ND}

rates are effectively uncorrelated with the true rates (Figure 2). Additionally, the performance of

the different tip rate metrics for multi-regime phylogenies is not sensitive to the source of the

simulated phylogenies (Figure S6). We found that BAMM substantially outperformed all other

metrics on datasets from studies that independently assessed BAMM's performance (Figure S6:

Moore et al. 2016; Meyer & Wiens 2017). Tip rates were also generally but more weakly

correlated with true net diversification rates, with the exception of λ_{ND}, which was uncorrelated

with true rates for diversity-dependent trees, presumably because this metric equally weights the

full depth of the tree (Figure S7).

In terms of mean per-tip error, λ_{BAMM} consistently outperformed the other metrics for

multi-regime, diversity-dependent and evolving rates trees (Figures 3, S8). Error in λ_{BAMM}

increased as a function of rate heterogeneity for evolving rate phylogenies, but was largely

independent of the magnitude of rate heterogeneity for the other scenarios. λ_{DR} generally

exhibited greater error than λ_{BAMM}, and this error increased as a function of the level of

heterogeneity for both the evolving rates and multi-regime trees. Error in λ_{DR} was generally

invariant to the number of rate regimes for the diversity-dependent scenarios. However, λ_{DR}

tended to have greater error than tip estimates from a simple model that assumes no variation in

rates through time or among lineages (λ_{CRBD}; all tips assigned the tree-wide CRBD rate). λ_{ND}

performed somewhat similarly to λ_{DR} for constant-rate and evolving rates trees, but worse for

diversity-dependent trees. Error in λ_{TB} increased with increasing rate heterogeneity for constant-

rate and evolving rates trees, but was relatively unaffected by rate heterogeneity in diversity-

dependent trees (Figure S9). However, error for this metric was far greater than for all other tip

metrics.

<u>Effects of regime size on performance</u>

Both metrics of performance assessment – the Pearson correlation and OLS slope –

generally increased with increasing regime size (Figure 4). This was found to be true for all tip

rate metrics, although λ_{TB} and λ_{ND} never achieved high performance. λ_{DR} tended to perform

better than other metrics when small rate regimes were included (e.g., 10 tips or fewer);

however, the slope between estimated and true rates was greater than 1 across the majority of

minimum regime sizes, indicating that λ_{DR} overestimates speciation rates (see also Figure S3).

Similar patterns were observed for net diversification rates with λ_{DR}, but the magnitude of the

overestimation was greater than for speciation (Figure S10). λ_{BAMM}, in contrast, approached a

slope of 1 when estimating speciation rates and slightly underestimated net diversification rates

(regimes with > 30 tips: OLS slope = 0.96 for λ, 0.87 for r).

Absolute error in regime mean tip rates was lowest for λ_{DR} and λ_{BAMM}, regardless of the

size of the rate regime (Figure 5). BAMM's ability to accurately estimate tip rates improved with

regime size, whereas absolute error was relatively consistent across regime sizes for λ_{DR} for

regimes greater than 10 species. We also found that λ_{DR} slightly outperformed λ_{BAMM} for small

rate regimes.

Note that, in Figures 4 and 5, each rate regime is treated as a single data point. Rate

regimes of sizes 1000, 100, and 1 tip are equivalent under this method of error assessment.

Figure 4 assesses how well these methods estimate rates for individual regimes, regardless of the

size of those regimes. In contrast, Figures 1-3 ask how well these methods perform at estimating

rates for a given tip.

Discussion

We assessed several tip rate metrics and confirmed that these are more accurate

estimators of the rate of speciation, rather than net diversification (Figures 1, 4, S7, S10). This

distinction was especially pronounced at high extinction fractions, where the rate of lineage

turnover is high, and rates of speciation and net diversification have the potential to be more

divergent. These results are consistent with supplemental analyses performed in Jetz et al. 2012.

It is also important to note that recent extinction will have a much greater influence on these

metrics than extinction events deeper in time (Quental & Marshall 2011). Net diversification rate

is a critical determinant of species richness, yet this quantity is potentially independent of the

underlying rate of speciation. Misinterpretation of tip rate metrics could therefore lead to highly

misleading perspectives on large-scale diversity dynamics. As we demonstrate (Figures 1, S2,

S3), tip rate metrics (λ_{ND}, λ_{DR}) provide relatively little information about net diversification, and

high values of these metrics are fully consistent with equilibrial models of speciation where the

true net diversification rate is zero. Thus, λ_{DR} and λ_{ND} should not be used to support claims about

the dynamics of species richness or net diversification *per se* without independent evidence

bearing on plausible levels of extinction.

In terms of accuracy, we found that BAMM performed better than non-model-based

metrics across all datasets we considered: estimated tip rates were most highly correlated with

true tip rates, and mean per-tip error in rates was lower across a range of rate-variable simulation

scenarios. This performance is likely to be at least partially due to the inclusion of extinction in

the BAMM inference model. BAMM is expected to perform well for phylogenies with discrete

shifts in diversification rates as this type of rate variation is most consistent with BAMM's

assumptions (Rabosky 2014; Mitchell & Rabosky 2016; Rabosky, Mitchell & Chang 2017;

Mitchell, Etienne & Rabosky 2018). However, BAMM performed surprisingly well for the

evolving rates phylogenies, which conform poorly to the assumptions of the inference model. In

these trees, the rate of speciation changes continuously under a diffusion process, and as a result,

the phylogeny exhibits rate heterogeneity without discrete rate shifts.

On evolving rates phylogenies, λ_{BAMM} performed better than λ_{DR} (Figure 2; Spearman's ρ

for $\lambda_{BAMM} = 0.83$, ρ for $\lambda_{DR} = 0.62$), despite the fact that λ_{DR} does not rely on the detection of

distinct rate regimes to estimate tip rates (Figure 5). λ_{BAMM} also exhibited the lowest mean per-

tip error across varying levels of rate heterogeneity (Figure 3).

Why do λ_{BAMM} and λ_{DR} exhibit such striking differences in performance across the

simulation scenarios considered here? To illustrate the differences between inference under these

metrics, we compared true tip rates to λ_{BAMM} and to λ_{DR} on a simulated birth-death tree with a

single rate shift (Figure 6), as well as on one evolving rates tree simulated for this study (Figure

7). It is clear that if BAMM has the statistical power to detect true rate shifts, then it will perform

well under rate shift scenarios. In contrast, λ_{DR} tracks true rate shifts but exhibits high sample

variance. With an evolving rates tree (Figure 7), the simulation model is very different from the

inference model in BAMM. However, it conservatively places rate shifts in order to

accommodate rate heterogeneity that is spread across the phylogeny under a rather different

model of rate variation. λ_{DR} also broadly tracks the overall pattern of the true rates, but the

variance in the corresponding estimates is so high that performance is negatively affected. If we

calculate mean (absolute) per-tip error in λ_{BAMM} and $\lambda_{DR,}$ the error is relatively similar between

λ_{BAMM} and λ_{DR}, but the variance in per-tip error for λ_{DR} is higher. Overall, BAMM exhibited

substantially lower error than λ_{DR} under precisely this scenario (Figure 3).

Thus, although BAMM is conservative in the estimation of tip rates relative to λ_{DR}, the

method exhibits lower overall error. It appears that λ_{DR} can recover more subtle rate

heterogeneity relative to BAMM (see Rabosky, Mitchell & Chang 2017 for discussion of power

in BAMM), but this apparent power advantage comes at the cost of increased variance (error) in

the resulting estimates. Remarkably, on a per-tip basis, we find that a simple constant-rate birth-

death process (λ_{CRBD}) frequently yields tip estimates with lower median error and less error

variance than those obtained with λ_{DR} (Figure 3), despite the simplifying (and incorrect)

assumption that rates are identical across all tips in a given tree. For example, across all multi-

regime simulations (Figure 3), λ_{CRBD} point estimates were more accurate than the corresponding

λ_{DR} point estimates for 84% of trees in the simulations; for λ_{BAMM}, the λ_{CRBD} estimates were

more accurate for a much smaller fraction of the total (36%). Similar results were noted for

diversity-dependent (λ_{CRBD} more accurate than 98% of λ_{DR} estimates, versus 15% of λ_{BAMM}

estimates) and evolving rates trees (λ_{CRBD} more accurate than 93% of λ_{DR} estimates, versus 36%

of λ_{BAMM} estimates). Given that λ_{DR} can and does track true heterogeneity in speciation rate

(Figures 6, 7), this pattern suggests that the metric is especially sensitive to the stochastic

variation in branch lengths that can emerge even when all tips have the same underlying

speciation rate.

Regardless of the performance summaries presented in this article, important questions

remain with respect to how well tip rate metrics can estimate the true rate of speciation from

empirical phylogenies. The phylogenies analyzed in this study were simulated under idealized

processes and neglect potential biases and sources of uncertainty that are present in real datasets.

For example, if the process of speciation takes time to complete, as is generally believed to be

the case (i.e., the protracted speciation process; Rosindell et al. 2010; Etienne and Rosindell

2012), then the most recent speciation events may still be on-going at the present and typical

species-level molecular phylogenies may fail to recognize these events. This will lead to an

overestimation of terminal branch lengths, as some terminal branches potentially include

incipient species. A related bias might arise due to incomplete taxon sampling, which

disproportionately affects the length of terminal (or otherwise recent) branch lengths (Pybus &

Harvey 2000). Likewise, variation in taxonomic practice across a phylogeny might lead to

spurious rate variation, particularly if different species concepts are used, or if some clades in the

phylogeny – but not others – have been subject to population genetic analysis or screens for

cryptic species diversity. Additionally, it has been shown that BAMM and other methods may

fail to infer accurate speciation rate dynamics if the phylogeny is in diversity decline – that is,

when extinction rates increase towards the present and ultimately exceed speciation rates

(Quental & Marshall 2011; Burin et al. 2018). A major, if obvious, caveat in the interpretation of

tip rates is that they apply to recent speciation rates and are necessarily limited with respect to

inferences about historical variation in speciation rate.

The greater the importance of the terminal branches in tip rate metrics, the greater the

impact these biases might have on tip rate estimates. On one end of the spectrum, metrics such as

λ_{TB} will be very sensitive to such biases as they rely exclusively on terminal branch lengths.

Such approaches may retain utility as summary statistics (e.g., Bromham, Hua & Cardillo 2016),

but we found that λ_{TB} exhibited the greatest amount of error in estimating speciation rates. On

the other end of the spectrum, a metric like λ_{ND} would be minimally impacted as this metric is

attempting to capture an average speciation rate over an entire root-to-tip path and does not

upweight the contribution of recent branch lengths. λ_{DR} is likely somewhere in the middle of this

spectrum, as it gives decreasing weight to branches towards the root. λ_{BAMM} is potentially

sensitive to such issues as well, although it may be possible to analytically correct for some

biases in the mechanics of the model itself (e.g., Rosindell et al. 2010; Etienne and Rosindell

2012).

Potential empirical biases aside, tip rates present a number of practical advantages in the

study of diversification rate variation. First, tip rates can be summarized and compared across

non-monophyletic assemblages of species (Jetz et al. 2012; Kennedy et al. 2016; Belmaker &

Jetz 2015; Oliveira et al. 2016; Quintero & Jetz 2018; Rabosky et al. 2018), making it possible to

summarize rate characteristics of entire communities or regional assemblages of species. Second,

estimation of rates at the present should be more robust to the influence of extinction, as

extinction can erase the history of lineage splitting deeper in the phylogeny (Nee et al. 1994;

Nee, May & Harvey 1994; Rabosky & Lovette 2008). Third, tip-specific rates can be paired with

species-specific trait values or geographic attributes in order to test potential trait- or geography-

dependent speciation rates (Freckleton, Phillimore & Pagel 2008; Jetz et al. 2012, Rabosky &

Goldberg 2017; Harvey & Rabosky 2017). Tip rates make it possible to relax strong assumptions

of rate homogeneity within character states, which are inherent to certain trait-dependent models,

including BiSSE and GeoSSE (Maddison, Midford & Otto 2007; Goldberg, Lancaster & Ree

2011; Ng & Smith 2014). Recent work has provided a conceptually rich and robust interpretive

framework for SSE models that does not assume rate-constancy within character states (Beaulieu

& O'Meara 2016; Caetano, O'Meara & Beaulieu 2018), but tip rates nonetheless can provide an

important check on results obtained with SSE models by providing a direct means of visualizing

the relationship between branch lengths and character states (Bromham, Hua & Cardillo 2016;

Hua & Bromham 2016; Harvey & Rabosky 2017). Visual inspection of data in this fashion has

the potential to reduce false positives by calling attention to potential outliers and other sources

of model inadequacy (Maddison & FitzJohn 2014; Rabosky & Goldberg 2015). A final

advantage for non-model-based tip rates, especially λ_{DR}, is that they can profitably be applied to

extremely large phylogenies: there are few computational limits to using them on phylogenies

with tens of thousands of tips or more, in contrast to formal model-based approaches for which

BAMM, HiSSE (Hidden State Speciation and Extinction; Beaulieu & O'Meara 2016), and other

methods are poorly suited. This computational efficiency also lends itself to more readily

accounting for phylogenetic uncertainty, because tip rate metrics can rapidly be computed across

posterior distributions of phylogenies and averaged (for example, see Jetz et al. 2012; Rabosky et

al. 2018).

In summary, tip rates offer a number of theoretical and practical advantages, particularly

in the study of associations between traits and diversification. We found that λ_{BAMM}

outperformed other metrics evaluated in this study and proved to be relatively accurate, even

under diversification scenarios that depart from the BAMM inference model. λ_{DR}

underperformed in comparison to λ_{BAMM}, but in many cases still did reasonably well, particularly

for small rate regimes. Despite our performance results, λ_{DR} is likely to remain a useful tool in

the study of trait- and geography-dependent diversification (Rabosky & Goldberg 2017; Harvey

& Rabosky 2017).

Acknowledgements

We thank Jonathan Mitchell for help with compiling the simulation datasets evaluated in this

study. We thank Francisco Henao Diaz, Matt Pennell, Brian O'Meara, the O'Meara Lab, and an

anonymous reviewer for comments on the manuscript. We also thank Michael Grundler, as well

as all other members of the Rabosky and Davis Rabosky labs at the University of Michigan for

thoughtful discussion. This work was supported in part by a University of Michigan Rackham

Predoctoral Fellowship (P.O.T.) and by a Fellowship from the David and Lucile Packard

Foundation (D.L.R.).

Data accessibility

All data and code necessary to repeat the analyses have been made available through the Dryad

digital data repository at XXXXXX.

Author contributions

P.O.T. and D.L.R. designed the project. P.O.T. assembled the datasets and performed all

analyses. P.O.T. and D.L.R. wrote the manuscript. Both authors contributed critically to

subsequent drafts and approved the final publication.

References

Alfaro, M.E., Faircloth, B.C., Harrington, R.C., Sorenson, L., Friedman, M., Thacker, C.E.,
Oliveros, C.H., Černý, D. & Near, T.J. (2018). Explosive diversification of marine fishes at
the Cretaceous–Palaeogene boundary. *Nature Ecology & Evolution*, **2**, 688–696.

Alfaro, M.E., Santini, F., Brock, C., Alamillo, H., Dornburg, A., Rabosky, D.L., Carnevale, G. &
Harmon, L.J. (2009). Nine exceptional radiations plus high turnover explain species
diversity in jawed vertebrates. *Proceedings of the National Academy of Sciences*, **106**,
13410–13414.

Barker, F.K., Burns, K.J., Klicka, J., Lanyon, S.M. & Lovette, I.J. (2013). Going to extremes:
contrasting rates of diversification in a recent radiation of new world passerine birds.
Systematic Biology, **62**, 298–320.

Beaulieu, J.M. & O'Meara, B.C. (2016). Detecting Hidden Diversification Shifts in Models of
Trait-Dependent Speciation and Extinction. *Systematic Biology*, **65**, 583–601.

Beaulieu, J.M. & O'Meara, B.C. (2015). Extinction can be estimated from moderately sized

molecular phylogenies. *Evolution*, **69**, 1036–1043.

Belmaker, J. & Jetz, W. (2015). Relative roles of ecological and energetic constraints,

diversification rates and region history on global species richness gradients. *Ecology Letters*,

18, 563–571.

Bromham, L., Hua, X. & Cardillo, M. (2016). Detecting Macroevolutionary Self-Destruction

from Phylogenies. *Systematic Biology*, **65**, 109–127.

Burin, G., Alencar, L.R.V., Chang, J., Alfaro, M.E. & Quental, T.B. (2018). How Well Can We

Estimate Diversity Dynamics for Clades in Diversity Decline? *Systematic Biology*, **105**, 1–

17.

Caetano, D.S., O'Meara, B.C. & Beaulieu, J.M. (2018). Hidden state models improve state-

dependent diversification approaches, including biogeographical models. *Evolution*.

Cai, T., Fjeldså, J., Wu, Y., Shao, S., Chen, Y., Quan, Q., Li, X., Song, G., Qu, Y., Qiao, G. &

Lei, F. (2017). What makes the Sino-Himalayan mountains the major diversity hotspots for

pheasants? *Journal of Biogeography*, **45**, 640–651.

Claramunt, S. (2010). Discovering exceptional diversifications at continental scales: the case of

the endemic families of Neotropical suboscine passerines. *Evolution*, **64**, 2004–2019.

Coyne, J.A. & Orr, H.A. (2004). *Speciation*. Sinauer, Cambridge.

Etienne, R.S. & Haegeman, B. (2012). A Conceptual and Statistical Framework for Adaptive

Radiations with a Key Role for Diversity Dependence. *The American Naturalist*, **180**, E75–

E89.

Etienne, R.S. & Rosindell, J. (2012). Prolonging the Past Counteracts the Pull of the Present:

Protracted Speciation Can Explain Observed Slowdowns in Diversification. *Systematic*

Biology, **61**, 204–213.

FitzJohn, R.G. (2010). Quantitative traits and diversification. *Systematic Biology*, **59**, 619–633.

FitzJohn, R.G., Maddison, W.P. & Otto, S.P. (2009). Estimating Trait-Dependent Speciation and

Extinction Rates from Incompletely Resolved Phylogenies. *Systematic Biology*, **58**, 595–

611.

Freckleton, R.P., Phillimore, A.B. & Pagel, M. (2008). Relating Traits to Diversification: A

Simple Test. *The American Naturalist*, **172**, 102–115.

Goldberg, E.E., Lancaster, L.T. & Ree, R.H. (2011). Phylogenetic inference of reciprocal effects

between geographic range evolution and diversification. *Systematic Biology*, **60**, 451–465.

Gomes, A.C.R., Sorenson, M.D. & Cardoso, G.C. (2016). Speciation is associated with changing

ornamentation rather than stronger sexual selection. *Evolution*, **70**, 2823–2838.

Harvey, M.G. & Rabosky, D.L. (2017). Continuous traits and speciation rates: Alternatives to

state-dependent diversification models (N. Cooper, Ed.). *Methods in Ecology and Evolution*,

12, 751–10.

Hua, X. & Bromham, L. (2016). PHYLOMETRICS: an R package for detecting

macroevolutionary patterns, using phylogenetic metrics and backward tree simulation.

Methods in Ecology and Evolution, **7**, 806–810.

Jablonski, D. (2008). Species selection: Theory and data. *Annual Review of Ecology, Evolution,*

and Systematics, **39**, 501–524.

Jetz, W., Thomas, G.H., Joy, J.B., Hartmann, K. & Mooers, A.O. (2012). The global diversity of

birds in space and time. *Nature*, **491**, 444–448.

Kay, K., Voelckel, C., Yang, J.Y., Hufford, K.M., Kaska, D.D. & Hodges, S.A. (2006). Floral

characters and species diversification. *Ecology and Evolution of Flowers* (eds L. Harder & S.

Barrett), pp. 311–325. Oxford.

Kennedy, J.D., Borregaard, M.K., Jønsson, K.A., Holt, B., Fjeldså, J. & Rahbek, C. (2016). Does

the colonization of new biogeographic regions influence the diversification and

accumulation of clade richness among the Corvides (Aves: Passeriformes)? *Evolution*, 38–

50.

Maddison, W.P. & FitzJohn, R.G. (2014). The Unsolved Challenge to Phylogenetic Correlation

Tests for Categorical Characters. *Systematic Biology*, **64**, 127–136.

Maddison, W.P., Midford, P.E. & Otto, S. (2007). Estimating a binary character's effect on

speciation and extinction. *Systematic Biology*, **56**, 701.

Marin, J. & Hedges, S.B. (2016). Time best explains global variation in species richness of

amphibians, birds and mammals. *Journal of Biogeography*, **43**, 1069–1079.

Meyer, A.L.S. & Wiens, J.J. (2017). Estimating diversification rates for higher taxa: BAMM can

give problematic estimates of rates and rate shifts. *Evolution*, **72**, 39–53.

Mitchell, J.S. & Rabosky, D.L. (2016). Bayesian model selection with BAMM: effects of the

model prior on the inferred number of diversification shifts. *Methods in Ecology and*

Evolution, **8**, 37–46.

Mitchell, J.S., Etienne, R.S. & Rabosky, D.L. (2018). Inferring Diversification Rate Variation

From Phylogenies With Fossils. *Systematic Biology*.

Mittelbach, G.G., Schemske, D.W., Cornell, H.V., Allen, A.P., Brown, J.M., Bush, M.B.,

Harrison, S.P., Hurlbert, A.H., Knowlton, N., Lessios, H.A., McCain, C.M., McCune, A.R.,

McDade, L.A., McPeek, M.A., Near, T.J., Price, T.D., Ricklefs, R.E., Roy, K., Sax, D.F.,

Schluter, D., Sobel, J.M. & Turelli, M. (2007). Evolution and the latitudinal diversity

gradient: speciation, extinction and biogeography. *Ecology Letters*, **10**, 315–331.

Moen, D. & Morlon, H. (2014). Why does diversification slow down? *Trends in Ecology and*

Evolution, **29**, 190–197.

Moore, B.R., Höhna, S., May, M.R., Rannala, B. & Huelsenbeck, J.P. (2016). Critically

evaluating the theory and performance of Bayesian analysis of macroevolutionary mixtures.

Proceedings of the National Academy of Sciences, **113**, 9569–9574.

Near, T.J., Dornburg, A., Kuhn, K.L., Eastman, J.T., Pennington, J.N., Patarnello, T., Zane, L.,

Fernandez, D.A. & Jones, C.D. (2012). Ancient climate change, antifreeze, and the

evolutionary diversification of Antarctic fishes. *Proceedings of the National Academy of*

Sciences, **109**, 3434–3439.

Nee, S., Holmes, E.C., May, R.M. & Harvey, P.H. (1994a). Extinction Rates Can Be Estimated

From Molecular Phylogenies. *Philosophical Transactions of the Royal Society B: Biological*

Sciences, **344**, 77–82.

Nee, S., May, R.M. & Harvey, P.H. (1994b). The reconstructed evolutionary process.

Philosophical Transactions of the Royal Society B: Biological Sciences, **344**, 305–311.

Nee, S., Mooers, A. & Harvey, P.H. (1992). Tempo and Mode of Evolution Revealed From

Molecular Phylogenies. *Proceedings of the National Academy of Sciences*, **89**, 8322–8326.

Ng, J. & Smith, S.D. (2014). How traits shape trees: new approaches for detecting character

state-dependent lineage diversification. *Journal of Evolutionary Biology*, **27**, 2035–2045.

Oliveira, B.F., Machac, A., Costa, G.C., Brooks, T.M., Davidson, A.D., Rondinini, C. &

Graham, C.H. (2016). Species and functional diversity accumulate differently in mammals.

Global Ecology and Biogeography, **25**, 1119–1130.

Pybus, O.G. & Harvey, P.H. (2000). Testing macro-evolutionary models using incomplete

molecular phylogenies. *Proceedings of the Royal Society of London. Series B: Biological*

Sciences, **267**, 2267–2272.

Quental, T.B. & Marshall, C.R. (2011). The Molecular Phylogenetic Signature of Clades in

Decline. *PLoS ONE*, **6**, e25780–9.

Quintero, I. & Jetz, W. (2018). Global elevational diversity and diversification of birds. *Nature*,

555, 246–250.

Rabosky, D.L. (2014). Automatic detection of key innovations, rate shifts, and diversity-

dependence on phylogenetic trees. *PLoS ONE*, **9**, e89543.

Rabosky, D.L. (2010). Extinction rates should not be estimated from molecular phylogenies.

Evolution, **64**, 1816–1824.

Rabosky, D.L. & Goldberg, E.E. (2017). FiSSE: A simple nonparametric test for the effects of a

binary character on lineage diversification rates. *Evolution*, **106**, 13410–11.

Rabosky, D.L. & Goldberg, E.E. (2015). Model Inadequacy and Mistaken Inferences of Trait-
Dependent Speciation. *Systematic Biology*, **64**, 340–355.

Rabosky, D.L. & Lovette, I.J. (2008). Explosive evolutionary radiations: decreasing speciation or
increasing extinction through time? *Evolution*, **62**, 1866–1875.

Rabosky, D.L., Chang, J., Title, P.O., Cowman, P.F., Sallan, L., Friedman, M., Kaschner, K.,
Garilao, C., Near, T.J., Coll, M. & Alfaro, M.E. (2018). An inverse latitudinal gradient in
speciation rate for marine fishes. *Nature*, **559**, 392–395.

Rabosky, D.L., Donnellan, S.C., Grundler, M. & Lovette, I.J. (2014). Analysis and Visualization
of Complex Macroevolutionary Dynamics: An Example from Australian Scincid Lizards.
Systematic Biology, **63**, 610–627.

Rabosky, D.L., Grundler, M., Anderson, C., Title, P.O., Shi, J.J., Brown, J.W., Huang, H. &
Larson, J.G. (2014). BAMMtools: an R package for the analysis of evolutionary dynamics
on phylogenetic trees. *Methods in Ecology and Evolution*, **5**, 701–707.

Rabosky, D.L., Mitchell, J.S. & Chang, J. (2017). Is BAMM Flawed? Theoretical and Practical
Concerns in the Analysis of Multi-Rate Diversification Models. *Systematic Biology*, **66**,
477–498.

Rabosky, D.L., Title, P.O. & Huang, H. (2015). Minimal effects of latitude on present-day
speciation rates in New World birds. *Proceedings of the Royal Society of London. Series B:*
Biological Sciences, **282**, 20142889.

Redding, D.W. & Mooers, A.O. (2006). Incorporating Evolutionary Measures into Conservation
Prioritization. *Conservation Biology*, **20**, 1670–1678.

Ricklefs, R.E. (2006). Global variation in the diversification rate of passerine birds. *Ecology*, **87**,

2468–2478.

Rohde, K. (1992). Latitudinal Gradients in Species Diversity: The Search for the Primary Cause.

Oikos, **65**, 514–527.

Rosindell, J., Cornell, S.J., Hubbell, S.P. & Etienne, R.S. (2010). Protracted speciation revitalizes

the neutral theory of biodiversity. *Ecology Letters*, **13**, 716–727.

Sanderson, M.J. & Donoghue, M.J. (1996). Reconstructing shifts in diversification rates on

phylogenetic trees. *Trends in Ecology and Evolution*, **11**, 15–20.

Schluter, D. (2016). Speciation, Ecological Opportunity, and Latitude. *The American Naturalist*,

187, 1–18.

Silvestro, D., Schnitzler, J. & Zizka, G. (2011). A Bayesian framework to estimate

diversification rates and their variation through time and space. *BMC Evolutionary Biology*,

11, 311.

Stadler, T. (2011). Simulating Trees with a Fixed Number of Extant Species. *Systematic Biology*,

60, 676–684.

Steel, M. & Mooers, A. (2010). The expected length of pendant and interior edges of a Yule tree.

Applied Mathematics Letters, **23**, 1315–1319.

Zink, R.M., Klicka, J. & Barber, B. (2004). The tempo of avian diversification during the

Quaternary. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **359**,

215–220.

 Figures/Captions

662 **Table 1.** Summary of simulated phylogenies used in this study.

simulation model	number of trees	tree size	regime number	source
single-regime, constant-rate birth-death	100	100	1	Mitchell & Rabosky 2016
single- and multi-regime, constant-rate birth-death	100	51-148	1-6	Moore et al. 2016
single- and multi-regime, constant-rate birth-death	400	10-4296	1-67	Rabosky, Mitchell & Chang 2017
multi-regime, constant-rate birth-death	20	939-3708	11	Meyer & Wiens 2017
single- and multi-regime, constant-rate birth-death	188	4-3955	1-73	Mitchell, Etienne & Rabosky 2018
single-regime, constant-rate birth-death, lambda uniform	1000	100	1	this study
single-regime, constant-rate birth-death, net diversification uniform	1000	100	1	this study
pure birth root regime, 1-4 discrete shifts to diversity-dependent regimes	1200	54-882	1-5	Rabosky 2014; Mitchell & Rabosky 2016
speciation rate evolves via diffusion process	1200	25-1208	1	Rabosky 2010; Beaulieu & O'Meara 2015; Rabosky 2016; this study

Figure 1. Mean absolute error in λ (top) and *r* (bottom) for three different tip rate metrics, across a range of relative extinction rates. For BAMM, the estimated speciation and net diversification rates are presented in the top and bottom panels, respectively. Absolute error of zero implies perfect accuracy. Inset plots show error in λ with truncated y-axis scale to facilitate comparison among metrics. All tip rate metrics track λ more accurately than they track *r*. See Figure S4 for λ_{TB}, which performed much worse than the other metrics.

Figure 2. True tip rates (λ_{TRUE}) in relation to estimated tip rates. Tip rates were compared

separately for different major categories of phylogeny simulations (rows) and are plotted

separately by inference method (columns). Plotting region is restricted to the 99th percentile of

true rates, but Spearman correlations between true and estimated rates (lower right of each figure

panel) are based on the full range of the data. Colors indicate the density of points in the scatter

plots. The horizontal gaps in λ_{ND} for diversity-dependent trees are an artefact of all trees having

the same crown age. λ_{BAMM} exhibited the strongest correlation with true rates for all simulation

categories.

Figure 3. Mean per-tip absolute error in speciation rates as a function of the magnitude of rate

heterogeneity in each simulated phylogeny. Results are presented separately for different

categories of rate variation (Table 1); left column shows estimates from a constant-rate birth-

death model for reference. The boxes and whiskers represent the $0.25 - 0.75$, and the $0.05 - 0.95$

quantile ranges, respectively. In some cases, λ_{ND} and λ_{DR} had more error than a simple CRBD

model with no variation in tip rates. λ_{BAMM} had the least amount of error across all amounts of

rate heterogeneity. See Figure S9 for λ_{TB}.

Figure 4. Performance of tip rate metrics as a function of regime size, including Pearson

correlation (a) and OLS regression slope (b) for mean rates with respect to λ_{TRUE}. λ_{DR} and λ_{BAMM}

outperform the other metrics when summarized in this fashion, although λ_{DR} tends to

overestimate the rate of speciation. The x-axis denotes the minimum regime size across which

performance was summarized. For example, x = 20 corresponds to the correlations and slopes

computed for all regimes with 20 or more tips; a value of x = 1 is the corresponding results for

all regimes. The OLS slope for λ_{TB} is not visible as it ranges between 7 and 9.

720 **Figure 5.** Mean per-regime absolute error in relation to true rate regime size, as binned into 10

721 size categories. The boxes and whiskers represent the 0.25 – 0.75, and the 0.05 – 0.95 quantile

722 ranges, respectively. Perfectly estimated rates have an error of zero. λ_{DR} and λ_{BAMM} exhibit the

723 least error when averaged by regimes, and λ_{DR} does slightly better for small clades (10-clade

724 median error 0.07 for λ_{DR}, and 0.08 for λ_{BAMM}).

Figure 6. Relationship between λ_{TRUE}, λ_{BAMM}, and λ_{DR} for a simulated phylogeny containing a

single rate shift (orange circle). Subplots to the right of the tree illustrate true and estimated rates

for each tip (left) and corresponding absolute error (right). Asterisks at the bottom denote mean

per-tip error in tip rate metrics. Mean per-tip error is relatively low and similar between λ_{DR} and

λ_{BAMM}, but the sample variance in λ_{DR} tip rates is high. In this example, the variance in absolute

per-tip error in λ_{DR} is 0.002 versus 0.0003 for λ_{BAMM}. On average, λ_{DR} tends to either

overestimate or underestimate rates relative to λ_{BAMM}, even if the mean per-tip error is relatively

low for both metrics.

Figure 7. Relationship between λ_{DR}, λ_{BAMM}, and λ_{DR} for a phylogeny simulated under an

"evolving rates" model, such that the speciation rate itself varies under a diffusion model. See

Figure 6 for additional details. Neither metric is particularly well equipped to infer the true rate

variation in this case. However λ_{BAMM}'s conservative estimates are still more accurate relative to

λ_{DR}, which is negatively impacted by high variance in tip rates. Here, variance in absolute per-tip

error in λ_{DR} is 0.012 versus 0.003 for λ_{BAMM}.

simulations for evaluating speciation rate

simulations for evaluating net diversification rate

Figure S1. Details of the phylogeny simulations designed to evaluate the performance of the four tip metrics in terms of speciation rate and net diversification rate. From the top row, it is clear that when λ is sampled uniformly with respect to ε, the distribution of r is not uniform: the mean, range and variance in r decrease dramatically as ε increases. The reverse is true for the distribution of λ when r is sampled uniformly with respect to ε (bottom row). Our simulation design ensures that λ and r are sampled from identical uniform distributions with respect to ε and ensures comparability of the resulting error estimates.

Figure S2. Root-mean-square error (RMSE) in λ (top) and *r* (bottom) for three different tip rate

metrics, across a range of relative extinction rates. For BAMM, the estimated speciation and net

diversification rates are presented in the top and bottom panels, respectively. Error of zero

implies perfect accuracy. Inset plots show error in λ with truncated y-axis scale to facilitate

comparison among metrics. All tip rate metrics track λ more accurately than they track *r*.

Figure S3. Log proportional accuracy in λ (top) and *r* (bottom) for different tip rate metrics,

across a range of relative extinction rates. For BAMM, the estimated net diversification rate is

presented. Proportional error of 0 implies perfect accuracy. Inset plots reveal greater detail in

error for λ to ease metric comparison. All tip metrics track λ much more accurately than they

track *r*, and λ$_{BAMM}$ does so with the least amount of error. See Figure S4 for λ$_{TB}$.

Figure S4. Mean absolute error in λ (top) and r (bottom) for λ_{TB}, with λ_{DR} and λ_{BAMM} on the

same scale for comparison. For BAMM, the estimated net diversification rate is presented. λ_{TB}

more accurately tracks λ and than r, but the amount of error is an order of magnitude greater than

for other metrics.

Figure S5. True tip rates (top row: λ_{TRUE}, bottom row: r_{TRUE}) in relation to λ_{TB}. Tip rates were compared separately for different major categories of phylogeny simulations (rows). Plotting region is restricted to the 99th percentile of true rates, but Spearman correlations between true and estimated rates (lower right of each figure panel) are based on the full range of the data. Colors indicate the density of points in the scatter plots. λ_{TB} is a largely unbiased but noisy measure of true speciation rate.

Figure S6. True tip rates (λ_{TRUE}) in relation to estimated tip rates from multi-regime

phylogenies, as inferred from the best two tip rate metrics, λ_{DR} and λ_{BAMM}. Data are separated by

source, to confirm that patterns described in the main text are not driven by any one simulation

study. Spearman's correlation is presented in the bottom right corner. Colors indicate the density

of points in the scatter plots. Regardless of the dataset, λ_{BAMM} performs noticeably better than

λ_{DR}.

Figure S7. True net diversification tip rates (r_{TRUE}) in relation to estimated tip rates. Tip rates

were compared separately for different major categories of phylogeny simulations (rows).

Plotting region is restricted to the 99th percentile of true rates, but Spearman correlations

between true and estimated rates (lower right of each figure panel) are based on the full range of

the data. Colors indicate the density of points in the scatter plots. The horizontal gaps in λ_{ND} for

diversity-dependent trees are an artefact of all trees having the same crown age. Relative

performance comparison aside, correlations with r_{TRUE} are lower than with λ_{TRUE} (Figure 2).

Figure S8. Root-mean-square error (RMSE) in speciation rates as a function of the magnitude of

rate heterogeneity in each simulated phylogeny. Results are presented separately for different

categories of rate variation (Table 1); left column shows estimates from a constant-rate birth-

death model for reference. The boxes and whiskers represent the $0.25 - 0.75$, and the $0.05 - 0.95$

quantile ranges, respectively. In some cases, λ_{ND} and λ_{DR} had more error than a simple CRBD

model with no variation in tip rates. λ_{BAMM} had the least amount of error across all amounts of

rate heterogeneity.

Figure S9. Mean per-tip absolute error in λ_{TB} as a function of the magnitude of rate

heterogeneity in each simulated phylogeny. λ_{DR} and λ_{BAMM} are included on the same scale for

comparison. Results are presented separately for different categories of rate variation (Table 1).

The boxes and whiskers represent the $0.25 - 0.75$, and the $0.05 - 0.95$ quantile ranges,

respectively. Error in λ_{TB} generally increases with increasing rate heterogeneity, and this error is

substantially greater than error in other tip rate metrics.

Figure S10. Performance of tip rate metrics as a function of regime size, including Pearson

correlation (a) and OLS regression slope (b) for mean rates with respect to r_{TRUE}. λ_{DR} and r_{BAMM}

outperform the other metrics when summarized in this fashion, although λ_{DR} overestimates the

rate of net diversification (more so than it overestimated λ_{TRUE}, Figure 4). The x-axis denotes the

minimum regime size across which performance was summarized. For example, x = 20

corresponds to the correlations and slopes computed for all regimes with 20 or more tips; a value

of x = 1 is the corresponding results for all regimes. The OLS slope for λ_{TB} is not visible as it

ranges between 7 and 9.