Apomixis-related genes identified from a coexpression network in *Paspalum*

notatum, a Neotropical grass

Apomixis-related genes identified from a coexpression network in *Paspalum notatum*

Fernanda A. de Oliveira[1], Bianca B. Z. Vigna[2], Carla C. da Silva[1], Alessandra P. Fávero[2],

Frederico de P. Matta[2], Ana L. S. Azevedo[3], Anete P. de Souza[1,4,*]

[1]Centro de Biologia Molecular e Engenharia Genética (CBMEG), Universidade Estadual de

Campinas (UNICAMP), Campinas, SP, Brazil

[2]Embrapa Pecuária Sudeste, São Carlos, SP, Brazil

[3]Embrapa Gado de Leite, Juiz de Fora, MG, Brazil

[4]Departmento de Biologia Vegetal, Instituto de Biologia, UNICAMP, Campinas, SP, Brazil

E-mail addresses:

f.ancelmo.o@gmail.com, bianca.vigna@embrapa.br, silvacbio@yahoo.com.br,

alessandra.favero@embrapa.br, frederico.matta@embrapa.br, ana.azevedo@embrapa.br,

anete@unicamp.br

[*]**Corresponding author**

E-mail: anete@unicamp.br (AP)

Telephone number: +55 19 3521-1132

Abstract

Apomixis is a highly desirable trait in modern agriculture, due to the maintenance of characteristics of the mother plant in the progeny. However, incorporating it into breeding programs requires a deeper knowledge of its regulatory mechanisms. *Paspalum notatum* is considered a good model for such studies because it exhibits both sexual and apomictic cytotypes, facilitating the performance of comparative approaches. Therefore, we used comparative transcriptomics between contrasting *P. notatum* cytotypes to identify novel candidate genes involved in the regulation of the expression of this phenotype. We assembled and characterized a transcriptome from leaf and inflorescence from apomictic tetraploids and sexual diploids/tetraploids of *P. notatum* accessions, and then assembled a coexpression network based on pairwise correlation between transcripts expression profiles. We identified genes exclusively expressed in each cytotype and differentially expressed genes between pairs of cytotypes. Gene ontology enrichment analyses were performed for the interpretation of data. We *de novo* assembled 114,306 of reference transcripts. 536 novel candidate genes for the control of apomixis were detected through statistical analyses of expression data, contains in this set, the interactions among genes potentially linked to the apomixis-controlling region, differentially expressed, several genes also already reported in the literature and their neighbors transcriptionally related in the coexpression network. The reference transcriptome obtained in this study represents a robust set of expression data for *P. notatum*. Additionally, novel candidate genes identified in this work represent a valuable resource for future grass breeding programs.

Author Summary

Clonal mode of reproduction by seeds is termed apomixis, which results from the failure of gamete formation (meiosis) and fertilization in the sexual female reproductive pathway. The manipulation of seeds production genetically identical to the mother plant bears great promise for agricultural applications, however clarification regarding gene interactions involved in reproductive process is needed. *Paspalum* is considered a model genus for the analysis of apomixis mechanisms. Here, we describe an overall analysis of the expression profiles of *Paspalum notatum* transcripts in response to changes in reproductive mode (sexual to apomictic), which allowed us to identify several candidate apomixis genes. Among these, we found genes potentially associated with the apomixis control region, in addition to genes already described in the literature for *Paspalum*, which highlights the representativeness of assembled transcriptome. For the first time in the literature, we explored the main biological processes involved in controlling the expression of apomictic reproduction based on co-regulatory networks of candidate apomixis genes.

Introduction

Apomixis is a mode of asexual reproduction through seeds in which the plants produced offspring is genetically identical to the female parent [1]. In apomixis, a non-reduced cell undergoes developmental pathways different from those of sexual cells. The multiple trajectories are classified as sporophytic or gametophytic according to the developmental origin of the cell from which the embryo is derived [2-4]. In sporophytes, the embryo develops from a somatic cell of the ovule through numerous mitotic divisions. In gametophytic apomixis, a chromosomally unreduced embryo sac can be formed either from the megaspore mother cell (diplospory) or from a nearby nucellar cell (apospory) without fertilization (parthenogenesis) through a process termed apomeiosis. Endosperm formation is required to produce a viable seed [5,6]. Apomixis is widely distributed among angiosperms, among which Poaceae represents the family with the largest number of apomictic genera, with the 47 apomictic species of *Paspalum* standing out in particular [7-9].

The polyploid nature of all apomicts provides a challenge for genetic and genomic analysis [10]. Polyploidization is known to cause immediate and extensive genomic changes, including sequence rearrangements and/or elimination, changes in DNA methylation, and the loss of balance in gene expression [11,12]. Apomixis is frequently correlated with polyploidy and might have arisen through deregulation of the sexual developmental pathway, due to the increase in number of genomic complements, through a mechanism regulated by both genetic and epigenetic components [13-16].

Paspalum notatum Flüggé, also known as bahiagrass, belongs to the genus *Paspalum,*

which exhibits numerous characteristics that make it a very interesting system for the study of

apomixis [8]. This species is considered an agamic complex that includes different ploidy levels

and reproductive modes in which diploid cytotypes ($2n=2\times=20$) are sexual and self-

incompatible, whereas polyploids ($3\times$, $4\times$, $5\times$, and $6\times$) are self-compatible pseudogamous

apomicts [17]. No $4\times$ sexual cytotypes have been found in nature, but they have been obtained

artificially [18,19]. The inheritance of apomixis in *Paspalum* is controlled by a single complex

dominant locus that confers apospory, i.e., epigenetically controlled parthenogenesis [20], with

the capacity to form endosperm with a maternal excess genome contribution ratio of 4:1

(maternal:paternal) [8]. The apomixis-controlling region (ACR) is small compared to other

apomictic systems [8], does not demonstrate recombination and conserves a relatively narrow

region that is linked to apomixis among the distinct species [21,22]. Comparative mapping of the

ACR has shown synteny with a portion of the long arm of chromosome 12 of rice in *P. simplex*,

P. malacophyllum and *P. procurrens* [21-23]. In *P. notatum* the ACR shows synteny with

regions of rice chromosomes 2 and 12, suggesting the presence of chromosomal rearrangements

[21,24-26].

Apomixis presents potential significance for agriculture, allowing the maintenance of

heterosis in hybrid progeny. Therefore, this mode of reproduction has become the subject of

exhaustive cytoembryological, cytogenetic and molecular analyses [10,25]. Some key genes

associated with the components of apomixis and the isolation of some sequence candidates have

already been discovered [6,12,14,27-41]. However, insights into the genetic mechanisms

underlying asexual reproduction in natural apomicts are still needed to develop a stable and

universal apomixis system to be applied in breeding programs [39]. Given the complexity of this

trait, an understanding of the genomic structure of the apomictic locus is likely to be an essential

prerequisite for manipulation of the sexual pathway in model plants or economically important

crops [15]. However, the suppression of recombination events around the ACR limits forward

genetic strategies to isolate the mechanism triggering apomixis by map-based [31]. In this

context, identify and validate genes that are differentially expressed in apomicts and to conduct

more meticulous investigations aiming to detect their effects in phenotype, have garnered

extreme interest in the study of apomixis [31,40].

RNA sequencing (RNA-seq) is the most effective method for simultaneously predicting

new transcripts and identifying differentially expressed genes in distinct tissues, genotypes,

abiotic conditions or developmental stages [42,43]. Conversely, considering the large amount of

data generated from RNA-seq, new approaches that efficiently extract meaningful associations

from highly multivariate datasets are needed [44]. The construction of coexpression networks

from gene expression data using pairwise correlation metrics provides us with valuable

information on alterations in biological systems in response to differential gene expression

patterns [44-46].

The objective of this study was to identify candidate genes possibly involved in the

regulation of the expression of apomixis in *P. notatum*. To reach this goal, we used RNA-seq

technology to obtain a reference transcriptome and comprehensively analyze gene expression

profiles in leaves and florets from apomictic and sexual genotypes with different ploidy levels in

P. notatum. Our study revealed several differentially expressed genes among the analyzed

genotypes, and a coexpression network analysis and Gene Ontology (GO) annotation of

transcripts allowed us to investigate the main biological processes of candidate genes potentially

linked to the ACR. These candidate genes may be used to further explore and clarify the

mechanisms regulating apomixis in forage grasses.

Results

Verification of the DNA content and mode of reproduction of *P.

notatum **accessions**

The cytoembryological analysis confirmed the expected mode of reproduction of the

accessions based on the literature (Tables 1 and 2). The 2C DNA content of all plants from

accessions BGP_22 and BGP_306 was half the 2C DNA content from all plants from BGP_30,

BGP_34, BGP_115, and BGP_216. Therefore, all evaluated plants had DNA contents

compatible with the described ploidy levels of their respective accessions in the literature (Tables

1 and 2).

Table 1. *Paspalum notatum* **Accessions Evaluated in this Study.**

BGP	FORMER	REPRODUCTION	ORIGIN	COLLECTOR*	CHROMOSOME

CODE	CODE	MODE			NUMBER
306		Sexual	Candói (PR), Brazil	VLiGu 14829	2n=2×=20 [47]
	BRA-024236				
22	BRA-006173	Sexual	Bagé (RS), Brazil	VGnMaBd 9607	2n=2×=20 [48,49]
216	BRA-019470	Sexual**	Corrientes, Argentina	Q 3664	2n=4×=40 [50]
30	BRA-006467	Apomictic	Alegrete (RS), Brazil	VMrFrLw 9747	2n=4×=40 [51]
34	BRA-006513	Apomictic	Uruguaiana (RS), Brazil	VMrFrLw9782	2n=4×=40 [51,52]
115	BRA-010006	Apomictic	Laguna (SC), Brazil	VDBdSv 10137	2n=4×=40 [51]

* Collectors: Bd, I. I. Boldrini; D, M. Dall'Agnol; Fr, J. M. O. Freitas; Gn, J. O. N. Gonçalves;

Gu, A. Guglieri; Li, L. Essi; Lw, H. M. Longhi-Wagner; Ma, M. C. Assis; Mr, C. O. C. Moraes;

Q, C. L. Quarín; Sv, G. P. da Silva; V, J. F. M. Valls; LAT, latitude (decimal degrees); LONG,

longitude (decimal degrees). ** Facultative apomictic with a high level of sexuality (> 70%),

according to Quarín et al. [53].

Table 2. DNA Contents and Modes of Reproduction of the Bahiagrass Accessions

Evaluated in this Study.

Accession	DNA content		Mode of

(BGP code)	Plants	2C (pg)	Δ2C among plants	Accession mean 2C (pg)	reproduction
30	BGP30_1[1]	3.2			
	BGP30_2[1]	3.1			
	BGP30_3	2.9	0.4	3.20	facultative apomixis
	BGP30_4	2.8			
	BGP30_5[1,3]	3.0			
34	BGP34_1[1,2,3]	2.9			
	BGP34_2	2.9			
	BGP34_3[1]	2.5	0.5	2.80	facultative apomixis
	BGP34_4	2.7			
	BGP34_5[1]	3.0			
115	BGP115_1	2.7			
	BGP115_2[1,3]	2.9			
	BGP115_3[1]	2.6	0.3	2.68	obligate apomixis
	BGP115_4	2.6			
	BGP115_5[1]	2.7			
216	BGP216_1[1,3]	2.6			
	BGP216_2[1]	2.7			
	BGP216_3	2.9	0.6	2.82	sexual
	BGP216_4	3.2			
	BGP216_5[1]	2.7			
22	BGP22_1	1.5	0.3	1.52	sexual
	BGP22_2	1.7			

	BGP22_3[1,2,3]	1.4			
	BGP22_4[1]	1.4			
	BGP22_5[1]	1.5			
	BGP306_1[1,3]	1.5			
	BGP306_2	1.5			
306	BGP306_3	1.4	0.1	1.44	sexual
	BGP306_4[1]	1.4			
	BGP306_5[1]	1.4			

[1]Indicates plants with foliar RNA evaluated; [2]Indicates plants with inflorescence RNA evaluated;

[3]Indicates plants with DNA evaluated.

RNA-seq analysis and *de novo* transcriptome assembly

After removing barcodes sequences and low-quality and contaminant reads, 313,198,097 high-quality 72-bp paired-ends reads from floret and leaf tissues were used to assemble a *P. notatum* reference transcriptome. A total of 203,808 transcripts were assembled, out of which a set of 114,306 non-redundant transcripts were filtered (56.08% of all transcripts) (Table 3). Final transcripts in the reference transcriptome had an average length of 750.51 bp with an N50 of 906 bp and GC percentage of 47.60%. According to the length distribution of non-redundant transcripts, we obtained 21,585 (18.88%) transcripts that were longer than 1 kb, a size range that commonly confers a high annotation rate. More than half of the total annotated transcripts were > 500 bp in length (Fig S1).

Table 3. Statistics for the Assembled Transcriptome of *Paspalum notatum*.

	All assembled transcripts	Non-redundant transcripts
Total	203,808	114,306
Total assembled bases	219,683,637	85,787,466
Average length (bp)	1,077.90	750.51
N50 length (bp)	1,599	906
Percent GC (%)	46.76	47.60

The Bowtie aligner mapped 94.39% of sequenced reads onto the assembled transcripts.

73.78% mapped to the non-redundant transcripts set. Out of these, 82.97% and 83.48% of the

sequencing reads belonged to sexual and apomictic samples, respectively, showing similar

representation within the final transcriptome. The BUSCO analysis included 956 conserved

single-copy plant orthologs; the *P. notatum* transcriptome showed high assembly completeness,

with 664 (69%) complete sequences (532 as a single copy and 132 as multiple copies) and 169

(17%) fragmented sequences. One hundred and twenty-three (12%) BUSCO plant orthologs

were not identified in the reference *P. notatum* transcriptome.

Transcriptome annotation

Among the 114,306 unigenes, 51,939 (45.44%) were similar to proteins from the NCBI

Nr database, 34,750 (30.40%) to proteins from the UniProtKB/Swiss-Prot, and 54,568 (47.74%)

to proteins from Phytozome.

The GO database was then used to retrieve the GO identifiers for all possible transcripts.

Overall, 21,537 (41.47%) transcripts were assigned to 4,614 GO terms in three main categories:

2,510 biological process, 1,486 molecular function, and 618 cellular components. The KEGG

annotation was possible for 9,221 annotated transcripts belonging to 129 pathways. Among

these, the most represented were Purine Metabolism Pathway (1,079 members), followed by

Metabolism of Thiamine (907 members) and Starch and Sucrose Metabolism (336 members).

Overall, 33,873 transcripts remained without hits after searching against all protein databases.

We then performed a search for the presence of open reading frames (ORFs), which were found

in 15,107 transcripts, 2,347 of which were complete. These sequences that showed ORFs without

annotation require further investigation since they may represent genes that have not yet been

described and possibly new genes that may be unique to *P. notatum*.

Expression levels of genes and identification of DEGs

The sequenced reads were realigned against the 114,306 transcripts of the reference

transcriptome to estimate the expression level of the transcripts in each genotype. Most identified

transcripts had expression levels of FPKM ≤ 100 (110,621 transcripts), whereas only 49

transcripts had high expression levels (FPKM ≥ 400), in these, most of them involved in

photosynthesis process, house-keeping genes and response to stress. The remaining transcripts

were not mapped and thus, their expression levels could not be determined. A comparison of the

110,656 transcripts that showed different levels of expression in the two tissues of the sequenced

samples revealed genes that were expressed in the genotypes of only one phenotypic class and

that were not expressed in the others (Fig 1), which were considered "unique" transcripts. Thus,

19,304 unique transcripts were found exclusively in the 4×APO group, 2,173 in the 2×SEX

group, and 217 in the 4×SEX group. All transcripts expressed in florets were also expressed in

leaves; there were no unique transcripts expressed in the florets from either diploid sexual or

tetraploid apomictic samples (Fig 1).

Fig 1. Venn diagram showing the distribution of *P. notatum* transcripts. Expression level of

transcripts from two tissues (floret and leaf) and each phenotypic class (sexual diploid, sexual

tetraploid, and apomictic tetraploid).

For pairwise differential expression analyses of all three phenotypic classes, we

considered only leaf samples because they expressed all representative transcripts from the

transcriptome assembly and had three clones for each genotype, maximizing the accuracy and

reliability of the differential gene expression analysis. However, within the same phenotypic

class, different expression patterns were observed among genotypes, mainly in 4×APO samples

(Fig S2). Thus, to reduce the effects of the differences in expression caused by genotype, we

performed a two-step pipeline. In the first step, we identified 72,318 transcripts that were equally

expressed among all 4×APO genotypes and 47,069 equally expressed transcripts in the 2×SEX

genotypes. This initial analysis was not necessary for the 4×SEX because only one genotype was

analyzed. From these, we selected 28,969 transcripts that were expressed in all three phenotypic

classes. In the second step of the differential expression analysis, pairwise comparisons of 28,969

transcripts allowed the identification of 2,072 DEGs between 2×SEX and 4×APO, which

included 772 and 1,173 overexpressed transcripts, respectively. There were 1,173 DEGs between

2×SEX and 4×SEX, including 308 and 798 overexpressed transcripts, respectively, and 1,317

were identified as DEGs between 4×APO and 4×SEX, 618 and 661 of which were

overexpressed, respectively. Next, we inferred the enriched functions for the identified DEGs

that were over- or underexpressed (Figs S5-S7).

Among the identified DEGs, we isolated a set of 510 transcripts that were consistently

differentially expressed between apomictic and sexual samples independently of the ploidy level

(diploid or tetraploid). We identified GO terms enriched in this set of transcripts, which belonged

to 26 biological processes (BP), 13 molecular functions (MF), and three cellular components

(CC) (Fig 2). We considered these transcripts as potentially involved in the determination of the

reproductive mode.

Fig 2. Enriched ontology terms of 510 transcripts differentially expressed in the *P. notatum*

transcriptome. Transcripts consistently differentially expressed between apomictic and sexual

samples.

qRT-PCR validation of RNA-seq data

We selected a subset of 18 DEGs detected in-silico between apomictic and sexual

samples to confirm the expression profiles through an independent technique, qRT-PCR. Five

equally expressed transcripts detected in-silico showed desirable expression stability and

coefficient of variance in qRT-PCR reactions (Table S1), out of which one was selected to be an

internal control for qRT-PCR verification of 2×SEX vs. 4×APO analyses (01RefGen-*Pnot*) and

one for the verification of 4×APO vs. 4×SEX analyses (04RefGen-*Pnot*).

Out of 18 primer pairs designed for target DEGs, eight (Table S2) were used for qRT-

PCR validation of RNA-seq data. The qRT-PCR analyses revealed consistent results compared

to those detected through RNA-seq (Figs S3 and S4), emphasizing the power of the latter

technique. Two primers pairs were not evaluated in qRT-PCR but showed interesting results in

terms of DNA amplification. The first was designed from a transcript that was exclusively

expressed in apomict samples in RNA-seq analyses (i.e., showing a zero expression value in the

sexual samples). This transcript did not amplify using genomic DNA from sexual samples as

PCR templates. For the second primer pair, although genomic DNA amplification occurred in all

samples, when complementary DNA (cDNA) was used as PCR template, only apomictic

samples presented amplicons. The expression level of this transcript estimated from RNA-seq of

the same samples was very low in 2×SEX samples. These results suggest that the first transcript,

expressed only in apomictic cytotypes, is exclusive to the genomes of these samples, whereas the

second transcript was poorly expressed or silenced in sexual samples.

Detailed search for potential apomixis genes

We detected 1,612 transcripts that showed high similarity to the ACR in the *P. notatum*

reference transcriptome. Out of these, 1,356 transcripts aligned to corresponded rice

chromosome 2, and 256 aligned to rice chromosome 12. Curiously, 40 of these transcripts were

part of the set identified as unique to apomictic samples, nine were unique to sexual diploids, and

six were unique to sexual tetraploid samples expression. Moreover, considering the analyses of

differential expression, 16 putative ACR transcripts were differentially expressed in 2×SEX vs.

4×APO, 20 DEGs in 4×APO vs. 4×SEX, and 17 DEGs in 2×SEX vs. 4×SEX. All these findings,

especially this set of 108 unique or differentially expressed transcripts, represent a valuable set of

genes that deserve to be carefully investigated to determine their roles in apomixis and determine

whether they are effectively genetically within the ACR in *P. notatum*.

The BLAST results in Table S3 show the similarity scores of the *P. notatum* transcripts to

previously *Paspalum* sequences associated to the apomictic mode of reproduction, revealing that

our transcriptome contains genes already found to be involved in asexual reproductive

development in *Paspalum*.

270 Transcripts coexpression network

271 We identified 879,481 connections among 53,262 transcripts from RNA-seq data (Fig 3).

272 Transcripts were grouped into 642 clusters according to their correlated pattern of expression

273 level. In the network, the unique transcripts from each phenotypic class tended to form

274 coexpression modules (Fig 3B-D). Despite this subdivision, the relevant biological relationships

275 of these transcripts with all remaining transcripts can be recovered in a fully integrated network.

Fig 3. Coexpression network for *Paspalum notatum*. (A) Coexpression network of all analyzed transcripts; the more centralized genes in light gray are those common to all genotypes analyzed with non-differential expression. Node color denotes the differentially expressed transcripts between the phenotypic classes: purple (2×SEX vs. 4×APO), yellow (2×SEX vs. 4×SEX), and dark green (4×APO vs. 4×SEX). The networks of exclusively expressed transcripts for each phenotypic class are highlighted: (B) transcripts unique to sexual diploids (blue); (C) transcripts unique to sexual tetraploids (green); and (D) transcripts unique to apomictic tetraploids (red). Edges denote interaction strength. Circular nodes represent transcripts.

We identified a direct correlation among the 108 differentially expressed and/or unique putative ACR genes and their first neighbors in the gene expression network. Thus, we retrieved a sub-network composed of 536 strongly correlated transcripts (Fig 4). We used GO enrichment analysis to summarize the main putative functions of this set of transcripts, which included 43 biological processes (BP), 31 molecular functions (MF) and 17 cellular components (CC) (Fig. 5).

Fig 4. Gene expression sub-network for 536 *Paspalum notatum* transcripts. The 108 differentially expressed and/or unique putative ACR transcripts are presented in this sub-network, along with their first neighbors. The color patterns are the same as those used in the complete coexpression network in Fig 3.

Fig 5. Enriched biological processes in the 536 transcripts in the gene expression sub-network.

Discussion

We constructed a *P. notatum* reference transcriptome using the next-generation sequencing approaches, RNA-seq, to investigate the gene expression changes associated with apomictic and sexual reproduction. RNA-seq enabled the assembly of a non-redundant transcriptome, containing 114,306 transcripts from florets and leaves from six distinct genotypes with different ploidy levels and reproductive modes. Robust metrics indicated that the transcriptome was a quality assembly with a high degree of integrity, which further expands the bahiagrass transcriptome database.

Even though we did not sequence inflorescences samples at different developmental stages (premeiosis, late premeiosis/meiosis, postmeiosis and anthesis) to search for candidate apomixis-linked genes, differential expression analyses between leaf samples from sexual and apomictic cytotypes allowed us to identify DEGs that may be representative of the mode of reproduction and not dependent of the level of ploidy. We isolated expression patterns of phenotypic classes, aimed both at removing transcripts whose expression could be related to the effect of genotype and at identifying transcripts expressed in all genotypes within the same phenotypic class.

GO classification of the assembled transcripts into 4,614 known terms, was used to perform functional enrichment analyses of candidate apomixis genes. Based on a set of 510 DEGs detected between apomictic and sexual, among the enriched GO terms, we emphasize "positive regulation of miRNA metabolic process (GO:2000630)" and "regulation of primary

miRNA processing (GO:2000636)", both of which either activate or increase the frequency, rate,

or extent of miRNA production. miRNA, in turn, is directly related to "DNA metabolic process

(GO:0006259)", which decreases the rate of gene expression (negative regulation of gene

expression - GO:0010629) through an epigenetic RNA-based gene silencing process (RNA-

directed DNA methylation - GO:0080188). Five transcripts were mainly involved in these

processes and showed higher expression levels in apomict samples in comparison to sexual

samples; these transcripts correspond to a pseudoARR-B transcription factor; pre-mRNA

splicing factor, putative (*Ricinus communis*); hypothetical protein SORBIDRAFT_05g016770;

uncharacterized protein, LOC100501330 (*Zea mays*); and an uncharacterized protein

LOC103641690. The GO analysis reveals the significant enrichment of terms associated with the

regulation of gene expression by epigenetic silencing among transcripts showing higher

expression levels among apomictic samples in comparison to sexual samples of the bahiagrass.

This result is consistent with the growing body of evidence that suggests that apomixis arises

from deregulation of the sexual pathway, where epigenetic mechanisms play a significant role in

at least some elements of apomictic development [8,15,54].

These DEGs represent a set of candidate genes that, together with the 19,304 transcripts

exclusively expressed in apomictic samples, deserve further investigation. In apospory, gene

expression occurs at specific stages of apomictic development such as apomeiosis,

parthenogenesis, and endosperm development. Increased or decreased expression of some genes

during these specific stages may hinder the analysis of differential expression between sexual

and apomictic genotypes [40,55]. Nonetheless, we believe that the identification of DEGs as

performed herein, using selected transcripts with detectable expression pattern among all

genotypes of all phenotypic classes, may have minimized this influence. At the same time, our

approach increased the potential for discovery of candidate genes involved in apomictic process

and not only in a single step. Thus, future reverse genetics experiments based on qRT-PCR and

in situ hybridization could be useful to identify the role of specific genes in the whole apomixis

process.

Functionally related genes tend to be transcriptionally coordinated [56,57]. Therefore, the

construction of transcripts coexpression network provided a powerful resource, for the

identification of transcripts that are coexpressed with unique and differentially expressed genes,

despite showing undetectable differences in expression between samples. The sub-network

containing 536 candidate coexpressed transcripts associated with the expression of apomixis is

an example of how we can recover the relevant biological relationships of genes of interest using

the transcriptome sequencing approaches. By starting on a smaller scale of transcripts that are

DEGs and/or unique genes and possibly integrated into the ACR and combining the information

by adding their nearest neighbors, we can obtain a broader view of the processes involved in the

regulation of candidate genes. Significantly enriched BP were mainly associated with plant

reproduction, for instance, "male gamete generation (GO: 0048232)"; "sister chromatid

segregation (GO: 0000819)"; "meiosis I (GO: 0007127)"; "resolution of meiotic recombination

intermediates (GO: 0000712)"; "regulation of cytokinesis (GO: 0032465)"; "mitotic cytokinesis

(GO: 0000281)"; "spindle assembly (GO: 0051225)"; "mitotic spindle organization (GO:

0007052)"; and "mitotic DNA integrity checkpoint (GO: 0044774)". In addition, the sub-

network was enriched in epigenetic processes such as "macromolecule methylation (GO:

0043414)"; "histone-serine phosphorylation (GO: 0035404)"; "regulation of protein phosphatase

type 2A activity (GO: 0034047)"; and "negative regulation of MAP kinase activity (GO:

0043407)". Transcripts from this sub-network were associated with reproductive processes and

the regulation of epigenetic changes by modulating histones. In particular, MAP kinase activity

was negatively regulated. The mitogen-activated protein 3-kinase (MAP3K) gene family has

been identified as differentially expressed in apomictic and sexual flowers of *P. notatum* [31]

and in flowers of *P. simplex* [14], and it might play a role in parthenogenetic development of the

embryo in both species [8]. Recently, an essential gene to the formation of unreduced embryo

sacs in *P. notatum* was identified from the characterization of a MAP3K retrieved in previous

transcriptomic surveys [58].

The size of the ACR in *Paspalum* is relatively small compared with other apomictic

systems [8], but the lack of recombination makes it difficult to study this region by map-based

cloning [31]. Thus, the discovery of DEGs that are potentially located within this region is

extremely valuable for understanding the complex regulatory network of gene–gene interaction.

These DEGs could also be used for future manipulation of the apomixis trait, which has

outstanding importance in agricultural biotechnology.

In the genus *Paspalum,* the first approach for understanding the molecular basis of

apomixis was applied by Pessino et al. [27]. This approach led to the identification of three small

sequences that are highly expressed during early megagametophyte development in apomictic

plants through differential display experiments in the inflorescences of sexual and aposporous *P.*

notatum. Here, we recovered the reported sequences as a single representative transcript (Table

S3) similar to a kinesin motor protein, involved in the biological process of microtubule-based

movement. The kinesin motor protein has also been reported by other authors as differentially

expressed between apomictic and sexual plants [29,31,59]. Interestingly, the transcript assembled

in this work also showed differential expression levels between apomictic and sexual samples.

Since then, several other candidate genes have been reported using *P. notatum* and *P. simplex*

species [14,27,30,31] and more recently, the first approach using RNA-seq was published for

bahiagrass based on expression profiling of apomictic and sexual flowers [40]. Some of these

candidate genes had been investigated in more detail [35,37-39]. However, many questions

remain unanswered, and further research is needed to define the relationships between the

structure, the position, and the function of the known apomixis-linked genes [8]. Furthermore,

based on the set of genes selected through differential expression patterns, we recovered

previously published sequences in our *P. notatum* transcriptome.

One of the candidate genes previously reported in the literature is n20gap-1, a lorelei-like

P. notatum gene [31,35] encoding a GPI-anchored protein that supposedly plays a role in the

final stages of the apomixis developmental cascade. Laspina et al. [31] previously reported this

sequence as linked to the chromosomal locus governing apospory at a genetic distance of 22 cM.

Here, one transcript similar to this gene present in the set of transcripts potentially integrated

within the ACR, corresponding to rice chromosome 2. Additionally, we found transcripts,

including some that were differentially expressed, that aligned with *P. notatum* sequences

characterized by Podio et al. [38], corresponding to *Somatic Embryogenesis Receptor-Like*

Kinase (*SERK*), in addition to other numerous transcripts with the same annotation. The

candidate SERK-like genes play crucial roles in somatic embryogenesis in angiosperms and have

been reported as related to apomixis. Albertini et al. [59] analyzed two members of this protein

family, namely *PpSERK1* and *PpSERK2*, and found that *PpSERK1* expression levels was high

during premeiosis and decreased during the meiosis and post-meiotic stages, whereas *PpSERK2*

expression was high from premeiosis to anthesis in the nucellar cells of aposporous genotypes.

The authors proposed that the expression pattern of *PpSERK* in *Poa pratensis* was compatible

with its role in the specification of aposporous initials. In *P. notatum*, the expression of two

different members of the *SERK* family (*PnSERK1* and *PnSERK2*) was observed, and *PnSERK2*

displayed a spatial expression pattern similar to that reported for the *PpSERKs*, which are

expressed in nucellar cells at meiosis in the apomictic genotype [38]. We also identified

transcripts that showed similarity to other interesting candidate genes related to apospory, such

as the PnTgs1-like gene that encodes a trimethylguanosine synthase-like protein, which plays a

fundamental role in nucellar cell fate, as its diminished expression is correlated with initiation of

the apomictic pathway in plants [37]. Additionally, two transcripts were similar to the sequence

of *Ps*ORC3, which seems to play an active role in mechanisms repressing sexuality in apomictic

P. simplex, acting in the development of apomictic seeds, which deviate from the canonical 2

(maternal):1 (paternal) genome ratio [39]. The recovery of these previously published sequences

demonstrates the representative and informative nature of assembled transcriptome. Indeed, this

outcome reaffirms that high-throughput sequencing technology enhances our understanding of

global RNA expression.

In this study, we applied RNA-seq technology and bioinformatics methods to assemble a

useful transcriptome and identify differences between apomictic and sexual reproduction. Our

results reveal DEGs, genes exclusively expressed in apomicts or sexuals, of which are in

potential association with the ACR genomic region, including the apo locus. The validation of

genes from this set of candidates may enable valuable insights to the understanding of apospory.

Moreover, these genes may be considered to screen for molecular markers linked to apomixis in

P. notatum, which will be crucial to boost breeding programs for apomictic forage grasses.

Materials and methods

Plant materials collections and RNA extraction

The six accessions of *P. notatum* used in this study belong to the Germplasm Bank of

Paspalum, maintained by Embrapa Pecuária Sudeste, São Carlos, SP, Brazil (22° 01'S and

47°54' W; 856 m above sea level) (Table 1). The accessions were chosen based on their origins

(different ecotypes), genetic dissimilarity [60] and ploidy levels and reproduction modes.

Tetraploid *P. notatum* is apomictic, whereas the diploid is sexual. The Q3664 (BGP 216)

accession is a hybrid from a cross between a sexual colchicine-induced tetraploid (PT-2) and a

white-stigma apomictic tetraploid of *P. notatum* [50]. The Q3664 is characterized as a facultative

apomictic accession with a high level of sexuality ($> 70\%$) [53].

For this study, five clones (biological replicates) of each accession were planted in 8 L

pots, filled with 1:1 soil/vermiculite and grown in a greenhouse under the same environmental

conditions. The climate is humid subtropical (according to the Köppen-Geiger classification

system), with annual average low and high temperatures of 15.3 and 27.0°C, respectively, and

total rainfall of 1,422.8 mm, occurring mainly during the spring and summer seasons [61].

Young leaf samples from three clones of each accession were collected during summer

(December). One clone each of the accessions BGP34 (apomictic tetraploid) and BGP22 (sexual

diploid) presented inflorescences, which were also collected. All leaf and floret samples were

immediately placed in liquid nitrogen and subsequently stored at -80°C until RNA extraction.

Total RNA was isolated according to Oliveira et al. [62]. RNA integrity was assessed in a 1%

agarose denaturing gel and quantified using a NanoVue Plus spectrophotometer (GE Healthcare

Life Sciences, Little Chalfont, UK).

Verification of the mode of reproduction of *P. notatum* accessions

through cytoembryological analysis

The mode of reproduction of the plants was confirmed using the embryo sac clarification

method proposed by Young et al. [63], with minor modifications. Inflorescences at anthesis

(when the embryo sac is fully developed) were collected and fixed in FAA (95% ethanol,

distilled water, glacial acetic acid, formalin 40%, 40:13:3:3 v/v) for 24 h at room temperature.

Afterward, the FAA was replaced with 70% ethanol and the samples were stored at 4°C. Ovaries

were extracted by dissection under a stereoscopic microscope and stored in 70% ethanol. The

embryo sacs were clarified by replacing 70% alcohol with the following series of solutions: 85%

ethanol; absolute ethanol; ethanol:methyl salicylate (1:1); ethanol:methyl salicylate (1:3); 100%

methyl salicylate (two times). The samples remained in each solution for 24 h. Observations

were carried out with an Axiophot microscope (Carl Zeiss, Jena, Germany) using the differential

interference contrast (DIC) microscopy technique. A total of 100 embryo sacs per accession were

evaluated.

Verification of the ploidy level of *P. notatum* accessions through flow

cytometry

Flow cytometry analyses were performed to confirm the exact ploidy level of each plant

used in the experiment and results were compared to the literature data. Approximately 5 mg of

young leaf tissue from each plant was used (Table 1). *Pisum sativum* cv. Ctirad samples

(2C=9.09 pg), were used as an internal control [64,65]. Samples were triturated in a petri dish

containing 800 µL of LB01 buffer (0.45425 g TRIS, 0.186125 g NaEDTA, 0.0435 g-spermine,

0.29225 g NaCl, 1.491 g KCl, and 250 μL of Triton X-100 in 250 mL of distilled water, 7.5 pH,

0.11% v/v of β-mercaptoethanol) to obtain a nuclear suspension [66]. The nuclear suspension

was filtered through a mesh of 40 μm and incubated at room temperature for 5 min, followed by

the addition of 25 μL of propidium iodide and 25 μL of RNase. For each sample, at least 10,000

nuclei were analyzed. Samples were analyzed with a FACSCalibur flow cytometer (Becton

Dickinson, New Jersey, USA). Histograms were obtained in Cell Quest software and analyzed in

Flowing Software (available at http://www.flowingsoftware.com). Only histograms with both

peaks (sample and standard) of approximately the same height were considered. The nuclear

DNA index (pg) of the plants was estimated based on the value of the G1 peak as an internal

reference. We calculated the mean value of C per biological replicate and accession and the

difference between the highest and lowest values (Δ) observed in replicates of each accession.

RNA-seq library construction, Illumina sequencing, and data

quality control

The cDNA libraries were constructed from each RNA sample (18 leaf and two floret

libraries) using specific barcodes and the TruSeq RNA Sample Preparation Kit (Illumina Inc.,

San Diego, CA, USA) following manufacturer's instructions. Library quality was confirmed

using the Agilent 2100 Bioanalyzer (Agilent Technologies, Palo Alto, CA, USA) and quantified

via quantitative real-time PCR (qPCR) (Illumina protocol SY-930-10-10). Clustering was

conducted using a TruSeq Paired-End Cluster Kit on a cBot (Illumina Inc., San Diego, CA,

USA). Paired-end sequencing was performed on the Illumina Genome Analyzer IIx Platform

with TruSeq SBS 36-Cycle kits (Illumina, San Diego, CA, USA) following manufacturer's

specifications. RNA-seq was performed with floret libraries of the accessions BGP22

(diploid/sexual) and BGP34 (tetraploid/apomictic) in two separate lanes, without biological

replicates. The remaining 18 leaf libraries from six different accessions (Table 1), each with

three independent biological replicates (clones), were distributed randomly in the flow cell, with

three libraries per lane.

Raw data was converted to FastQ files containing 72-bp reads. Quality control was

performed using the NGS QC Toolkit v2.3.3 [67]. Initially, high-quality reads (Phred quality

score $\geq$ 20 in at least 75% of bases) and reads with more than 60 bases were selected.

Subsequently, reads were trimmed at the 3' end for the removal of barcodes. All reads were

deposited in the NCBI Short Read Archive (SRA) under accession number SRP150615.

Transcriptome assembly and completeness assessment

High-quality reads from leaves and florets were *de novo* assembled into a reference

transcriptome for *P. notatum*, performed with Trinity program v2.0.2 [68] using default settings.

Contigs redundancy was minimized through the selection of the first Butterfly sequence from

each Chrysalis component, which is considered the most representative contig. To assess the

sequenced reads-support of the assembled contigs, the high-quality short reads were mapped

back to the transcriptome using the Bowtie2 sequence aligner v2.2.5 [69]. The Benchmarking

Universal Single-Copy Orthologs (BUSCO) tool, an approach for the assessment of conserved

orthologs among plant species in a set of sequences [70], was used to estimate the completeness

of the transcriptome assembly.

Transcriptome annotation

All transcripts were aligned to proteins from the the NCBI non-redundant (Nr) database,

from the UniProtKB/Swiss-Prot and from the grass protein database from Phytozome v9.0, using

the BLASTX algorithm with an e-value cutoff of 1e-06. Gene Ontology (GO) [71] terms were

retrieved for transcripts showing similarity to sequences from NCBI Nr or UniProtKB databases

using Blast2GO software [72]. We used REVIGO [73] to summarize and visualize GO terms

sets. Pathways were assigned to metabolic pathways from the Kyoto Encyclopedia of Genes and

Genomes (KEGG) database [74].

Estimation of transcript expression levels

The expression levels of the transcripts were estimated using the FPKM method

(expected number of fragments per kilobase of transcript sequence per million base pairs

sequenced). Read counts for each transcript from each sample (Table 1) was obtained using

Bowtie v2-2.1.0 and normalized using the RSEM software (RNA-seq by expectation

maximization) [75]. A Venn diagram for the visualization of the amount of exclusive and shared

transcripts among samples was created using the online platform available at

http://bioinformatics.psb.ugent.be/webtools/Venn/. Principal component analysis (PCA) was

used to assess expression patterns of sequenced genotypes.

Differential expression analysis

EBSeq [76] was used to identify differentially expressed genes (DEGs) using FPKM

values, at a false discovery rate (FDR) ≤ 0.05. Transcripts with a log2 fold change (FC) ≥ 1.5 in

transcript abundance were regarded as overexpressed. GO enrichment analysis was carried out

using R software (version 3.3.1) and the 'goseq' 1.24.0 Bioconductor package [77]. P-values

were subjected to Bonferroni correction, and we considered adjusted p-values ≤ 0.05 as enriched.

Resulting enriched GO terms were summarized using REVIGO [73].

Genotypes were grouped according to their phenotypic classes as sexual diploids

(2×SEX), sexual tetraploids (4×SEX) and apomictic tetraploids (4×APO) to be compared in

pairwise differential expression analyses. A two-step pipeline was used for the identification of

the DEGs (Fig 6). In the first step, the objective was to select a list of equally expressed

transcripts among genotypes belonging to the same phenotypic class. This procedure allowed the

detection of gene expression patterns related with characteristics that were shared among

genotypes, independently of physiological or developmental particularities. Thus, EBSeq was

used to estimate the pairwise posterior probabilities of a transcript being equally express between

genotypes: sexual diploids (accessions: BGP22 and BGP306), and apomictic tetraploids

(accessions: BGP30, BGP34 and BGP115). Transcripts that presented PPEE ≥ 0.95 were kept for

the second step of the analysis. The second step consisted of another round of pairwise

differential expression analyses between phenotypic classes: (i) 2×SEX vs. 4×APO; (ii) 2×SEX

vs. 4×SEX; (iii) 4×APO vs. 4×SEX.

Fig 6. Workflow of differential expression analyses used in this study. We realized pairwise

differential expression analyses of all three phenotypic classes (apomictic tetraploid "4×APO",

sexual diploid "2×SEX" and sexual tetraploid "4×SEX"). Each phenotypic class had different

genotype with three clones. In the first step, the objective was to select a list of equally expressed

transcripts among genotypes belonging to the same phenotypic class. EBSeq was used to

estimate the pairwise posterior probabilities of a transcript being equally express (PPEE)

between genotypes of the same phenotypic class: apomictic tetraploid (A×B; A×C; B×C) and

sexual diploid (D×E). Transcripts that presented PPEE ≥ 0.95 were kept for the second step of

the analysis. The second step consisted of another round of pairwise differential expression

analyses between phenotypic classes: (i) 4×APO vs. 2×SEX; (ii) 4×APO vs. 4×SEX; (iii) 2×SEX

vs. 4×SEX. The objective was to select a list of transcripts that presented pairwise posterior

probabilities of being differentially expressed (PPDE ≥ 0.95). Additionally, we identified a list of

transcripts that were consistently differentially expressed between sexual and apomictic samples,

independently of the ploidy level (diploid or tetraploid).

Quantitative reverse transcription PCR (qRT-PCR) validation of

differential expression results

To verify the reliability and accuracy of transcriptome data and differential expression

analyses pipeline, 18 DEGs were randomly selected for quantification through reverse

transcription PCR (qRT-PCR). Eight transcripts showing similar expression patterns based on

FPKM values were used as internal reaction controls. Primer sets for controls and targets were

designed using Primer3Plus software [78]. All primer pairs were initially tested in regular PCR

reactions using genomic DNA as a template from the same genotypes and clones used in RNA-

seq. Only primers that amplified the genomic DNA of all genotypes were used in the following

qRT-PCR amplification efficiency tests. Primer pairs with an amplification efficiency of 90-

110% and R^2> 0.99 were used for relative expression analyses.

Total RNA (500 ng) was used for cDNA synthesis using the QuantiTect Reverse

Transcription Kit (Qiagen Inc., Chatsworth, CA, USA). qRT-PCR were performed using a

CFX384 Real-Time PCR Detection System with iTaq Universal SYBR Green Supermix (Bio-

Rad Laboratories Inc., Hercules, CA, USA), according to the manufacturer's instructions. The

reaction conditions were 95°C (10 min) and then 40 cycles of 95°C (30 s) and 60°C (1 min). All

experiments were performed using technical triplicate, and no-template controls were included.

The specificity of amplicons was confirmed through the analysis of the melting curve. The

baseline and Cq (quantification cycle) values were automatically determined, and the expression

values were estimated using the ΔΔCt method implemented by the CFX Manager 2.1 software

(Bio-Rad Laboratories, Inc., USA). Reference genes were selected according to their expression

stability (M < 0.5) and coefficient of variance (CV < 0.25) among samples. Mann-Whitney U-

tests were performed to estimate statistical significance of distribution differences between

samples.

Search for putative apomixis genes

To perform a detailed comparative analysis of *P. notatum* transcripts obtained in this

study with their syntenic counterparts on rice chromosomes that are conservatively linked to

apomixis in *Paspalum* species [8], we aligned all *P. notatum* transcripts to rice transcripts using

the BLASTN tool and selected those that presented putative homology to genes in the

highlighted area of rice chromosomes 2 and 12. Then, we selected only *P. notatum* transcripts

present in the ACR, which were delimited by a set of molecular markers completely linked to the

apospory locus in this species. Thus, the region encompassed the C1069 and C996 markers for

rice chromosome 12 and between C560 and C932 markers for chromosome 2 [8,21,24-26,79].

We also compared the *P. notatum* reference transcriptome with sequences reported for the genus

Paspalum, potentially associated with apomixis. The sequences were used as queries for

BLASTN search in the 114,306 transcripts database, with an e-value cutoff of 1e-05.

Construction of a transcripts coexpression networks

To generate coexpression network, we used expression values in FPKM of all assembled

transcripts from floret and leaf tissues. Transcripts showing null values for most of the replicates

were excluded to reduce noise and eliminate residuals in the analysis. We calculated an all-

versus-all coexpression network matrix using the Pearson correlation coefficient cutoff of ≥ 0.8.

The highest reciprocal rank (HRR) method proposed by Mutwil et al. [80] was used to select

only the strongest edges, considering an HRR limit of 30. The Heuristic Cluster Chiseling

Algorithm (HCCA) was applied to partition the networks into manageable clusters, with three-

step node vicinity networks (NVN) [80]. The interactive coexpression network was visualized

using the Cytoscape 3.4.0 software [81].

Acknowledgments

623   We are grateful to Maria Augusta C. Horta and Natalia F. Murad for their help in

624   generating the coexpression networks and Mariana V. Cruz for critically reading the manuscript.

References

1. Asker S, Jerling L. Apomixis in plants. Boca Raton, FL: CRC Press; 1992.

2. Nogler GA. Gametophytic apomixis. In: Johri BM, editor. Embryology of angiosperms.
   Berlin: Springer; 1984. pp. 475-518.

630
631

3. Koltunow AM. Apomixis: embryo sacs and embryos formed without meiosis or
   fertilization in ovules. Plant Cell. 1993;5: 1425-1437.

632
633

4. Conner JA, Goel S, Gunawan G, Cordonnier-Pratt MM, Johnson VE, Liang C, et al.
   Sequence analysis of bacterial artificial chromosome clones from the apospory-specific
   genomic region of *Pennisetum* and *Cenchrus*. Plant Physiol. 2008;147: 1396-1411.

634
635
636

5. Conner JA, Gunawan G, Ozias-Akins P. Recombination within the apospory specific
   genomic region leads to the uncoupling of apomixis components in *Cenchrus ciliaris*.
   Planta. 2013;238: 51-63.

637
638
639

6. Conner JA, Mookkan M, Huo H, Chae K, Ozias-Akins P. A parthenogenesis gene of
   apomict origin elicits embryo formation from unfertilized eggs in a sexual plant. Proc
   Natl Acad Sci U S A. 2015;112: 11205-11210.

640
641
642

7. Carman JG. Asynchronous expression of duplicate genes in angiosperms may cause
   apomixis, bispory, tetraspory, and polyembryony. Biol J Linn Soc. 1997;61: 51-94.

643
644

8. Ortiz JPA, Quarin CL, Pessino SC, Acuña C, Martínez EJ, Espinoza F, et al. Harnessing
   apomictic reproduction in grasses: what we have learned from *Paspalum*. Ann Bot.
   2013;112: 767-787.

645
646
647

9. Hojsgaard D, Klatt S, Baier R, Carman JG, Hörandl E. Taxonomy and biogeography of

apomixis in *Angiosperms* and associated biodiversity characteristics. CRC Crit Rev Plant

Sci. 2014;33: 414-427.

10. Ozias-Akins P. Apomixis: developmental characteristics and genetics. CRC Crit Rev

Plant Sci. 2006;25: 199-214.

11. Adams KL, Wendel JF. Polyploidy and genome evolution in plants. Curr Opin Plant

Biol. 2005;8: 135-141.

12. Cervigni GDL, Paniego N, Pessino S, Selva JP, Díaz M, Spangenberg G, et al. Gene

expression in diplosporous and sexual *Eragrostis curvula* genotypes with differing ploidy

levels. Plant Mol Biol. 2008;67: 11-23.

13. Spillane C, Curtis MD, Grossniklaus U. Apomixis technology development—virgin

births in farmers' fields? Nat Biotechnol. 2004;22: 687-691.

14. Polegri L, Calderini O, Arcioni S, Pupilli F. Specific expression of apomixis-linked

alleles revealed by comparative transcriptomic analysis of sexual and apomictic

Paspalum simplex Morong flowers. J Exp Bot. 2010;61: 1869-1883.

15. Calderini O, Donnison I, Polegri L, Panara F, Thomas A, Arcioni S, et al. Partial isolation

of the genomic region linked with apomixis in *Paspalum simplex*. Mol Breeding.

2011;28: 265-276.

16. Selva JP, Pessino SC, Meier MS, Echenique VC. Identification of candidate genes related

to polyploidy and/or apomixis in *Eragrostis curvula*. Am J Plant Sci. 2012;03: 403-416.

17. Burton GW. The method of reproduction in common bahia grass, *Paspalum notatum1*. J

Am Soc Agron. 1948;40: 443-452.

18. Martínez EJ, Urbani MH, Quarin CL, Ortiz JPA. Inheritance of apospory in bahiagrass,

Paspalum notatum. Hereditas. 2001;135: 19-25.

19. Quarin CL, Urbani MH, Blount AR, Martinez EJ, Hack CM, Burton GW, et al.

Registration of Q4188 and Q4205, sexual tetraploid germplasm lines of bahiagrass. Crop

Sci. 2003;43: 745-746.

20. Podio M, Cáceres ME, Samoluk SS, Seijo JG, Pessino SC, Ortiz JPA, et al. A

methylation status analysis of the apomixis-specific region in *Paspalum* spp. suggests an

epigenetic control of parthenogenesis. J Exp Bot. 2014;65: 6411-6424.

21. Pupilli F, Martinez EJ, Busti A, Calderini O, Quarin CL, Arcioni S. Comparative

mapping reveals partial conservation of synteny at the apomixis locus in *Paspalum spp*.

Mol Genet Genomics. 2004;270: 539-548.

22. Hojsgaard DH, Martínez EJ, Acuña CA, Quarin CL, Pupilli F. A molecular map of the

apomixis-control locus in *Paspalum procurrens* and its comparative analysis with other

species of *Paspalum.* Theor Appl Genet. 2011;123: 959-971.

23. Pupilli F, Labombarda P, Caceres ME, Quarín CL, Arcioni S. The chromosome segment

related to apomixis in *Paspalum simplex* is homoeologous to the telomeric region of the

long arm of rice chromosome 12. Mol Breed. 2001;8: 53-61.

24. Martínez EJ, Hopp HE, Stein J, Ortiz JPA, Quarin CL. Genetic characterization of

apospory in tetraploid *Paspalum notatum* based on the identification of linked molecular

markers. Mol Breed. 2003;12: 319-327.

25. Stein J, Pessino SC, Martínez EJ, Rodriguez MP, Siena LA, Quarin CL, et al. A genetic

map of tetraploid *Paspalum notatum* Flügge (bahiagrass) based on single-dose molecular

markers. Mol Breed. 2007;20: 153-166.

26. Podio M, Rodríguez MP, Felitti S, Stein J, Martínez EJ, Siena LA, et al. Sequence

characterization, *in silico* mapping and cytosine methylation analysis of markers linked to

apospory in *Paspalum notatum*. Genet Mol Biol. 2012;35: 827-837.

27. Pessino SC, Espinoza F, Martínez EJ, Ortiz JPA, Valle EM, Quarín CL. Isolation of

cDNA clones differentially expressed in flowers of apomictic and sexual *Paspalum*

notatum. Hereditas. 2001;134: 35-42.

28. Rodrigues JCM, Cabral GB, Dusi DMA, de Mello LV, Rigden DJ, Carneiro VTC.

Identification of differentially expressed cDNA sequences in ovaries of sexual and

apomictic plants of *Brachiaria brizantha*. Plant Mol Biol. 2003;53: 745-757.

29. Albertini E, Marconi G, Barcaccia G, Raggi L, Falcinelli M. Isolation of candidate genes

for apomixis in *Poa pratensis* L. Plant Mol Biol. 2004;56: 879-894.

30. Calderini O, Chang SB, de Jong H, Busti A, Paolocci F, Arcioni S, et al. Molecular

cytogenetics and DNA sequence analysis of an apomixis-linked BAC in *Paspalum*

simplex reveal a non pericentromere location and partial microcolinearity with rice. Theor

Appl Genet. 2006;112: 1179-1191.

31. Laspina NV, Vega T, Seijo JG, González AM, Martelotto LG, Stein J, et al. Gene

expression analysis at the onset of aposporous apomixis in *Paspalum notatum*. Plant Mol

Biol. 2008;67: 615-628.

32. Sharbel TF, Voigt ML, Corral JM, Thiel T, Varshney A, Kumlehn J, et al. Molecular

signatures of apomictic and sexual ovules in the *Boechera holboellii* complex. Plant J.

2009;58: 870-882.

33. Yamada-Akiyama H, Akiyama Y, Ebina M, Xu Q, Tsuruta S, Yazaki J, et al. Analysis of

expressed sequence tags in apomictic guineagrass (*Panicum* maximum). J Plant Physiol.

2009;166: 750-761.

34. Sharbel TF, Voigt ML, Corral JM, Galla G, Kumlehn J, Klukas C, et al. Apomictic and

sexual ovules of *Boechera* display heterochronic global gene expression patterns. Plant

Cell. 2010;22: 655-671.

35. Felitti SA, Seijo JG, González AM, Podio M, Laspina NV, Siena L, et al. Expression of

lorelei-like genes in aposporous and sexual *Paspalum notatum* plants. Plant Mol Biol.

2011;77: 337-354.

36. Okada T, Hu Y, Tucker MR, Taylor JM, Johnson SD, Spriggs A, et al. Enlarging cells

initiating apomixis in *Hieracium praealtum* transition to an embryo sac program prior to

entering mitosis. Plant Physiol. 2013;163: 216-231.

37. Siena LA, Ortiz JPA, Leblanc O, Pessino S. PNTGS1-like expression during

reproductive development supports a role for RNA methyltransferases in the aposporous

pathway. BMC Plant Biol. 2014;14: 297.

38. Podio M, Felitti SA, Siena LA, Delgado L, Mancini M, Seijo JG, et al. Characterization

and expression analysis of SOMATIC EMBRYOGENESIS RECEPTOR KINASE

(SERK) genes in sexual and apomictic *Paspalum notatum*. Plant Mol Biol. 2014;84: 479-

495.

39. Siena LA, Ortiz JPA, Calderini O, Paolocci F, Cáceres ME, Kaushal P, et al. An

apomixis-linkedORC3-like pseudogene is associated with silencing of its functional

homolog in apomictic *Paspalum simplex*. J Exp Bot. 2016;67: 1965-1978.

40. Ortiz JPA, Revale S, Siena LA, Podio M, Delgado L, Stein J, et al. A reference floral

transcriptome of sexual and apomictic *Paspalum notatum.* BMC Genomics. 2017;18:

318.

41. Bocchini M, Galla G, Pupilli F, Bellucci M, Barcaccia G, Ortiz JPA, et al. The vesicle

trafficking regulator PN_SCD1 is demethylated and overexpressed in florets of apomictic

Paspalum notatum genotypes. Sci Rep. 2018;8: 3030.

42. Huang W, Guo Y, Du W, Zhang X, Li A, Miao X. Global transcriptome analysis

identifies differentially expressed genes related to lipid metabolism in Wagyu and

Holstein cattle. Sci Rep. 2017;7: 5278.

43. Liang R, Han B, Li Q, Yuan Y, Li J, Sun D. Using RNA sequencing to identify putative

competing endogenous RNAs (ceRNAs) potentially regulating fat metabolism in bovine

liver. Sci Rep. 2017;7: 6396.

44. Miao X, Luo Q, Zhao H, Qin X. Co-expression analysis and identification of fecundity-

related long non-coding RNAs in sheep ovaries. Sci Rep. 2016;6: 39398.

45. de la Fuente A. From 'differential expression' to 'differential networking' – identification

of dysfunctional regulatory networks in diseases. Trends Genet. 2010;26: 326-333.

46. Gov E, Arga KY. Interactive cooperation and hierarchical operation of microRNA and

transcription factor crosstalk in human transcriptional regulatory network. IET Syst Biol.

2016;10: 219-228.

47. Dahmer N, Schifino-Wittmann MT, Dall'Agnol M, Castro BD. Cytogenetic data for

Paspalum notatum Flügge accessions. Sci Agric. 2008;65: 381-388.

48. Pagliarini MS, Takayama SY, de Freitas PM, Carraro LR, Adamowski EV, Silva N, et al.

Failure of cytokinesis and 2n gamete formation in Brazilian accessions of *Paspalum*.

Euphytica. 1999;108: 129-135.

49. Pagliarini MS, Carraro LR, Freitas PM, Adamowski EV, Batista LAR, Valls JFM.

Cytogenetic characterization of Brazilian *Paspalum* accessions. Hereditas. 2001;135: 27-

34.

50. Forbes I, Burton GW. Cytology of Diploids, natural and induced tetraploids, and intra-

species hybrids of bahiagrass, *Paspalum notatum* Flugge. Crop Sci. 1961;1: 402-406.

51. Pozzobon MT, Valls JFM. Chromosome number in germplasm accessions of *Paspalum*

notatum (Gramineae). Braz J Genet. 1997;20: 29-34.

52. Adamowski EDV, Pagliarini MS, Bonato ABM, Batista LAR, Valls JFM. Chromosome

numbers and meiotic behavior of some *Paspalum* accessions. Genet Mol Biol. 2005;28:

773-780.

53. Quarin CL, Burson BL, Burton GW. Cytology of intra- and interspecific hybrids between

two cytotypes of *Paspalum notatum* and *P. cromyorrhizon*. Bot Gaz. 1984;145: 420-426.

54. Brukhin V. Molecular and genetic regulation of apomixis. Russ J Genet. 2017;53: 943-

964.

55. Schmidt A, Schmid MW, Klostermeier UC, Qi W, Guthörl D, Sailer C, et al.

Apomictic and sexual germline development differ with respect to cell cycle,

transcriptional, hormonal and epigenetic regulation. PLoS Genet. 2014;10: e1004476.

56. Stuart JM, Segal E, Koller D, Kim SK. A gene-coexpression network for global

discovery of conserved genetic modules. Science. 2003;302: 249-255.

57. Persson S, Wei H, Milne J, Page GP, Somerville CR. Identification of genes required for

cellulose synthesis by regression analysis of public microarray data sets. Proc Natl Acad

Sci U S A. 2005;102: 8633-8638.

58. Mancini M, Permingeat HR, Colono C, Siena LA, Pupilli F, Azzaro C, et al. The

MAP3K-coding QUI-GON JINN (QGJ) gene is essential to the formation of unreduced

embryo sacs in *Paspalum*. Front. Plant Sci. 2018. doi: 10.3389/fpls.2018.01547.

59. Albertini E, Marconi G, Reale L, Barcaccia G, Porceddu A, Ferranti F, et al. *SERK* and

APOSTART. Candidate genes for apomixis in *Poa pratensis*. Plant Physiol. 2005;138:

2185-2199.

60. Cidade FW, Vigna BBZ, de Souza FHD, Valls JFM, Dall'Agnol M, Zucchi MI, et al.

Genetic variation in polyploid forage grass: assessing the molecular genetic variability in

the *Paspalum* genus. BMC Genet. 2013;14: 50.

61. CEPAGRI. (2010) Centro de Pesquisas Meteorológicas e Climáticas Aplicadas à

Agricultura. Available from: http://www.cpa.unicamp.br/o-cepagri.html Cited 17 August

2016.

62. Oliveira RR, Viana AJC, Reátegui ACE, Vincentz MGA. Short communication an

efficient method for simultaneous extraction of high-quality RNA and DNA from various

plant tissues. Genet Mol Res. 2015;14: 18828-18838.

63. Young BA, Sherwood RT, Bashaw EC. Cleared-pistil and thick-sectioning techniques for

detecting aposporous apomixis in grasses. Can J Bot. 1979;57: 1668-1672.

64. Dolezel J, Sgorbati S, Lucretti S. Comparison of three DNA fluorochromes for flow

cytometric estimation of nuclear DNA content in plants. Physiol Plant. 1992;85: 625-631.

65. Greilhuber J, Temsch EM, Loureiro JCM. Nuclear DNA content measurements. In:

Doležel J, Greilhuber J, Suda. J, editors. Flow cytometry with plant cells: analysis of

genes, chromosomes and genomes. Weinheim, Germany: Wiley-VCH Verlag GmbH and

Co.; 2007. pp. 67-101.

66. Dolezel J. Flow cytometry, its application and potential for plant breeding. In: Lelley T,

editor. Current topics in plant cytogenetics related to plant improvement. Vienna: WUV-

Universitätsverlag; 1997. pp. 80-90.

67. Patel RK, Jain M. NGS QC toolkit: a toolkit for quality control of next generation

sequencing data. PLoS One. 2012;7: e30619.

68. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-length

transcriptome assembly from RNA-Seq data without a reference genome. Nat Biotechnol.

2011;29: 644-652.

69. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment

of short DNA sequences to the human genome. Genome Biol. 2009;10: R25.

70. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO:

assessing genome assembly and annotation completeness with single-copy orthologs.

Bioinformatics. 2015;31: 3210-3212.

71. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology:

tool for the unification of biology. Nat Genet. 2000;25: 25-29.

72. Conesa A, Gotz S, Garcia-Gomez JM, Terol J, Talon M, Robles M. Blast2GO: a

universal tool for annotation, visualization and analysis in functional genomics research.

Bioinformatics. 2005;21: 3674-3676.

73. Supek F, Bošnjak M, Škunca N, Šmuc T. REVIGO summarizes and visualizes long lists

of gene ontology terms. PLoS One. 2011;6: e21800.

74. Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids

Res. 2000;28: 27-30.

75. Li B, Dewey CN. RSEM: accurate transcript quantification from RNA-Seq data with or

without a reference genome. BMC Bioinformatics. 2011;12: 323.

76. Leng N, Dawson JA, Thomson JA, Ruotti V, Rissman AI, Smits BMG, et al. EBSeq: an

empirical Bayes hierarchical model for inference in RNA-seq experiments.

Bioinformatics. 2013;29: 1035-1043.

77. Young MD, Wakefield MJ, Smyth GK, Oshlack A. Gene ontology analysis for RNA-seq:

accounting for selection bias. Genome Biol. 2010;11: R14.

78. Untergasser A, Nijveen H, Rao X, Bisseling T, Geurts R, Leunissen JAM. Primer3plus,

an enhanced web interface to Primer3. Nucleic Acids Res. 2007;35: W71-W74.

79. Labombarda P, Busti A, Caceres ME, Pupilli F, Arcioni S. An AFLP marker tightly

linked to apomixis reveals hemizygosity in a portion of the apomixis-controlling locus in

Paspalum simplex. Genome. 2002;45: 513-519.

80. Mutwil M, Usadel B, Schutte M, Loraine A, Ebenhoh O, Persson S. Assembly of an

interactive correlation network for the *Arabidopsis* genome using a Novel Heuristic

Clustering algorithm. Plant Physiol. 2010;152: 29-43.

81. Shannon P. Cytoscape: a software environment for integrated models of biomolecular

interaction networks. Genome Res. 2003;13: 2498-2504.

Supporting Information

S1 Fig. Length distribution of the number of non-redundant transcripts successful annotated. Comparison of the all non-redundant transcripts and the number of transcripts annotated in the NCBI non-redundant protein database by size range.

S2 Fig. Principal component analysis (PCA) according to the FPKM values of *P. notatum* transcriptome. PCA analysis of leaf and floret transcriptomes for all genotypes and clones used in RNA-seq.

S3 Fig. Histograms of gene expression obtained by qRT-PCR. qRT-PCR validation showing the relative expression patterns of 4 genes that are differentially expressed between tetraploid apomictic "4×APO" (red box) and diploid sexual "2×SEX" (blue box). *P < 0.05; **P < 0.01; ***P < 0.001: statistically significant differences in gene expression between phenotypic classes compared using the Mann-Whitney U-test.

S4 Fig. Histograms of gene expression obtained by qRT-PCR. qRT-PCR validation showing the relative expression patterns of 4 genes that are differentially expressed between tetraploid apomictic "4×APO" (red box) and tetraploid sexual "4×SEX" (green box). *P < 0.05; **P < 0.01; ***P < 0.001: statistically significant differences in gene expression between the phenotypic classes compared using the Mann-Whitney U-test.

S5 Fig. Functional classification of enriched overexpressed DEGs in the 2×SEX vs. the 4×APO comparison. Gene Ontology biological process (blue boxes), Gene Ontology cellular component (yellow boxes), and Gene Ontology molecular function (orange boxes). a) Categories

enriched in overexpressed transcripts in 2×SEX and b) categories enriched in overexpressed

transcripts in 4×APO.

S6 Fig. Functional classification of enriched overexpressed DEGs in the 2×SEX vs. 4×SEX

comparison. Gene Ontology biological process (blue boxes), Gene Ontology cellular component

(yellow boxes), and Gene Ontology molecular function (orange boxes). a) Categories enriched in

overexpressed transcripts in 2×SEX and b) categories enriched in overexpressed transcripts in

4×SEX.

S7 Fig. Functional classification of enriched overexpressed DEGs in the 4×APO vs. 4×SEX

comparison. Gene Ontology biological process (blue boxes), Gene Ontology cellular component

(yellow boxes), and Gene Ontology molecular function (orange boxes). a) categories enriched in

overexpressed transcripts in 4×APO and b) categories enriched in overexpressed transcripts in

4×SEX.

S1 Table. Primer sequences and amplicons of the candidate reference genes evaluated in

this study.

S2 Table. Quantitative RT-PCR primer sequences.

S3 Table. BLAST search results for sequences of *P. notatum* from the literature against the

assembled transcriptome.