1

5 **The Evolution of SlyA/RovA Transcription Factors**

6 **From Repressors to Counter-Silencers in *Enterobacteriaceae***

10 W. Ryan Will[1], Peter Brzovic[2], Isolde Le Trong[3], Ronald E. Stenkamp[2,3],
11 Matthew B. Lawrenz[4], William W. Navarre[5*], Kara Main-Hester[5], Virginia L. Miller[6],
12 Stephen J. Libby[1#†] and Ferric C. Fang[1,5†]
13

Departments of [1]Laboratory Medicine, [2]Biochemistry, [3]Biological Structure and [5]Microbiology,
University of Washington, Seattle, WA 98195

[4]Department of Microbiology and Immunology and the Center for Predictive Medicine for
Biodefense and Emerging Infectious Diseases, University of Louisville School of Medicine,
Louisville, KY 40202

[6]Departments of Microbiology and Immunology, and Genetics, University of North Carolina
School of Medicine, Chapel Hill, NC 27599

*Present address, University of Toronto, ON, Canada M5S 1A8

#Corresponding author. E-mail address: fcfang@uw.edu

†Contributed equally to this paper

Abstract

Gene duplication and subsequent evolutionary divergence have allowed conserved proteins

to develop unique roles. The MarR family of transcription factors (TFs) has undergone extensive

duplication and diversification in bacteria, where they act as environmentally-responsive

repressors of genes encoding efflux pumps that confer resistance to xenobiotics, including many

antimicrobial agents.   We have performed structural, functional, and genetic analyses of

representative members of the SlyA/RovA lineage of MarR TFs, which retain their ancestral

functions, including repression of their own expression and that of divergently-transcribed multi-

drug efflux pumps, as well as allosteric inhibition by aromatic carboxylate compounds. In contrast

to other MarR family members, SlyA and RovA have acquired the ability to counter-silence

horizontally-acquired genes, which has greatly facilitated the evolution of *Enterobacteriaceae* by

horizontal gene transfer. SlyA/RovA TFs in different species have independently evolved novel

regulatory circuits to provide the enhanced levels of expression required for their new role. This

demonstrates the ability of TFs to acquire new functions as a result of evolutionary divergence of

cis-regulatory sequences.

Introduction

47   As organisms adapt to new or changing environments, their regulatory networks must

evolve to ensure that individual genes are appropriately expressed in response to environmental

signals (1). An important mechanism for the evolution of conserved essential proteins, including

transcription factors (TFs), is gene duplication, which allows the subsequent diversification of

gene and protein families and the development of new functions (2). More than 50% of bacterial

genes are believed to have descended from original duplication events (3-5), providing a broad

foundation from which bacteria can evolve complex and adaptive traits.

54   The MarR family is an ancient family of TFs, predating the divergence of archaea and

bacteria (6). It has undergone extensive gene duplication events, with recent estimates suggesting

that bacteria encode an average of seven MarR TFs per genome (7). MarR TFs typically function

as environmentally-responsive repressors of genes encoding efflux pumps that export xenobiotics,

including many antimicrobial agents. The prototypical MarR protein of *Escherichia coli* represses

a single operon, *marRAB*, which encodes a transcriptional activator (MarA) required for the

expression of the AcrAB efflux pump, which in turn confers resistance to β-lactams, quinolones,

and tetracyclines (8-10). MarR is allosterically regulated by many small molecules, in particular

small aromatic carboxylate compounds such as salicylate, which induce a structural change that

reduces the affinity of MarR for DNA (9, 11-13) and derepresses the expression of its cognate

promoters. MarR TFs are defined by the presence of a winged helix-turn-helix (wHTH) DNA-

binding domain (14). Dimerization allows MarR TFs to recognize palindromic sequences via the

α4 recognition helix, which makes sequence-specific contacts with the major groove, while the

wing makes sequence-independent contacts via the minor groove (15).

The SlyA lineage of MarR TFs originally resulted from a MarR family gene duplication.

SlyA has been best characterized in *Salmonella enterica* serovar Typhimurium, where it serves

primarily to upregulate virulence genes (16-18). Although this contrasts with the classical

repressive role of MarR TFs, work in our and other laboratories has demonstrated that SlyA

positively regulates genes by a counter-silencing mechanism, in which repression of AT-rich

promoters by the histone-like nucleoid-associated protein H-NS is relieved by SlyA (16, 19). SlyA

cooperatively remodels the H-NS-DNA complex in concert with the response regulator PhoP (19,

20), which is activated by conditions found within phagosomal compartments, including low

Mg^{2+}(21), acidic pH (22), and cationic antimicrobial peptides (23). SlyA orthologs, represented by

Hor, Rap, and RovA in *Pectobacterium* (*Erwinia*), *Serratia*, and *Yersinia,* respectively (24), are

conserved in every species of *Enterobacteriaceae*, even including endosymbionts such as *Sodalis*

glossinidius (25), which have undergone extensive gene loss and degenerative evolution (26). This

high degree of conservation suggests that the SlyA lineage occupies an essential role in the

regulatory network organization of *Enterobacteriaceae*. Although conclusive mechanistic

evidence to demonstrate that other SlyA orthologs function as counter-silencers has not yet been

obtained, existing evidence is strongly suggestive of counter-silencing, as several are known to

up-regulate horizontally-acquired traits, which are generally repressed by H-NS, in a number of

species, including *Yersinia* spp. (27-29)*, Dickeya dadantii* (30), *Pectobacterium carotovorum* (24)

and *Shigella flexneri* (31).

We sought to understand the evolutionary transition of the SlyA/RovA TF lineage from

the ancestral function of MarR family TFs as environmentally-responsive and dedicated repressors

of small regulons to counter-silencers of extensive networks of horizontally-acquired genes. To do

this, we performed structural and comparative analyses of representative members of the SlyA

lineage to identify the evolutionary changes that allowed SlyA to adopt its new role. Here we show

that SlyA has retained the ability to undergo conformational changes in response to aromatic

carboxylates, regulate gene expression in an environmentally-responsive manner, and repress the

expression of a linked drug efflux system. We further show that SlyA/RovA promoters have

evolved to support the higher levels of expression that are required for counter-silencing. Finally,

we show that linked efflux pumps are not conserved in some *Enterobacteriaceae,* even though

SlyA/RovA TFs have been evolutionarily retained, suggesting that these regulators have been

conserved not due to their primordial role in regulating antimicrobial resistance but rather as a

consequence of their counter-silencing function, which is essential to maintain the regulated

expression of horizontally-acquired genes in *Enterobacteriaceae*.

Results

Salicylate-mediated inhibition of SlyA activity. As environmentally-responsive repressors

whose conformation and regulatory actions are modulated by small aromatic carboxylates (11,

13), MarR family TFs are inhibited by salicylate *in vitro* (11). To determine whether the SlyA

lineage has retained ancestral MarR characteristics, we performed *in vitro* transcription assays

(IVTs) of *slyA* and the divergently transcribed *ydhIJK* efflux pump operon. Supercoiled plasmid

DNA containing the *slyA-ydhIJK* region was incubated with RNA polymerase (RNAP) and

increasing SlyA concentrations in the presence or absence of salicylate. SlyA repressed *slyA*

transcription approximately 5.3-fold, while *ydhI* transcription was inhibited ~19-fold (Fig. 1A, B).

The addition of 2mM sodium salicylate reduced SlyA-mediated repression to 2.8-fold and 3.2-

fold, respectively, indicating that the sensitivity to aromatic carboxylates observed in classical

MarR TFs has been retained by SlyA.

We then confirmed that the *ydhIJK* operon encodes a functional antimicrobial efflux pump

by growing wildtype, *slyA*, *ydhIJK*, and *slyA ydhIJK* mutant strains in the presence of an aromatic

carboxylate with antimicrobial activity, fusaric acid (Fig. 1C). Both the *ydhIJK* and *slyA ydhIJK*

mutant strains exhibited delayed growth in the presence of fusaric acid relative to the wildtype

strain, suggesting that *ydhIJK* confers fusaric acid resistance in wildtype *S*. Typhimurium.

Conversely, *slyA* mutants exhibit improved growth in the presence of fusaric acid, as predicted

when *ydhIJK* is derepressed. Collectively, these observations indicate that the *S*. Typhimurium

SlyA TF has retained ancestral functions characteristic of the MarR family.

To determine whether salicylate inhibits SlyA-mediated counter-silencing as well as

repression, expression of the counter-silenced *pagC* gene was measured by qRT-PCR in the

presence or absence of 2mM sodium salicylate (Fig. 1D). Wildtype cultures grown in salicylate

phenocopied a *slyA* mutant strain, with a >14-fold reduction in *pagC* expression, indicating that

salicylate is a general allosteric inhibitor of SlyA function, most likely inducing a structural change

that reduces affinity for DNA as described in other MarR TFs (9, 14). Unexpectedly, SlyA retained

an ability to interact with DNA upstream of the *ydhI* promoter even in the presence of salicylate

(Fig. S3), but this interaction may be non-specific, as salicylate inhibited SlyA interaction with a

20 bp DNA segment centered near the -35 position of the *ydhI* TSS, which exhibited the strongest

binding by SlyA.

To determine whether salicylate-mediated inhibition correlates with structural changes in

SlyA, ^{1}H,^{15}N-TROSY NMR spectra of *S*. Typhimurium SlyA in the presence or absence of

salicylate were collected. The apo-SlyA spectrum is well dispersed (Fig. 2), with ~85% of the

expected resonances observed. However, wide variation among individual resonances with respect

to peak width and intensity may signify [1] weak non-specific interactions between SlyA dimers,

[2] varying rates of exchange of amide protons with solvent, or [3] conformational exchange.

Previous studies of MarR found that apo-MarR is also highly disordered (14), suggesting that the

observed spectral characteristics of apo-SlyA can be ascribed to the presence of multiple

conformational states in solution. Addition of salicylate to SlyA induces chemical shift

perturbations throughout the spectrum and nearly 90% of expected resonances are now observed.

The large-scale changes in chemical shifts show that backbone amides throughout the protein are

stabilized in a different environment relative to apo-SlyA. These observations are consistent with

ligand binding to SlyA dimers inducing global structural changes, likely stabilizing a single protein

conformation in solution, a conformation that is no longer able to interact with specific high-

affinity DNA binding sites.

Crystal structure of salicylate-SlyA. To further analyze the mechanism of allosteric inhibition

of SlyA, the structure of the salicylate-SlyA co-crystal was determined. Studies by other groups

have previously determined the structure of apo-SlyA and the SlyA bound to DNA (32),

demonstrating that SlyA is similar in overall structure to other MarR proteins, consisting of six

alpha helices (Fig. 3). Helices α1, α5, and α6 make up the dimerization domain, while α3 and α4,

along with the wing region between α4 and α5 comprise the wHTH DNA-binding domain. These

two domains are separated by α2. Dolan *et al.* (32) previously observed that the two recognition

helices of the apo-SlyA dimer are only ~15Å apart, in a closed conformation. During interaction

with a high-affinity binding site, the helices move a significant distance to accommodate the 32Å

distance between major grooves.

SlyA formed large crystals in the presence of 75mM sodium salicylate (Fig. S4). However,

we were unable to obtain usable crystals of apo-SlyA. Diffraction data set and refinement statistics

are summarized in Tables S1 and S2, respectively. The two SlyA molecules in the asymmetric unit

form two different SlyA dimers in this crystal form. Space group symmetry operations generate

the other subunit in each dimer. The dimers are very similar in structure, and further discussion

will focus on the dimer formed by polypeptide chain A. We were unable to observe electron

density for the tips of the wings, so these regions are absent from our structural model. Difference

electron density maps (|Fo|-|Fc|) identified two salicylate molecules bound per SlyA monomer at

sites referred to as Site I and Site II (Fig. 3A-C). Salicylate molecules interact with these sites via

hydrophobic interactions with their aromatic rings, while the carboxylate and hydroxyl groups are

positioned to interact with the solvent. Site I is composed of residues from α2, α3, and α4, as well

as I58 in the loop between α3 and α4, and is well positioned to sterically inhibit DNA binding. It

should be noted that the residue numbers in the deposited PDB file are not in register with the

residue numbers in this text which are based on alignment of SlyA orthologs. Comparison with

apo-SlyA and SlyA-DNA structures (32) indicates that this salicylate molecule causes the α4

recognition helix to rotate by ~35° around its axis, disrupting specific contacts with the DNA major

groove. Site II is formed by residues from both dimer subunits, almost completely sequestering

the salicylate molecule from the solvent. The buried polar groups of salicylate interact with S7'

and R14' in α1 of one subunit and R17 in α1 and H38 in α2 from the other subunit. A third

salicylate binding site was observed on the surface of each subunit of the dimer. However, this site

is adjacent to SlyA residues involved in crystal packing contacts and may not be biologically

relevant.

Evolutionary analysis of the SlyA TF lineage. To identify other variables that might have

contributed to the evolution of the SlyA lineage, we performed an evolutionary analysis of

representative SlyA proteins throughout the *Enterobacteriaceae*. Phylogenetic analysis reveals

four clusters of SlyA orthologs (Fig. 4A). SlyA orthologs in enteric pathogens including *E. coli*,

S. Typhimurium, and *S. flexneri* form cluster I. Orthologs in more distantly related pathogens such

as *Yersinia* spp. and *Proteus mirabilis* make up cluster II, while orthologs in insect endosymbionts

such as *S. glossinidius* and plant pathogens such as *D. dadantii* make up clusters III and IV,

respectively. Despite having significantly diverged from cluster I, orthologs in both clusters II and

IV are known to function as pleiotropic regulators (27, 30). This suggests that the SlyA lineage

began to diverge from the ancestral function of MarR TFs prior to the divergence of

Enterobacteriaceae from other bacteria. To determine if allosteric inhibition is likely to be

conserved throughout the SlyA lineage, we aligned the sequences of SlyA orthologs from different

bacterial species to analyze the conservation of residues involved in salicylate binding (Fig. S5,

Table S3). Cluster I exhibited complete conservation of the salicylate-binding residues. Cluster II

exhibited the greatest variation among the free-living species, with both the *Proteus mirabilis* and

Y. pseudotuberculosis proteins exhibiting an H38Y mutation, and the *P. mirabilis* protein encoding

an additional S7A mutation, either of which would alter the second salicylate binding site. Some

members of cluster IV (*P. carotovorum*) also exhibited a mutation in site II (I20V). The orthologs

from the endosymbiotic species of cluster III also exhibited a unique set of polymorphisms, with

all examined species exhibiting a V35I mutation (sites I and II), and the *Wigglesworthia*

glossinidia protein containing an A10S mutation (site II).

As the ancestral function of MarR TFs is the negative regulation of genes encoding drug

efflux pumps, a function that is conserved in *S.* Typhimurium, we examined *slyA* orthologs

throughout the *Enterobacteriaceae* for linkage to the *ydhIJK* efflux operon (Fig. 4B). Linkage is

conserved among the enteric pathogens of cluster I, although some, such as *S. flexneri*, no longer

encode the efflux pump YdhK. However, linkage to the *ydhIJK* operon is no longer conserved in

clusters II, III, or IV, as the entire operon has been lost in nearly all of the species examined. The

ydhIJK locus does not appear to have been exchanged for another drug efflux system. Rather, the

region upstream of *slyA* is highly variable, resulting in the replacement of *ydhIJK* by a wide variety

of sequences (Fig. S6). Sequences homologous to the proximal portion of the *ydhI* gene are

detectable even in species that have lost functional coding sequences (*e.g., Y. pseudotuberculosis*,

D. dadantii). The retention of *slyA* orthologs outside of cluster I despite the loss of the linked efflux

system suggests that proteins in the SlyA lineage have evolved to serve a function distinct from

that of classical MarR family TFs. SlyA is convergently transcribed with *slyB*, encoding a putative

lipoprotein. Although this linkage appears to be more strongly conserved, with *slyB* present in all

examined species with the exception of *W. glossinidia*, the significance of this conservation is

unclear, as the function of *slyB* is unknown.

Despite being closely related, *rovA* in *Y. pseudotuberculosis* and *Y. pestis* (data not shown)

is not linked to a functional *ydhIJK* operon, whereas *ydhIJK* is retained in *Y. enterocolitica*. This

indicates that *ydhIJK* was not lost by the former species until after the divergence of *Yersinia* from

the other *Enterobacteriaceae*. Other species of *Enterobacteriaceae* are likely to have lost *ydhIJK*

more recently, as each became adapted to its own specific niche. This appears to represent an

example of parallel evolution, with multiple species independently losing *ydhIJK* as their own

SlyA orthologs adapted to new roles.

Functional characterization of a distantly related SlyA ortholog. To understand how the SlyA

lineage adapted to its emergent role as an important pleiotropic regulatory protein in the

Enterobacteriaceae, we compared the functions of SlyA from *S*. Typhimurium (SlyA$_{STM}$) and

RovA of *Y. pseudotuberculosis* and *Y. enterocolitica,* a relatively divergent ortholog that also

functions as a counter-silencer (27-29, 33). RovA of *Y. pseudotuberculosis* exhibits 76% identity

with SlyA$_{STM}$ (Fig. 4A, Fig. S5). RovA is essential for virulence in *Y. enterocolitica*, and *Y. pestis*

(27, 34) and has been suggested to function both as an activator, interacting with RNAP (35), and

as a counter-silencer, alleviating H-NS-mediated repression (28, 29, 33). Notably, direct activation

has not been demonstrated for any other member of the SlyA lineage, and the most direct evidence

to suggest that RovA functions as an activator is derived from IVT studies using small linear

fragments of DNA as a template (35). We have previously shown that small linear fragments are

not necessarily representative of physiological regulatory events in intact cells and can generate

spurious results in IVT assays (16). Genetic analysis of the *inv* gene, which is positively regulated

by RovA, in an *hns* mutant strain of *E. coli* also suggested that RovA functions as both a counter-

silencer and an activator. However, these experiments failed to consider the potential contribution

of the H-NS paralog, StpA, which is up-regulated in *hns* mutants and can provide partial

complementation (36, 37), which complicates the interpretation of regulatory studies in an *hns*

mutant strain.

We attempted to corroborate the published findings by performing IVT analysis of *inv*

expression in the presence of RovA. However, expression of *inv* exhibited only a very modest

(~1.5-fold) increase following the addition of RovA (Fig. 5A), which became saturated at a 20 nM

concentration; these results are not supportive of direct *inv* activation by RovA. In contrast, RovA

was confirmed to function as an auto-repressor, like other MarR family TFs, as IVT analysis

demonstrated a 4-fold decrease in *rovA* expression following the addition of RovA protein, with

the effect reaching saturation at a 500 nM concentration (Fig. 5B). RovA has also retained the

ability to respond to salicylate (Fig. 5C), despite the presence of an H38Y mutation in the second

salicylate-binding pocket (Table S3), as 5mM salicylate completely inhibited RovA-mediated

repression, similar to our observations with SlyA. It is also notable that RovA has retained

salicylate sensitivity despite the loss of the linked YdhIJK efflux pump in *Y. pseudotuberculosis*.

To prove that RovA functions as a counter-silencer in *Y. pseudotuberculosis*, we measured *inv*

expression in wild-type and *rovA* mutant strains expressing H-NST from enteropathogenic *E. coli*

(H-NST$_{EPEC}$). Mutations in *hns* cannot be generated in *Yersinia* as *Yersinia* spp. carry a single

essential *hns* gene, unlike many other members of the *Enterobacteriaceae* which encode *hns*-like

genes such as *stpA*, which are able to partially compensate for the loss of *hns*. H-NST$_{EPEC}$ is a

truncated *hns* homolog that has been demonstrated to function as a dominant negative form of H-

NS by binding and inhibiting the activity of wild-type H-NS protein (38). The expression of *inv* in

rovA mutant bacteria was approximately four-fold lower than in wild-type cells (Fig. 5D).

However, *inv* expression was fully restored upon inhibition of H-NS by *hnsT*$_{EPEC}$, demonstrating

that RovA functions solely as a counter-silencer of the *inv* gene. The *rovA* and *slyA* genes are

capable of complementing each other in both *S.* Typhimurium and *Y. pseudotuberculosis* when

expressed *in trans* (Fig. 5E, F), up-regulating both *inv* and *pagC*, further indicating that RovA

functions as a counter-silencer, like SlyA. Together, these observations demonstrate that RovA

has retained the ancestral characteristic of environmentally-responsive repression exhibited by

other members of the MarR TF family, despite being one of the most divergent members of the

SlyA lineage. However, in contrast to MarR TFs outside of the SlyA lineage, it is also able to

function as a counter-silencer of horizontally-acquired genes, as exemplified by *inv*.

SlyA counter-silencing requires high expression levels. It is interesting to note that SlyA does

not play a major regulatory role in *Escherichia coli,* the most well-studied member of the

Enterobacteriaceae. SlyA is only known to regulate two *E. coli* genes, *hlyE* and *fimB* (39-41),

despite exhibiting a high degree of homology (91% identity) to SlyA in *S.* Typhimurium. To

better understand the different roles of SlyA in the regulatory hierarchy of *E. coli* and *S.*

Typhimurium, we performed a comparative genetic analysis of the *S.* Typhimurium ($slyA_{STm}$)

and *E. coli* ($slyA_{Eco}$) alleles. In allelic exchange experiments, $slyA_{STm}$ was swapped with $slyA_{Eco}$

to determine the effect on the expression of the counter-silenced *S.* Typhimurium *pagC* gene

(Fig. 6A). We observed that $SlyA_{Eco}$ is able to counter-silence the expression of *pagC* similarly

to $SlyA_{STm}$, suggesting that the diminished role of SlyA in *E. coli* is not attributable to

differences in protein sequence. This suggested that the importance of different SlyA lineage

proteins within their respective regulatory networks may be determined by differences in levels

of expression. To test this hypothesis, *slyA* promoter-ORF chimeras were constructed and

assessed for their ability to counter-silence *pagC*. Chimeric constructs in the chromosome were

unstable, so counter-silencing was instead assessed in a *slyA* mutant carrying pKM05, a plasmid

with the $slyA_{STM}$ ORF transcribed by the *E. coli* promoter (P$slyA_{Eco}$), or pKM07, a plasmid with

the $slyA_{STM}$ ORF transcribed by its native promoter (P$slyA_{STM}$). Although *pagC* expression was

similar with either construct under non-inducing conditions, expression was approximately 10-

fold lower with *slyA* expressed from P$slyA_{Eco}$ under inducing conditions (Fig. 6B), suggesting

that the diminished role of SlyA in *E. coli* is at least partially attributable to differences in *slyA*

expression levels in *S.* Typhimurium and *E. coli*. Although *slyA* expression driven by P$slyA_{Eco}$ is

only slightly lower under non-inducing conditions, it is approximately 3-fold greater when

driven by P$slyA_{STM}$ under inducing conditions (Fig. 6C), suggesting that P$slyA_{Eco}$ is less

responsive to environmental signals associated with virulence gene expression in *Salmonella*,

and thus is unable to regulate *slyA* expression in a manner appropriate for counter-silencing.

Parallel experiments in *Y. pseudotuberculosis* revealed that *rovA* is also more strongly expressed

from its native promoter (P*rovA*$_{Yps}$) than from a chimera expressing *rovA* from P*slyA*$_{Eco}$ (Fig.

S7).

Discussion

The ancient MarR family of transcription factors is represented throughout the bacterial

kingdom and in many archaeal species (42). This study sought to understand how the SlyA/RovA

lineage of MarR TFs in the *Enterobacteriaceae* evolved to acquire the novel function of counter-

silencing. Our observations demonstrate that SlyA proteins have retained vestiges of their ancestral

functions, e.g., the environmentally-responsive repression of small molecule efflux systems, while

acquiring an ability to act as pleiotropic counter-silencers of horizontally-acquired genes. The

latter role has been facilitated by the parallel evolution of cis-regulatory elements that support

higher levels of gene expression.

SlyA, like other MarR family TFs, is subject to allosteric inhibition by small aromatic

carboxylate compounds such as salicylate, which bind and stabilize the SlyA dimer in a

conformation unfavorable for DNA binding (Fig. 2, Fig. S3). However, the specific structure and

arrangement of the effector binding sites varies significantly among the MarR family. Our

structural data indicate that SlyA binds a total of six salicylate molecules per dimer, although two

of these are not likely to be biologically relevant (Fig. 3). Although the general structure and

architecture of the salicylate-SlyA complex is similar to complexes formed by other MarR TFs,

including MarR (14) and MTH313 from *Methanobacterium thermautrophicum* (43), the various

TFs differ significantly in their specific interactions with salicylate. MarR binds a total of four

salicylate molecules per dimer, whereas MTH313 binds only two (Fig. 3D, E). All four salicylate

binding sites in MarR flank the wHTH domain and are partially exposed to the solvent, whereas

none of these sites correspond to the salicylate binding sites of SlyA. MTH313 binds salicylate

asymmetrically at sites similar to sites I and II of SlyA, binding only one salicylate molecule per

monomer. It is possible that the structural variability in the MarR family with respect to salicylate

binding reflects an inherent evolutionary flexibility and ligand promiscuity. The MarR family

sensor region may have evolved to interact with a variety of small molecules, and salicylate may

simply represent a promiscuous probe for potential interactions. Alternatively, recent studies

indicate that the true ligand of MarR may be copper, which oxidizes a cysteine residue (C80) to

promote disulfide bond formation between MarR dimers, resulting in subsequent tetramerization

and inhibition of DNA binding (44). The copper atom is thought to be liberated from membrane-

associated proteins upon oxidative stress induced by bactericidal xenobiotic agents. As almost all

characterized MarR TFs have a reactive cysteine residue corresponding to a region near Site II of

SlyA, cysteine oxidation, which has also been described in the MarR family TF OhrR (45), may

be a common inhibitory mechanism. Indeed, SlyA contains a cysteine residue that is conserved

throughout the *Enterobacteriaceae* at position 82 (Fig. S5).

A phylogenetic analysis of SlyA orthologs in *Enterobacteriaceae* demonstrates that the

SlyA lineage is strongly conserved, even in endosymbiotic species exhibiting significant genome

loss (Fig. 4, Fig. S5). This suggests that SlyA has a central and essential role in the transcriptional

regulatory networks of these species. For example, a recent analysis of the endosymbiont *W.*

glossinidia found that *slyA* is subject to significant evolutionary constraints (46). This conservation

does not appear to be due to its role as a regulator of antimicrobial resistance, as *slyA* does not

exhibit significant linkage to the efflux pump operon *ydhIJK*, outside of the enteric pathogens in

cluster I. Rather, SlyA proteins appear to function predominantly as pleiotropic counter-silencers,

facilitating the integration of horizontally-acquired genes, including virulence genes, into existing

regulatory networks. A general counter-silencing role has been suggested in multiple

enterobacterial species including *E. coli* (18, 41), *Salmonella* (16, 19, 20) and *Shigella* (31) in

cluster I, *Yersinia* in cluster II, and *Dickeya* (30, 47), *Serratia* and *Pectobacterium* (*Erwinia*) (24)

in cluster IV. Although similar evidence does not currently exist for the endosymbionts in cluster

III, this may simply reflect the limited genetic analyses that have been performed in these species.

To understand the evolutionary adaptations to accommodate counter-silencing by the

SlyA/RovA lineage, we compared three orthologs, $slyA_{STM}$, $slyA_{Eco}$ and *rovA*. The $slyA_{STM}$ and

rovA genes are essential for *Salmonella* and *Yersinia* virulence, whereas $slyA_{Eco}$ plays a negligible

role in *E. coli*, despite exhibiting a much higher degree of similarity to $slyA_{STM}$ than does *rovA*.

These differing roles are attributable to differences in expression, as $SlyA_{Eco}$ is able to function as

a counter-silencer in *S*. Typhimurium when its expression is driven by the *S*. Typhimurium

promoter (Fig. 6). We also observed that $slyA_{Eco}$ transcription is significantly reduced in medium

with low Mg^{2+} concentrations, a condition associated with the *Salmonella*-containing vacuole

(SCV) of macrophages, in which SlyA cooperates with the response regulator PhoP to counter-

silence virulence genes necessary for intracellular survival (16). This suggests that the diminished

role of $slyA_{Eco}$ in *E. coli* may result from its inability to respond to appropriate environmental cues.

This is further reinforced by the observation that RovA of *Yersinia* spp. is able to function as a

counter-silencer in *S*. Typhimurium when expressed *in trans* from an inducible promoter, and that

the *rovA* promoter of *Y. pseudotuberculosis* drives *rovA* transcription more strongly than P$slyA_{Eco}$

in a chimeric strain (Fig. S7). We conclude that the ability of a given SlyA ortholog to serve as a

counter-silencer is contingent on its level and pattern of expression. Notably, another group

recently demonstrated a significant expansion of the SlyA regulon in *E. coli* when $slyA_{Eco}$ is

overexpressed, with 30 operons exhibiting regulation by SlyA, 24 of which are also repressed by

H-NS (48), indicating that SlyA$_{Eco}$ is capable of functioning as a counter-silencer but is not

expressed under the appropriate conditions. In *S*. Typhimurium, the appropriate conditions are

those associated with the SCV. However, SlyA orthologs in other species such as the plant

pathogens of cluster IV and the endosymbionts of cluster III are likely to require expression under

vastly different conditions corresponding to their environmental niches. The loss of the

divergently-transcribed *ydhIJK* operon in *Y. pseudotuberculosis* and other enteric species may be

a consequence of genetic alterations to enhance *rovA* expression as well as the redundancy of drug

efflux pumps. An additional possibility is that enhanced *slyA/rovA* expression might result in

hyper-repression of *ydhIJK*, which would negate the usefulness of the pump to the cell. Even the

plant pathogens of cluster IV, which are most likely to encounter small phenolic compounds in the

plant environment (49), have failed to retain *ydhIJK*. It is also notable that the *slyA* orthologs in

the four species (*E. coli*, *S*. Typhimurium, *Y. enterocolitica*, *Y. pseudotuberculosis*) in which the

transcriptional start sites have been characterized initiate transcription at different positions (29,

50-52), indicating that each species has evolved its cis-regulatory circuit independently (Fig. S6).

Previous studies have demonstrated that regulatory evolution can promote adaptation to

new niches (53). The SlyA/RovA TF lineage provides a unique example of parallel regulatory

evolution to achieve a common functional objective. Throughout the *Enterobacteriaceae*, the

associated *ydhIJK* pump genes have been repeatedly lost, yet their regulators have been retained,

presumably to facilitate the evolution of an appropriately responsive regulatory circuit to enable

counter-silencing. This suggests that SlyA/RovA proteins possess intrinsic features that predispose

them for a counter-silencing role, perhaps the abilities to respond to environmental stimuli and to

recognize a variety of AT-rich target DNA sequences. Studies are underway to characterize these

features and to determine their contribution to the evolutionary capacity of *Enterobacteriaceae*.

Materials and methods.

Bacterial strains and general reagents. All oligonucleotides and plasmids used in this study are

described in Table S4 and Table S5, respectively. Unless otherwise indicated, bacteria were grown

in Luria Bertani (LB) broth with agitation. *Salmonella enterica* serovar Typhimurium strains were

constructed in the ATCC 14028s background and grown at 37°C, unless otherwise indicated. The

14028s *slyA* mutant strain was described previously (16). *Yersinia pseudotuberculosis* YPIII (54)

and YP107 (55) (a gift from P. Dersch, Helmholtz Centre for Infection Research) were used as

the wildtype and *rovA* strains, respectively, and grown at 24°C. A *ydhIJK* deletion mutant was

constructed using the λ-Red recombinase system (56) and oligonucleotides WNp318 and

WNp319. A *slyA ydhIJK* strain was generated by introducing the *slyA::Cm* cassette from 14028s

slyA::Cm (16) to 14028s *ydhIJK* via P22HTint-mediated transduction. To exchange the wild-type

S. Typhimurium *slyA* coding sequence with the *E. coli* allele, *S*. Typhimurium *slyA* was replaced

with a *thyA* cassette, via FRUIT (57), using the oligonucleotides STM-slyA-targ-F and STM-slyA-

targ-R . The *slyA* coding sequence from *E. coli* K-12 was then amplified using the primers STM-

Eco-slyA-F and STM-Eco-slyA-R, which include 40 and 41 bases, respectively, from the regions

flanking the *S*. Typhimurium *slyA* coding sequence. This fragment was electroporated into the

slyA::thyA mutant strain and the resulting transformants were plated on minimal media containing

trimethoprim, as described for the FRUIT method (57), to select for replacement with the *E. coli*

slyA allele.

Cloning. A 2249 bp region containing both *slyA* and *ydhI* was amplified by PCR using the primers

SlyAreg-F and SlyAreg-R. The resulting fragment was digested with *Bam*HI and *Hin*dIII and

ligated into the low-copy IVT scaffold vector pRW20 (16) to generate pRW39. An IVT target

containing the 2379 bp *rovA* region from *Y. pseudotuberculosis* was generated using the primers

BamHI-rovA-F and EcoRI-rovA-R. The resulting fragment and pRW20 were both digested with

*Bam*HI and *Eco*RI and ligated together to construct pRW54. An IVT target containing the 2708

bp *inv* region from *Y. pseudotuberculosis* was generated via PCR using the primers BamHI-inv-F

and EcoRI-inv-R. The resulting fragment and pRW20 were both digested with *Bam*HI and *Eco*RI

and ligated together to construct pRW55. To perform *rovA* and *slyA* complementation studies, both

genes were cloned into the arabinose-inducible expression vector pBAD18 (58). The *rovA* gene

was amplified using the oligonucleotides EcoRI-rovA-F and KpnI-rovA-R. The *slyA* gene was

amplified using the oligonucleotides EcoRI-slyA-F and KpnI-slyA-R. Both fragments were

digested with *Eco*RI and *Kpn*I, and ligated into pBAD18, generating pRW58 (*slyA*) and pRW59

(*rovA*). pRW60 was constructed by cloning an *N*-terminal 6×His-tagged copy of *rovA* into

pTRC99A. The *rovA* gene was amplified from YPIII genomic DNA by PCR using the

oligonucleotides 6HisRovA-F and 6HisRovA-R. The resulting fragment and pTRC99a were

digested with *Nco*I and *Bam*HI, and ligated together. An *E. coli* P$_{slyA}$-S. Typhimurium *slyA* coding

sequence chimera was constructed using overlapping PCR. The *slyA-ydhIJK* intergenic region was

amplified from *E. coli* K12 genomic DNA using primers KMp178 and KMp206, and from *S.*

Typhimurium 14028s using oligonucleotides KMp177 and KMp206. The corresponding *slyA* ORF

from *S*. Typhimurium was amplified with primers KMp207 and KMp181. The ORF and promoter

segments were amplified along with 40bp of complementary overlapping sequence. Products from

the ORF and promoter reactions were mixed 1:1 and amplified using primers KMp178 (*E. coli*) or

KMp177 (*S*. Typhimurium) and KMp181 to amplify the promoter/ORF chimeras. The resulting

fragment and pTH19Kr (59) were digested with *Bam*HI and *Hin*dIII and ligated together,

generating pKM05 (*E. coli* promoter) and pKM07 (*S*. Typhimurium promoter). Plasmids were

transformed into S. Typhimurium 14028s *slyA* for gene expression analysis. The *hnsT* coding

sequence was amplified from E2348/69 genomic DNA using the oligonucleotides EcoRI-hnsT-F

and HindIII-hnsT-R. Both pBAD18 and the resulting PCR fragment were digested with *Eco*RI and

*Hin*dIII, agarose gel-purified, and ligated together to generate pRW57.

Fusaric acid resistance assays. Cultures were grown overnight in LB broth at 37°C, then diluted

1:100. Thirty microliters of this dilution were added to 270 µl of LB containing 30 µg/ml freshly

prepared fusaric acid in 100-well BioScreen plates (GrowthCurves, Piscataway, NJ). Cultures

were grown with continuous maximum shaking at 37°C and regular OD_{600} measurements were

taken on a BioScreen C MBR. Fresh 20 mg/ml stock solutions of fusaric acid were prepared in

DMF (dimethyl formamide).

NMR analysis of SlyA-salicylate. SlyA protein for NMR analysis was prepared as an *N*-terminal

6×His-tagged protein from cells grown in M9 minimal medium supplemented with ^{15}N-

ammonium chloride (Cambridge Isotope Labs, Tewksbury MA). SlyA protein expression was

induced by the addition of 2 mM IPTG (isopropyl-ß-D-thiogalactopyranoside) and the protein

purified to homogeneity using Ni-affinity chromatography as previously described (20), except

that samples were dialyzed in 50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 0.1mM EDTA, and 3 mM

DTT following purification. Protein and ligand solutions for NMR experiments were prepared in

25 mM sodium phosphate, 150 mM NaCl buffer at pH 7.0 containing 10% D_2O. NMR spectra

were collected on a Bruker DMX 500hHz spectrometer (Bruker, Billerica MA) on samples

equilibrated at 35°C and consisting of 250-350 µM ^{15}N-labeled His-SlyA in the absence or

presence of 2 mM sodium salicylate. Spectra were processed using NMR-Pipe (60) and analyzed

using NMR-View (61).

SlyA-salicylate crystallization. SlyA was over-expressed and purified for crystallization as

previously described (20), except that samples were dialyzed in 50 mM Tris-HCl, pH 8.0, 150 mM

NaCl, 0.1 mM EDTA, and 3 mM DTT following purification. Cryo I and II sparse matrix

crystallization screens (Emerald Biosystems, Bainbridge Island, WA) were used to determine

initial conditions for His-SlyA crystal formation by sitting-drop vapor diffusion in 24-well crystal

trays. Equal volumes (4µL) of SlyA and crystallization solutions were mixed before plates were

sealed and kept at room temperature. Crystals appeared within 4-7 days in 20% PEG 300, 10%

glycerol, 0.1 M phosphate/citrate buffer, pH 4.2, 0.2M ammonium sulfate (condition #14).

Subsequent crystal growth was performed using lab-made 20% PEG 400, 10% glycerol, 0.1 M

phosphate/citrate buffer, pH 4.2, and 0.2 M ammonium sulfate. Crystallization of SlyA with

sodium salicylate was achieved by adding dilutions of a 2M sodium salicylate aqueous stock

solution to the protein-crystallization solution mixture. SlyA-salicylate crystals appeared within

7-10 days. A single crystal appeared at a concentration of 75 mM sodium salicylate and grew to

a maximum size of 500µ x 300µ x 150µ (Fig. S4). This crystal was used for X-ray diffraction

experiments.

476   The crystal was frozen at 100°K in its crystallization solution for diffraction data

collection on GM/CA-CAT beamline 23-ID-D at the Advanced Photon Source. The space group

for the crystals is $P2_12_12$ with two SlyA molecules in the asymmetric unit. The diffraction data

were processed with HKL2000 (62). Dataset statistics are shown in Table S1. The crystal structure

of the salicylate complex of SlyA was solved using a model of a previously investigated structure

of apo-SlyA (unpublished data). That apo-structure was solved using the molecular replacement

program, MOLREP (63) with a search model generated by applying Swiss-Model (64) and the

SlyA amino acid sequence to PDB entry 2FBH (65). The structural model for the salicylate

complex was refined using REFMAC-5 (66) in the CCP4 suite (67). Rfree (68) was calculated

using 5% of the data in the test set. A high-resolution limit of 2.3Å was applied for the refinement,

consistent with standards appropriate when the structure was solved. This is the resolution at

which Rmerge for the data set drops below 0.40. XtalView (69) and Coot (70) were used to

examine sigma A weighted |Fo|-|Fc| and 2|Fo|-|Fc| electron density maps (71). MOLSCRIPT(72),

and Raster3D (73) were used to produce structural figures for this paper. Table S2 contains

refinement statistics for the structure. Coordinates and structure factors have been deposited in the

Protein Data Bank with identifier 3DEU.

6×His-RovA purification. RovA was purified using the same protocol as described previously

for SlyA (16). However, overexpression cultures were grown at 24°C to OD_{600}=0.5. IPTG was

added to a final concentration of 1mM, and cultures were incubated at 24°C for an additional four

h before the cells were harvested by centrifugation and cell pellets stored at -80°C for the

subsequent purification of RovA.

In vitro **transcription.** IVT assays were performed essentially as previously described (16) with

the following modifications. All SlyA IVT assays were performed at 37°C. All RovA IVT assays

were performed at 24°C. Where indicated, sodium salicylate was added to IVT reactions prior to

the addition of SlyA or RovA. All oligonucleotides and templates used in IVT reactions are

indicated in Table S6.

RNA isolation and qRT-PCR. RNA was purified using Trizol (Life Technologies, Carlsbad CA)

according to the manufacturer's protocols. cDNA was generated using the QuantiTect Reverse

Transcription Kit (Qiagen, Hilden, Germany), and quantified in a BioRad CFX96 (BioRad,

Hercules CA), using SYBR Green Master Mix (74).

For the analysis of *pagC* expression under inducing conditions, *S*. Typhimurium cultures

were grown to early stationary phase ($OD_{600}{\approx}2.0$) at 37°C in LB broth, then washed three times in

N-minimal medium containing either 10 µM (inducing) or 10 mM $MgCl_2$ (non-inducing). Cultures

were re-suspended in the appropriate N-minimal medium and incubated for an additional 30 min

at 37°C before cells were harvested for RNA purification studies. For the complementation of *slyA*

with *slyA* or *rovA in trans*, overnight cultures were diluted to 0.05 OD_{600} and grown for two h at

24°C before adding arabinose to a final concentration of 0.02% w/v. Cultures were grown for an

additional six h before harvesting RNA.

Y. pseudotuberculosis strains YPIII and YP107 were diluted from overnight cultures to

0.05 OD_{600} in LB and grown with shaking at 24°C. For the inhibition of H-NS via H-NST$_{EPEC}$

over-expression, and complementation of *rovA* with *rovA* or *slyA in trans*, YPIII and YP107

cultures carrying either pBAD18 or pRW57 were grown for two hrs before arabinose was added

to a final concentration of 0.02% w/v. Cultures were grown for an additional six hrs before

harvesting RNA. For *slyA*/*rovA* complementation studies, rpoAYS primers targeting *rpoA* were

used as loading controls, as *rpoA* is sufficiently conserved between the two species as to allow use

of the same primers.

Acknowledgments

The National Institutes of Health provided support to FCF (AI39557, AI44486,

AI118962, AI112640) and SJL (A148622). GM/CA@APS has been funded in whole or in part

with Federal funds from the National Cancer Institute (ACB-12002) and the National Institute of

General Medical Sciences (AGM-12006). This research used resources of the Advanced Photon

Source, a U.S. Department of Energy (DOE) Office of Science User Facility operated for the

DOE Office of Science by Argonne National Laboratory under Contract No. DE-AC02-

06CH11357. We thank Eric Larson for helpful discussions and suggestions. We also thank

Catherine Eakin and Ponni Rajagopal for assistance with protein purification and crystal growth,

Petra Dersch for kindly providing *Y. pseudotuberculosis* YPIII and YP107, and Ralph Isberg for

providing *Y. pseudotuberculosis* IP32953. Finally, we thank Dr. Rachel Klevit for use of her

NMR facilities to collect spectra of SlyA-salicylate complexes.

Figure 1. SlyA retains the ancestral functions of MarR family TFs. To determine whether

SlyA is an autorepressor like other members of the MarR family, *in vitro* transcription (IVT)

assays were performed on supercoiled template DNA containing *slyA* and the divergently

expressed *ydhI* gene in the presence of increasing SlyA concentrations (A). Reactions were

performed with or without 2mM sodium salicylate to determine whether SlyA is inhibited by

small aromatic carboxylate compounds like other MarR TFs. (IVT analysis of SlyA-mediated

regulation of *pagC* is shown in Fig. S1.) To determine whether SlyA exhibits a dose-dependent

response to salicylate, IVT assays were performed on *ydhI* in the presence of 2μM SlyA and

increasing concentrations of sodium salicylate (B). Wildtype, *slyA*, *ydhIJK*, and *slyA ydhIJK*

cultures were grown in the presence of 30 μg/ml fusaric acid and cell density (OD_{600}) measured

over time to determine if *ydhIJK* encodes a functional anti-microbial efflux pump (C). Data

represent the mean (solid line) of three independent experiments, each consisting of three

replicates. Dashed lines represent the SD. (Growth curves in the absence of fusaric acid is shown

in Fig. S2.) To determine whether salicylate also inhibits SlyA counter-silencing activity, *pagC*

transcripts were quantified by qRT-PCR from early stationary phase ($OD_{600} \approx 2.0$) cultures in

minimal N-medium containing 10 μM $MgCl_2$ in the presence or absence of 2 mM sodium

salicylate (D). Transcript levels are normalized to *rpoD*, and data represent the mean $\pm$ SD; n$\geq$3.

Figure 2. SlyA undergoes significant structural alterations upon salicylate binding. To

assess salicylate-induced SlyA structural changes, ^{1}H,^{15}N-TROSY NMR spectroscopy was

performed on 0.3mM uniformly-labelled ^{15}N-SlyA in the absence (black) or presence (red) of

2mM sodium salicylate. (Differential DNA Footprinting Analysis, or DDFA, of the SlyA-DNA

interaction in the presence or absence of salicylate is shown in Fig. S3.)

563   **Figure 3. Structure of the SlyA dimer bound to salicylate.** The crystal structure of the SlyA

564   dimer with salicylate bound at sites I and II was determined at a resolution of 2.3Å (A). Residues

565   responsible for coordinating bound salicylates are shown for sites I (B) and II (C). The structures

566   of salicylate-bound MarR (D; PDB entry 1JGS) and MTH313 (E; PDB entry 3BPX) dimers are

567   shown for comparison. (An image of the SlyA-salicylate crystal is shown in Fig. S4.)

Figure 4. The SlyA TF lineage in *Enterobacteriaceae*. A phylogram of the SlyA TF lineage

was generated using PhyML (75), and *slyA* lineage genes from representative members of

Enterobacteriaceae were aligned (A). The *slyA* genes fall into four clusters: cluster I consists of

most enteric species, including *S*. Typhimurium and *E. coli*; cluster II contains more distantly

related pathogenic species from *Proteus* and *Yersinia*; cluster III includes endosymbiotic species

such as *S. glossinidius* and *W. glossinidia*; cluster IV contains plant pathogens such as

Pectobacterium (Erwinia) and *Dickeya*. (A multiple sequence alignment of these SlyA orthologs

can be found in Fig. S5.) Regions flanking *slyA* were examined for genetic linkage to the *ydhIJK*

efflux locus (B). The *slyA* gene is highlighted in red. The efflux pump encoded by *ydhK* is only

found in a small subset of species, suggesting that SlyA has evolved to serve a role independent

of *ydhIJK* repression in *Enterobacteriaceae*. (A multiple sequence alignment of the region

upstream of the *slyA* coding sequence is shown in Fig. S6.)

Figure 5. Environmentally-responsive repressing and counter-silencing functions are

conserved in the SlyA TF lineage. To determine whether RovA functions as a counter-silencer

in *Y. pseudotuberculosis*, transcription of the Rov-regulated *inv* gene was quantified by qRT-

PCR in wildtype or *rovA* mutant strains expressing *hnsT*$_{EPEC}$, which inhibits H-NS (A). Although

inv expression is decreased in a *rovA* mutant strain, expression is fully restored by the inhibition

of H-NS, indicating that RovA functions as a counter-silencer. IVT assays of the *inv* regulatory

region in the presence of increasing RovA concentrations detected only a minimal impact on

transcription levels, suggesting that RovA is unlikely to function as a classical activator (B). IVT

assays of *rovA* in the presence of increasing RovA concentrations of RovA demonstrate that

RovA functions as an autorepressor (C). RovA remains sensitive to inhibition by small aromatic

carboxylate molecules, as increasing concentrations of sodium salicylate inhibited RovA-

mediated *rovA* repression in IVT assays. Reciprocal complementation studies were performed by

providing both *rovA* and *slyA in trans* expressed on the arabinose-inducible pBAD18 vector in

rovA or *slyA* mutant *Y. pseudotuberculosis* and *S.* Typhimurium strains, respectively, and

measuring transcription of *inv* and *pagC* via qRT-PCR. Transcript levels are normalized to *rpoA*

in both species, and data represent the mean ± SD; n=3.

Figure 6. *S*. Typhimurium SlyA is a more effective counter-silencer than *E. coli* SlyA due to

higher expression levels. Expression of the SlyA counter-silenced *pagC* gene was measured by

qRT-PCR in *S.* Typhimurium strains encoding wildtype *slyA*, *slyA$_{Eco}$*, or mutant *slyA* under either

inducing (10 µM MgCl$_2$) or non-inducing (10mM MgCl$_2$) conditions (A). Expression of *pagC*

(B) and *slyA* (C) was compared in *S.* Typhimurium strains in which *slyA* transcription was driven

by either the wildtype (WT) or *E. coli* (P*slyA$_{Eco}$*) *slyA* promoter. Transcript levels are normalized

to *rpoD* and data represent the mean ± SD; n=3. (A comparative analysis of a P*slyA$_{Eco}$*-*rovA*

chimera is shown in Fig. S7.)

References

1. Fang FC, Frawley ER, Tapscott T, Vázquez-Torres A. 2016. Discrimination and Integration of Stress Signals by Pathogenic Bacteria. Cell Host Microbe 20:144-153.
2. Ohno S. 1970. Evolution by Gene Duplication. Springer, New York.
3. Brenner SE, Hubbard T, Murzin A, Chothia C. 1995. Gene duplications in *H. influenzae*. Nature 378:140.
4. Teichmann SA, Park J, Chothia C. 1998. Structural assignments to the *Mycoplasma genitalium* proteins show extensive gene duplications and domain rearrangements. Proc Natl Acad Sci U S A 95:14658-63.
5. Teichmann SA, Babu MM. 2004. Gene regulatory network growth by duplication. Nat Genet 36:492-6.
6. Pérez-Rueda E, Collado-Vides J, Segovia L. 2004. Phylogenetic distribution of DNA-binding transcription factors in bacteria and archaea. Comput Biol Chem 28:341-50.
7. Grove A. 2017. Regulation of Metabolic Pathways by MarR Family Transcription Factors. Comput Struct Biotechnol J 15:366-371.
8. Seoane AS, Levy SB. 1995. Characterization of MarR, the repressor of the multiple antibiotic resistance (*mar*) operon in *Escherichia coli*. J Bacteriol 177:3414-9.
9. Martin RG, Rosner JL. 1995. Binding of purified multiple antibiotic-resistance repressor protein (MarR) to *mar* operator sequences. Proc Natl Acad Sci U S A 92:5456-60.
10. George AM, Levy SB. 1983. Amplifiable resistance to tetracycline, chloramphenicol, and other antibiotics in *Escherichia coli*: involvement of a non-plasmid-determined efflux of tetracycline. J Bacteriol 155:531-40.
11. Cohen SP, Levy SB, Foulds J, Rosner JL. 1993. Salicylate induction of antibiotic resistance in *Escherichia coli*: activation of the *mar* operon and a *mar*-independent pathway. J Bacteriol 175:7856-62.
12. Wilkinson SP, Grove A. 2006. Ligand-responsive transcriptional regulation by members of the MarR family of winged helix proteins. Curr Issues Mol Biol 8:51-62.
13. Alekshun MN, Levy SB. 1999. The *mar* regulon: multiple resistance to antibiotics and other toxic chemicals. Trends Microbiol 7:410-3.
14. Alekshun MN, Levy SB, Mealy TR, Seaton BA, Head JF. 2001. The crystal structure of MarR, a regulator of multiple antibiotic resistance, at 2.3 A resolution. Nat Struct Biol 8:710-4.
15. Zhu R, Hao Z, Lou H, Song Y, Zhao J, Chen Y, Zhu J, Chen PR. 2017. Structural characterization of the DNA-binding mechanism underlying the copper(II)-sensing MarR transcriptional regulator. J Biol Inorg Chem 22:685-693.
16. Will WR, Bale DH, Reid PJ, Libby SJ, Fang FC. 2014. Evolutionary expansion of a regulatory network by counter-silencing. Nat Commun 5:5270.
17. Will WR, Navarre WW, Fang FC. 2015. Integrated circuits: how transcriptional silencing and counter-silencing facilitate bacterial evolution. Curr Opin Microbiol 23:8-13.
18. Lithgow JK, Haider F, Roberts IS, Green J. 2007. Alternate SlyA and H-NS nucleoprotein complexes control *hlyE* expression in *Escherichia coli* K-12. Mol Microbiol 66:685-98.
19. Perez JC, Latifi T, Groisman EA. 2008. Overcoming H-NS-mediated transcriptional silencing of horizontally acquired genes by the PhoP and SlyA proteins in *Salmonella enterica*. J Biol Chem 283:10773-83.

654   20.   Navarre WW, Halsey TA, Walthers D, Frye J, McClelland M, Potter JL, Kenney LJ,
655   Gunn JS, Fang FC, Libby SJ. 2005. Co-regulation of *Salmonella enterica* genes required
656   for virulence and resistance to antimicrobial peptides by SlyA and PhoP/PhoQ. Mol
657   Microbiol 56:492-508.
658   21.   García Véscovi E, Soncini FC, Groisman EA. 1996. Mg^{2+} as an extracellular signal:
659   environmental regulation of *Salmonella* virulence. Cell 84:165-74.
660   22.   Prost LR, Daley ME, Le Sage V, Bader MW, Le Moual H, Klevit RE, Miller SI. 2007.
661   Activation of the bacterial sensor kinase PhoQ by acidic pH. Mol Cell 26:165-74.
662   23.   Bader MW, Sanowar S, Daley ME, Schneider AR, Cho U, Xu W, Klevit RE, Le Moual
663   H, Miller SI. 2005. Recognition of antimicrobial peptides by a bacterial sensor kinase.
664   Cell 122:461-72.
665   24.   Thomson NR, Cox A, Bycroft BW, Stewart GS, Williams P, Salmond GP. 1997. The *rap*
666   and *hor* proteins of *Erwinia*, *Serratia* and *Yersinia*: a novel subgroup in a growing
667   superfamily of proteins regulating diverse physiological processes in bacterial pathogens.
668   Mol Microbiol 26:531-44.
669   25.   Dale C, Maudlin I. 1999. *Sodalis* gen. nov. and *Sodalis glossinidiu*s sp. nov., a
670   microaerophilic secondary endosymbiont of the tsetse fly *Glossina morsitans morsitans*.
671   Int J Syst Bacteriol 49 Pt 1:267-75.
672   26.   Akman L, Rio RV, Beard CB, Aksoy S. 2001. Genome size determination and coding
673   capacity of *Sodalis glossinidius*, an enteric symbiont of tsetse flies, as revealed by
674   hybridization to *Escherichia coli* gene arrays. J Bacteriol 183:4517-25.
675   27.   Cathelyn JS, Crosby SD, Lathem WW, Goldman WE, Miller VL. 2006. RovA, a global
676   regulator of *Yersinia pestis*, specifically required for bubonic plague. Proc Natl Acad Sci
677   U S A 103:13514-9.
678   28.   Cathelyn JS, Ellison DW, Hinchliffe SJ, Wren BW, Miller VL. 2007. The RovA regulons
679   of *Yersinia enterocolitica* and *Yersinia* pestis are distinct: evidence that many RovA-
680   regulated genes were acquired more recently than the core genome. Mol Microbiol
681   66:189-205.
682   29.   Heroven AK, Nagel G, Tran HJ, Parr S, Dersch P. 2004. RovA is autoregulated and
683   antagonizes H-NS-mediated silencing of invasin and *rovA* expression in *Yersinia
684   pseudotuberculosis*. Mol Microbiol 53:871-88.
685   30.   Haque MM, Kabir MS, Aini LQ, Hirata H, Tsuyumu S. 2009. SlyA, a MarR family
686   transcriptional regulator, is essential for virulence in *Dickeya dadantii* 3937. J Bacteriol
687   191:5409-18.
688   31.   Weatherspoon-Griffin N, Wing HJ. 2016. Characterization of SlyA in *Shigella flexneri*
689   Identifies a Novel Role in Virulence. Infect Immun 84:1073-1082.
690   32.   Dolan KT, Duguid EM, He C. 2011. Crystal structures of SlyA protein, a master
691   virulence regulator of *Salmonella*, in free and DNA-bound states. J Biol Chem
692   286:22178-85.
693   33.   Ellison DW, Miller VL. 2006. H-NS represses inv transcription in *Yersinia enterocolitica*
694   through competition with RovA and interaction with YmoA. J Bacteriol 188:5101-12.
695   34.   Revell PA, Miller VL. 2000. A chromosomally encoded regulator is required for
696   expression of the *Yersinia enterocolitica inv* gene and for virulence. Mol Microbiol
697   35:677-85.

35.   Tran HJ, Heroven AK, Winkler L, Spreter T, Beatrix B, Dersch P. 2005. Analysis of
RovA, a transcriptional regulator of *Yersinia pseudotuberculosis* virulence that acts
through antirepression and direct transcriptional activation. J Biol Chem 280:42423-32.
36.   Shi X, Bennett GN. 1994. Plasmids bearing hfq and the *hns*-like gene *stpA* complement
hns mutants in modulating arginine decarboxylase gene expression in *Escherichia coli*. J
Bacteriol 176:6769-75.
37.   Sonden B, Uhlin BE. 1996. Coordinated and differential expression of histone-like
proteins in *Escherichia coli*: regulation and function of the H-NS analog StpA. EMBO J
15:4970-80.
38.   Williamson HS, Free A. 2005. A truncated H-NS-like protein from enteropathogenic
Escherichia coli acts as an H-NS antagonist. Mol Microbiol 55:808-27.
39.   McVicker G, Sun L, Sohanpal BK, Gashi K, Williamson RA, Plumbridge J, Blomfield
IC. 2011. SlyA protein activates *fimB* gene expression and type 1 fimbriation in
Escherichia coli K-12. J Biol Chem 286:32026-35.
40.   Wyborn NR, Stapleton MR, Norte VA, Roberts RE, Grafton J, Green J. 2004. Regulation
of *Escherichia coli* hemolysin E expression by H-NS and *Salmonella* SlyA. J Bacteriol
186:1620-8.
41.   Corbett D, Bennett HJ, Askar H, Green J, Roberts IS. 2007. SlyA and H-NS regulate
transcription of the *Escherichia coli* K5 capsule gene cluster, and expression of *slyA* in
Escherichia coli is temperature-dependent, positively autoregulated, and independent of
H-NS. J Biol Chem 282:33326-35.
42.   Perera IC, Grove A. 2010. Molecular mechanisms of ligand-mediated attenuation of
DNA binding by MarR family transcriptional regulators. J Mol Cell Biol 2:243-54.
43.   Saridakis V, Shahinas D, Xu X, Christendat D. 2008. Structural insight on the mechanism
of regulation of the MarR family of proteins: high-resolution crystal structure of a
transcriptional repressor from *Methanobacterium thermoautotrophicum*. J Mol Biol
377:655-67.
44.   Hao Z, Lou H, Zhu R, Zhu J, Zhang D, Zhao BS, Zeng S, Chen X, Chan J, He C, Chen
PR. 2014. The multiple antibiotic resistance regulator MarR is a copper sensor in
Escherichia coli. Nat Chem Biol 10:21-8.
45.   Hong M, Fuangthong M, Helmann JD, Brennan RG. 2005. Structure of an OhrR-*ohrA*
operator complex reveals the DNA binding mechanism of the MarR family. Mol Cell
20:131-41.
46.   Sabater-Muñoz B, Toft C, Alvarez-Ponce D, Fares MA. 2017. Chance and necessity in
the genome evolution of endosymbiotic bacteria of insects. ISME J 11:1291-1304.
47.   Zou L, Zeng Q, Lin H, Gyaneshwar P, Chen G, Yang CH. 2012. SlyA regulates type III
secretion system (T3SS) genes in parallel with the T3SS master regulator HrpL in
Dickeya dadantii 3937. Appl Environ Microbiol 78:2888-95.
48.   Curran TD, Abacha F, Hibberd SP, Rolfe MD, Lacey MM, Green J. 2017. Identification
of new members of the *Escherichia coli* K-12 MG1655 SlyA regulon. Microbiology
163:400-409.
49.   Nicholson R, Hammerschmidt R. 1992. Phenolic-compounds and their role in disease
resistance. Annual Review of Phytopathology 30:369-389.
50.   Salgado H, Peralta-Gil M, Gama-Castro S, Santos-Zavaleta A, Muñiz-Rascado L, García-
Sotelo JS, Weiss V, Solano-Lira H, Martínez-Flores I, Medina-Rivera A, Salgado-Osorio
G, Alquicira-Hernández S, Alquicira-Hernández K, López-Fuentes A, Porrón-Sotelo L,

Huerta AM, Bonavides-Martínez C, Balderas-Martínez YI, Pannier L, Olvera M,
Labastida A, Jiménez-Jacinto V, Vega-Alvarado L, Del Moral-Chávez V, Hernández-
Alvarez A, Morett E, Collado-Vides J. 2013. RegulonDB v8.0: omics data sets,
evolutionary conservation, regulatory phrases, cross-validated gold standards and more.
Nucleic Acids Res 41:D203-13.
51.   Kröger C, Colgan A, Srikumar S, Händler K, Sivasankaran SK, Hammarlöf DL, Canals
R, Grissom JE, Conway T, Hokamp K, Hinton JC. 2013. An infection-relevant
transcriptomic compendium for *Salmonella enterica* Serovar Typhimurium. Cell Host
Microbe 14:683-95.
52.   Lawrenz MB, Miller VL. 2007. Comparative analysis of the regulation of *rovA* from the
pathogenic yersiniae. J Bacteriol 189:5963-75.
53.   Osborne SE, Walthers D, Tomljenovic AM, Mulder DT, Silphaduang U, Duong N,
Lowden MJ, Wickham ME, Waller RF, Kenney LJ, Coombes BK. 2009. Pathogenic
adaptation of intracellular bacteria by rewiring a *cis*-regulatory input function. Proc Natl
Acad Sci U S A 106:3982-7.
54.   Bölin I, Norlander L, Wolf-Watz H. 1982. Temperature-inducible outer membrane
protein of *Yersinia pseudotuberculosis* and *Yersinia enterocolitica* is associated with the
virulence plasmid. Infect Immun 37:506-12.
55.   Quade N, Mendonca C, Herbst K, Heroven AK, Ritter C, Heinz DW, Dersch P. 2012.
Structural basis for intrinsic thermosensing by the master virulence regulator RovA of
Yersinia. J Biol Chem 287:35796-803.
56.   Datsenko KA, Wanner BL. 2000. One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. Proc Natl Acad Sci U S A 97:6640-5.
57.   Stringer AM, Singh N, Yermakova A, Petrone BL, Amarasinghe JJ, Reyes-Diaz L,
Mantis NJ, Wade JT. 2012. FRUIT, a scar-free system for targeted chromosomal
mutagenesis, epitope tagging, and promoter replacement in *Escherichia coli* and
Salmonella enterica. PLoS One 7:e44841.
58.   Guzman LM, Belin D, Carson MJ, Beckwith J. 1995. Tight regulation, modulation, and
high-level expression by vectors containing the arabinose P_{BAD} promoter. J Bacteriol
177:4121-30.
59.   Hashimoto-Gotoh T, Yamaguchi M, Yasojima K, Tsujimura A, Wakabayashi Y,
Watanabe Y. 2000. A set of temperature sensitive-replication/-segregation and
temperature resistant plasmid vectors with different copy numbers and in an isogenic
background (chloramphenicol, kanamycin, *lacZ, repA*, *par*, *polA*). Gene 241:185-91.
60.   Delaglio F, Grzesiek S, Vuister GW, Zhu G, Pfeifer J, Bax A. 1995. NMRPipe: a
multidimensional spectral processing system based on UNIX pipes. J Biomol NMR
6:277-93.
61.   Johnson BA, Blevins RA. 1994. NMR View: A computer program for the visualization
and analysis of NMR data. J Biomol NMR 4:603-14.
62.   Otwinowski Z, Minor W. 1997. Processing of X-ray diffraction data collected in
oscillation mode. Macromolecular Crystallography, Pt a 276:307-326.
63.   Vagin A, Teplyakov A. 1997. *MOLREP*: an Automated Program for Molecular
Replacement. J Appl Cryst 30:1022-1025.
64.   Schwede T, Kopp J, Guex N, Peitsch MC. 2003. SWISS-MODEL: An automated protein
homology-modeling server. Nucleic Acids Res 31:3381-5.

65.   Lunin V, Evdokimova E, Kudritska M, Osipiuk J, Joachimiak A, Edwards A, Savchenko
790   A. 2005. The crystal structure of transcriptional regulator PA3341. Midwest Center for
791   Structural Genomics, RCSB PDB.
792   66.   Murshudov GN, Vagin AA, Dodson EJ. 1997. Refinement of macromolecular structures
793   by the maximum-likelihood method. Acta Crystallogr D Biol Crystallogr 53:240-55.
794   67.   Collaborative Computational Project Nm. 1994. The CCP4 suite: programs for protein
795   crystallography. Acta Crystallogr D Biol Crystallogr 50:760-3.
796   68.   Brünger AT. 1993. Assessment of phase accuracy by cross validation: the free R value.
797   Methods and applications. Acta Crystallogr D Biol Crystallogr 49:24-36.
798   69.   McRee DE. 1999. XtalView/Xfit--A versatile program for manipulating atomic
799   coordinates and electron density. J Struct Biol 125:156-65.
800   70.   Emsley P, Cowtan K. 2004. Coot: model-building tools for molecular graphics. Acta
801   Crystallogr D Biol Crystallogr 60:2126-32.
802   71.   RJ R. 1986. Improved Fourier coefficients for maps using phases from partial structures
803   with errors. Acta Cryst A42:140-149.
804   72.   Kraulis P. 1991. MOLSCRIPT: a program to produce both detailed and schematic plots
805   of protein structures. J Appl Cryst 24:946-950.
806   73.   Merritt EA, Bacon DJ. 1997. Raster3D: photorealistic molecular graphics. Methods
807   Enzymol 277:505-24.
808   74.   Aparicio O, Geisberg JV, Sekinger E, Yang A, Moqtaderi Z, Struhl K. 2005. Chromatin
809   immunoprecipitation for determining the association of proteins with specific genomic
810   sequences *in vivo*. Curr Protoc Mol Biol Chapter 21:Unit 21.3.
811   75.   Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New
812   algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
813   performance of PhyML 3.0. Syst Biol 59:307-21.