**Quantitatively predicting optimal antibiotic dose levels from drug-target binding**

*Fabrizio Clarelli[1], Adam Palmer[2], Bhupender Singh[1], Merete Storflor[1], Silje Lauksund[1], Ted Cohen[3], Sören Abel[1,4,#], Pia Abel zur Wiesch[1,3,#,*]* [ORCID: https://orcid.org/0000-0001-9420-9005]

1. Department of Pharmacy, Faculty of Health Sciences, University of Tromsø, Tromsø, Norway.
2. Laboratory of Systems Pharmacology, Harvard Medical School, Boston, MA, USA.
3. Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New Haven, Connecticut, United States of America.
4. Centre for Molecular Medicine Norway, Nordic EMBL Partnership, Oslo, Norway.
#  Authors contributed equally
*  e-mail: pia.z.wiesch@uit.no

**Abstract**

Improved predictions of antibiotic efficacy can inform the development of new antibiotics and extend the effectiveness of existing drugs and thereby help combatting the global antibiotic resistance crisis. We describe a computational model (COMBAT- COmputational Model of Bacterial Antibiotic Target-binding) that leverages accessible biochemical parameters to quantitatively predict the antimicrobial effects of antibiotics based on their drug-target affinity. We validate our model with MICs of a range of quinolone antibiotics in clinical isolates demonstrating that antibiotic efficacy can be predicted from drug-target binding ($R^2>0.9$). Conversely, we experimentally demonstrate that changes in drug-target binding can be predicted from antibiotic efficacy with 92-94% accuracy by exposing bacteria overexpressing target molecules to ciprofloxacin. To test the generality of COMBAT, we predict target molecule occupancy at MIC from antimicrobial action with 90% accuracy for a different antibiotic class, the beta-lactam ampicillin. Finally, we predict antibiotic concentrations that can select for resistance due to novel resistance mutations. COMBAT provides a framework to inform optimal antibiotic dose levels that maximize efficacy and minimize the rise of resistant mutants.

**Introduction**

The rise of antibiotic resistance represents an urgent public health threat. In order to effectively combat the spread of antibiotic resistance, we must optimize the use of existing drugs and develop new drugs that are effective against drug-resistant strains. Accordingly, methods to improve antibiotic dose levels to i) maximize efficacy against susceptible strains and ii) minimize resistance evolution play a key role in our defense against antibiotic resistant pathogens.

It is noteworthy that dosing strategies for treatment of susceptible strains (e.g., dosing level[1],

dosing frequency[2], and treatment duration[3-5]) have recently been substantially improved, even for

antibiotic treatments that have been standard of care for decades. This suggests that there likely

remains significant room for optimization in our antibiotic treatment regimens. It also highlights

the difficulty in identifying optimal dosing levels for new antibiotics. Indeed, optimizing dosing

is one of the biggest challenges in drug development. Typically, time-consuming trial-and-error

approaches are used and each failed drug candidate makes this process more expensive[6].

It is even more challenging to optimize dose levels to minimize the emergence of antibiotic

resistance, both for existing and novel antibiotics. There remains substantial debate about which

dosing strategies best prevent the emergence of resistant mutants during treatment[7-9]. In this

context, a useful concept that links antibiotic concentrations with resistance evolution is the

resistance selection window (mutant selection window) that ranges from the lowest

concentration at which the resistant strain grows faster than the wild-type, usually well below the

wild-type minimum inhibitory concentration (MIC), to the MIC of the resistant strain[10-12].

Antibiotic concentrations above the resistance selection window safeguard against *de novo*

resistance emergence. Antibiotic concentrations below the resistance selection window do not

kill the susceptible strain, but also do not favor the resistant strain and therefore do not promote

emergence of resistance. The latter may be preferable if one cannot dose above the MIC of the

resistant strain due to toxicity or solubility limits. To limit resistance emergence, it is therefore

important to identify the resistance selection window and optimize dosing accordingly.

Limitations in our knowledge of how antibiotic treatment regimens affect bacterial populations

contribute to the need for lengthy and expensive trial-and-error approaches, with the sheer

number of possible dosing regimens making it difficult to identify an optimal regimen. We argue

that this knowledge gap is a major limitation for the improvement of dosing regimens of existing

drugs and a real obstacle for the development of new antibiotics[13,14].

Pharmacodynamic models that can make predictions of bacterial killing and selection on the

basis of drug-target interactions offer new promise to inform rational antibiotic dosing

practices[15]. Recently described models that include drug-target binding have been useful in

gaining a better qualitative understanding of complicated drug effects, such as post-antibiotic

effects, inoculum effects, and bacterial persistence[15-18]. However, to speed the development of

new antibiotics or to inform practices which minimize resistance, we require quantitative

predictions for antibiotics or resistant bacterial strains that do not exist yet. Models which permit

quantitative predictions of changes in drug efficacy as a function of modification of antibiotic

molecules (i.e. new drugs) or novel resistance mutations would be invaluable. Such tools would

advance our general mechanistic understanding of antibiotic action, could guide dosing trials of

new drugs, and suggest better dosing of existing drugs.

In this report, we describe a mechanistic computational modeling framework (COMBAT-

COmputational Model of Bacterial Antibiotic Target-binding) that allows us to predict drug

effects based solely on accessible biochemical parameters describing drug-target interaction.

These parameters can be determined early in drug development. We use this framework to

investigate how changes in drug target binding, either due to improvements in existing

antibiotics or due to resistance mutations in bacteria, affect antibiotic efficacy. We first show that

COMBAT accurately predicts bacterial susceptibility as a function of drug-target binding and,

conversely, allows inference of these biochemical parameters on the basis of observed patterns of

bacterial growth suppression or killing. We then use COMBAT to predict the susceptibility of

newly arising resistant variants based on the molecular mechanism of resistance and determine

the resistance selection window.

**Results**

**Quinolone target affinities correlate with antibiotic efficacy**

To investigate how biochemical changes in antibiotic action modifies bacterial susceptibility, we

explored how the affinity of antibiotics to their target affects the MIC. We compared the MICs of

quinolones, an antibiotic class in which individual antibiotics have a wide range of affinities to

their target, gyrase ($K_D \sim 10^{-4}$ - $10^{-7}$ M) but are of similar molecular sizes and have a similar mode

of action[19]. This choice allowed us to isolate the effects of differences in drug-target affinity on

the MIC.

We obtained binding affinities of quinolones to their gyrase target in *Escherichia coli* from

previous studies[20-24]. We then retrieved MIC data for several quinolones from clinical

Enterobacteriaceae isolates collected before 1990[25], i.e., before the widespread emergence of

quinolone resistance[19]. We assume that quinolone affinities obtained from clinical

Enterobacteriaceae isolates collected before the emergence of resistance correspond to those

measured in wild-type *E. coli*.

To make qualitative predictions of MICs, we employed a simplified model based on the

assumptions that i) drug-target binding occurs much more quickly than bacterial replication, ii)

the antibiotic concentration remains constant and iii) that during the 18 hours of an MIC assay,

the concentration gradient of the drug inside and outside the cell has equilibrated. Under these

assumptions, the MIC can be expressed as

$MIC = K_D \frac{f_c}{1-f_c}$                                             (1)

where $K_D$ represents the affinity constant and $f_c$ the fraction of the target bound at the MIC[26].

Accordingly, this model predicts that the MIC is linearly correlated with $K_D$.

Fig. 1 shows the correlations between drug-target affinities and MICs for seven quinolones and

clinical isolates of 11 different Enterobacteriaceae species. We observed a significant ($p < 0.018$)

linear correlation between MIC and $K_D$ in all species, confirming the qualitative model

prediction.

**A quantitative model to predict antibiotic efficacy**

While it was encouraging that our model can qualitatively predict MIC changes, our aim was to

quantitatively predict antibiotic treatment performance. The simplified model assumes that the

binding kinetics are much faster than bacterial replication, which may not be true in all cases. To

expand the generalizability of the model, we extended the modeling framework to allow that

bacterial replication may occur in a similar time frame as drug-target binding events.

The full model (COMBAT- COmputational Model of Bacterial Antibiotic Target-binding)

describes the binding and unbinding of antibiotics to their targets and predicts how such binding

dynamics affects bacterial replication and death (Fig. 2a). In previous work linking drug-target

binding kinetics with bacterial replication[18], we described a population of bacteria with $\theta$ target

molecules per cell with a system of $\theta + 1$ (bacteria with 0, 1, …, $\theta$ bound target molecules)

ordinary differential equations (ODEs). This system increases in complexity with the number of

target molecules and makes fitting the model to data computationally too demanding for most

settings. To simplify this prior approach, we developed new mathematical models based on

partial differential equations (PDEs), where a single equation describes all bacteria

simultaneously. The sum of bacteria within all target occupancy states over time can be

described by a time kill curve (Fig. 2b), during which the bacterial population is characterized by

the distribution of bacterial cells with different levels of target occupancies at each time-step

(Fig. 2c). This curve can be visualized as a two-dimensional surface in a three-dimensional

coordinate system where the number of bacteria is represented on the z-axis, the percent of

bacteria with the fraction of bound target molecules on the x-axis, and time on the y-axis (Fig.

2d).

Antibiotic action is described by rates of binding ($k_f$) and unbinding ($k_r$) to bacterial target

molecules (Fig. 2a, e). The binding of an antibiotic to a target results in the formation of an

antibiotic-target molecule complex $x$, where $x$ ranges between 0 and $\theta$.

COMBAT consists of two mass balance equations: equation 2 describing bacterial numbers as a

function of bound targets and time and equation 3 describing antibiotic concentration as a

function of time (Methods section).

$$\frac{\partial B(x,t)}{\partial t} + \overbrace{\frac{\partial}{\partial x}(v_B(x,t)B(x,t))}^{\text{Binding kinetics}} = \overbrace{-r(x)B(x,t)F_{lim}(t) + S_B(x,t)F_{lim}(t)}^{\text{Replication and its effects on binding}} - \overbrace{\delta(x)B(x,t)}^{\text{Death}} \quad (2)$$

The term for binding kinetics is given in brown, the term for replication in blue and the term for death in red.

$$\frac{dA(t)}{dt} = -\hat{k}_f A(t) \int_0^\theta (\theta - x)B(x,t)dx + k_r \int_0^\theta xB(x,t)dx \qquad (3)$$

where $v_B = v_f - v_r$, $v_f = \hat{k}_f A(t)(\theta - x)$ and $v_r = k_r x$. $v_B$, $v_f$, and $v_r$ can be seen as a generalized velocity $v = \frac{dx}{dt}$.

Equation 4 (part of the replication term in equation 2) describes how daughter cells inherit bound target molecules from the mother cell during replication:

$$S_B(x,t) = 2\int_x^\theta h(x,z)r(z)B(z,t)dz \,; \, \forall x \in [0,\theta] \qquad (4)$$

Equation 5 (part of the replication term in equation 2) is a logistic growth model describing reduced bacterial replication as the carrying capacity is approached:

$$F_{lim} = \left(1 - \frac{\int_0^\theta B(x,t)dx}{K}\right) \qquad (5)$$

**Model fit to ciprofloxacin time-kill data**

We used the quinolone ciprofloxacin to quantitatively fit bacterial time-kill curves, since this is a commonly used antibiotic for which binding parameters have been directly measured.

Supplementary Tab. S1 gives an overview of the known parameters used for fitting;

Supplementary Tab. S2 gives the parameters resulting from our fit.

The functional relationship between the levels of bacterial replication and death on the fraction

of bound target molecules is extremely hard to obtain experimentally. We therefore treated the

relationships between the fraction of bound target and bacterial replication and death as free

parameters in our model fitting. Ciprofloxacin is considered to have both bacteriostatic and

bactericidal action (mixed action)[27,28], and we fitted functions for a monotonically decreasing

replication and a monotonically increasing killing with each successively bound target molecule

(see Methods & Supplementary Fig. S1).

Overall, we found that COMBAT could fit the time-kill curves well ($R^2 = 0.93$, Fig. 3a). Fig. 3b

shows the predicted bacterial replication $r(x)$ and death as a function of target occupancy $\delta(x)$

based on the fit obtained in Fig. 3a. After model calibration, we simulated bacterial replication

during exposure to different antibiotic concentrations for 18 h. For this simulation, positive

values indicate an increase in the number of bacteria, and negative values indicate a decrease in

the number of bacteria. We estimated a MIC of 0.0139 mg/L (Fig. 3c), a value that is within the

range of MIC determinations for wt *E. coli* (0.01 mg/L, 0.015 mg/L, 0.017 mg/L and 0.023 mg/L

[11,29-31]).

**Accurate prediction of target overexpression from time-kill data**

Having shown that COMBAT can quantitatively fit experimental data on antibiotic action within

biologically plausible parameters, we continued to test the predictive ability of the model. Given

our hypothesis that modifications in antibiotic-target interactions lead to predictable changes in

bacterial susceptibility, we experimentally induced changes in the antibiotic-target interaction of

ciprofloxacin in *E. coli*. We then quantified these biochemical changes by fitting COMBAT to

corresponding time-kill curves and compared them to the experimental results. Ciprofloxacin

acts on gyrase $A_2B_2$ tetramers[19]. We used an *E. coli* strain for which both gyrase A and gyrase B

are under the control of a single inducible promoter ($P_{lacZ}$), such that the amount of gyrase $A_2B_2$

tetramer can be experimentally manipulated[32]. We measured net growth rates for this strain at

different ciprofloxacin concentrations in the presence of 10 μM isopropyl β-D-1-

thiogalactopyranoside (IPTG; mild overexpression) and 100 μM IPTG (strong overexpression)

and compared it to the wild-type in the absence of the inducer (Fig. 4a).

Like previously reported, we find that increasing gyrase content makes *E. coli* more susceptible

to ciprofloxacin[32]. We fitted net growth rates allowing the target molecule content, i.e. gyrase

$A_2B_2$, to vary. We assumed that the only change between the different conditions was the amount

of target. We further assumed that the relationship between bound target and bacterial replication

or death did not differ between the control strain containing a mock plasmid (no IPTG) and the

experiments with overexpression (Fig. 4b, between 0 % and 100 %). Finally, we assumed that

the maximal kill rate at very high antibiotic concentrations was accurately measured in our

experiments and forced the function describing bacterial death through the measured value when

all target molecules are bound. We found the best fit for a 1.31x increase in $GyrA_2B_2$ target

molecule content for bacteria grown in the presence of 10 μM IPTG and a 2.02x increase in

$GyrA_2B_2$ target molecule content for those grown in the presence of 100 μM IPTG.

We subsequently tested these predictions experimentally by analyzing Gyrase A and B content

by western blot Fig. 4c; Supplementary Fig. S2). Using realistic association and dissociation

rates for biological complexes[33], we predicted a range of functional tetramers based on the

relative amount of Gyrase A and B proteins (Fig. 4d). Supplementary Tab. S3 details the

individual measurements, and the procedure to estimate tetramers is provided in the methods

section. We found that the observed overexpression was very close to our theoretical prediction,

with 1.43x [95 % CI 1.19-1.81] overexpression (model prediction = 1.31x overexpression) in the

presence of 10 μM IPTG and 2.15x [95 % CI 1.73-2.87] overexpression in the presence of

100 μM IPTG (model prediction = 2.02x overexpression).

**Accurate prediction of target occupancy at MIC from time-kill data**

Next, we tested whether COMBAT can be applied to the action of the beta-lactam ampicillin, a

very different antibiotic with a distinct mode of action from quinolones. Using published

pharmacodynamic data of *E. coli* exposed to ampicillin[31] also allowed us to compare COMBAT

predictions to established pharmacodynamic approaches. Most of the biochemical parameters for

ampicillin binding to its target, penicillin-binding proteins (PBPs), have been determined

experimentally (Supplementary Tab. S1). Ampicillin is believed to act as a bactericidal drug[34],

and this mode of action is supported by findings from single-cell microscopy[26]. We therefore

assume that ampicillin binding does not affect bacterial replication. In order to model the

consumption of beta-lactams at target inhibition and eventual target recovery, we made small

adjustments to equation 13 (see Methods, description of beta-lactam action).

We fitted COMBAT to published time-kill curves of *E. coli* exposed to ampicillin (Fig. 5a).

Again, COMBAT provides a good fit to the experimental data between 0 min and 40-60 min.

After that time, observed bacterial killing showed a characteristic slowdown at high ampicillin

concentrations which is often attributed to persistence[18] (Fig. 5a). For the sake of simplicity, we

chose to omit bacterial population heterogeneity in this work and therefore cannot describe

persistence, even though COMBAT can be adapted to capture this phenomenon[18]. Because

ampicillin acts in an entirely bactericidal manner, we assume a constant replication rate (see

Methods & Supplementary Fig. S1) and fitted bacterial death as a function of target binding,

$\delta(x)$ (Fig. 5b, fitted parameters in Tab. S4). Fig. 5c shows the predicted net growth rate over a

range of drug concentrations. We estimated a MIC of 2.6 mg/L. This MIC is based on the

Clinical & Laboratory Standards Institute definition of the MIC determined at 18 h. The original

source of the MIC, which was based on experimental data and a pharmacodynamic model[31]

determined an MIC of 3.4 mg/L at 1 h. If we change our prediction to 1 h, our estimated MIC is

3.32 mg/L, which is within 2.5 % of the reported value[31].

Having established that COMBAT can also adequately capture the pharmacodynamics of

ampicillin, we next tested whether we can estimate experimentally determined target occupancy

at the MIC. Our estimated mean occupancy considering both living and dead bacteria is 89 %

(Fig. 5b), a value within previously reported experimental estimates from *Staphylococcus aureus*

(84-99 %)[35].

**Sensitivity of antibiotic efficacy to parameters of drug-target binding**

It is possible to vary all parameters in COMBAT and explore their effect. We used this to test

how hypothetical chemical changes to ampicillin or ciprofloxacin would affect antibiotic

efficacy (Supplementary Fig. S3-S11). These changes could reflect either bacterial resistance

mutations or modifications of the antibiotics themselves. We predict that changes in drug-target

affinity, $K_D$, have more profound effects than changes in target molecule content, bacterial

reaction to increasingly bound target (i.e. $\delta(x)$ and $r(x)$), or changes in target molecule content.

We also predict that the individual binding rates $k_r$ and $k_f$, and not just the ratio of these terms,

the $K_D$, are important factors in efficiency. The faster a drug binds, the more efficient we

predicted it will be. One intuitive explanation for the observation that $k_f$ drives efficacy is that a

slow binding fails to rapidly interfere with bacterial replication, which may allow for the

production of additional target molecules and thereby reduce the ratio of free antibiotic to target

molecules.

**Forecasting the resistance selection window**

Finally, we illustrate how COMBAT can be used to explore how the molecular mechanisms of

resistance mutations affect antibiotic concentrations at which resistance can emerge, i.e., the

resistance selection window. We compared predicted net growth rates as a function of

ciprofloxacin concentrations for a wild-type strain and an archetypal resistant strain. For this

analysis, we assumed that the resistant strain has a 100x slower drug-target binding rate (i.e.

~100x increased MIC, realistic for novel point mutations[36]) and that the maximum replication

rate of the resistant strain is 85 % of the wild type strain[37]. We then predicted the antibiotic

concentrations at which resistance would be selected. Interestingly, when comparing COMBAT

to previous pharmacodynamics models (Fig. 5), we observed that estimates of replication rates

depend on the selected time frame (Fig. 6a). When the timeframe for MIC determination is set to

18 h as defined by CLSI[38], the "competitive resistance selection window", i.e., the concentration

range below the MIC of both strains where the resistant strain is fitter than the wild type, ranges

from 0.002 mg/L to 0.014 mg/L for ciprofloxacin (Fig. 6a) and 1 mg/L to 2.6 mg/L for

ampicillin (Supplementary Fig. S12), respectively. This corresponds well with previous

observations that ciprofloxacin resistance is selected for well below MIC[11]. However, when

measuring after 15 min or 45 min, the results are substantially different. The reason for this is

illustrated in Fig. 6b. COMBAT reproduces non-linear time kill curves where bacterial

replication continues until sufficient target is bound to result in a negative net growth rate. This

compares well with experimental data around MIC in Fig. 3a and 5a. In Fig. 6b, we show model

predictions for ciprofloxacin concentrations corresponding to a zero net growth (i.e. same

population size) after 15 min, 45 min and 18 h ($MIC_{Resistant; 15 min}$, $MIC_{Resistant; 45 min}$,

$MIC_{Resistant; 18 h}$). In all cases, the bacterial population first increases and then decreases slowly.

This may have consequences for the selection of resistant strains. Fig. 6c illustrates how the

resistance selection windows depending on the observed time frame. This suggests that even at

concentrations above the 18 h MIC of the resistant strain, there may be initial growth of the

resistant strain. In this case, the resistant strain could continue growing at concentration of up to

7 mg/L ciprofloxacin at 15 min, even though the MIC at 18 h is 1.27 mg/L.

**Discussion**

Optimizing dosing levels of antibiotics is important for maximizing drug efficacy against wild-

type strains as well as for minimizing the rise of resistant mutants. The determination of optimal

dosing strategies typically requires expensive empirical studies; the need for such studies arises

in part from our currently limited capacity to predict how antibiotics will affect bacteria at a

given concentration. In fact, drug attrition is mainly due to insufficient predictions of efficacy

(pharmacodynamics) rather than pharmacokinetics[6]. For optimizing drug development and for

minimizing resistance, we need quantitative predictions for antibiotics or resistant bacterial

strains that do not exist yet. The ability to accurately predict MICs on the basis of biochemical

parameters and, more generally, to define antibacterial activity across a range of drug

concentrations, would allow us to estimate antibiotic efficacy for novel compounds or against not

yet emerged resistant strains[15,39]. Recent studies have reported methods to predict MICs from

whole genome sequencing data[40,41]. However, these methods require transfer of prior knowledge

on how the resistance mutations affect MICs in other organisms. There are no methods that could

predict *a priori* how chemical changes to an antibiotic structure or novel resistance mutations

affect bacterial growth at a given antibiotic concentration.

Here, we accurately predict antibiotic action on the basis of accessible biochemical parameters of

drug-target interaction. Our computational model, COMBAT provides a framework to predict

the efficacy of compounds based on drug-target affinity, target number, and target occupancy.

These parameters may change both when improving antibiotic lead structures as well as when

bacteria evolve resistance. Importantly, they can be measured early in drug development and

may even be a by-product of target-based drug discovery[42]. When these data are available,

COMBAT makes only one assumption: that the rate of bacterial replication decreases and/or the

rate of killing increases with successive target binding. While fitting, we allow this relationship

to be gradual or abrupt and select the best fit. This means we do not model specific molecular

mechanisms down-stream of drug-target binding, but their effects are subsumed in the functions

that connect the kinetic of drug-target binding to bacterial replication and death.

In previous work, for example on antipsychotics[16], antivirals[17] and antibiotics[15,18], models of

drug-target binding kinetics have been used to improve our qualitative understanding of

pharmacodynamics. Our study substantially advances this work by making accurate quantitative

predictions across antibiotics and bacterial strains when measurable biochemical characteristics

change. This is possible because COMBAT employs an elegant mathematical approach, based

on partial differential equations, that makes it computationally feasible to fit the model to a large

range of data. Importantly, we are not only able to predict antibiotic action from biochemical

parameters, but can also vice versa use COMBAT to accurately predict biochemical changes

from observed patterns of antibiotic action. We have confirmed the excellent predictive power of

COMBAT with clinical data as well as experiments with antibiotics with very different

mechanisms of action. This gives us confidence that biochemical parameters are major

determinants of antibiotic action in bacteria and that COMBAT helps to make rational decisions

about antibiotic dosing.

In drug development, our mechanistic modeling approach provides insight into which chemical

characteristics of drugs may be useful targets for modification. For example, our sensitivity

analyses indicate that antibiotics with a similar affinity but faster binding inactivate bacteria

more quickly and therefore prevent replication and production of more target molecules, which

would change the ratio of antibiotic to target. Furthermore, because e.g. antibiotic binding and

unbinding rates can be determined early in the drug development process, such insight can help

the transition to preclinical and clinical dosing trials. This may contribute to reducing bottlenecks

between these phases of drug development and thereby save money and time.

Avoiding antibiotic concentrations that select for resistance is challenging because the precise

concentrations are only known after extensive experiments have been performed that identify the

MIC of (nearly) all possibly emerging resistant mutants. Predicting the resistance selection

windows of novel resistant mutants on the basis of biologically plausible changes in drug-target

binding would allow us to better assess what drug concentrations need to be achieved to avoid

selection of resistance. This approach offers new promise to assess resistance risks prior to

characterizing the majority of resistance mutations and thereby reduce the failure rates of

candidate compounds late in the drug development process when resistance is observed in

patients and substantial resources have been invested.

Our approach also offers insight into determinants of the resistance selection window. Rather

than determining the resistance selection window for a comprehensive collection of possibly

arising resistance mutations in each bacteria-drug pair, it would be attractive to build

transferrable knowledge that allows estimating the resistance selection window. In concordance

with a recent meta-analysis of experimental data[43], our sensitivity analyses predict that changes

in drug target binding and unbinding have a greater impact on the MIC than changes in target

molecule content or down-stream processes. Thus, a more comprehensive characterization of the

binding parameters of spontaneous resistant mutants would allow an overview of the maximal

biologically plausible levels of resistance that can arise with one mutation. Dosing above this

level should then safeguard against resistance. This is especially useful for compounds for which

it is difficult to saturate the mutational target for resistance, or for safeguarding against resistance

to newly introduced antibiotics for which we do not yet have a good overview of resistance

conferring mutations. If toxicity, solubility or other constraints do not allow dosing above the

MIC of expected resistant strains, COMBAT can predict the concentration range at which

resistance is less strongly selected. This could guide decisions on treating with low versus high

doses, which is currently controversially debated[7,8]. Good quantitative estimates on the dose-

response relationship of new drugs would also help defining the therapeutic window, i.e. the

range of drug concentrations at which the drug is effective but not yet toxic.

Our quantitative work can help to identify optimal dosing strategies at constant antibiotic

concentrations for homogeneous bacterial populations. These measures are commonly used to

assess antibiotic efficacy. In addition, previous work has demonstrated that drug-target binding

models can qualitatively describe antibiotic efficacy over the fluctuating concentrations that

actually occur in patients[26,44]. They can also explain complicated phenomena such as biphasic

kill curves, the post-antibiotic effect, or the inoculum effect[15,18,45] that often complicate the

clinical phase of drug development. COMBAT has similar characteristics that allow capturing

these complex phenomena. Therefore, employing COMBAT may be useful for guiding drug

development to maximize antibiotic efficacy and minimize *de novo* resistance evolution.

**Methods**

*Mathematical model*

COMBAT incorporates the binding and unbinding of antibiotics to their targets and describes

how target binding affects bacterial replication and death. This work extends the model

developed in[18]. COMBAT consists of a system of two mass balance equations: one PDE for

bacteria (describing replication and death as a function of both time and target binding) and one

ODE for antibiotic molecules (describing the concentrations as function of time).

In the most basic version of COMBAT, we ignored differences between extracellular and

intracellular antibiotic concentrations and only followed the total antibiotic concentration $A$,

assuming that the time needed for drug molecules to enter bacterial cells is negligible. We model

ciprofloxacin (to which there is a limited diffusion barrier[46]) and ampicillin (where the target is

not in the cytosol, even though the external membrane in gram negatives has to be crossed to

reach PBPs). We therefore believe that this assumption is justified in wild-type *E. coli*. This

basic version of COMBAT is therefore more accurate for describing antibiotic action where the

diffusion barrier to the target is weak.

*Binding kinetics*

We describe the action of antibiotics as a binding and unbinding process to bacterial target

molecules[18]. For simplicity, we assume a constant number of available target molecules $\theta$. The

binding process is defined by the formula $A + T \rightleftharpoons x$, where the intracellular antibiotic

molecules $A$ react with target molecules T at a rate $k_f$ and form an antibiotic-target molecule

complex x, where values for x range between 0 and $\theta$. If the reaction is reversible, the complex

dissociates with a rate $k_r$.

In[18], the association and dissociation terms are described by the following terms

$$\frac{dB_i(t)}{dt} = \overbrace{\hat{k}_f A(t)\big((\theta - i + 1)B_{i-1}(t) - (\theta - i)B_i(t)\big)}^{\text{Association term}} -$$

$$\underbrace{k_r\big(iB_i(t) - (i+1)B_{i+1}(t)\big)}_{\text{Dissociation term}}; \ i\epsilon[0,\theta] \tag{6}$$

where $\hat{k}_f = \frac{k_f}{V_{tot} n_A}$, $k_f$ is the association rate, $V_{tot}$ is the volume in which the experiment is

performed, $n_A$ is Avogadro's number, $k_r$ is the dissociation rate, $B_i$ is the number of bacteria with

$i$ bound targets, and $\theta$ is the total number of targets. Green denotes the association term, while

the dissociation term is in orange.

This approach requires the use of a large number of ordinary differential equations, $(\theta + 1)$ for

the bacterial population and one for the antibiotic concentration. To generalize this approach, we

assume that the variable of bound targets is a real number $x \in \mathcal{R}$. Under this continuity

assumption, we consider the bacterial cells as a function of $x$ and the time $t$, thereby reducing the

total number of equations to two.

Under the continuity approximation $(x \in \mathcal{R})$, we can rewrite the binding kinetics in the form

$$\frac{\partial B(x,t)}{\partial t} = \overbrace{\frac{\partial}{\partial x}\left(\hat{k}_f A(t)(\theta - x)B(x,t)\right)}^{\text{Association term}} - \overbrace{\frac{\partial}{\partial x}\left(k_r x B(x,t)\right)}^{\text{Dissociation term}} \qquad (7)$$

or simply

$$\frac{\partial B(x,t)}{\partial t} = \frac{\partial}{\partial x}\left(v_f(x,t)B(x,t) - v_r(x,t)B(x,t)\right) \qquad (8)$$

where $v_f = \hat{k}_f A(t)(\theta - x)$ and $v_r = k_r x$ can be considered as two velocities, i.e., the derivative

of the bound targets with respect to the time $\frac{dx}{dt}$. Green denotes the association term, while the

dissociation term is in orange.

*Replication rate*

We assume that the replication rate of bacteria, $r(x)$, is dependent on the number of bound target

molecules $x$. The function $r(x)$ is a monotonically decreasing function of $x$, such that fewer

bacteria replicate as more target is bound. $r(0)$ is the maximum replication rate, corresponding to

the replication rate of bacteria in absence of antibiotics. Thus, $r(x)$ describes the bacteriostatic

action of the antibiotics, i.e., the effect of the antibiotic on bacterial replication.

*Carrying capacity*

Replication ceases as the total bacterial population approaches the carrying capacity $K$. At that

point, the replication term of the equation is

$$\frac{\partial B(x,t)}{\partial t} = r(x)B(x,t)\frac{K - \int_0^\theta B(x,t)dx}{K} = r(x)B(x,t)F_{lim} \qquad (9)$$

where $F_{lim} = \frac{K - \int_0^\theta B(x,t)dx}{K}$ is the replication-limiting term due to the carrying capacity $K$, and

$0 \leq F_{lim} \leq 1$.

*Distribution of target molecules upon division*

We assume that the total number of target molecules doubles at replication, such that each

daughter cell has the same number as the mother cell. We also assume that the total number of

drug-target complexes is preserved in the replication and that the distribution of $x$ bound target

molecules of the mother cell to its progeny is described by a hypergeometric sampling of $n$

molecules from $x$ bound and $2\theta - x$ unbound molecules. Under the continuity assumption, we

generalize the concept of hypergeometric distribution. Because the hypergeometric distribution

is a function of combinations and because a combination is defined as function of factorials, we

can use $\Gamma$ functions in place of factorials and redefine a continuous hypergeometric distribution

as a function of $\Gamma$ functions. A $\Gamma$ function is

$\Gamma(\zeta) = \int_0^\infty x^{\zeta-1} e^{-x} dx; \ Re(\zeta) > 0$         (10)

where $\zeta$ is a complex number. In this way, the distribution can be expressed as a probability

density function of continuous variables. The amount of newborn bacteria is given by the term

$r(x)B(x,t)F_{lim}(t)$. We assume that bound target molecules are distributed randomly between

mother and daughter cells, with each of them inheriting 50% upon division on average. This

means that twice the amount of newborn cells must be redistributed along $x$ to account for the

random distribution process. For example, if a mother cell with 4 bound targets divides, we have

two daughter cells, each with a number of bound targets between 0 and 4 (their sum has to be 4),

following the generalized hypergeometric distribution. For simplicity, we define $S(x,t)$ to be a

function related to the replication rate that depends on the number of bacteria with a number of

bound target molecules ranging between $x$ and $\theta$, their specific replication rate $r(x)$, and the

fraction of their daughter cells expected to inherit $x$ antibiotic-target complexes $h(x,z)$:

$S(x,t) = 2 \int_x^\theta h(x,z)r(z)B(z,t)\, dz$         (11)

*Death rate*

The death rate function $\delta(x)$ depends on the number of bound target molecules. The function

$\delta(x)$ is assumed to be a monotonically increasing function of $x$, where $\delta(\theta)$ is the maximum

death rate, when all targets in the bacteria have been bound by antibiotics. The shape of this

function describes the bactericidal action of the antibiotic.

*Bacteriostatic and bactericidal effects*

We consider several potential functional forms of the relationship between the percentage of

bound targets and replication and death rates, because the exact mechanisms how target

occupancy affects bacteria is unknown (Supplementary Fig. S1). We use a sigmoidal function

that can cover cases ranging from a linear relationship to a step function. When the inflection

point of a sigmoidal function is at 0 % or 100 % target occupancy, the relationship can also be

described by an exponential function. We assume that replication in bactericidal and death in

bacteriostatic drugs is independent of the amount of bound target. With sufficient experimental

data, the replication rate $r(x)$ and/or the death rate $\delta(x)$ can be obtained by fitting COMBAT to

time-kill curves of bacterial populations after antibiotic exposure. The sigmoidal shape of $r(x)$

and $\delta(x)$ can be written as:

$$r(x) = \frac{r_0}{1+e^{\gamma_r(x-x_{rth})}}; \; \delta(x) = \frac{d_{max}}{1+e^{-\gamma_d(x-x_{dth})}} \hspace{2cm} (12)$$

where $x_{rth}$ is the replication rate threshold, $x_{dth}$ is the death rate threshold, and both represent the

point where the sigmoidal function reaches ½ of its maximum. $\gamma_r$ and $\gamma_d$ represent the shape

parameters of the replication and death rate functions, respectively. These factors determine the

steepness around the inflection point. When they are extreme, the relationship approaches a

linear or a step function.

*Full equation describing bacterial population*

Putting these components together, the full equation describing a bacterial population is:

$$\frac{\partial B(x,t)}{\partial t} + \overbrace{\frac{\partial}{\partial x}\Big(\textcolor{green}{v_f(x,t)B(x,t)} - \textcolor{orange}{v_r(x,t)B(x,t)}\Big)}^{\text{Binding kinetics}} =$$

$$\overbrace{\textcolor{blue}{-r(x)B(x,t)F_{lim}(t) + S_B(x,t)F_{lim}(t)}}^{\text{Replication and its effects on binding}} - \overbrace{\textcolor{red}{\delta(x)B(x,t)}}^{\text{Death}} \qquad (13)$$

where $B(x,t)$ is the number of bacteria. As in equations 2, 6, 7 and 8, green denotes the binding

term, orange the unbinding term (together the binding kinetics is given in brown), blue the term

describing bacterial replication and red the term describing bacterial death.

*Equation describing antibiotic concentration*

The free antibiotic concentration results from mass conservation, i.e., all antibiotic molecules

associating with their target are subtracted and all dissociating antibiotic molecules are added.

Equation 3 in the results section describes the dynamics of the antibiotic concentration.

*Description of beta-lactam action*

Beta-lactams acetylate their target molecules (PBPs) and thereby inhibit cell wall synthesis. The

acetylation of PBPs consumes beta-lactams. However, PBPs can recover through deacetylation.

We modified the term of drug-target dissociation in the equation describing antibiotic

concentrations (equation 3), and set the unbinding rate $k_r = 0$. To reflect the recovery of target

molecules, we substituted the dissociation rate $k_r$ in the equation describing the bacterial

population with the deacetylation rate $k_a$, as described in[26].

*Initial and boundary conditions*

At $t = 0$, we assume that all bacteria have zero bound targets ($x = 0$), and the initial

concentration of bacteria is $B(x, 0) = 0, x > 0$, and $B(0,0) = B_0$.

At the boundaries of the partial differential equation ($x = 0, x = \theta$), we specify that the outgoing

velocities are zero. For $x = 0$, i.e. no bound target molecules, the unbinding velocity $v_r(0, t) =$

0, and in $x = \theta$, i.e. all targets are bound, the binding velocity $v_f(\theta, t) = 0$. When the

replication term at $x = 0$ and the death term at $x = \theta$ are known, we can solve the partial

differential equation with two ordinary differential equations at the boundaries. They are similar

to the equations at $x = 0$ and at $x = \theta$ described by Abel zur Wiesch et al.[18], but taking into

account that x is a continuous variable instead of a natural number.

*Numerical schemes*

To solve our system of differential equations, we used a first-order upwind scheme. Specifically,

we used the spatial approximation $u_-^f = \frac{u(i)-u(i-1)}{\Delta x}$ for the binding term ($v_f > 0$) and the spatial

approximation $u_+^f = \frac{u(i+1)-u(i)}{\Delta x}$ for the unbinding term ($v_r < 0$). For the time approximation of

both the PDEs and the ODEs, we used the forward approximation $\frac{\Delta B}{\Delta t} = \frac{B^{n+1}-B^n}{\Delta t}$[47]. We also

verified that the Courant-Friedrichs-Lewy condition is satisfied. For fitting the experimental data

of bacteria exposed to ciprofloxacin and ampicillin, we used the particle swarm method

("particleswarm" function in Matlab, MathWorks software).

*Concentrations of gyrase $A_2B_2$ tetramers*

We assumed that gyrases A and B first homo-dimerize to $A_2$ and $B_2$, respectively, which in turn

bind to each other to form the tetramer TR[48]. The following system of equations describes their

binding kinetics:

$$
552 \quad
\begin{cases}
\frac{dA}{dt} = -2k_1 A^2 + 2k_{-1} A_2 \\
\frac{dB}{dt} = -2k_2 B^2 + 2k_{-2} B_2 \\
\frac{dA_2}{dx} = k_1 A^2 - k_{-1} A_2 - k_3 A_2 B_2 + k_{-3} TR \\
\frac{dB_2}{dt} = k_2 B^2 - k_{-2} B_2 - k_3 A_2 B_2 + k_{-3} TR \\
\frac{dTR}{dt} = k_3 A_2 B_2 - k_{-3} TR
\end{cases}
\tag{14}
$$

$$
553 \quad
\begin{cases}
A + A \underset{k_{-1}}{\overset{k_1}{\rightleftharpoons}} A_2 \\
B + B \underset{k_{-2}}{\overset{k_2}{\rightleftharpoons}} B_2 \\
A_2 + B_2 \underset{k_{-3}}{\overset{k_3}{\rightleftharpoons}} TR
\end{cases}
\tag{15}
$$

First, we calibrated the model to ensure that we obtain the correct number of gyrase $A_2 B_2$

tetramers (~100) per wild type bacterial cell[49,50]. This results in an average of each 206 gyrase A

and B monomers. Because the association and dissociation rates of the dimers and tetramers are

unknown, we sampled $10^4$ sets of six parameters in equation 14 ($k_{-3}, \dots k_3$) in a Latin hypercube

approach from a biologically plausible range where the association rates are between $10^7$ - $10^9$

$M^{-1} s^{-1}$ and the dissociation rates between $10^{-3}$ - $10^{-1}$ $s^{-1}$ [33]. This results in $10^4$ estimates for each of

the six experimental replicates quantifying gyrase A and B (Fig. 4, Supplementary Fig. S2,

Supplementary Tab. S3).

***Experimental methods***

*Strains, growth conditions and strain construction*

*Escherichia coli* strain BW25113[51] (SoA2740) was transformed with plasmids pCA24N-SC101-

gyrAB[32] and pCA24N-SC101-ΔP-YFP[32] using electroporation, resulting in strains

BW25113/pCA24N-SC101-gyrAB (SoA3329) and BW25113/pCA24N-SC101-ΔP-YFP

(SoA3330), respectively. pCA24N-SC101-gyrAB encodes the *E. coli gyrAB* genes under control

of the IPTG inducible LacZ promoter. pCA24N-SC101-ΔP-YFP encodes a promoterless copy of

YFP and was used as a control. Bacteria were grown at 30°C on either LB agar or in LB broth,

both supplemented with 10 µg/mL chloramphenicol (Cm) and 10 µM (mild induction) or 100

µM (strong induction) of isopropyl β-D-1-thiogalactopyranoside (IPTG) (43714 5X, VWR

Chemicals) when necessary.

*Time-kill curves*

Overnight cultures of BW25113 or SoA3329 and SoA3330 were diluted 1:1000 in pre-warmed

LB or LB-Cm and LB-Cm-IPTG, respectively, and grown with shaking to $OD_{600}$ ~0.5. A 1:3

dilution series of ciprofloxacin was made and added to the cultures at indicated concentrations.

Additional cultures without antibiotics and with a very high concentration of ciprofloxacin

(2.187 mg/L) were used to determine the minimal and maximal kill rate, respectively. Samples

were taken immediately prior to addition of the antibiotic and in ~20 min intervals or after 45

min, respectively. Samples were washed once in phosphate buffered saline (PBS) before colony

forming units (CFUs) were determined for each sample by plating a 1:10 dilution series in PBS

on LB agar plates.

*GyrAB quantification*

To quantify the relative amount of GyrAB, samples of SoA3329 and SoA3330 were collected

after 45 min of drug treatment as described above. An equal number of cells corresponding to 1

mL culture at $OD_{600}$ = 1 were harvested by centrifugation. Pelleted cells were lysed at room

temperature for 20 min using B-PER bacterial protein extraction reagent (90078, Thermo

Scientific) supplemented with 100 µg/mL lysozyme, 5 units/mL DNaseI (all part of B-PER™

with Enzymes Bacterial Protein Extraction Kit, 90078, Thermo Scientifc) and 100 µM/mL

PMSF (52332, Calbiochem). Samples were stored at -80°C until further use.

Samples were heated to 70°C for 10 min after addition of 1x Bolt sample reducing agent (B0009,

Life Technologies) and 1x fluorescent compatible sample buffer (LC2570, Invitrogen). Proteins

in whole-cell lysates were separated on 4-15 % Mini-Protean TGX Precast gels (456-1085, Bio-

Rad) and transferred to 0.2 µm Nitrocellulose membranes (1704158, Bio-Rad).

Membranes were blocked in Odyssey blocking buffer-TBS (927-50000, Li-Cor) for at least one

hour at room temperature. Primary antibodies raised against GyrA (Rabbit α-Gyrase A, PA005,

Inspiralis), GyrB (Rabbit α-Gyrase B , PB005, Inspiralis), and CRP (Mouse α-E. coli CRP,

664304, Nordic Biosite antibodies) were diluted 1:250, 1:250, and 1:2,000 in Odyssey blocking

buffer-TBS, respectively. The blocked membranes were incubated with the appropriate primary

antibodies overnight at 4°C, washed 4x for 15 min each in TBS-T solution (Tris buffered saline

supplemented with Tween20: 0.138 M sodium chloride, 0.0027 M potassium chloride, 0.1 %

Tween20, pH 8.0 at 25°C), and incubated for 2 h at room temperature with fluorescent labelled

secondary antibodies (1:10,000 of IRDye® 680RD Goat anti-Mouse IgG, P/N 925-68070, Li-

Cor and 1:5000 of IRDye® 800CW Goat anti-Rabbit IgG, P/N 925-32211, Li-Cor) in Odyssey

blocking buffer-TBS. Finally, the membranes were washed 4x for 15 min each in TBS-T

solution and imaged at 700 nm and 800 nm using a Li-Cor Odyssey Sa scanning system.

Band intensities were quantified from unmodified images using the record measurement tool of

Photoshop CS6, normalized to the CRP loading control after background subtraction, and

reported relative to SoA3330. For clarity, the "levels" tool of Photoshop CS6 was used to

enhance the contrast of shown Western blot images.

**Data Availability**

Computer code will be available at https://www.abel-zur-wiesch-lab.com/.

**References**
620

**Acknowledgements**

We thank Giovanni Montani, Forrest W. Crawford, Roberto Natalini, Klaus Harms, Christoph

Zimmer, Angelo Vannozzi, Jingyi Liang, Vi Tran, Antal Martinecz and Rafal Mostowy for

helpful discussions and feedback on the manuscript. We would like to acknowledge R. Regoes

for providing previously published raw data. This work was funded by Bill and Melinda Gates

Foundation Grant OPP1111658 (to T.C. & P.AzW.), Research Council of Norway (NFR) Grant

262686 (to P.AzW.) and 249979 (to S.A.), and Helse-Nord Grant 14796 (to S.A.).

**Author Contributions**

P.AzW. designed the study. F.C. developed the mathematical models. A.P., B.S., M.S., and S.A.

designed the experiments. A.P., B.S., M.S., and S.L. performed the experiments. F.C., A.P., B.S.,

M.S., T.C., S.A., and P.AzW. analyzed the data. T.C., S.A., and P.AzW. wrote the manuscript.

**Competing Interests statement**

The authors declare no competing interests.

**Fig. 1| Clinical data confirm linear correlation between MICs and affinities of quinolones to gyrase.** We analyzed MIC and drug-target affinity data from 11 Enterobacteriaceae isolates and seven different quinolones. The x-axes show the affinities ($K_D$), and the y-axes show the MICs, both in mol/L. The adjusted $R^2$ and p-value of each correlation are given. In cases where there was more than one $K_D$ value reported in the literature, we used the mean for this analysis. The tested MIC values are the median of several clinical isolates described previously[25].

**Fig. 2| Illustration of modeling approach. a**, Schematic illustration of binding kinetics (adapted

from[52]). The grey triangles depict the drug target molecules, and the orange circles represent

antibiotic molecules within bacteria. The arrows indicate individual binding and unbinding

events of the antibiotic to its target molecule in the cell. $\hat{k}_f$ is the adjusted forward reaction rate,

$k_r$ is the reverse reaction rate, $A$ is the concentration of antibiotics inside the bacterium, $x$ is the

number of bound targets, $\theta$ is the number of targets and $B_x$ is the number of bacteria with $x$ bound

targets. **b**, Modeled sample time-kill curve, in which the sum of bacteria in all binding states (i.e.,

the entire population of living bacteria) is followed over time after exposure to antibiotics. The

vertical dotted lines indicate the time points depicted in (**c**); 1 min (grey), 14 min (yellow), and

80 min (purple). **c**, The percentage of bound antibiotic targets in the bacterial population at

indicated time points. **d**, Illustration of how the partial differential equation describes the

bacterial population as a surface in a three-dimensional coordinate system, the dimensions of

which represent percent bound target (x-axis), time (y-axis), and number of bacteria (z-axis). The

three time points shown in (**c**) represent two-dimensional cross-sections at different points of the

y-axis. **e**, Overview of used parameters and functions.

**Fig. 3| Model predictions for the MIC and the bacteriostatic and bactericidal effects of ciprofloxacin. a**, Model fit to experimental time-kill curves. The points indicate the experimental data of three independent replicates, and the lines indicate the model fit. Each color indicates a ciprofloxacin concentration as reported in the figure. **b**, The blue line indicates the bacteriostatic effect ($r(x)$, replication rate) of ciprofloxacin and the red line the bactericidal effect ($\delta(x)$, death rate) as a function of the number of bound targets predicted by the model fit in (**a**).

666    The values of the fitted parameters are listed in Supplementary Tab. S2. **c**, The net growth rate as

667    determined by the slope of a line connecting the initial bacterial density and the final bacterial

668    density of a time-kill curve at 18 h on a logarithmic scale, is given as function of the drug

669    concentration (blue). The dotted horizontal line indicates zero net growth, and the intersection

670    with the blue line predicts the MIC (0.0139 mg/mL).

**Fig. 4| Prediction of relative antibiotic target molecule content from time-kill curves. a**,

Dose-response curves of *E. coli* expressing *gyrA* and *gyrB* under the same IPTG-inducible

promoter (SoA3329) grown in the presence of 10 μM IPTG (mild overexpression; red) and 100

μM IPTG (strong overexpression; yellow). A control strain (SoA3330), which expresses wild-

type GyrAB levels and contains a mock plasmid, is grown in the absence of inducer (blue). The

x-axis indicates the ciprofloxacin concentration, and the y-axis indicates the fold change in

colony forming units over time. The dotted lines indicate experimental data, and the solid lines

indicate the model fit. The best model fit was obtained for relative target molecule contents of

131 % (mild overexpression) and 202 % (strong overexpression) relative to the control strain

(WT). **b**, Death rates of *E. coli* expressing different levels of GyrAB. The colors represent

GyrAB expression conditions as in (**a**). The x-axis shows the percentage of bound antibiotic

target normalized to the control strain; the y-axis shows the death rate $\delta(x)$. Each line represents

the best fit for $\delta(x)$. **c**, Western blot analysis of GyrA&B in the strains/conditions shown in (**a**).

CRP (cAMP receptor protein) was used as loading control. A representative example of six

replicates is shown; see Supplementary Fig. S2 for full blots. **d**, comparison of theoretical

prediction (from (**b**), solid colors) and $GyrA_2B_2$ tetramer levels estimated from relative GyrA&B

monomer levels (quantified in (**c**), translucent colors). For the experimental measurements, the

bars indicate the mean, and the whiskers represent the 95 % confidence interval.

**Fig. 5| Model prediction of MIC and target occupancy at MIC for ampicillin. a**, Model fit to

previously published time-kill curves[31]. The points represent experimental data, and the lines

represent the fit of the model. Each color indicates a single ampicillin concentration, as described

in the legend. **b**, Replication (blue) and death (red) rates as a function of the number of bound

targets predicted by the model fit in (**a**). The black line indicates the predicted distribution of

target occupancies in a bacterial population (both living and dead cells) exposed to ampicillin at

697    the MIC for 18 h. **c**, The net growth rate, as determined by the slope of a line connecting the

698    initial bacterial density and the bacterial density at 18 h on a logarithmic scale predicted from the

699    model fit in (**a**), is shown as function of the drug concentration (blue). The dotted horizontal line

700    indicates zero net growth, and the intersection with the blue line predicts the MIC (2.6 mg/mL).

**Fig. 6| Predicted mutation selection windows for *E. coli* exposed to ciprofloxacin. a**, The

drug concentration of ciprofloxacin is shown on the x-axes, and the average bacterial net growth

rate in the first 15 min (grey panel), 45 min (yellow panel), and 18 h (purple panel) of exposure

is given on the y-axes. The blue line represents the wild-type strain based on the fits shown in

Fig. 3, and the red line represents a strain with a hypothetical resistance mutation that decreases

the binding rate ($k_f$) 100-fold and imparts a 15 % fitness cost. The horizontal dotted line indicates

no net growth. The vertical dotted line indicates where the resistant strain becomes more fit than

the wild-type, the solid vertical line indicates the MIC of the wild-type, and the dashed vertical

line indicates the MIC of the resistant strain. **b**, Modeled time kill curves of the resistant strain

for ciprofloxacin concentrations at which there is no growth at 15 min (grey; $MIC_{15\ min} = 7$

mg/L), 45 min (yellow; $MIC_{45\ min} = 3$ mg/L) and 18 h (purple; $MIC_{18\ h} = 1.27$ mg/L). The

horizontal dotted line indicates the initial population size; the vertical dotted lines represent the

time points at which the initial and final population size is the same. **c**, The mutation selection

window depends on the time at which bacterial growth is observed. The x-axis shows the

observed time at which replication rates were determined, the y-axis shows ciprofloxacin

concentrations. The dotted curve shows the ciprofloxacin concentration at which the resistant

becomes fitter than the WT ($Fitness_{Resistant} > Fitness_{WT}$), the solid line the MIC of the WT

($MIC_{WT}$), and the dashed line the MIC of the resistant strain ($MIC_{Resistant}$). The area between the

dotted and dashed line indicates the competitive resistance selection window.

1      Boeree, M. J. *et al.* A dose-ranging trial to optimize the dose of rifampin in the treatment of
tuberculosis. *Am J Respir Crit Care Med* **191**, 1058-1065, doi:10.1164/rccm.201407-1264OC
(2015).
2      Lan, A. J., Colford, J. M. & Colford, J. M., Jr. The impact of dosing frequency on the efficacy
of 10-day penicillin or amoxicillin therapy for streptococcal tonsillopharyngitis: A meta-
analysis. *Pediatrics* **105**, E19 (2000).
3      Roord, J. J., Wolf, B. H., Gossens, M. M. & Kimpen, J. L. Prospective open randomized
study comparing efficacies and safeties of a 3-day course of azithromycin and a 10-day
course of erythromycin in children with community-acquired acute lower respiratory tract
infections. *Antimicrob Agents Chemother* **40**, 2765-2768 (1996).
4      Van Deun, A., Salim, M. A., Das, A. P., Bastian, I. & Portaels, F. Results of a standardised
regimen for multidrug-resistant tuberculosis in Bangladesh. *Int J Tuberc Lung Dis* **8**, 560-567
(2004).
5      WHO. The shorter MDR-TB regimen.
http://www.who.int/tb/Short_MDR_regimen_factsheet.pdf?ua=1 (2016).
6      Waring, M. J. *et al.* An analysis of the attrition of drug candidates from four major
pharmaceutical companies. *Nat Rev Drug Discov* **14**, 475-486, doi:10.1038/nrd4609 (2015).
7      Kouyos, R. D. *et al.* The path of least resistance: aggressive or moderate treatment? *Proc Biol*
*Sci* **281**, 20140566, doi:10.1098/rspb.2014.0566 (2014).
8      Read, A. F., Day, T. & Huijben, S. The evolution of drug resistance and the curious
orthodoxy of aggressive chemotherapy. *Proc Natl Acad Sci U S A* **108 Suppl 2**, 10871-10877,
doi:10.1073/pnas.1100299108 (2011).
9      Colijn, C. & Cohen, T. How competition governs whether moderate or aggressive treatment
minimizes antibiotic resistance. *Elife* **4**, doi:10.7554/eLife.10559 (2015).
10     Drlica, K. & Zhao, X. Mutant selection window hypothesis updated. *Clin Infect Dis* **44**, 681-
688, doi:10.1086/511642 (2007).
11     Gullberg, E. *et al.* Selection of resistant bacteria at very low antibiotic concentrations. *PLoS*
*Pathog* **7**, e1002158, doi:10.1371/journal.ppat.1002158 (2011).
12     Drlica, K. The mutant selection window and antimicrobial resistance. *J Antimicrob Chemother*
**52**, 11-17, doi:10.1093/jac/dkg269 (2003).
13     Sommer, M. O. A., Munck, C., Toft-Kehler, R. V. & Andersson, D. I. Prediction of
antibiotic resistance: time for a new preclinical paradigm? *Nat Rev Microbiol* **15**, 689-696,
doi:10.1038/nrmicro.2017.75 (2017).

14 Muliaditan, M., Davies, G. R., Simonsson, U. S. H., Gillespie, S. H. & Della Pasqua, O. The
implications of model-informed drug discovery and development for tuberculosis. *Drug*
*Discov Today* **22**, 481-486, doi:10.1016/j.drudis.2016.09.004 (2017).
15 Walkup, G. K. *et al.* Translating slow-binding inhibition kinetics into cellular and in vivo
effects. *Nat Chem Biol* **11**, 416-423, doi:10.1038/nchembio.1796 (2015).
16 Sykes, D. A. *et al.* Extrapyramidal side effects of antipsychotics are linked to their association
kinetics at dopamine D2 receptors. *Nat Commun* **8**, 763, doi:10.1038/s41467-017-00716-z
(2017).
17 Shen, L. *et al.* A critical subset model provides a conceptual basis for the high antiviral
activity of major HIV drugs. *Sci Transl Med* **3**, 91ra63, doi:10.1126/scitranslmed.3002304
(2011).
18 Abel zur Wiesch, P. *et al.* Classic reaction kinetics can explain complex patterns of antibiotic
action. *Sci Transl Med* **7**, doi:ARTN 287ra73
10.1126/scitranslmed.aaa8760 (2015).
19 Aldred, K. J., Kerns, R. J. & Osheroff, N. Mechanism of quinolone action and resistance.
*Biochemistry* **53**, 1565-1574, doi:10.1021/bi5000564 (2014).
20 Shen, L. L. & Pernet, A. G. Mechanism of inhibition of DNA gyrase by analogues of
nalidixic acid: the target of the drugs is DNA. *Proc Natl Acad Sci U S A* **82**, 307-311 (1985).
21 Shen, L. L. *et al.* Mechanism of inhibition of DNA gyrase by quinolone antibacterials: a
cooperative drug--DNA binding model. *Biochemistry* **28**, 3886-3894 (1989).
22 Andriole, V. T. <<The>> *quinolones*. 3rd edn, (Academic Press, 2000).
23 Jungkind, D. L. & American Society for Microbiology Eastern Pennsylvania Branch.
*Antimicrobial resistance a crisis in health care.* (Plenum Press, 1995).
24 Kampranis, S. C. & Maxwell, A. The DNA gyrase-quinolone complex. ATP hydrolysis and
the mechanism of DNA cleavage. *J Biol Chem* **273**, 22615-22626 (1998).
25 Siporin, C., Heifetz, C. L. & Domagala, J. M. *The New generation of quinolones*. (M. Dekker,
1990).
26 Abel Zur Wiesch, P., Clarelli, F. & Cohen, T. Using Chemical Reaction Kinetics to Predict
Optimal Antibiotic Treatment Strategies. *PLoS Comput Biol* **13**, e1005321,
doi:10.1371/journal.pcbi.1005321 (2017).
27 Elliott, T. S., Shelton, A. & Greenwood, D. The response of Escherichia coli to
ciprofloxacin and norfloxacin. *J Med Microbiol* **23**, 83-88, doi:10.1099/00222615-23-1-83
(1987).
28 Silva, F., Lourenco, O., Queiroz, J. A. & Domingues, F. C. Bacteriostatic versus bactericidal
activity of ciprofloxacin in Escherichia coli assessed by flow cytometry using a novel far-red
dye. *J Antibiot (Tokyo)* **64**, 321-325, doi:10.1038/ja.2011.5 (2011).
29 Garoff, L., Yadav, K. & Hughes, D. Increased expression of Qnr is sufficient to confer
clinical resistance to ciprofloxacin in Escherichia coli. *J Antimicrob Chemother* **73**, 348-352,
doi:10.1093/jac/dkx375 (2018).
30 Sulavik, M. C. *et al.* Antibiotic susceptibility profiles of Escherichia coli strains lacking
multidrug efflux pump genes. *Antimicrob Agents Chemother* **45**, 1126-1136,
doi:10.1128/AAC.45.4.1126-1136.2001 (2001).
31 Regoes, R. R. *et al.* Pharmacodynamic functions: a multiparameter approach to the design of
antibiotic treatment regimens. *Antimicrob Agents Chemother* **48**, 3670-3676,
doi:10.1128/AAC.48.10.3670-3676.2004 (2004).
32 Palmer, A. C. & Kishony, R. Opposing effects of target overexpression reveal drug
mechanisms. *Nat Commun* **5**, 4296, doi:10.1038/ncomms5296 (2014).
33 Fontana, W. Lecture Notes: Continuous-Time Monte-Carlo of Reaction Systems.

34    Ocampo, P. S. *et al.* Antagonism between bacteriostatic and bactericidal antibiotics is
prevalent. *Antimicrob Agents Chemother* **58**, 4573-4582, doi:10.1128/AAC.02463-14 (2014).
35    Chambers, H. F., Sachdeva, M. J. & Hackbarth, C. J. Kinetics of penicillin binding to
penicillin-binding proteins of Staphylococcus aureus. *Biochem J* **301 ( Pt 1)**, 139-144 (1994).
36    Al-Emran, H. M. *et al.* Detection of a Novel gyrB Mutation Associated With
Fluoroquinolone-Nonsusceptible Salmonella enterica serovar Typhimurium Isolated From a
Bloodstream Infection in Ghana. *Clin Infect Dis* **62 Suppl 1**, S47-49, doi:10.1093/cid/civ790
(2016).
37    Vogwill, T. & MacLean, R. C. The genetic basis of the fitness costs of antimicrobial
resistance: a meta-analysis approach. *Evol Appl* **8**, 284-295, doi:10.1111/eva.12202 (2015).
38    Jorgensen, J. H. & Ferraro, M. J. Antimicrobial susceptibility testing: a review of general
principles and contemporary practices. *Clin Infect Dis* **49**, 1749-1755, doi:10.1086/647952
(2009).
39    Tonge, P. J. Drug-Target Kinetics in Drug Discovery. *ACS Chem Neurosci* **9**, 29-39,
doi:10.1021/acschemneuro.7b00185 (2018).
40    Metcalf, B. J. *et al.* Using whole genome sequencing to identify resistance determinants and
predict antimicrobial resistance phenotypes for year 2015 invasive pneumococcal disease
isolates recovered in the United States. *Clin Microbiol Infect* **22**, 1002 e1001-1002 e1008,
doi:10.1016/j.cmi.2016.08.001 (2016).
41    Eyre, D. W. *et al.* WGS to predict antibiotic MICs for Neisseria gonorrhoeae. *J Antimicrob
Chemother* **72**, 1937-1947, doi:10.1093/jac/dkx067 (2017).
42    Payne, D. J., Gwynn, M. N., Holmes, D. J. & Pompliano, D. L. Drugs for bad bugs:
confronting the challenges of antibacterial discovery. *Nat Rev Drug Discov* **6**, 29-40,
doi:10.1038/nrd2201 (2007).
43    van der Putten, B. C. *et al.* Quantifying the contribution of four resistance mechanisms to
ciprofloxacin minimum inhibitory concentration in Escherichia coli: a systematic review.
372086, doi:10.1101/372086 %J bioRxiv (2018).
44    Copeland, R. A. The drug-target residence time model: a 10-year retrospective. *Nat Rev Drug
Discov* **15**, 87-95, doi:10.1038/nrd.2015.18 (2016).
45    Srimani, J. K., Huang, S., Lopatkin, A. J. & You, L. Drug detoxification dynamics explain the
postantibiotic effect. *Mol Syst Biol* **13**, 948, doi:10.15252/msb.20177723 (2017).
46    Cramariuc, O. *et al.* Mechanism for translocation of fluoroquinolones across lipid
membranes. *Biochim Biophys Acta* **1818**, 2563-2571, doi:10.1016/j.bbamem.2012.05.027
(2012).
47    Hirsch, C. 1 online resource (Butterworth-Heinemann,, Oxford, 2007).
48    Wu, J. *et al.* The dimer state of GyrB is an active form: implications for the initial complex
assembly and processive strand passage. *Nucleic Acids Res* **39**, 8488-8502,
doi:10.1093/nar/gkr553 (2011).
49    Maier, T. *et al.* Quantification of mRNA and protein and integration with protein turnover in
a bacterium. *Mol Syst Biol* **7**, 511, doi:10.1038/msb.2011.38 (2011).
50    Malmstrom, J. *et al.* Proteome-wide cellular protein concentrations of the human pathogen
Leptospira interrogans. *Nature* **460**, 762-765, doi:10.1038/nature08184 (2009).
51    Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes in
Escherichia coli K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**, 6640-6645,
doi:10.1073/pnas.120163297 (2000).
52    Martinecz, A. & Abel Zur Wiesch, P. Estimating treatment prolongation for persistent
infections. *Pathog Dis* **76**, doi:10.1093/femspd/fty065 (2018).