Parallel Evolution in the Emergence of Highly Pathogenic Avian Influenza A Viruses

Marina Escalera-Zamudio[1][†][*], Michael Golden[1][†], Bernardo Gutiérrez[1], Julien Thézé[1], Jeremy Russell Keown[2], Loic Carrique[2], Thomas A. Bowden[2] and Oliver G. Pybus[1,3][*]

1. Department of Zoology, Oxford University, Parks Rd Oxford, OX1 3PS, UK

2. Division of Structural Biology, Wellcome Centre for Human Genetics, Oxford, OX3 7BN, UK

3. Royal Veterinary College, London, UK.

[†]MEZ and MG contributed equally to this paper.
[*]Corresponding authors:
marina.escalerazamudio@zoo.ox.ac.uk; oliver.pybus@zoo.ox.ac.uk

other email addresses:
michael.golden@zoo.ox.ac.uk
bernardo.gutierrez@zoo.ox.ac.uk
julien.theze@zoo.ox.ac.uk
jeremy@strubi.ox.ac.uk
loic@strubi.ox.ac.uk
thomas.bowden@strubi.ox.ac.uk

ABSTRACT (145 words)

Parallel molecular evolution and adaptation are important phenomena commonly observed in

viruses. Here we exploit parallel molecular evolution to understand virulence evolution in

avian influenza viruses (AIV). Highly-pathogenic AIVs independently evolve from low-

pathogenic ancestors via acquisition of a polybasic cleavage sites (pCS). Why some AIV

lineages but not others evolve in this way is unknown. We hypothesise that the parallel

emergence of highly-pathogenic AIV may be facilitated by permissive or compensatory

mutations occurring across the AIV genome. We combined phylogenetic, statistical and

structural approaches to discover parallel mutations in AIV genomes associated with the

highly-pathogenic phenotype. Parallel mutations were screened using a new statistical test of

mutation-phenotype association and further evaluated in the contexts of positive selection

and protein function. The mutational panel we present reveals new links between virulence

evolution and other viral traits and raises the possibility of predicting aspects of AIV

evolution.

INTRODUCTION (5066 words)

Avian Influenza A viruses (AIVs) infect bird populations worldwide [1]. While most AIVs

cause only a mild-to-asymptomatic infection, highly-pathogenic (HP) avian influenza,

restricted to the H5 and H7 subtypes of AIV, can cause extremely high mortality (up to 100%)

in domestic bird populations [1]. Both low-pathogenic (LP) and HP viruses can cause severe

disease in humans following avian-to-human transmission, and there are potential instances

of human-to-human transmission [2,3]. Recently, the emergence of H7N9 AIV in China caused

>1500 infections and 615 deaths in people between 2013 and early 2018 [4]. Low virulence

AIVs have evolved into HP strains on multiple occasions, resulting in geographically and

temporally distinct HP outbreaks that generate sizeable economic losses to the poultry

industry, and pose an increased risk for avian-to-human spread [5][6]. Therefore, it is important

to understand the mechanisms driving the evolution of virulence in AIVs.

The virulence of an infectious disease is, in general, a complex trait [7][8][9]. However,

the AIV HP phenotype is one of the simplest and best characterized molecular markers of

virulence, and is therefore an attractive system in which to study virulence evolution. The HP

phenotype is conferred by a polybasic cleavage site (pCS) within the virus haemagglutinin

(HA) protein. A pCS is the outcome of one or more insertion mutations resulting in multiple

basic amino acids, which renders the HA protein a target for broadly-expressed host

proteases, facilitating systemic virus spread within avian hosts [10][11]. The pCS insertion has

occurred several times during the evolutionary history of AIVs. Over ten different pCS

sequences have been described for HP H5NX viruses since 1961, and over fifteen pCS have

been described for HP H7NX viruses since 1963 [12]. The evolutionary pathways that lead to

the independent, parallel evolution of pCSs and the emergence of the HP phenotype in

different AIV lineages are not well understood.

Parallel molecular evolution describes repeated evolutionary changes leading to the

same phenotype or genotype in independent populations, and can result from adaptation by

natural selection to similar selective pressures or ecological niches [13,14]. RNA viruses exhibit

high rates of mutation and replication, large population sizes, consistently varying

environments, and strong selective pressures. These factors, combined with small genome

sizes, which may restrict the number of mutational pathways leading to a given genotype,

may result in parallel molecular evolution being common in RNA viruses [15].

We hypothesize that the parallel evolution of the HP genotype from LP ancestors in

AIV subtypes H5 and H7 may have a genetic basis, and may be therefore associated with

permissive or compensatory secondary mutations occurring elsewhere in the virus genome.

Such mutations could arise prior to or immediately after the emergence of a pCS, perhaps to

compensate for possible detrimental effects on the stability of the HA protein [16,17]. Little is

known about the genetic predisposition of AIV to acquire a pCS (except for RNA structural

studies of the cleavage site region itself [18]), nor about the role of mutations elsewhere in the

virus genome favouring the evolution to a HP phenotype.

Here we use a combination of analytical methods from evolutionary and structural

biology to detect parallel mutations across the AIV genome that are associated with the

evolution of the HP phenotype (hereafter termed *HP-cluster associated parallel mutations, or*

HAPMs). We developed, validated and implemented a new phylogenetic model of binary trait

evolution that tests whether genotype-to-phenotype associations observed across multiple

lineages are statistically significant. This approach explicitly incorporates the phylogenetic

correlation among observations and thus is less susceptible to error than standard statistical

tests [19]. We further investigated if the HAPMs detected had evolved under positive selection,

and assessed their possible effects on protein function and/or stability. We identify numerous

HAPMs among HP AIV lineages of same viral subtype, many of which are also associated

with other viral processes, including antigenic escape, host-response modulation, and

replication efficiency. We discuss how the mutational panel presented here can provide a

starting point for future molecular studies on the evolution of the HP phenotype.

METHODS (1524)

74   We developed a new analysis strategy to detect parallel genetic changes associated

75 with the LP-to-HP evolution of AIV (summarized in Figure 1). For convenience, we refer to

76 parallel genetic changes as HAPMs (*HP-cluster associated parallel mutations*). The list of

77 HAPMs identified for each virus subtype was further analysed to identify those most likely to

78 be functionally and selectively relevant using the following criteria: *i)* they exhibit a

79 statistically-supported association with the HP phenotype, *ii*) they show evidence of being

80 evolving under positive selection, *iii)* they are predicted to be stabilizing to protein structure,

81 and *iv)* they have known or predicted functional relevance, derived from structural analysis.

83 **1. Large-scale phylogenetic analysis**

84   All available coding sequences for all genome segments of H7NX and H5NX viruses,

85 from all hosts and geographical regions, were downloaded from the Influenza Virus

86 Database. Sequences were excluded if (i) they were >200nt shorter than the full coding

87 sequence length, (ii) they represented identical sequences, or (iii) they contained >10%

88 ambiguities. Sequences were aligned using MAFFT [20] and preliminary large-scale

89 phylogenies for each genome segment were estimated using RAxML under a general time

90 reversible nucleotide substitution model with gamma distributed among-site rate variation

91 (GTR+G). Branch support was assessed using a bootstrap approach with 100 replicates [21].

92 HP strains were identified by (i) detecting >1 basic residue insertions within the consensus

93 cleavage site in the HA protein alignments [12] and (ii) by confirming HP status with case

94 reports in the literature [5]. Phylogenetic clusters of HP sequences were identified as (i)

95 monophyletic and well supported (bootstrap value >80) clusters that, (ii) contain >2 HP

96 sequences, (iii) descend from LP ancestors in all genome segments, and (iii) have emerged

97 independently from other HP clusters, and are thus distinct in time and space.

99 **2. Phylogenetically-informed subsampling**

To render further analyses computationally feasible, we sub-sampled the

abovementioned datasets in a phylogenetically-informed manner. Subsampling was

undertaken so as to retain the mutational pathways along internal tree branches that lead

from LP ancestors to HP clusters, whilst reducing overall alignment sizes. For each genome

segment, we retained (i) selected HP clusters, (ii) their immediate LP ancestors in the tree,

(iii) up to 250 randomly-selected LP reference sequences, in order to maintain overall tree

topology, and (iv) a group of ancestral sequences for rooting purposes. Sub-sampled

datasets were re-aligned and screened for recombination using the RDP, GENECOV and

BOOTSCAN components of the RDP3 package [22]. No evidence for recombination was

detected by three or more methods under a Bonferroni corrected p-value<0.001. Prior to

estimation of final phylogenies, HA alignments were edited to remove the cleavage site and

NA alignments were edited to remove hypervariable regions with poor alignment confidence.

Trees were estimated from the sub-sampled alignments using RAxML under the GTR+G

model and bootstrapped with 100 replicates [21]. Accession numbers and alignment sizes are

provided in Supplementary File 1. The varying sizes and/or missing HP clusters in some

alignments are due to the lack of complete genomes for all strains in the public database.

3. Detecting HAPMs occurring across HP clusters

To screen for HAPMs within AIV genomes, all mutations (deletions, insertions and

non-synonymous variable sites) present in ≥1% of the sequences in each sub-sampled

alignment were coded as a discrete taxon trait and subjected to ancestral state

reconstruction, using the maximum likelihood framework implemented in the "*ape*" R

package [23]. The resulting phylogenetic reconstructions were inspected visually to identify

HAPMs shared by multiple HP clusters. HAPMs were identified if: (i) they occurred within 2

or more HP clusters, (ii) they were present in >60% of the sequences within the HP clusters,

and (iii) they were not observed on internal branches anywhere else in the phylogeny

(Supplementary File 2).

To illustrate in detail the evolutionary association between HAPMs and AIV HP

clusters, we selected three HAPMS for each virus subtype for ancestral state reconstruction

using Bayesian molecular clock phylogenetic analysis. These six examples were chosen to

illustrate the range of evolutionary dynamics exhibited by HAPMs, and because they are all

significantly associated with the HP phenotype (see next section). For each site, we

estimated the posterior probability of amino acid states at tree nodes ancestral to HP clusters

in the subsampled alignments, using the discrete trait evolution model in BEAST v1.8.4 [24].

Time-scaled phylogenies were estimated under a SRD06 nucleotide substitution model, a

strict molecular clock model and a Bayesian Skyline coalescent tree prior. Partitions were

generated for each site, and ancestral states/transition rates were estimated for each

mutation using an asymmetric continuous time Markov chain model [25,26]. Two MCMC runs

were computed for 100×10^6 states or until convergence was reached. Maximum clade

credibility (MCC) trees were summarized using TreeAnnotator, after removing 10% of the

runs as burn-in [24]. Additionally, we re-estimated the full evolutionary history of these HAPMs

by mapping the amino acid changes onto the large-scale ML phylogenies generated in step 1

(Supplementary File 3).

4. Testing for genotype-to-phenotype associations

We developed a new evolutionary model to statistically test for associations between

binary phenotypic traits and amino acid states in a phylogenetically-informed manner. We

used this model to determine which of the HAPMs identified above were significantly

associated with the HP phenotype. Our method serves the same purpose as a χ^2 test, but

correctly incorporates the phylogenetic history of the trait and phenotype of interest, as it is

known that failing to account for these phylogenetic correlations can lead to erroneous false

positives [19,27]. The model assumes that phenotypes and amino acid states are discrete traits

that change along phylogeny branches. Given an alignment, its phylogeny, and a set of

binary values for each sequence, the method proceeds to test if a specific amino acid state is

associated with the given phenotype. This is repeated for all amino acid states and all

codons. A coupling parameter λ is used to measure the strength of association between an

amino acid state and the phenotype state. The null hypothesis of zero association is

represented by $\lambda = 1$, and can be rejected in favour of a positive association (λ >1) using a

likelihood ratio test (LRT). The type I and II error rates of this test were evaluated using

simulations. A full mathematical description of the model and simulation details are provided

in Supplementary Text 5. We applied this model to our subsampled H5NX and H7NX

alignments. The H5NX alignments were further reduced by collapsing clade C8 to only its

most basal sequences, in order to avoid the overrepresentation of HP sequences.

5. Testing for diversifying positive selection

To identify positively-selected HAPMs, we used various *dN/dS* (ω) ratio estimation

methods. The branch-site model A (BSA) in CODEML [28] was used to identify positively-

selected sites (PSS) associated with specific phylogeny branches, namely the HP clusters.

The statistical significance of the BSA model was compared to a null hypothesis with an ω=1

and evaluated using a LRT. The datamonkey server [29] was also used to test for sites

evolving under pervasive and episodic positive selection under both site- (MEME: Mixed

Effects Model of Evolution; SLAC: Single-Likelihood Ancestor Counting and FEL: Fixed

Effects Likelihood) and branch-site models (BSREL: branch-site random effects likelihood)

[30,31,32,33]. Test significances were assessed using a LRT and subjected to Holm-Bonferroni

multiple test correction. Finally, PSS were compared to the HAPMs detected and to the

DEPS algorithm results (Directional Evolution of Protein Sequences using Maximum

Likelihood) [34] (Supplementary Table 1). The LRT results and lists of the PSSs identified are

provided in the Supplementary File 4.

6. Inferring mutation effects on protein stability and function

We undertook structural analyses to assess if the identified HAPMs might be

functionally relevant. To determine the location and potential functional implications of the

different HAPMs, we modelled protein 3D structures from existing structural data in the PDB

database (https://pdb101.rcsb.org/), using representative HP sequences from our datasets

as queries. For H7, we used the crystal structure of A/H7N2/New York/107/2003 AIV (PDB

ID:3M5G), whilst for H5 we used the crystal structure of A/Vietnam/30262III/04 AIV (PDB ID:

2IBX). The high-resolution structure PDB 2Q06 was used to model NS1, and PDB 2Z16 was

used to model M1. For the PB2/PB1/PA complex, the high-resolution structure from the bat

Influenza A/H17N10 virus (PDB 4WSB) was used as a template, as this is the only structural

data currently available for the complete influenza A virus polymerase complex [35]. For the

HA, M1, and NS1 proteins, we used the I-TASSER server [36] to generate models with the

mutations of interest, that were compared manually to the PDB files for validation. For the

polymerase complex, mutations were introduced to the superimposed 3D protein structures

using PyMOL (Schrödinger, LLC). Finally, to predict which HAPMs had a possible effect on

protein stability, we used the PIPS (Phylogenetic Inference of Protein Stability) program [37].

This program implements an informed Bayesian approach and estimates, for each amino

acid state, a value of free energy change (ddG) in terms of the thermodynamic stability of an

unfolding/refolding protein, weighted under a phylogenetic background. Negative ddG values

are predicted to be stabilizing for protein structure. Values close to zero are predicted to have

little effect or to be only weakly stabilizing, whilst positive values are predicted to destabilize

protein   structure   [37].

RESULTS

Large-scale phylogenetic analysis of independent HP clusters

Large-scale phylogenetic analysis confirmed that H7NX viruses are split into two major clades, the Eurasian and North American lineages (Supplementary Figure 1) [38]. H7 HP outbreaks occurred independently through time and space and correspond to distinct lineages that, in most cases, display a unique pCS motif [5,6] (Figure 2). LP H7NX viruses carry a consensus amino acid sequence -PEXPKGRGLF- at the HA cleavage site, whereas a minimal consensus of -PEXPKXnK/RnRGLF- is identified for HP viruses (Figure 2a) [39] [11]. Nine H7NX well-supported clusters and spatio-temporally distinct HP lineages were identified in the HA phylogeny (Figure 2b) and in trees of other genome segments (Supplementary Figs. 2,3). Minor inconsistencies were observed for cluster C1, which segregated into C1.1 and C1.2 in the PB2, PB1, PA, M, and NS1 trees (Supplementary Figure 2).

The large-scale phylogeny of H5NX viruses shows a basal clade of HP and LP H5NX viruses circulating in wild birds worldwide from the late 1960s to the early 2000s (Figure 2d, Supplementary Figure 1). This basal clade is ancestral to the large and long-lived HP H5N1 lineage (named C8 here). This clade emerged in 2003 in China and subsequently spread to Asia, Africa, Europe and America [1] and persists until today [5]. Within C8, there is a distinct group of HP H5NX viruses (H5N2, H5N6 and H5N8 subtypes) that circulates in both wild and domestic birds (Figure 2d). Eight H5NX HP clusters were identified as being spatio-temporally distinct and well-supported (Figure 2c). Most HP clusters displayed a unique pCS sequence, except for C8, reflecting the ongoing evolution of this long-lived lineage. As with the H7NX viruses, phylogenetic clustering of HP H5NX viruses was conserved within other genome segment phylogenies (Supplementary Figures 3,4).

Identifying HAPMs significantly associated with the HP phenotype

The number of amino acid sites that were polymorphic (i.e. variants present at >1% frequency) varied among virus subtypes and genome segments. For H7NX viruses, the

number of polymorphic sites per segment ranged from 21 (for M1) to 125 (for HA). Maximum

likelihood ancestral state reconstructions revealed 16 parallel genetic changes (HAPMs)

present in two or more H7NX HP lineages (Supplementary File 2). For H5NX viruses,

between 12 (for M1) and 141 (for HA) polymorphic sites were detected for each segment.

We identified 39 HAPMs present in two or more H5NX HP lineages (Supplementary File 2).

Of these, 17 occurred within sub-lineages of cluster C8 and thus were not considered further.

To focus attention on those HAPMs significantly associated with the HP phenotype,

we developed a new phylogenetic model to test the null hypothesis of no association

between mutations at each position and the HP phenotype (Supplementary Text 5). For

H7NX viruses, we found a strong association with the HP phenotype for 9 out of 16 HAPMs

(Table 1). For H5NX viruses, a strong association with the HP phenotype was supported for

12 out of 22 HAPMs (Table 1).

To illustrate the evolutionary dynamics of HAPMs, we chose a subset of six HAPMs

that were significantly associated with the HP phenotype and subjected them to molecular

clock phylogenetic analysis (Figure 3). We observed a variety of evolutionary patterns and in

general obtained high posterior probability values for ancestral amino acids at nodes

preceding the emergence of HP clusters. By way of illustration, consider site 355 of H7NX

PB2. At this site K is present in HP clusters C5 and C6, but not in lineages representing the

LP background (Figure 3a). A similar pattern is seen for mutation 627K in H5NX viruses

(Figure 3f). More extensive parallel evolution is exhibited by mutations 143T in H7 (Figure

3b) and 127L in H5 (Figure 3d), which are shared by multiple HP clusters, but again not

observed within LP lineages. Mutation 209N in H7NX NS1 is found within several HP

clusters, but is also common in LP sequences of the American lineage (Figure 3c). Lastly,

site 154 in H5 exhibits a more complex, step-wise evolutionary process, from N to I/L to Q,

reflecting the transition from LP to HP (Figure 3e). N is present only in LP viruses, I/L occurs

in HP viruses circulating prior to and during the early expansion of C8, and Q is only present

in more recent HP viruses from C8 (Figure 3e).

To facilitate computational analyses, we sub-sampled AIV alignments before

detecting HAPMs (see Method step 2). Thus, the original alignments before subsampling

may contain HP lineages that were not present after sub-sampling (Supplementary Figure 1).

Thus, we investigated whether if these lineages also contained any HAPMs significantly

associated with the HP phenotype, by mapping HAPM amino acid changes onto the full

phylogenetic trees (Supplementary File 3). Since these HP lineages were not used to define

HAPMs, mutations present in them constitute an independent test of our statistical approach.

We found that, in addition to the HP clusters used here, mutation 154L in H5 was fixed in two

independent HP H5N1 outbreaks in Vietnam from 2008-2013 [40], and in HP H5N2 viruses in

China circulating from 2009-2015 [41] (Supplementary File 3). We also observed that mutation

143T is present in H7N6 viruses isolated from an outbreak in Aichi prefecture, Japan, in 2009

[42] (Figure 3b; Supplementary Files 1,3). Although this was not a HP outbreak, these viruses

contained an additional arginine residue within the pCS (-PKGR to -PKRR), which triggered

concerns about whether this lineage might evolve the HP phenotype. Fortunately, this

lineage was successfully eradicated before any HP phenotype strains were observed [42].

Most HAPMs were not observed on phylogeny branches immediately ancestral to the

HP clusters, suggesting that these appear shortly before, during, or after the emergence of

the HP phenotype and making it difficult to discern if they are permissive or compensatory

change. However mutation A143T in H7 is present in LP viral sequences that are basal and

immediately ancestral to the emergence of the HP H7N1 outbreak in Italy 2000 [43]. This

observation supports the notion that at least some of the HAPMs identified here could be

permissive.

HAPMs associated with diversifying positive selection

Although parallel evolution is often caused by positive selection, parallel genetic

changes can, in theory, also arise by chance [14]. Consequently, we investigated whether if

some HAPMs were inferred as being positively selected sites (PSS; see Methods section 5).

Analyses under site- and branch-site models showed evidence for diversifying positive

selection at some HAPMs, and for some HP clusters (H7NX clusters C4 and C7, and H5NX

clusters C1, C2 and C4; Supplementary Table 1). As expected [15], dN/dS (ω) values for whole

genome segments were <1 in all cases (data not shown), and the overall proportion of PSS

detected was low (Supplementary Table 1, Supplementary File 4).

For the H7NX viruses, 4/16 HAMPs (corresponding to 4/9 of the HAPMs significantly

associated with the HP phenotype) were identified as PSS. For the H5NX viruses, only 2/22

HAPMs (corresponding to 2/12 of the HAPMs significantly associated with the HP

phenotype) were identified as PSS (see Supplementary Table 1 for details). At first

impression, this suggests that most HAPMs significantly associated with the HP phenotype

are not a direct consequence of positive selection, and may instead represent neutral

variation that has arisen through genetic hitchhiking [14,16]. However, if some HAPMs are

selectively neutral in the absence of a pCS (LP phenotype), but offer a positive effect when a

pCS is present (HP phenotype), then they may not be inferred as PSS using dN/dS methods

yet still represent a permissive/compensatory change. This is a consequence of dN/dS

estimation methods being conservative in detecting sites under positive selection in some

circumstances [44]. For example, mutation E627K in PB2, which was not inferred to be a PSS,

is known to have a positive effect on the adaptation of AIV to mammalian hosts and

increases replication efficacy [45]. The role of selection on whole HP lineages is less clear: for

example, experiments in turkey populations suggest that the HP and LP phenotypes of H7N1

viruses have indistinguishable transmission rates and basic reproductive numbers (R_0) [46].

Structural and protein stability analyses

Most of the HAPMs significantly associated with the HP phenotype had negative ddG

values, as assessed under a Bayesian approach for inferring the stability effects of individual

mutations from homologous protein sequences with a known phylogeny [37] (Figures 4 and 5).

Thus, they are predicted to be fairly stabilizing/neutral changes to protein structure. One

exception is E627K in H5NX PB2, which is predicted to be destabilizing. However, it has

been shown that other compensatory mutations in PB2 appear to improve the genetic stability of E627K [47].

Structural analysis suggests that several H7NX HAPMs significantly associated with the HP are functionally relevant (Figure 4). A143T in HA is a non-conservative amino acid change located in a solvent-accessible region within the 130-loop proximal to the receptor surface. This is within antigenic pocket A, contributing to the electrostatic environment needed for ligand binding. Although it has no direct interaction with glycans [48], mutation A143T induces glycosylation of residue 141 by adding a N-linked glycosylation motif. This N-linked residue produces steric masking of the antigenic site and may affect HA folding and stability [75]. Similarly, M/L271I in HA is located within the receptor binding subdomain [49,50]. R355K in PB2 is a conservative amino acid change within a solvent-exposed loop of the electrostatic surface of the cap binding domain. This site is in direct contact with first phosphate after the cap [51] and correlates with high pathogenicity of AIVs in mouse models [52]. D209N in NS1 is a conservative amino acid change located within a disordered tail interacting with importin alpha, a cellular host factor that promotes the entry of proteins to the cell nucleus (Figure 4), and may thus interfere with cell cycle regulation pathways [53].

HAMPs significantly associated with the HP phenotype in H5NX viruses were also found to be structurally relevant (Figure 5). Mutation F127L is a conservative amino acid change within the globular head of HA1, close to antigenic site 3 [54]. N154I/L is a non-conservative amino acid change within antigenic site 2, and involves the loss of the asparagine-linked glycosylation site [55]. Both these HAPMs correspond to variable sites related to escape mutants [54]. However, they are not crucial for the protective efficacy of H5 vaccines, suggesting that the loss of the saccharide linkage may have little impact on antigenic variation [50,55]. I167T is located in the surface-exposed region of the HA protein, and has been related to increased infectivity in the human lower respiratory tract in HP H5N1 viruses from Egypt [56]. S145L in HA is also located in the surface exposed region and has been linked to increased virulence in poultry [57]. Position 145 also undergoes occasional deletion (as observed in approximately 10% of sequences) and it has been argued that this

deletion is critical for the antigenic evolution of long-lived HPAIV clusters [75]. Mutation E627K

in PB2 is a well-characterised adaptation marker associated with enhanced transmission in

mammalian hosts [58], found frequently among H5NX viruses causing human infections [7]. Site

627 is surface exposed and located on the "627 domain" and is essential for viral RNA

replication and transcription in a cellular context [59] (Figure 5). Mutation A674T in PB2 is

located at the interface of the polymerase complex trimer and is predicted to increase

strength of the PB2 protein structure. In *in vivo* mouse models this mutation is a prerequisite

for the acquisition of E627K, as it stabilizes polymerase activity when co-occurring with 627K

[60], supporting the notion that this HAPM acts as a permissive mutation.

DISCUSSION

The occurrence of parallel molecular evolution in AIV genomes can be used as a

starting point for the identification of mutations associated with the evolution of the HP

phenotype. We report combined evolutionary and structural evidence to support the

hypothesis that some AIV parallel mutations are significantly associated with the HP

phenotype and may potentially act as permissive/compensatory changes to facilitate the

evolution of HP viruses. Previous studies have noted parallel genetic changes in

independent AIV lineages [61], but these were not linked to the emergence of the HP

phenotype. Similarly, specific instances of parallel genetic changes associated with the

transition from a LP to a HP phenotype giving rise to single H7N1 outbreaks in Italy have

also been described [43,62,63]. We provide the first systematic and statistically-rigorous

screening of AIV mutations associated with HP outbreaks. We demonstrate the effectiveness

of phylogenetically-informed hypothesis testing coupled with comparative evolutionary and

structural approaches to identify potentially important mutations exhibiting parallel evolution

in H5 and H7 AIV genomes.

Permissive/compensatory mutations associated with the HP phenotype are expected

to be favoured under a HP genetic background, and therefore occur repeatedly in HP

lineages. Mutations that occur at nodes preceding and coinciding with the emergence of

pCS-defined HP clusters (*e.g.* those that are present in the immediate LP ancestors and in

the HP cluster), may represent permissive changes that facilitate the subsequent evolution of

the HP phenotype. Conversely, mutations that occur only within HP clusters (after a pCS has

arisen), may be more likely to be compensatory changes.

However, sampling of AIV genomes is likely biased towards HP outbreaks, resulting

in a phylogenetic under-representation of the immediate LP ancestors of any given HP

lineage. This under-sampling of local LP strains makes it difficult to discern whether the

HAPMs correspond to permissive versus compensatory changes. Further molecular

experiments are needed to understand how the parallel mutations detected here actually

contribute to the evolution of the HP phenotype by, for example, stabilizing protein structure,

altering host-virus or virus-virus inter-molecular interactions, or by combining additively to

increase virulence. However, such experimental validation may take several years to

complete especially if HP AIV gain-of-function experiments will be required [64]. In the

meantime, we should continue to exploit fully the information revealed by parallel molecular

evolution, which represents a "natural" replicated experiment. This situation further highlights

the importance of systematic sampling and whole genome sequencing of both HP viruses

and their immediate LP ancestors, prior and during AIV outbreaks.

Some of the HAPMs identified here were previously identified either *in vitro* or *in silico*

as being associated with HP AIV (Table 1) [47,52,56-58,60]. Although this supports our results, it is

possible that the number, biological roles, and genomic distribution of mutations involved in

HP adaptation is currently underestimated. Notably, the HAPMs described here are not

shared among the H7NX and H5NX virus subtypes, yet they commonly occur within the

same genome segments (Table 1), suggesting that the evolution of the HP phenotype may

interact with other well-studied biological processes, such as antigenic escape (determined

by the HA protein), host-response modulation (regulated by PB2 and NS1) and replication

efficiency (PB2). Our findings are in agreement with previous studies showing that the

acquisition of a pCS may require coadaptation of the HA and other viral proteins [65].

Some of the mutations occurring close to antigenic sites may be indirectly associated

with differential patterns of immunity in varied host populations, defined by complex factors

such as animal age and sex distributions, previous exposure to infections/vaccines, or

differences in host genetic diversity [66]. Interestingly, mathematical modelling has suggested

that the sporadic population dynamics of HP strains may be determined, at least in part, by

their linkage to one or more specific antigenic variants, whose frequencies are driven by the

dynamics of host immunity, rather than by mutability [67,68]. Furthermore, the emergence of the

HP phenotype has been often associated with virus adaptation from wild to domestic bird

host populations [61]. The strength of this association is difficult to assess in nature given the

substantial bias (in both the literature and sequence databases) towards AIV infections and

HP outbreaks in poultry versus those in wild birds. However, using our new model of

mutation-trait association (Methods section 4; Supplementary Text 5), we found no

association between any of the detected HAMPs (or any other mutation) and the host

species (domestic versus wild bird; data not shown). This suggests that the parallel

mutations identified here are not directly linked to transmission among domestic bird

populations, although high population densities together with a limited genetic variability

within the host (as occurs in intense farming facilities), may facilitate virus transmission. If

acquiring a pCS does indeed reduce AIV transmission fitness in natural systems [46], then it is

interesting to speculate that both intensive farming and compensatory/permissive mutations

may act to facilitate the spread of a HP strain; the latter by increasing R_0 and the former by

lowering the threshold R_0 required to sustain transmission in a population.

Lastly, we propose that the set of HAPMs associated with HPAIV lineages identified

here (or any improvement to this set obtained in the future), may, with sufficient validation

and testing, function as an early detection system capable of highlighting lineages that pose

a greater risk of evolving the HP phenotype in the future. Importantly, the mutational panel

identified here could potentially benefit to AIV control and bird health even if the mechanistic

and functional actions of the mutations described are unknown. We suggest that circulating

AIV strains in domestic and wild bird populations that do not contain a pCS site, but which

exhibit a combination of the HAPMs identified here, are potential targets for more focussed

and targeted genomic and epidemiological surveillance.

AUTHOR CONTRIBUTIONS

MEZ and OGP designed research. MEZ performed research. MEZ and OGP analysed data.

MG developed the statistical model and analysed data. BG and JT analysed genomic data.

JRK and LC analysed structural data. TAB supervised structural data analysis. MEZ wrote

the manuscript, with comments from all authors.

ACKNOWLEDGEMENTS

MEZ is supported by an EMBO Long Term Fellowship (ALTF376-2017). MG, JT and OGP

are supported by the EU's Seventh Framework Programme (FP7/2007-2013)/European

Research Council (614725-PATHPHYLODYN). BG is supported by the 2017 Universities of

Academic Excellence Scholarship Program of the Secretariat for Higher Education, Science,

Technology and Innovation of the Republic of Ecuador (ARSEQ-BEC-003163-2017) and by

Universidad San Francisco de Quito. JRK and LC are supported by the Wellcome

Investigator award (200835/Z/16/Z). The Wellcome Centre for Human Genetics is supported

by grant 203141/Z/16/Z. TAB is supported by the Medical Research Council

(MR/L009528/1).

 FIGURE LEGENDS

Figure 1. Summary of analytical approach

A summary of our approach to identifying HAPMs (HP-cluster associated parallel mutations)

in AIV genomes. 1. Large-scale data set curation and phylogenetic analysis to detect

genotypes and phenotypes of interest (HP lineages highlighted in red). 2. Phylogenetically-

informed subsampling to reduce alignment size whilst retaining key mutational pathways

along internal branches. 3. Detecting HAPMs across multiple HP clusters using ancestral

state reconstructions (HAPMs shared among clusters represented by coloured dots) 4.

Statistical testing for genotype-to-phenotype associations. 5. Identifying HAPMs evolving

under positive selection using different dN/dS (ω) estimation methods. 6. Inferring the

possible effects of HAPMs on protein stability and function by inferring ddG values, and by

using existing structural data to infer interactions.

Figure 2. Independent spatio-temporal emergence of the HP phenotype

(a) Geographical occurrence of H7NX HP outbreaks. The year of circulation, virus subtype

and consensus sequence for the polybasic cleavage site (pCS) are shown for the HP

clusters analysed in this work. Countries of emergence are in red. (b) Phylogenetic

distribution of HP outbreaks in the H7NX phylogeny. Distinct well-supported HP clusters

within the HA tree are highlighted in red and named (C1-C9). Posterior probabilities and

bootstrap support values are shown for nodes of interest. (c) Geographical occurrence of

H5NX HP outbreaks, excluding C8 (clade 2.2) which is distributed worldwide. (d)

Phylogenetic distribution of HP outbreaks in the H5NX phylogeny. Distinct well-supported HP

clusters within the HA tree are highlighted in red and named (C1-C8). The H5NX HP viruses

within cluster C8 (clade 2.2) are highlighted in blue.

Figure 3. Time-scaled reconstruction of amino acid evolution at selected HAPMs

Maximum clade credibility trees are shown for six illustrative HAPMs (a-f) that exhibit a

significant associated with the HP phenotype. Amino acid changes at each of the six sites

were inferred using a Bayesian molecular clock phylogenetic approach. Branches are

coloured according to the inferred amino acid states at internal nodes (tip states not shown).

Ancestral nodes preceding the emergence of a given HAPM are highlighted by coloured

circles. The posterior probabilities for the amino acid state occurring at specified node are

shown. The HP clusters are highlighted by grey circles.

Figure 4. Functional relevance of HAPMs of H7NX viruses

Protein structure models for the HA (Hemagglutinin), PB2/PB1 (Polymerase complex) and

the NS1 (Non-structural protein) of the H7NX viruses. Functionally relevant HAPMs are

highlighted in red within the structural models; only those HAMPs that are significantly

associated with the HP phenotype are indicated. Estimated values for free energy change

(ddG) are indicated for these sites. Negative values or those close to zero can be interpreted

as fairly stabilizing/neutral mutations with respect to protein structure.

Figure 5. Functional relevance of HAPMs of H5NX viruses

Protein structure models for the HA (Hemagglutinin) and the PB2 polymerase subunit of the

H5NX viruses. See Figure 4 legend for further details.

TABLE 1. HAPMs significantly associated with the HP phenotype of H5 and H7 virus subtypes

H7NX							
HAPM	Segment	HP clusters	Num. HP clusters	λ [95% CI]	p-value	H7 numbering	Reference[*]
A143T [†]	HA	C2, C5, C6, C7 and C8	5	7.1 [3.3, 13.2]	0.00	125	67,68,69,75
D209N [†]	NS1	C1.1, C6 and C8	3	2.4 [1.0, 4.3]	0.03		
R355K [†]	PB2	C5 and C6	2	6.0 [2.8, 11.8]	0.00		57
M/L274I	HA	C4 and C8	2	6.2 [3.1, 11.5]	0.00	256	
V480I	PB2	C5 and C6	2	9.2 [4.3, 18.9]	0.00		
V438I	HA	C3 and C8	2	4.6 [2.4, 8.3]	0.01	423	
D384N	HA	C6 and C8	2	5.1 [2.6, 9.5]	0.02	369	
G154D	PB1	C2 and C6	2	7.7 [4.3, 13.2]	0.03		67
S152L	PB1	C6 and C9	2	3.1 [1.7, 5.2]	0.03		
H5NX							
HAPM	Segment	HP clusters	Num. HP clusters	λ [95% CI]	p-value	H5 numbering	Reference[*]
V113I	PB1	C2, C4 and C8	3	7.3 [3.7, 14.5]	0.01		
F127L[†]	HA	C3, C7 and C8	3	6.5 [3.3, 12]	0.03	111	59
N154I/L/Q[† §]	HA	C6, C7 and C8	3	See footnote	0.02	138	59
S145L/del[‡]	HA	C5 and C8	2	See footnote	0.01	129	62,75
A674T	PB2	C2 and C6	2	7.4 [3.9, 13.6]	0.00		65
M242I	HA	C1 and C8	2	7.4 [3.7, 13.7]	0.00	226	72
I167T	HA	C6 and C8	2	5.8 [3.0, 10.4]	0.03	151	61
E627K	PB2	C1 and C8	2	5.1 [2.7, 9.2]	0.04		
I451T	PB2	C2 and C8	2	3.8 [2.1, 6.8]	0.03		
R101K	M	C5 and C8	2	3.7 [1.7, 6.9]	0.02		
T204I	HA	C2 and C8	2	2.7 [1.4, 4.7]	0.03	188	
K379R	HA	C1 and C8	2	14.8 [7.4, 27.5]	0.00	363	

[†]Scored as positively selected site under different methods (see Supplementary Table 1 in the Supplementary Information).

[§] Individual lambda values for each site are as follow: for 154I 1.7 [1.0, 2.7], for 154L 3.2 [1.9, 5.2] and for 154Q: 3.4 [2.0, 5.8], all with a p-value of 0.00.

[‡] Individual lambda values for each site are as follow: for 145L 77.5 [69.5, 100.0] with a p-value of 0.00, and for 145del 7.0 [0.5, 14.7] with a p-value of 0.04.

[*] Where such HAPMs had been previously mentioned.

REFERENCES

Webster, R. G., Peiris, M., Chen, H. & Guan, Y. H5N1 outbreaks and enzootic influenza. *Emerg Infect Dis* **12**, 3-8, doi:10.3201/eid1201.051024 (2006).

Buxton Bridges, C. *et al.* Risk of influenza A (H5N1) infection among health care workers exposed to patients with influenza A (H5N1), Hong Kong. *J Infect Dis* **181**, 344-348, doi:10.1086/315213 (2000).

Du Ry van Beest Holle, M., Meijer, A., Koopmans, M. & de Jager, C. M. Human-to-human transmission of avian influenza A/H7N7, The Netherlands, 2003. *Euro Surveill* **10**, 264-268 (2005).

Lam, T. T. & Pybus, O. G. Genomic surveillance of avian-origin influenza A viruses causing human disease. *Genome Med* **10**, 50, doi:10.1186/s13073-018-0560-3 (2018).

Dhingra, M. S. *et al.* Geographical and Historical Patterns in the Emergences of Novel Highly Pathogenic Avian Influenza (HPAI) H5 and H7 Viruses in Poultry. *Front Vet Sci* **5**, 84, doi:10.3389/fvets.2018.00084 (2018).

Short, K. R. *et al.* One health, multiple challenges: The inter-species transmission of influenza A virus. *One Health* **1**, 1-13, doi:10.1016/j.onehlt.2015.03.001 (2015).

Hatta, M., Gao, P., Halfmann, P. & Kawaoka, Y. Molecular basis for high virulence of Hong Kong H5N1 influenza A viruses. *Science* **293**, 1840-1842, doi:10.1126/science.1062882 (2001).

Jackson, D., Hossain, M. J., Hickman, D., Perez, D. R. & Lamb, R. A. A new influenza virus virulence determinant: the NS1 protein four C-terminal residues modulate pathogenicity. *Proc Natl Acad Sci U S A* **105**, 4381-4386, doi:10.1073/pnas.0800482105 (2008).

Schmolke, M. *et al.* Differential contribution of PB1-F2 to the virulence of highly pathogenic H5N1 influenza A virus in mammalian and avian species. *PLoS Pathog* **7**, e1002186, doi:10.1371/journal.ppat.1002186 (2011).

Horimoto, T. & Kawaoka, Y. Reverse genetics provides direct evidence for a correlation of hemagglutinin cleavability and virulence of an avian influenza A virus. *J Virol* **68**, 3120-3128 (1994).

Abdelwhab, E. M. *et al.* Composition of the Hemagglutinin Polybasic Proteolytic Cleavage Motif Mediates Variable Virulence of H7N7 Avian Influenza Viruses. *Sci Rep* **6**, 39505, doi:10.1038/srep39505 (2016).

OFFLU. *OIE/FAO Network of expertise on animal influenza: Influenza A Cleavage Sites Report 2018*, <http://www.offlu.net/fileadmin/home/en/resource-centre/pdf/Influenza_A_Cleavage_Sites.pdf> (2018).

Stern, D. L. The genetic causes of convergent evolution. *Nat Rev Genet* **14**, 751-764, doi:10.1038/nrg3483 (2013).

Gutierrez, B., Escalera-Zamudio, M. & Pybus, O. G. Parallel molecular evolution and adaptation in viruses. *Current Opinion in Virology* **34**, 90-96, doi:10.1016/j.coviro.2018.12.006. (2019).

Pond, S. L., Murrell, B. & Poon, A. F. Evolution of viral genomes: interplay between selection, recombination, and other forces. *Methods Mol Biol* **856**, 239-272, doi:10.1007/978-1-61779-585-5_10 (2012).

Frost, S. D. W., Magalis, B. R. & Kosakovsky Pond, S. L. Neutral Theory and Rapidly Evolving Viral Pathogens. *Mol Biol Evol* **35**, 1348-1354, doi:10.1093/molbev/msy088 (2018).

Bloom, J. D., Gong, L. I. & Baltimore, D. Permissive secondary mutations enable the evolution of influenza oseltamivir resistance. *Science* **328**, 1272-1275, doi:10.1126/science.1187816 (2010).

Nao, N. *et al.* Genetic Predisposition To Acquire a Polybasic Cleavage Site for Highly Pathogenic Avian Influenza Virus Hemagglutinin. *MBio* **8**, doi:10.1128/mBio.02298-16 (2017).

Bhattacharya, T. *et al.* Founder effects in the assessment of HIV polymorphisms and HLA allele associations. *Science* **315**, 1583-1586, doi:10.1126/science.1131528 (2007).

Katoh, K., Misawa, K., Kuma, K. & Miyata, T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* **30**, 3059-3066 (2002).

Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312-1313, doi:10.1093/bioinformatics/btu033 (2014).

Martin, D. P. *et al.* RDP3: a flexible and fast computer program for analyzing recombination. *Bioinformatics* **26**, 2462-2463, doi:10.1093/bioinformatics/btq467 (2010).

23 Paradis, E. & Schliep, K. ape 5.0: an environment for modern phylogenetics and evolutionary
analyses in R. *Bioinformatics*, doi:10.1093/bioinformatics/bty633 (2018).
24 Drummond, A. J., Suchard, M. A., Xie, D. & Rambaut, A. Bayesian phylogenetics with BEAUti
and the BEAST 1.7. *Mol Biol Evol* **29**, 1969-1973, doi:10.1093/molbev/mss075 (2012).
25 Ferreira, M. A. R. & Suchard, M. A. Bayesian analysis of elapsed times in continuous-time
Markov chains. *Canadian Journal of Statistics* **36**, 355-368,
doi:https://dx.doi.org/10.1002/cjs.5550360302 (2008).
26 Lemey, P., Rambaut, A., Drummond, A. J. & Suchard, M. A. Bayesian phylogeography finds
its roots. *PLoS Comput Biol* **5**, e1000520, doi:10.1371/journal.pcbi.1000520 (2009).
27 Harvey, P. H. & Pagel, M. D. *The Comparative Method in Evolutionary Biology.* (Oxford
University Press, 1991).
28 Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**, 1586-1591,
doi:10.1093/molbev/msm088 (2007).
29 Delport, W., Poon, A. F., Frost, S. D. & Kosakovsky Pond, S. L. Datamonkey 2010: a suite of
phylogenetic analysis tools for evolutionary biology. *Bioinformatics* **26**, 2455-2457,
doi:10.1093/bioinformatics/btq429 (2010).
30 Kosakovsky Pond, S. L. & Frost, S. D. Not so different after all: a comparison of methods for
detecting amino acid sites under selection. *Mol Biol Evol* **22**, 1208-1222,
doi:10.1093/molbev/msi105 (2005).
31 Murrell, B. *et al.* Detecting individual sites subject to episodic diversifying selection. *PLoS*
Genet **8**, e1002764, doi:10.1371/journal.pgen.1002764 (2012).
32 Murrell, B. *et al.* Gene-wide identification of episodic selection. *Mol Biol Evol* **32**, 1365-1371,
doi:10.1093/molbev/msv035 (2015).
33 Smith, M. D. *et al.* Less is more: an adaptive branch-site random effects model for efficient
detection of episodic diversifying selection. *Mol Biol Evol* **32**, 1342-1353,
doi:10.1093/molbev/msv022 (2015).
34 Kosakovsky Pond, S. L., Poon, A. F., Leigh Brown, A. J. & Frost, S. D. A maximum likelihood
method for detecting directional evolution in protein sequences and its application to influenza
A virus. *Mol Biol Evol* **25**, 1809-1824, doi:10.1093/molbev/msn123 (2008).
35 Reich, S. *et al.* Structural insight into cap-snatching and RNA synthesis by influenza
polymerase. *Nature* **516**, 361-+, doi:10.1038/nature14009 (2014).
36 Yang, J. *et al.* The I-TASSER Suite: protein structure and function prediction. *Nat Methods* **12**,
7-8, doi:10.1038/nmeth.3213 (2015).
37 Bloom, J. D. & Glassman, M. J. Inferring stabilizing mutations from protein phylogenies:
application to influenza hemagglutinin. *PLoS Comput Biol* **5**, e1000349,
doi:10.1371/journal.pcbi.1000349 (2009).
38 Xu, Y. *et al.* Limited Antigenic Diversity in Contemporary H7 Avian-Origin Influenza A Viruses
from North America. *Sci Rep* **6**, 20688, doi:10.1038/srep20688 (2016).
39 Bosch, F. X., Garten, W., Klenk, H. D. & Rott, R. Proteolytic cleavage of influenza virus
hemagglutinins: primary structure of the connecting peptide between HA1 and HA2
determines proteolytic cleavability and pathogenicity of Avian influenza viruses. *Virology* **113**,
725-735 (1981).
40 Nguyen, T. *et al.* Characterization of a highly pathogenic avian influenza H5N1 virus
sublineage in poultry seized at ports of entry into Vietnam. *Virology* **387**, 250-256,
doi:10.1016/j.virol.2009.03.006 (2009).
41 Zhao, G. *et al.* Novel reassortant highly pathogenic H5N2 avian influenza viruses in poultry in
China. *PLoS One* **7**, e46183, doi:10.1371/journal.pone.0046183 (2012).
42 Uchida, Y. *et al.* Genetic characterization and susceptibility on poultry and mammal of H7N6
subtype avian influenza virus isolated in Japan in 2009. *Vet Microbiol* **147**, 1-10,
doi:10.1016/j.vetmic.2010.05.037 (2011).
43 Monne, I. *et al.* Emergence of a highly pathogenic avian influenza virus from a low-pathogenic
progenitor. *J Virol* **88**, 4375-4388, doi:10.1128/JVI.03181-13 (2014).
44 Anisimova, M. & Kosiol, C. Investigating protein-coding sequence evolution with probabilistic
codon substitution models. *Mol Biol Evol* **26**, 255-271, doi:10.1093/molbev/msn232 (2009).
45 Linster, M. *et al.* Identification, characterization, and natural selection of mutations driving
airborne transmission of A/H5N1 virus. *Cell* **157**, 329-339, doi:10.1016/j.cell.2014.02.040
(2014).
46 Saenz, R. A. *et al.* Quantifying transmission of highly pathogenic and low pathogenicity H7N1
avian influenza in turkeys. *PLoS One* **7**, e45059, doi:10.1371/journal.pone.0045059 (2012).

606	47	Long, J. S. *et al.* The effect of the PB2 mutation 627K on highly pathogenic H5N1 avian
607		influenza virus is dependent on the virus lineage. *J Virol* **87**, 9983-9996,
608		doi:10.1128/JVI.01399-13 (2013).

609	48	Yang, H., Chen, L. M., Carney, P. J., Donis, R. O. & Stevens, J. Structures of receptor
610		complexes of a North American H7N2 influenza hemagglutinin with a loop deletion in the
611		receptor binding site. *PLoS Pathog* **6**, e1001081, doi:10.1371/journal.ppat.1001081 (2010).

| 612 | 49 | Lu., J. *et al.* Molecular evolution, diversity and adaptation of H7N9 influenza A viruses in |
| 613 | | China. *Emerging Infectious Diseases* **24**, doi:DOI: 10.3201/eid2410.171063 (2018). |

614	50	Ha, Y., Stevens, D. J., Skehel, J. J. & Wiley, D. C. H5 avian and H9 swine influenza virus
615		haemagglutinin structures: possible origin of influenza subtypes. *EMBO J* **21**, 865-875,
616		doi:10.1093/emboj/21.5.865 (2002).

| 617 | 51 | Guilligay, D. *et al.* The structural basis for cap binding by influenza virus polymerase subunit |
| 618 | | PB2. *Nat Struct Mol Biol* **15**, 500-506, doi:10.1038/nsmb.1421 (2008). |

| 619 | 52 | Katz, J. M. *et al.* Molecular correlates of influenza A H5N1 virus pathogenesis in mice. *J Virol* |
| 620 | | **74**, 10807-10810 (2000). |

621	53	Melen, K. *et al.* Nuclear and nucleolar targeting of influenza A virus NS1 protein: striking
622		differences between different virus subtypes. *J Virol* **81**, 5995-6006, doi:10.1128/JVI.01714-06
623		(2007).

| 624 | 54 | Stevens, J. *et al.* Structure and receptor specificity of the hemagglutinin from an H5N1 |
| 625 | | influenza virus. *Science* **312**, 404-410, doi:10.1126/science.1124513 (2006). |

626	55	Hoffmann, E., Lipatov, A. S., Webby, R. J., Govorkova, E. A. & Webster, R. G. Role of specific
627		hemagglutinin amino acids in the immunogenicity and protection of H5N1 influenza virus
628		vaccines. *Proc Natl Acad Sci U S A* **102**, 12915-12920, doi:10.1073/pnas.0506416102 (2005).

629	56	Saleh, M. M., Selim, A., Shahera, A. A., El -Shahidy, M. S. & El-Trabilli, M. M. Molecular
630		characterization of recently isolated highly pathogenic avian influenza virus (H5N1) in Egypt.
631		*Nature and Science* **15** (2017).

| 632 | 57 | Perdue, M. L. & Suarez, D. L. Structural features of the avian influenza virus hemagglutinin |
| 633 | | that influence virulence. *Vet Microbiol* **74**, 77-86 (2000). |

634	58	Steel, J., Lowen, A. C., Mubareka, S. & Palese, P. Transmission of influenza virus in a
635		mammalian host is increased by PB2 amino acids 627K or 627E/701N. *PLoS Pathog* **5**,
636		e1000252, doi:10.1371/journal.ppat.1000252 (2009).

| 637 | 59 | Nilsson, B. E., Te Velthuis, A. J. & Fodor, E. Role of the PB2 627 Domain in Influenza A Virus |
| 638 | | Polymerase Function. *J Virol* **91**, doi:10.1128/JVI.02467-16 (2017). |

639	60	Lee, C. Y. *et al.* Prerequisites for the acquisition of mammalian pathogenicity by influenza A
640		virus with a prototypic avian PB2 gene. *Sci Rep* **7**, 10205, doi:10.1038/s41598-017-09560-z
641		(2017).

| 642 | 61 | Lebarbenchon, C. & Stallknecht, D. E. Host shifts and molecular evolution of H7 avian |
| 643 | | influenza virus hemagglutinin. *Virol J* **8**, 328, doi:10.1186/1743-422X-8-328 (2011). |

644	62	Fusaro, A. *et al.* Evolutionary trajectories of two distinct avian influenza epidemics:
645		Parallelisms and divergences. *Infect Genet Evol* **34**, 457-466,
646		doi:10.1016/j.meegid.2015.05.020 (2015).

647	63	Banks, J. *et al.* Changes in the haemagglutinin and the neuraminidase genes prior to the
648		emergence of highly pathogenic H7N1 avian influenza viruses in Italy. *Arch Virol* **146**, 963-973
649		(2001).

650	64	Casadevall, A. & Imperiale, M. J. Risks and benefits of gain-of-function experiments with
651		pathogens of pandemic potential, such as influenza virus: a call for a science-based
652		discussion. *MBio* **5**, e01730-01714, doi:10.1128/mBio.01730-14 (2014).

653	65	Veits, J. *et al.* Avian influenza virus hemagglutinins H2, H4, H8, and H14 support a highly
654		pathogenic phenotype. *Proc Natl Acad Sci U S A* **109**, 2579-2584,
655		doi:10.1073/pnas.1109397109 (2012).

| 656 | 66 | Hill, S. C. *et al.* Antibody responses to avian influenza viruses in wild birds broaden with age. |
| 657 | | *Proc Biol Sci* **283**, doi:10.1098/rspb.2016.2159 (2016). |

658	67	Wikramaratna, P. S., Pybus, O. G. & Gupta, S. Contact between bird species of different
659		lifespans can promote the emergence of highly pathogenic avian influenza strains. *Proc Natl*
660		*Acad Sci U S A* **111**, 10767-10772, doi:10.1073/pnas.1401849111 (2014).

| 661 | 68 | Thompson, C. P. *et al.* A naturally protective epitope of limited variability as an influenza |
| 662 | | vaccine target. *Nat Commun* **9**, 3859, doi:10.1038/s41467-018-06228-8 (2018). |

1 **Large-scale phylogenetic analysis:** Large-scale dataset compilation and initial phylogenetic analyses

2 **Phylogenetically-informed subsmapling:** to reduce tree size, yet retaining the mutational pathways of interest

3 **Detecting HAPMs occurring across HP clusters:** using ancestral state reconstruction (ML)

4 **Testing for genotype-to-phenotype associations:** under a model that considers the phylogenetic history of the trait and the phenotype

5 **Testing for diversifying positive selection:** selecting parallel genetic changes scored as PSS

6 **Inferring mutation effects on protein stability and function:** using a structural biology approach

H7NX

(a)

C9
Canda 2004-2005
H7N3
pCS: PENP*KQAY(X)KRMTR

C7
UK 1963
H7N3
pCS: PETP*KRRRR

C8
Mexico 2012-2016
H7N3
pCS: PENP*KDRKSRHRRTR

C4
Europe (UK, Sweden) 2008
H7N7
pCS: PEIP*KRRGR

C5
Europe
(Netherlands, Germany) 2003
H7N7
pCS: PEIP*KRRRR

C2
Italy1999-2000
H7N1
pCS: PEIP*KGSRVRR

C3
Italy 2013
H7N7
pCS: PETP*KRRERR

C6
Pakistan 1995-2004
H7N3
pCS: PETP*KRRKR

C1
China-Japan 2017
H7N9
pCS: PEVP*KRKRTAR

(b)

H5NX

(c)

C3
Mexico (Durango) 2005
H5N2
pCS: PKRK*TR

C2
Mexico (Puebla) 1994
H5N2
pCS: PQRK*RKTR

C1
USA (Texas) 1993
H5N2
pCS: PQRK*TR

C5
Ireland 1983
H5N8
pCS: PQRK*RKKR

C6
France 2015
H5N9
pCS: HQRR*KR

C7
Italy1997
H5N2
pCS: PQRR*RKKR

C4
South Africa 1960
H5N3
pCS: PQRE*TRRQKR

Note:
C8 (H5N1 and H5NX) correspond to clade 2.3.4.4, which is widely distributed worldwide.
Thus, it is not marked in the map.

(d)

H7

Head

S136N
ddG: -3.17

A143T
ddG: -3.78

M/L274I
ddG: 2.72

Stalk

V438I
ddG: 0.64

D384N
ddG: 2.98

Polymerase complex
PB2/PB1

V480I
ddG: 0.20

R355K
ddG: 0.26

G154D/S
ddG: -1.61

S152L
ddG: -1.22

PB1
PB2
RNA

NS1

Effector Domain

N127T
ddG: -2.52

V180I
ddG: -1.49

D209N
ddG: -2.40

D139N
ddG: -2.34

RNA Binding Domain

T80S
ddG: 0.21

H5

PB2

Head

Stalk

F127L
ddG: 2.40

N154I/L
ddG: -3.21

S145L
ddG: -1.74

I167T
ddG: 0.06

E627K
ddG: 3.58

A674T
ddG: 4.76

TABLE 1. HAPMs significantly associated with the HP phenotype of H5 and H7 virus subtypes

H7NX							
HAPM	**Segment**	**HP clusters**	**Num. HP clusters**	**λ [95% CI]**	**p-value**	**H7 numbering**	**Reference**[*]
A143T [†]	HA	C2, C5, C6, C7 and C8	5	7.1 [3.3, 13.2]	0.00	125	67,68,69,75
D209N [†]	NS1	C1.1, C6 and C8	3	2.4 [1.0, 4.3]	0.03		
R355K [†]	PB2	C5 and C6	2	6.0 [2.8, 11.8]	0.00		57
M/L274I	HA	C4 and C8	2	6.2 [3.1, 11.5]	0.00	256	
V480I	PB2	C5 and C6	2	9.2 [4.3, 18.9]	0.00		
V438I	HA	C3 and C8	2	4.6 [2.4, 8.3]	0.01	423	
D384N	HA	C6 and C8	2	5.1 [2.6, 9.5]	0.02	369	
G154D	PB1	C2 and C6	2	7.7 [4.3, 13.2]	0.03		67
S152L	PB1	C6 and C9	2	3.1 [1.7, 5.2]	0.03		
H5NX							
HAPM	**Segment**	**HP clusters**	**Num. HP clusters**	**λ [95% CI]**	**p-value**	**H5 numbering**	**Reference**[*]
V113I	PB1	C2, C4 and C8	3	7.3 [3.7, 14.5]	0.01		
F127L [†]	HA	C3, C7 and C8	3	6.5 [3.3, 12]	0.03	111	59
N154I/L/Q [† §]	HA	C6, C7 and C8	3	See footnote	0.02	138	59
S145L/del [‡]	HA	C5 and C8	2	See footnote	0.01	129	62,75
A674T	PB2	C2 and C6	2	7.4 [3.9, 13.6]	0.00		65
M242I	HA	C1 and C8	2	7.4 [3.7, 13.7]	0.00	226	72
I167T	HA	C6 and C8	2	5.8 [3.0, 10.4]	0.03	151	61
E627K	PB2	C1 and C8	2	5.1 [2.7, 9.2]	0.04		
I451T	PB2	C2 and C8	2	3.8 [2.1, 6.8]	0.03		
R101K	M	C5 and C8	2	3.7 [1.7, 6.9]	0.02		
T204I	HA	C2 and C8	2	2.7 [1.4, 4.7]	0.03	188	
K379R	HA	C1 and C8	2	14.8 [7.4, 27.5]	0.00	363	

[†]Scored as positively selected site under different methods (see Supplementary Table 1 in the Supplementary Information).

[§] Individual lambda values for each site are as follow: for 154I 1.7 [1.0, 2.7], for 154L 3.2 [1.9, 5.2] and for 154Q: 3.4 [2.0, 5.8], all with a p-value of 0.00.

[‡] Individual lambda values for each site are as follow: for 145L 77.5 [69.5, 100.0] with a p-value of 0.00, and for 145del 7.0 [0.5, 14.7] with a p-value of 0.04.

* Where such HAPMs had been previously mentioned.