Non-parametric polygenic risk prediction using partitioned GWAS summary statistics

Sung Chun[1,2,3,¶], Maxim Imakaev[1,2,3,¶], Nathan O. Stitziel[4,5,6,*], Shamil R. Sunyaev[1,2,3,7,*]

[1] Division of Genetics, Brigham and Women's Hospital, Boston, Massachusetts, 02115, USA

[2] Department of Medicine, Harvard Medical School, Boston, Massachusetts, 02115, USA

[3] Program in Medical and Population Genetics, Broad Institute, Cambridge, Massachusetts,

02142, USA

[4] Cardiovascular Division, Department of Medicine, Washington University School of Medicine,

Saint Louis, Missouri, 63110, USA

[5] Department of Genetics, Washington University School of Medicine, Saint Louis, Missouri,

63110, USA

[6] McDonnell Genome Institute, Washington University School of Medicine, Saint Louis,

Missouri, 63110, USA

[7] Department of Biomedical Informatics, Harvard Medical School, Boston, Massachusetts,

02115, USA

[¶] These authors contributed equally to this work.

[*] Corresponding authors

nstitziel@wustl.edu (NOS)

ssunyaev@rics.bwh.harvard.edu (SRS)

Abstract

Traditional methods for polygenic risk prediction of complex disease have historically

focused on the set of genetic markers associated with disease at a genome-wide level of

significance. Newer approaches have been developed to incorporate information from all markers

across the genome, however these methods typically require individual level genotypes or

depend on accurately specifying the underlying genetic architecture in order to optimally apply

shrinkage methods on estimated effect sizes. Here, we propose a novel partitioning-based risk

prediction model to achieve non-parametric shrinkage that does not require explicitly modeling

the genetic architecture. We start with a set of summary statistics in the form of SNP effect sizes

from a large GWAS cohort. We remove the correlation structure across summary statistics arising

due to linkage disequilibrium. Then, we partition the decorrelated summary statistics and

determine the appropriate shrinkage factors for each partition using an independent small cohort

with individual level data. We show that the resulting non-parametric shrinkage is equivalent to

applying piecewise linear interpolation on conditional mean effects. Based on simulated datasets,

we show that our new non-parametric shrinkage (NPS) can reliably correct for linkage

disequilibrium in summary statistics of dense 5 million genome-wide markers and considerably

outperforms a state-of-the-art method across wide range of genetic architecture.

1 Introduction

In complex trait genetics, the ability to predict phenotype from genotype is the ultimate

measure of our understanding of the allelic architecture underlying the heritability of a trait.

Complete understanding of the genetic basis of a trait should allow for predictive methods having

accuracies approaching the trait's broad sense heritability. In addition to improving our

fundamental understanding of basic genetics, phenotypic prediction has obvious practical utility,

ranging from crop and livestock applications in agriculture to estimating the genetic component of

risk for common human diseases in medicine. For example, a portion of the current guideline on

the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk focuses on

estimating a patient's risk of developing disease [1]; in theory, genetic predictors have the

potential to reveal a substantial proportion of this risk early in life (even before clinical risk factors

are evident) enabling prophylactic intervention for high-risk individuals.

The field of phenotypic prediction originated in plant and animal genetics (reviewed in

refs. [2] and [3]). The first approaches relied on "major genes" – allelic variants of large effect

sizes readily detectable by genetic linkage or association. Similarly, after the early results of

human genome-wide association studies (GWAS) became available, the first risk predictors in

humans were based on combining the effects of markers significantly and reproducibly

associated with the trait, typically those with association statistics exceeding a genome-wide level

of significance [4,5].

More recent studies have demonstrated that a multitude of small effect alleles, not

detectable at a genome-wide level of statistical significance using currently existing sample sizes,

play an important role in complex trait genetics [2,3,6]. This has led to increasing popularity of

highly polygenic and even infinitesimal models of the allelic architecture underlying complex traits

[7,8]. Under the assumption that numerous alleles of small effect are involved in the trait,

predictors based only on GWAS-variants are expected to have low accuracy because they do not

include most of the DNA variants underlying the trait. New methods of phenotype prediction

leverage this insight by fitting all variants simultaneously [8–13]. These methods include

extensions of BLUP [10,11], or Bayesian approaches that extend both shrinkage techniques and

random effect models [8]. Newer methods benefited from allowing for classes of alleles with

vastly different effect size distributions. However, these methods require individual level genotype

data that do not exist for large meta-analyses and are computationally expensive.

"Polygenic scores" [7,14–17] represent an alternative approach based on summary

statistics. These scores, that are generally additive over genotypes weighted by apparent effect

sizes exceeding a given *p*-value threshold, have the potential to incorporate many relevant

variants undetected by GWAS, albeit at the expense of including numerous variants unrelated to

the trait. In theory, the risk predictor based on conditional mean effects can achieve the optimal

accuracy of linear risk models regardless of underlying genetic architecture by properly down-

weighting noise introduced by non-causal variants [18]. In practice, however, implementing the

conditional mean predictor poses a dilemma. In order to estimate the conditional mean effects,

we need to know the underlying genetic architecture first, but the true architecture is unknown

and difficult to model accurately. The current methods circumvent this issue by deriving

conditional means under a highly simplified model of genetic architecture. However, the optimality

of resulting risk predictor is not guaranteed when the true underlying genetic architecture deviates

from the assumed prior. In particular, recent studies have revealed complex dependencies of

heritability on minor allele frequency (MAF) and local genomic features such as regulatory

landscape and intensity of background selections [19–23]. Several studies proposed to extend

polygenic scores by incorporating additional complexity into the parametric Bayesian models,

however, these methods have not been demonstrated with the genome-wise set of markers due

to computational challenges [17,24,25]. Recently, there have been growing interests in non-

parametric and semi-parametric approaches, such as those based on modeling of latent variables

and kernel-based estimation of prior or marginal distributions, however, thus far these models

cannot leverage summary statistics or account for the linkage disequilibrium (LD) structure in the

data [26–29].

Here, we propose a novel risk prediction approach called partitioning-based non-

parametric shrinkage (NPS). Without specifying a parametric model of underlying genetic

architecture, we aim to estimate the conditional mean effects directly from the data. Large GWAS

studies often provide estimated effect sizes at each SNP in a discovery cohort as summary

statistics. We will show that using a second small independent training cohort with available

genotype-level data, we can learn the shrinkage weights for apparent effect sizes from the

discovery GWAS. In this way, we can leverage the large sample size of the discovery cohort,

which is typically available only as summary statistics rather than individual genotypes. We

evaluate the performance of this new approach under a simulated genetic architecture of dense 5

million SNPs across the genome.

Results

Method Overview

11   We start from a simple observation that conditional mean effects can be approximated by

12   piece-wise linear interpolation in the absence of LD (Fig 1A). We can partition SNPs into K

13   disjoint intervals based on observed effect sizes ($\hat{\beta}_j$) and fit a linear function $f(\hat{\beta}_j) = \omega_k \hat{\beta}_j$ on

14   each interval of $k = 1, \dots, K$. Specifically, when x_{ij} is the genotype of individual i at marker j and

15   β_j is the true effect size at marker j, the predicted phenotype $\hat{y}_i$ based on conditional mean

16   effects $E[\beta_j \mid \hat{\beta}_j]$ can be interpolated as the following:

18
$$\hat{y}_i = \sum_{j=1}^{M} E[\beta_j \mid \hat{\beta}_j] x_{ij} \approx \sum_{j=1}^{M} \left(\sum_{k=1}^{K} \omega_k \hat{\beta}_j I(b_{k-1} < \hat{\beta}_j \leq b_k) \right) x_{ij} \qquad (1)$$

20   where b_{k-1} and b_k are partition boundaries and $I(\cdot)$ is an indicator function for partition k. This

21   equation can be further simplified by changing the order of summation as below:

23
$$= \sum_{k=1}^{K} \omega_k \left(\sum_{j \in \mathcal{S}_k} \hat{\beta}_j x_{ij} \right) = \sum_{k=1}^{K} \omega_k G_{ik} \qquad (2)$$

25   where $\mathcal{S}_k$ is the set of all markers assigned to partition k. If we define a partitioned risk score G_{ik}

26   to be a risk score of individual i calculated using only SNPs in partition k, ω_k becomes equivalent

to the per-partition shrinkage weight. Based on equation (2), we can utilize a small genotype-level training cohort to estimate ω_k by fitting phenotypes y_i with partitioned risk scores G_{ik} of training individual i. We do not need to specify the underlying distribution of true genetic effects β_j since ω_k can be directly learned from the data.

We rely on a small independent cohort with full genotype information to estimate the shrinkage weights. For quantitative traits, the per-partition shrinkage weights can be estimated by linear regression. For binary phenotypes such as case/control status, ω_k can be derived from a linear discriminant analysis (LDA)-based classifier in the *K*-dimensional feature space formed by partitioned risk scores G_{ik}. LDA guarantees the optimal accuracy of classifier when case and control subgroups follow multivariate normal distributions in the feature space. The partitioned risk scores of cases and controls indeed follow approximately normal distributions as far as each partition consists of an enough number of SNPs [30]. Furthermore, we can assume that the covariance of partitioned risk scores approximately equal between cases and controls and is a diagonal matrix because 1) G_{ik} of a single partition explains only a small fraction of phenotypic variation on the observed scale with typical GWAS sample sizes [31] and 2) simulations suggest that covariance across partitions is negligibly small despite the liability thresholding effect. Under these conditions, LDA-derived shrinkage weights can be independently estimated for each partition and simplify to:

$$\omega_k \approx 2 \frac{E[\,G_{ik}\,|\,y_i = 1] - E[\,G_{ik}\,|\,y_i = 0]}{var[\,G_{ik}\,|\,y_i = 1] + var[\,G_{ik}\,|\,y_i = 0]} \tag{3}$$

where the phenotype of an individual y_i is encoded in 0 or 1 depending on being a control or a case. Note that estimating ω_k is a far easier problem than estimating per-SNP effects. The training sample size is small but still larger than the number of parameters to estimate, namely *K*, whereas for $\hat{\beta}_j$, the GWAS sample size is considerably smaller than the number of markers.

In the presence of LD, we cannot apply piece-wise linear interpolation directly on observed effect sizes because the conditional mean effect at each SNP depends on genetic

effects of all adjacent SNPs in LD. To address the problem, we split the genome into ~2 Mbps

windows, and in each window, we project SNP genotypes into a decorrelated space by a linear

transformation obtained by the spectral decomposition of local LD matrix (Fig 1B, See Methods).

Specifically, when $\boldsymbol{D}$ is a local reference LD matrix obtained from training genotypes and

decomposes into $\boldsymbol{Q}\,\boldsymbol{\Lambda}\,\boldsymbol{Q}^T$, where $\boldsymbol{\Lambda}$ is a diagonal matrix of non-negative eigenvalues λ_j and $\boldsymbol{Q}$ is a

matrix of orthonormal eigenvalues q_j, the decorrelating projection is defined as:

8
$$\mathcal{P} = \boldsymbol{\Lambda}^{-1/2}\,\boldsymbol{Q}^T \qquad\qquad (4)$$

10   and for the axis of projection defined by q_j, the transformation $\mathcal{P}$ yields the projected genotype x_{ij}^P

11   and decorrelated effect estimate $\hat{\eta}_j$ as the following:

13
$$x_{ij}^P = \frac{1}{\sqrt{\lambda_j}} q_j^T x_i$$

14   and $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad (5)$

15
$$\hat{\eta}_j = \frac{1}{\sqrt{\lambda_j}} q_j^T \hat{\beta}$$

where x_i and $\hat{\beta}$ are m-dimensional vectors of genotypes and per-SNP estimated effects,

respectively, at all SNPs, and m is the number of SNPs in the window. Under the projection, both

projected genotypes and sampling noises of decorrelated effect estimates become uncorrelated

across different axes of projection (Supplementary Note). Although strictly speaking, the lack of

correlation does not imply independence, we apply the non-parametric shrinkage technique as if

x_{ij}^P and $\hat{\eta}_j$ are from unlinked markers.

Partitioning strategy

We use a higher number of partitions to better interpolate the mean conditional effect

curve. Since the full combinatorial optimization of partitioning cut-offs b_k is neither necessary or

practical, we place the cut-offs based on heuristics without optimizing them on individual dataset

(See Methods, Supplementary Note and Fig S1). Note that widely-used *p*-value thresholding

approach is a special case of non-parametric shrinkage with the number of partition *K* being 2.

When SNP effect sizes are measured relative to standardized genotypes, thresholding on *p*-

values of association is equivalent to thresholding on estimated SNP effect sizes.

In the presence of LD, non-parametric shrinkage requires double partitioning on both λ_j

and $\hat{\eta}_j$ because the conditional mean effects in the decorrelated space, i.e. $E\left[\eta_j \mid \hat{\eta}_j\right]$, depend not

only on $\hat{\eta}_j$ but also on eigenvalue λ_j of the projection vector. To provide intuition behind this, an

eigenvalue captures how many correlated SNPs are tracked by a projection vector, and we can

show that the scale of true underlying decorrelated effect sizes, namely $var\left[\eta_j\right]$, is proportional to

the eigenvalue λ_j (Supplementary Note). Thus, with increasing eigenvalues, the scale of

decorrelated effects becomes large relative to the sampling error $\frac{1}{N}$. Therefore, estimated effects

become more reliable. On the other hand, with decreasing eigenvalues, $\hat{\eta}_j$ becomes dominated

by sampling error. In addition, conditional mean effects depend on $\hat{\eta}_j$ as a function of

polygenicity. If we consider genetic architecture with a low proportion of causal SNPs, some of

projection vectors may not involve even a single causal SNP, and in this case, their true

decorrelated effect size η_j will be 0. Such non-causal projection vectors will be enriched among

small estimates of $\hat{\eta}_j$, thus of which conditional mean effects become attracted toward 0.

In the special case of infinitesimal genetic architecture in which all SNPs are causal with

normally distributed effect sizes, we already know analytically derived conditional mean effects

[17], therefore can cross-check the accuracy of our shrinkage weights ω_k estimated by non-

parametric shrinkage (Fig 2A-B, Supplementary Note). For this, we simulated a genome

consisted of 1,236 loci of 2 Mb each, of which LD structures were sampled from real human

haplotypes but each locus was treated unlinked from each other for simplicity (See Methods). The

ratio of discovery cohort size to number of markers were set to 1, and an independent cohort of

2,500 cases and 2,500 controls was set aside for training. To apply the non-parametric shrinkage,

we first partitioned decorrelated effect sizes into 10 subgroups on intervals of eigenvalues λ_j

while allocating equal amount of $\sum_j \lambda_j$ to each partition (Fig 2A). The per-partition shrinkage

weights ω_k trained by non-parametric shrinkage closely tracked the theoretical optimum in most

of the bins. However, interestingly, in the lowest and highest partitions of eigenvalues, $\mathcal{S}_{\lambda 1}$ and

$\mathcal{S}_{\lambda 10}$, the estimated shrinkage was significantly biased away from the optimal curve. The smallest

eigenvalues are too noisy to estimate with the reference LD panel. Therefore, it is correct to

down-weight ω_1 almost to 0. In case of partition $\mathcal{S}_{\lambda 10}$, it spans the widest interval of eigenvalues

but consists of the fewest number of SNPs. While it is ideal to apply a finer partitioning in this

interval so to better interpolate the theoretical curve, the total number of SNPs and independent

projection vectors in the genome are the fundamental limiting factor.

In case of infinitesimal architecture, theory predicts that per-partition shrinkage weights

depend only on eigenvalues and not on estimated effect sizes. To examine the robustness of

non-parametric shrinkage, we applied general 10-by-10 double partitioning on λ_j and $\hat{\eta}_j$ collected

under infinitesimal simulations. Similarly to eigenvalues, we placed partition boundaries on the

interval of $\hat{\eta}_j$ keeping each partition to have equal $\sum_j \hat{\eta}_j{}^2$. This is a heuristic approach to distribute

partition boundaries toward the intermediate range of $\hat{\eta}_j$ and not to over-partition the largest

values of $\hat{\eta}_j$. This approach works reliably well across a wide range of simulation conditions we

tested. In overall, the shrinkage weights estimated by double partitioning agree with the

theoretical expectation. The estimated conditional mean effects, interpolated with $\omega_k \hat{\eta}_j$, follow the

linear trajectory (Fig 2B for $\mathcal{S}_{\lambda 10}$, Fig S2 for other partitions of eigenvalues).

For non-infinitesimal genetic architecture, we do not have an analytic derivation of

conditional mean effects, therefore empirically estimated the conditional means using the true

underlying effects η_j and true LD structure of the population (See Methods). One percent of

SNPs were simulated to be causal with normally distributed effect sizes. The ratio between

number of discovery samples to number of markers was set to 1:1. As expected, the true

conditional mean dips for the lowest values of $\hat{\eta}_j$ but catches up toward no shrinkage ($\omega_k = 1$)

with increasing values of $\hat{\eta}_j$ (Fig 2C-D). A notable difference between the partitions of largest

eigenvalues $\mathcal{S}_{\lambda 10}$ (Fig 2C) and second smallest eigenvalues $\mathcal{S}_{\lambda 2}$ (Fig 2D) is that the true

conditional mean is very close to almost no shrinkage for large $\hat{\eta}_j$ in the former. This is because

the scale of true effects η_j is proportional to eigenvalues, therefore, with large enough λ_j, the

sampling error becomes relatively small and the estimated effect sizes more accurate. In all

partitions, conditional mean effects estimated by non-parametric shrinkage stayed very close to

the true conditional means (Fig S3 for eigenvalue partitions other than $S_{\lambda 2}$ and $S_{\lambda 10}$). The

observed consistency between estimated and true conditional means is not because of shared

underlying data. At each simulation run, we re-generated the entire dataset starting from fresh

sampling of true underlying genetic effects β_j under the same genetic architecture parameters.

Simulated benchmark

To benchmark the accuracy of non-parametric shrinkage, we simulated the genetic

architecture using the real LD structure of dense 5 million common SNPs from the 1000

Genomes Project (See Methods). For baseline simulations, we considered a simple mixture-

normal genetic architecture with the causal fraction of SNPs being 1% to 0.01% (50,000 to 500

SNPs). This is the model most commonly used as a Bayesian prior. With decreasing causal

fractions, the accuracy of non-parametric shrinkage improved in terms of both liability-scale R^2 as

well as Nagelkerke's R^2 on the observed scale (Table 1). With fewer causal SNPs, per-SNP

effect sizes increase making it easier to predict the phenotype.

We incrementally added more complexity to simulated genetic architecture starting from

the baseline model. First, we incorporated the dependency of heritability on minor allele

frequency (MAF) into the simulation. Because of this dependency, recent studies have reported

that low frequency SNPs contribute less heritability than previously expected under no

dependency [20]. Low-frequency SNPs tend to be captured by eigenvectors of small eigenvalues

and are challenging to handle with spectral decomposition. Lowering the overall heritability

contribution of low-frequency SNPs made non-parametric shrinkage predict more accurately

(Table 1).

Further, we introduced clumping of causal SNPs in open chromatins by making the

fraction of causal SNPs five-fold higher in DNase I hypersensitive sites (DHS), comprising 15% of

the genome, compared to the rest of genome. The prediction accuracy of non-parametric

shrinkage went down slightly compared to simulations of uniformly distributed causal SNPs but

still remained robust even if we did not explicitly account for DHS overlap in the current version of

non-parametric shrinkage (Table 1).

We evaluated the performance of non-parametric shrinkage vis-a-vis a comparable

successful parametric technique (Table 1). LDPred is the state-of-the-art parametric method,

which is similarly based on summary statistics estimated in large GWAS datasets and an

independent training set with individual-level data. As a Bayesian method, it is optimal for the

spike and slab allelic architecture. We found that our method resulted in more accurate

predictions than LDPred across a range of genome-wide simulations. This is seemingly surprising

given that the simulated allelic architectures are close to the LDPred assumptions. However, we

found that in most simulations, LDPred adopted the infinitesimal model irrespectively to the true

simulated regime pointing to the challenge of the computational optimization in the parametric

case. Notably, in a single simulation, where LDPred correctly selected a non-infinitesimal model

with a low causal fraction (0.01%), it achieved high prediction accuracy. This is reflected in Table

1 as a high confidence interval in one of the LDPred simulation sets. The simulations suggest that

the well optimized parametric model is capable to generate good predictions, but non-parametric

shrinkage is much more robust and does not suffer from optimization issues. This results in a

higher performance of NPS in practice. Overall, in our most realistic simulations incorporating

DHS clumping and MAF dependency, our method showed 1.20 and 1.38-fold higher accuracy in

Nagelkerke's R^2 compared to LDPred for causal fractions of 1% and 0.1% (P = 0.0007 and 0.001,

respectively).

Discussion

Understanding how phenotype maps to genotype has always been a central question of

basic genetics. With the explosive growth in the amount of training data, there is also a clear

prospect and enthusiasm for clinical applications of the polygenic risk prediction [32,33]. The

current reality is, however, that most large-scale GWAS datasets are available in the form of

summary statistics only. Nonetheless, data on a limited number of cases are frequently available

from epidemiological cohorts such as UK Biobank or from public repositories with a secured

access such as dbGaP. This motivated us to develop a method that is primarily based on

summary statistics but also benefits from smaller training data at the raw genotype resolution.

Although we heavily rely on the training data to construct a prediction model, the requirement for

out-of-sample training data is not unique for our method. Widely-used thresholding-based

polygenic scores and Bayesian parametric methods also need genotype-level data to optimize

their model parameters [17,34].

Human phenotypes vary in the degree of polygenicity [22], in the fraction of heritability

attributable to low-frequency variants [20] and in other aspects of allelic architecture [21,23]. Our

method is agnostic with respect to the underlying allelic architecture. This makes it potentially

equally applicable to a wide range of phenotypes.

A stream of recent publications point to a disproportionate contribution to heritability of

protein coding genes and regulatory regions highlighted by chromatin accessibility and epigenetic

modifications [19,35–38]. This heritability enrichment is often observed to be cell-type specific.

While in practice, we do not have the comprehensive annotations of functional genomics data

relevant to the phenotype of interest, incorporating available external annotations into the model

could increase the accuracy of polygenic risk prediction [24,25]. Although we do not explore this

direction in the present work, our method opens a clear perspective for the incorporation of the

functional data. The current scheme partitions summary statistics only based on the apparent

effect size, however, the same partitioning strategy can be easily applied to any functional

genomics variable.

Similarly to other methods, our method assumes that all datasets come from a

homogeneous population. It is well appreciated that polygenic risk models are not transferrable

between populations due to differences in allele frequencies and patterns of linkage

disequilibrium [39]. This problem should be addressed by the future work in the field.

26
27

Materials and Methods

Non-parametric shrinkage

The estimated effect sizes $\hat{\beta}_j$ at SNPs j = 1, ..., M are available as summary statistics

from a large discovery GWAS study of N samples. When the estimated effects were prepared as

per-allele effects, we converted them relative to standardized genotypes by multiplying

$\sqrt{2f_j(1-f_j)}$, where f_j is the allele frequency of SNP j in the discovery GWAS cohort. The logistic

per-allele effect sizes obtained from case/control GWAS were treated in the same way. In

addition, non-parametric shrinkage (NPS) requires a small independent genotype-level training

cohort in order to build the prediction model.

We processed the effect size estimates of discovery GWAS and training cohort of sample

size N' in non-overlapping windows of 4,000 SNPs each (~2.4 Mb in length). In each window,

given an N' × 4,000 standardized genotype matrix X, the raw reference LD matrix $D = \frac{1}{N'}X^T X$

was regularized in order to suppress sampling noise particularly in off-diagonal entries.

Specifically, pairwise LD was set to 0 if the SNPs were separated by > 500 kb or the absolute

value of estimated LD was smaller than $5/\sqrt{N'}$. Since the standard error of pairwise LD is

approximately $1/\sqrt{N'}$ under no correlation, we expect that on average, only 1.7 uncorrelated SNP

pairs escape the above regularization threshold per window. The regularized LD matrix D^* was

factorized into the following by spectral decomposition:

$$D^* = Q\,\Lambda\,Q^T$$

where Λ is a diagonal matrix of eigenvalues and Q is an orthonormal matrix of eigenvectors.

Since D^* is not non-negative semi-definite, Λ contained negative eigenvalues. Negative

eigenvalues were truncated along with those positive but smaller than 0.5 since they were

dominated by noise. Applying the decorrelating linear transformation $\mathcal{P}$ (equation 4), we obtained

decorrelated genotypes x_{ij}^P and decorrelated effect size estimates $\hat{\eta}_j$ for each projection defined

by eigenvalue λ_j and corresponding eigenvector q_j (equation 5), where i is an individual and j is

the index for decorrelating projection.

Although we chose the window size to be large enough to capture the majority of local LD

patterns, some LD structures, particularly near the edge, span across windows, which in turn

yield residual cross-window correlations. To eliminate such correlations, we applied LD pruning in

decorrelated space between adjacent windows. Specifically, we calculated Pearson correlations

$\rho_{jj'}$ between decorrelated genotypes x_j^P and $x_{j'}^P$, where j and j' are the axes of projection,

belonging to distinct windows. For the pairs with $|\rho_{jj'}| > 0.3$, we kept the one yielding larger

decorrelated effect sizes and eliminated the other.

Next, we merged decorrelated effect sizes $\hat{\eta}_j$ across all windows and defined the 10 × 10

double-partitioning boundaries on intervals of λ_j and $|\hat{\eta}_j|$. The eigenvalues were split to 10

intervals of λ_j, equally distributing $\sum_j \lambda_j$ across partitions. The partitions on eigenvalues are

denoted here by $\mathcal{S}_{\lambda 1}, \dots, \mathcal{S}_{\lambda 10}$ from the lowest to the highest. Each partition of eigenvalues was

sub-partitioned on intervals of $|\hat{\eta}_j|$, equally distributing $\sum_j \hat{\eta}_j^2$ across partitions. The partition

boundaries of $|\hat{\eta}_j|$ were defined separately for each partition of eigenvalues because the

distribution of $|\hat{\eta}_j|$ is dependent on λ_j.

In each sub-partition k, we calculated a partitioned risk score G_{ik} of training individual i as

the following:

17
$$G_{ik} = \sum_{j \in \mathcal{S}_k} \hat{\eta}_j \, x_{ij}^P$$

Given the phenotype y_i of each individual in training cohort, we estimated per-partition shrinkage

weights ω_k by linear discriminant analysis (LDA) using the equation (3). After re-weighting each

$\hat{\eta}_j$ by shrinkage weight of corresponding partition, we converted the decorrelated effect sizes

back to the original SNP effect space. This back-conversion has to be done window-by-window

using all $\hat{\eta}_j$ belonging to each window. Specifically, when a given window has m decorrelated

effects, $\hat{\eta}_{j_o}, \hat{\eta}_{j_o+1}, \dots, \hat{\eta}_{j_o+m-1}$, we applied the following back-transformation:

24
$$\boldsymbol{Q} \, \boldsymbol{\Lambda}^{-1/2} \begin{pmatrix} \omega_{k(j_0)} \, \hat{\eta}_{j_o} \\ \vdots \\ \omega_{k(j_0+m-1)} \, \hat{\eta}_{j_o+m-1} \end{pmatrix}$$

where $k(j)$ stands for the partition such that $j \in \mathcal{S}_k$ (Supplementary Note). The resulting re-

weighted SNP effect sizes is relative to standardized genotypes and can be converted to per-

allele effects by correcting for allele frequencies.

Because the accuracy of decorrelating projection declines near the edge of windows, the

overall performance of NPS is affected by the placement of window boundaries relative to

locations of strong genome-wide association peaks. To alleviate such dependency, we repeated

the same NPS procedure shifting by 1,000, 2,000, and 3,000 SNPs and took the average

reweighted effect sizes across four NPS runs.

Simulation with LD confined to each locus

To show that shrinkage weights estimated by NPS approximate conditional mean effects

in the decorrelated space, we simulated genetic architecture with SNPs in LD. The LD matrix was

calculated using the genotypes of the 1000 Genomes Project CEU panel (2n=198) for a total of

101,296 SNPs, which were obtained by 10-fold down-sampling of Illumina HumanHap550

genotyping SNPs. SNPs with minor allele frequency (MAF) < 5% were filtered out. The genome

was broken down to 2 Mb loci, and SNP-poor loci with less than 40 SNPs were excluded. Overall,

a total of 1,236 loci with 82 SNPs on average were used for this simulation. Since the raw LD

matrix was calculated with the reference LD panel of a small sample size, we suppressed

spurious long-range LD by setting the LD between SNPs separated by > 500 kb to 0. For

simplicity, we confined LD structure to each locus and disallowed LD spanning across loci.

We considered two genetic architectures: infinitesimal model with normally distributed

effect sizes and non-infinitesimal model for which 1% SNPs are casual with normally distributed

effect sizes. The discovery GWAS summary statistics were directly sampled from the following *m*-

dimensional multivariate normal distribution one locus each time:

25
$$\hat{\beta} \sim N(\mu = \beta \boldsymbol{D}, \Sigma = \frac{1}{N}\boldsymbol{D})$$

where $\hat{\beta}$ and β are *m*-dimensional vectors of true and estimated effect sizes of the locus, *D* is a

local LD matrix, and *N* is the discovery GWAS sample size. *N* was set to equal to the genome-

wide number of markers *M*. The standardized genotypes of training cohort were also generated

using a multivariate normal distribution:

$$X_i \sim N(\mu = 0, \Sigma = \boldsymbol{D})$$

where X_i is an *m*-dimensional genotype vector of individual i. We generated genotypes for 50,000

individuals and simulated phenotypes under a liability threshold model with the heritability h^2 of

0.5 and prevalence of 5%. By down-sampling controls, we assembled a training case/control

cohort of 2,500 cases and 2,500 controls.

For NPS, each locus was treated as a single analysis window without averaging over

sliding windows. The true underlying LD matrix $\boldsymbol{D}$ was hidden, and NPS had to estimate the

reference LD from the training cohort. For infinitesimal model, the theoretically optimal shrinkage

weight ω_k^o is known (Supplementary Note) [17]:

$$\omega_k^o = \lambda_j / (\lambda_j + \frac{M}{Nh^2})$$

For non-infinitesimal model, we do not have analytically known optimal shrinkage, therefore

instead empirically estimated it by regressing conditional mean effects against $\hat{\eta}_j$ with the fixed

intercept of 0 as follows:

$$E\left[\eta_j \mid \hat{\eta}_j \in \mathcal{S}_k\right] \sim \omega_k^o \hat{\eta}_j + 0$$

Then, we took the average of ω_k^o over 40 runs of simulations under the same genetic architecture

parameters. $E\left[\eta_j \mid \hat{\eta}_j \in \mathcal{S}_k\right]$ was estimated by taking the sample mean of true decorrelated effects

η_j in each partition $\mathcal{S}_k$. We calculated η_j by:

$$\eta_j = \sqrt{\lambda_j}\, \boldsymbol{q}_j^T \beta$$

using the true genetic effects β and spectral decomposition of true population LD matrix. The

shrinkage weights estimated by NPS are not in the same absolute scale as theoretical values.

Thus, we rescaled NPS estimates to the same scale as theoretical values by multiplying by a

global scaling factor. The accuracy of prediction model is not affected by global rescaling. For

infinitesimal model, the global scaling factor was $\sqrt{\sum_j (\omega_k^o \hat{\eta}_j)^2 / \sum_j (\omega_k \hat{\eta}_j)^2}$. For non-infinitesimal

model, it was obtained by regressing ω_k^o against estimated ω_k with fixed intercept of 0 by fitting

$\omega_k^o \sim \omega_k + 0$ across 100 overall partitions.

Simulation of genetic architecture with dense genome-wide markers

For simulated benchmarks, we generated genetic architecture with dense ~5 million

genome-wide markers from the 1000 Genomes Project. We kept only SNPs with MAF > 5% and

Hardy-Weinberg equilibrium test *p*-value > 0.001. We used EUR panel (2n=808) to populate LD

structures in simulated genetic data. Due to the limited sample size of LD panel, we regularized

the LD matrix by Schur product with a tapered banding matrix [40]. Briefly, we preserved the raw

pairwise LD for SNP pairs less than 150 kb apart, and then proportionally tapered them to 0 for

SNP pairs up to 300 kb apart, and finally set to 0 for distances over 300 kb. The resulting

regularized LD matrix $\boldsymbol{D^*}$ is not positive semi-definite. Thus, we approximated it to a positive

semi-definite LD matrix $\boldsymbol{D^o}$ constructed by:

$$\boldsymbol{D^o} = \boldsymbol{Q}\,\boldsymbol{\Lambda'}\boldsymbol{Q}^T + 0.01\,\boldsymbol{I}$$

where $\boldsymbol{D^*} = \boldsymbol{Q}\,\boldsymbol{\Lambda}\,\boldsymbol{Q}^T$ by spectral decomposition and $\boldsymbol{\Lambda'}$ is the matrix of eigenvalues $\boldsymbol{\Lambda}$ with negative

values masked out to 0.

Next, we generated genotypes across the entire genome simulating the genome-wide

patterns of LD. We assume that the standardized genotypes follow a multivariate normal

distribution (MVN). Since we assume that LD travels no farther than 300 kb, as far as we simulate

genotypes in blocks of length greater than 300kb, we can simulate the entire chromosome

without losing any LD patterns by utilizing a conditional multivariate normal distribution as the

following. The genotypes for the first block of 2,500 SNPs were sampled directly out of $N(\mu =$

$0, \Sigma = \boldsymbol{D}^o_{(1)})$. From the next block, we sampled the genotypes of 1,250 SNPs (average 0.75 Mb in

length) each, conditional on the genotypes of previous 1,250 SNPs. When the genotype of block l

is x_l and the LD matrix spanning block l and $l + 1$ is split into submatrices as the following:

$$\begin{pmatrix} \boldsymbol{D}^o_{(l)} & \boldsymbol{D}^o_{l,l+1} \\ \boldsymbol{D}^o_{l+1,l} & \boldsymbol{D}^o_{(l+1)} \end{pmatrix}$$

then, the genotype of next block $l + 1$ follows a conditional multivariate normal distribution as:

1. $$X_{l+1}|X_l = x_l \sim N\big(\mu = \mathbf{D}^o_{l+1,l}(\mathbf{D}^o_l)^{-1}x_l, \Sigma = \mathbf{D}^o_{(l+1)} - \mathbf{D}^o_{l+1,l}(\mathbf{D}^o_l)^{-1}\mathbf{D}^o_{l,l+1}\big)$$

2. After the genotype of entire chromosome was generated in this way, the standardized genotype

3. values were converted to allelic genotypes by taking the highest Nf_j^2 and lowest $N(1-f_j)^2$

4. genotypes as homozygotes and the rest as heterozygotes under the assumption of Hardy-

5. Weinberg equilibrium. N is the number of simulated samples, and f_j is the allele frequency of

6. SNP j.

7.   We simulated three different sets of genetic architecture: mixture normal, MAF

8. dependency and DNase I sensitive sites (DHS). The mixture normal is a spike-and-slab

9. architecture in which a fraction of SNPs have normally distributed causal effects β_j as below:

10. $$\beta_j \sim pN(0,1) + (1-p)\delta_0$$

11. where p is the fraction of causal SNPs (1, 0.1 or 0.01%) and δ_0 is a point mass at the effect size

12. of 0. For the MAF-dependent model, we extended the mixture normal so that the scale of causal

13. effect sizes varied across SNPs in proportion to $\big(f_j(1-f_j)\big)^\alpha$ with $\alpha = -0.25$ [20] as follows:

14. $$\beta_j \sim p\, N\left(0, \big(f_j(1-f_j)\big)^\alpha\right) + (1-p)\delta_0$$

15. Finally, for the DHS model, we further extended the MAF-dependent architecture to exhibit

16. clumping of causal SNPs within DHS peaks. Fifteen per cents of simulated SNPs were located in

17. the master DHS sites that we downloaded from the ENCODE project. We assumed a five-fold

18. higher causal fraction in DHS (p_{DHS}) compared to the rest of genome in order to simulate the

19. enrichment of per-SNP heritability in DHS reported in the previous study [19]. Specifically, β_j was

20. sampled from the following distribution:

21. $$\beta_j \sim \begin{cases} p_{DHS}\, N\left(0, \big(f_j(1-f_j)\big)^\alpha\right) + (1-p_{DHS})\delta_0 & \text{if SNP } j \text{ is in DHS} \\ \frac{1}{5}\, p_{DHS} N\left(0, \big(f_j(1-f_j)\big)^\alpha\right) + \left(1 - \frac{1}{5}p_{DHS}\right)\delta_0 & \text{otherwise} \end{cases}$$

22.   In each genetic architecture, we simulated phenotypes for discovery, training and

23. validation populations of 100,000, 50,000 and 50,000 samples, respectively, using a liability

24. threshold model of the heritability of 0.5 and prevalence of 0.05. In the discovery population, we

25. obtained GWAS summary statistics by testing for the association with the total liability instead of

case/control status. This is computationally easier than to generate a case/control GWAS cohort

of a realistic sample size because of the low prevalence. With the prevalence of 0.05, statistical

power of quantitative trait association studies using the total liability is roughly similar to those of

dichotomized case/control GWAS studies of same sample sizes [41]. For the training dataset, we

assembled a cohort of 2,500 cases and 2,500 controls by down-sampling controls out of the

simulated population of 50,000 samples. The validation population was used to evaluate the

accuracy of prediction model in terms of R^2 of the liability explained and Nagelkerke's R^2 to

explain case/control outcomes.

Benchmark evaluation of LDPred

11   We benchmarked LDPred on the exactly same datasets tested for NPS. The window size

was set to 1,670 SNPs in order to split the genome into 3,000 windows as recommended [17].

LDPred requires to optimize the causal fraction parameter using the training cohort available as

individual-level genotype data. We tested the casual SNP fractions of 1, 0.3, 0.1, 0.03, 0.01,

0.003, 0.001, 0.0003 and 0.0001 and selected the model parameter yielding the highest

prediction accuracy in the training cohort. This model was benchmarked in the validation cohort.

For fair comparison, we did not generate complementary alleles in simulations since they were

filtered out by LDPred automatically.

19
20

References

1. Stone NJ, Robinson JG, Lichtenstein AH, Bairey Merz CN, Blum CB, Eckel RH, et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. Circulation. 2014;129: S1-45. doi:10.1161/01.cir.0000437738.63853.7a

2. Goddard ME, Hayes BJ. Mapping genes for complex traits in domestic animals and their use in breeding programmes. Nat Rev Genet. 2009;10: 381–391. doi:10.1038/nrg2575

3. Falke KC, Glander S, He F, Hu J, de Meaux J, Schmitz G. The spectrum of mutations controlling complex traits and the genetics of fitness in plants. Curr Opin Genet Dev. 2013;23: 665–671. doi:10.1016/j.gde.2013.10.006

4. Ripatti S, Tikkanen E, Orho-Melander M, Havulinna AS, Silander K, Sharma A, et al. A multilocus genetic risk score for coronary heart disease: case-control and prospective cohort analyses. Lancet. 2010/10/26. 2010;376: 1393–1400. doi:S0140-6736(10)61267-6 [pii]10.1016/S0140-6736(10)61267-6

5. Wacholder S, Hartge P, Prentice R, Garcia-Closas M, Feigelson HS, Diver WR, et al. Performance of common genetic variants in breast-cancer risk models. N Engl J Med. 2010; doi:10.1056/NEJMoa0907727

6. Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, et al. Common SNPs explain a large proportion of the heritability for human height. Nat Genet. Nature Publishing Group; 2010;42: 565–9. doi:10.1038/ng.608

7. Chatterjee N, Wheeler B, Sampson J, Hartge P, Chanock SJ, Park JH. Projecting the performance of risk prediction based on polygenic analyses of genome-wide association studies. Nat Genet. 2013;45: 400–5, 405e1–3. doi:10.1038/ng.2579

8. Zhou X, Carbonetto P, Stephens M. Polygenic modeling with bayesian sparse linear mixed models. Visscher PM, editor. PLoS Genet. Public Library of Science; 2013;9: e1003264. doi:10.1371/journal.pgen.1003264

9. Abraham G, Tye-Din JA, Bhalala OG, Kowalczyk A, Zobel J, Inouye M. Accurate and

robust genomic prediction of celiac disease using statistical learning. PLoS Genet. 2014;10: e1004137. doi:10.1371/journal.pgen.1004137

10. Golan D, Rosset S. Effective Genetic-Risk Prediction Using Mixed Models. Am J Hum Genet. Elsevier; 2014;95: 383–393. doi:10.1016/j.ajhg.2014.09.007

11. Speed D, Balding DJ. MultiBLUP: improved SNP-based prediction for complex traits. Genome Res. 2014;24: 1550–1557. doi:10.1101/gr.169375.113

12. Zhang Z, Ober U, Erbe M, Zhang H, Gao N, He J, et al. Improving the accuracy of whole genome prediction for complex traits using the results of genome wide association studies. PLoS One. 2014;9: e93017. doi:10.1371/journal.pone.0093017

13. Weissbrod O, Geiger D, Rosset S. Multikernel linear mixed models for complex phenotype prediction. Genome Res. 2016;26: 969–979. doi:10.1101/gr.201996.115

14. Purcell SM, Wray NR, Stone JL, Visscher PM, O'Donovan MC, Sullivan PF, et al. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. Nature. 2009;460: 748–52. doi:10.1038/nature08185

15. Stahl E a, Wegmann D, Trynka G, Gutierrez-Achury J, Do R, Voight BF, et al. Bayesian inference analyses of the polygenic architecture of rheumatoid arthritis. Nat Genet. Nature Publishing Group; 2012;44: 483–9. doi:10.1038/ng.2232

16. Shi J, Park JH, Duan J, Berndt ST, Moy W, Yu K, et al. Winner's Curse Correction and Variable Thresholding Improve Performance of Polygenic Risk Modeling Based on Genome-Wide Association Study Summary-Level Data. PLoS Genet. 2016;12: e1006493. doi:10.1371/journal.pgen.1006493

17. Vilhjalmsson BJ, Yang J, Finucane HK, Gusev A, Lindstrom S, Ripke S, et al. Modeling Linkage Disequilibrium Increases Accuracy of Polygenic Risk Scores. Am J Hum Genet. 2015;97: 576–592. doi:10.1016/j.ajhg.2015.09.001

18. Goddard ME, Wray NR, Verbyla K, Visscher PM. Estimating Effects and Making Predictions from Genome-Wide Marker Data. Stat Sci. 2009;24: 517–529. doi:10.1214/09-STS306

19. Gusev A, Lee SH, Trynka G, Finucane H, Vilhjálmsson BJ, Xu H, et al. Partitioning

heritability of regulatory and cell-type-specific variants across 11 common diseases. Am J

Hum Genet. Elsevier; 2014;95: 535–52. doi:10.1016/j.ajhg.2014.10.004

20. Speed D, Cai N, Johnson MR, Nejentsev S, Balding DJ. Reevaluation of SNP heritability

in complex human traits. Nat Genet. Nature Publishing Group; 2017;49: 986–992.

doi:10.1038/ng.3865

21. Gazal S, Finucane HK, Furlotte NA, Loh PR, Palamara PF, Liu X, et al. Linkage

disequilibrium-dependent architecture of human complex traits shows action of negative

selection. Nat Genet. Nature Publishing Group; 2017;49: 1421–1427. doi:10.1038/ng.3954

22. Boyle EA, Li YI, Pritchard JK. An Expanded View of Complex Traits: From Polygenic to

Omnigenic. Cell. Elsevier; 2017;169: 1177–1186. doi:10.1016/j.cell.2017.05.038

23. Zeng J, de Vlaming R, Wu Y, Robinson MR, Lloyd-Jones LR, Yengo L, et al. Signatures of

negative selection in the genetic architecture of human complex traits. Nat Genet. Nature

Publishing Group; 2018;50: 746–753. doi:10.1038/s41588-018-0101-4

24. Hu Y, Lu Q, Powles R, Yao X, Yang C, Fang F, et al. Leveraging functional annotations in

genetic risk prediction for human complex diseases. PLoS Comput Biol. 2017;13: 1–16.

doi:10.1371/journal.pcbi.1005589

25. Hu Y, Lu Q, Liu W, Zhang Y, Li M, Zhao H. Joint modeling of genetically correlated

diseases and functional annotations increases accuracy of polygenic risk prediction. PLoS

Genet. 2017;13: 1–22. doi:10.1371/journal.pgen.1006836

26. Zeng P, Zhou X. Non-parametric genetic prediction of complex traits with latent Dirichlet

process regression models. Nat Commun. Springer US; 2017;8: 1–11.

doi:10.1038/s41467-017-00470-2

27. Efron B. Empirical bayes estimates for large-scale prediction problems. J Am Stat Assoc.

2009;104: 1015–1028. doi:10.1198/jasa.2009.tm08523

28. So HC, Sham PC. Improving polygenic risk prediction from summary statistics by an

empirical Bayes approach. Sci Rep. 2017;7: 1–11. doi:10.1038/srep41262

29. Gianola D, Fernando RL, Stella A. Genomic-Assisted Prediction of Genetic Value with

Semiparametric Procedures. Genetics. 2006;173: 1761–1776.

doi:10.1534/genetics.105.049510

30.   Wray NR, Yang J, Goddard ME, Visscher PM. The genetic interpretation of area under the

ROC curve in genomic profiling. Schork NJ, editor. PLoS Genet. Public Library of Science;

2010;6: e1000864. doi:10.1371/journal.pgen.1000864

31.   Dudbridge F. Power and predictive accuracy of polygenic risk scores. PLoS Genet.

2013;9: e1003348. doi:10.1371/journal.pgen.1003348

32.   Khera A V, Chaffin M, Aragam KG, Emdin CA, Klarin D, Haas M, et al. Genome - wide

polygenic score to identify a monogenic risk - equivalent for coronary disease. bioRxiv.

2017; doi:10.1101/218388

33.   Riglin L, Collishaw S, Richards A, Thapar AK, Maughan B, O'Donovan MC, et al.

Schizophrenia risk alleles and neurodevelopmental outcomes in childhood: a population-

based cohort study. The Lancet Psychiatry. Elsevier Ltd; 2017;4: 57–62.

doi:10.1016/S2215-0366(16)30406-0

34.   Wray NR, Yang J, Hayes BJ, Price AL, Goddard ME, Visscher PM. Pitfalls of predicting

complex traits from SNPs. Nat Rev Genet. Nature Publishing Group; 2013;14: 507–15.

doi:10.1038/nrg3457

35.   Maurano MT, Humbert R, Rynes E, Thurman RE, Haugen E, Wang H, et al. Systematic

localization of common disease-associated variation in regulatory DNA. Science.

2012;337: 1190–5. doi:10.1126/science.1222794

36.   Farh KK-H, Marson A, Zhu J, Kleinewietfeld M, Housley WJ, Beik S, et al. Genetic and

epigenetic fine mapping of causal autoimmune disease variants. Nature. Nature

Publishing Group; 2014; doi:10.1038/nature13835

37.   Trynka G, Sandor C, Han B, Xu H, Stranger BE, Liu XS, et al. Chromatin marks identify

critical cell types for fine mapping complex trait variants. Nat Genet. Nature Publishing

Group; 2013;45: 124–30. doi:10.1038/ng.2504

38.   Finucane HK, Bulik-Sullivan B, Gusev A, Trynka G, Reshef Y, Loh P-R, et al. Partitioning

heritability by functional annotation using genome-wide association summary statistics.

Nat Genet. 2015;47: 1228–1235. doi:10.1038/ng.3404

39. Martin AR, Gignoux CR, Walters RK, Wojcik GL, Neale BM, Gravel S, et al. Human

Demographic History Impacts Genetic Risk Prediction across Diverse Populations. Am J

Hum Genet. Elsevier; 2017;100: 635–649. doi:10.1016/j.ajhg.2017.03.004

40. Cai TT, Zhang CH, Zhou HH. Optimal rates of convergence for covariance matrix

estimation. Ann Stat. 2010;38: 2118–2144. doi:10.1214/09-AOS752

41. Yang J, Wray NR, Visscher PM. Comparing apples and oranges: Equating the power of

case-control and quantitative trait association studies. Genet Epidemiol. 2010;34: 254–

257. doi:10.1002/gepi.20456

1 Table 1. Simulated benchmark comparison of non-parametric shrinkage (NPS) and LDPred.

Simulated genetic architecture	Fraction of causal SNPs	Method	R^2 Liability	R^2 Nagelkerke	Fold improvement over LDpred R^2_L	R^2_{Nag}
Mixture normal	1 %	NPS	0.122	0.081	1.12	1.10
	0.1 %	NPS	0.139	0.101	1.35	1.34
	0.01 %	NPS	0.234	0.162	2.49	2.54
	1 %	LDPred	0.109	0.074		
	0.1 %	LDPred	0.103	0.075		
	0.01 %	LDPred	0.094	0.064		
+ MAF dependency α = -0.25	1 %	NPS	0.131	0.095	1.28	1.25
	0.1 %	NPS	0.161	0.113	1.31	1.30
	0.01 %	NPS	0.247	0.170	1.93	1.89
	1 %	LDPred	0.102	0.077		
	0.1 %	LDPred	0.123	0.087		
	0.01 %	LDPred	0.128	0.090		
+ 5-fold higher causal fraction in DHS	1% (3 in DHS, 0.6 outside)	NPS	0.122 +/- 0.003	0.082 +/- 0.006	**1.19 ***	**1.20 ***
	0.1% (0.3 in DHS, 0.06 outside)	NPS	0.153 +/- 0.013	0.109 +/- 0.009	**1.38 ***	**1.38 ***
	0.01% (0.03 in DHS, 0.006 outside)	NPS	0.241 +/- 0.004	0.167 +/- 0.003	1.10	1.09
	1% (3 in DHS, 0.6 outside)	LDPred	0.103 +/- 0.006	0.068 +/- 0.008		
	0.1% (0.3 in DHS, 0.06 outside)	LDPred	0.111 +/- 0.012	0.079 +/- 0.011		
	0.01% (0.03 in DHS, 0.006 outside)	LDPred	0.219 +/- 0.332	0.153 +/- 0.247		

3 Non-parametric shrinkage (NPS) is more robust and accurate than the state-of-the-art parametric method (LDPred). The number of markers was 5,012,500.

4 The GWAS sample size was 100,000. Prediction models were optimized in the training cohort of 2,500 cases and 2,500 controls. R^2 of prediction was

5 measured in the validation cohort of 50,000 samples. The 95% confidence intervals were estimated from three replicate simulation runs. Simulations of mixture

6 normal and MAF dependency were not replicated. The star(*) indicates a significant improvement in R^2 (paired t-test).

1 **Figure Legends**
2
Figure 1. Overview of non-parametric shrinkage (NPS).

(A) For unlinked markers, NPS partitions the estimated effect sizes ($\hat{\beta}_j$) into K subgroups at cut-

offs of $b_0, b_1, \ldots, b_K$. Partitioned risk scores G_{ik} are calculated for each partition k and individual i

using an independent genotype-level training cohort. The per-partition shrinkage weights ω_k are

determined by the separation of G_{ik} between training cases and controls. (B) For markers in LD,

genotypes and estimated effects are decorrelated first by a linear projection $\mathcal{P}$ in non-overlapping

windows of ~ 2.4 Mb in length, and then NPS is applied to the data. The size of black dots

indicates genotype frequencies in the training cohort. Before projection, genotypes are correlated

between SNP 1 and 2 by LD, and thus sampling noise of estimated effects ($\hat{\beta}_j \mid \beta_j$) is also

correlated between SNPs. The projection $\mathcal{P}$ neutralizes both correlation structures. The axes of

projection are marked by red dashed lines.

Figure 2. Per-partition shrinkage weights estimated by non-parametric shrinkage (NPS)

approximate the conditional mean effects in the decorrelated space.

(A) NPS shrinkage weights ω_k (red line) compared to the theoretical optimum (black line),

$\lambda_j/(\lambda_j + \frac{M}{Nh^2})$, under infinitesimal architecture. The partition of largest eigenvalues $\mathcal{S}_{\lambda 10}$ is marked

by grey box. (B) Conditional mean effects estimated by NPS (red line) in sub-partitions of $\mathcal{S}_{\lambda 10}$ by

$\hat{\eta}_j$ under infinitesimal architecture. The theoretical line (black) is the average over all λ_j in $\mathcal{S}_{\lambda 10}$.

(C-D) Conditional mean effects estimated by NPS (red line) in sub-partitions of $\mathcal{S}_{\lambda 10}$ (C) and $\mathcal{S}_{\lambda 2}$

(D) on intervals of $\hat{\eta}_j$ under non-infinitesimal architecture with the causal SNP fraction of 1%. The

true conditional means (black) were estimated over 40 simulation runs. (All) The mean NPS

shrinkage weights (red line) and their 95% CIs (red shade) were estimated from 5 replicates.

Grey vertical lines indicate partitioning cut-offs. No shrinkage line (green) indicates $\omega_k = 1$. The

number of markers *M* is 101,296. The discovery GWAS size *N* equals to *M*. The heritability *h²* is

0.5.

Fig 1

A

Estimated per-SNP effects from discovery GWAS ($\hat{\beta}_j$)

$\downarrow$

Partition SNPs into K subgroups:
$$\mathcal{S}_k = \{\, j : b_{k-1} < |\hat{\beta}_j| < b_k \,\}$$

$\downarrow$

Partitioned risk scores: $G_{ik} = \sum_{j \in \mathcal{S}_k} \hat{\beta}_j x_{ij}$

$\downarrow$

Estimate per-partition mean weights: ω_k

$\downarrow$

$$G_i = \sum_k \omega_k \sum_{j \in \mathcal{S}_k} \hat{\beta}_j x_{ij}$$

B

Decorrelating linear projection $\mathcal{P}$

Individual SNPs		Decorrelated space
genotypes X	$X^{P^T} := \mathcal{P} X^T$	X^P
$cov(X) = \mathbf{\Sigma}$		$cov(X^P) = \mathbf{I}$
estimated effects $\hat{\beta}$	$\hat{\eta} := \mathcal{P}\hat{\beta}$	$\hat{\eta}$
$cov(\hat{\beta} \mid \beta) = \frac{1}{N}\mathbf{\Sigma}$		$cov(\hat{\eta} \mid \beta) = \frac{1}{N}\mathbf{I}$

Fig 2

Supplementary Note for

Non-parametric polygenic risk prediction using partitioned GWAS summary statistics

Decorrelating projection

We split the genome into L non-overlapping windows of m SNPs each. An individual window is large enough to capture the majority of linkage disequilibrium (LD) patterns except near the edge. For the sake of simplicity, we assume that LD is confined to each window and there exists no LD across windows.

In genomic window l, let X_l be an $N \times m$ genotype matrix of a discovery cohort and X'_l be an $N' \times m$ genotype matrix of a training cohort. The sample sizes of discovery and training cohorts are N and N', respectively. The genotypes are standardized to the mean of 0 and variance of 1. Let $\hat{\beta}_l$ be an m-dimensional vector of apparent effect sizes at all SNPs from the discovery GWAS. $\hat{\beta}_l$ is also defined with respect to the standardized genotypes. Let β_l be an m-dimensional vector of true underlying genetic effects at all SNPs in window l. For convenience, we omit the subscript l when it is clear from the context.

The LD matrix D is given by $D = \frac{1}{N} X^T X$. Let us assume for a moment that D has full rank. In this case, D is symmetric and positive semi-definite, thus can be factorized by spectral decomposition into the following form:

$$D = Q \, \Lambda \, Q^T$$

where Q is an orthonormal matrix of eigenvectors and Λ is a diagonal matrix of positive eigenvalues. The extension to rank-deficient LD matrix is straight-forward and will be discussed in later section.

When the discovery cohort has the infinite sample size, the true genetic effects β can be directly recovered from the summary statistics $\hat{\beta}$ by correcting for LD:

$$\beta = D^{-1} \hat{\beta} = Q\Lambda^{-1}Q^T\hat{\beta} = Q\Lambda^{-\frac{1}{2}}\left(\Lambda^{-\frac{1}{2}}Q^T\hat{\beta}\right) \qquad (Eq\ S1)$$

We call the linear transformation $\mathcal{P} := \Lambda^{-\frac{1}{2}} Q^T$ a *decorrelating projection* because of the reasons presented in the next section. The inverse transformation $Q\Lambda^{-\frac{1}{2}}$ will retrieve SNP effects back from the decorrelated space defined by $\mathcal{P}$. Note that the sample size does not have to be infinite to order for $\mathcal{P}$ to be well-defined.

By applying $\mathcal{P}$ on summary statistics, we obtain *decorrelated effect size estimates $\hat{\eta}$*, and by applying $\mathcal{P}$ on genotypes, we obtain *projected genotypes X'^P* as follows:

$$\hat{\eta} := \mathcal{P}\hat{\beta}$$

$$(X'^P)^T := \mathcal{P}X'^T$$

Distribution of projected genotypes

Let X'_i be an m-dimensional genotype vector of individual i in window l. Then, X'_i follows the following multivariate normal distribution:

$$X'_i \sim N(\mathbf{0}, \boldsymbol{D})$$

Since the projected genotype X'^P_i is derived by multiplying $\mathcal{P}$ on X'_i by definition (Eq S2), it also follows a multivariate normal distribution. Specifically, the distribution of X'^P_i is:

$$X'^P_i \sim N\left(\Lambda^{-\frac{1}{2}} \boldsymbol{Q}^T \mathbf{0}, \left(\Lambda^{-\frac{1}{2}} \boldsymbol{Q}^T \right) \boldsymbol{D} \left(\Lambda^{-\frac{1}{2}} \boldsymbol{Q}^T \right)^T \right)$$

$$= N\left(\mathbf{0}, \Lambda^{-\frac{1}{2}} \boldsymbol{Q}^T \boldsymbol{Q}\Lambda\boldsymbol{Q}^T \boldsymbol{Q}\Lambda^{-\frac{1}{2}} \right) = N(\mathbf{0}, \boldsymbol{I})$$

since $\boldsymbol{D} = \boldsymbol{Q}\Lambda\boldsymbol{Q}^T$ and $\boldsymbol{Q}^T\boldsymbol{Q} = \boldsymbol{I}$. The projected genotypes are now decorrelated with the covariance of $\boldsymbol{I}$.

Distribution of decorrelated effect size estimates

In the discovery GWAS, the estimated effect sizes $\hat{\beta}$ are calculated by linear regression as below:

$$\hat{\beta} = \frac{1}{N}X^T y$$

where y is an N-dimensional phenotype vector. For convenience, we assume that y is standardized to the mean of 0 and variance of 1. At this time, we treat genotypes as fixed variables and model the genetic effects β and residuals ϵ as random. Since $y = X\beta + \epsilon$,

$$\hat{\beta} = \frac{1}{N}X^T(X\beta + \epsilon) = \boldsymbol{D}\beta + \frac{1}{N}X^T\epsilon$$

where the residual ϵ follows an N-dimensional multivariate normal distribution $N(\mathbf{0}, \sigma_e^2\boldsymbol{I})$. In an individual window, the genetic effects explain only a small fraction of phenotypic variation,

therefore $\sigma_e^2 \approx \text{var}(y) = 1$. The distribution of sampling noise in $\hat{\beta}$, namely the distribution of $\hat{\beta}$ given β, follows:

$$\hat{\beta} \mid \beta \sim N(\boldsymbol{D}\beta + \frac{1}{N}\boldsymbol{X}^T\boldsymbol{0}, \frac{\sigma_e^2}{N^2}\boldsymbol{X}^T\boldsymbol{I}\boldsymbol{X})$$
$$\approx N(\boldsymbol{D}\beta, \frac{1}{N}\boldsymbol{D})$$

since $\boldsymbol{D} = \frac{1}{N}\boldsymbol{X}^T\boldsymbol{X}$. Since the decorrelated effect size estimate $\hat{\eta}$ is a product of $\mathcal{P}$ and $\hat{\beta}$ by definition (Eq S2), the distribution of $\hat{\eta}$ given β also follows a multivariate normal distribution:

$$\hat{\eta} \mid \beta \sim N\left(\boldsymbol{\Lambda}^{-\frac{1}{2}}\boldsymbol{Q}^T\boldsymbol{D}\beta, \frac{1}{N}\boldsymbol{\Lambda}^{-\frac{1}{2}}\boldsymbol{Q}^T\boldsymbol{D}\left(\boldsymbol{\Lambda}^{-\frac{1}{2}}\boldsymbol{Q}^T\right)^T\right)$$

$$= N\left(\boldsymbol{\Lambda}^{-\frac{1}{2}}\boldsymbol{Q}^T\boldsymbol{Q}\boldsymbol{\Lambda}\boldsymbol{Q}^T\beta, \frac{1}{N}\boldsymbol{\Lambda}^{-\frac{1}{2}}\boldsymbol{Q}^T\boldsymbol{Q}\boldsymbol{\Lambda}\boldsymbol{Q}^T\boldsymbol{Q}\boldsymbol{\Lambda}^{-\frac{1}{2}}\right)$$

$$= N\left(\boldsymbol{\Lambda}^{\frac{1}{2}}\boldsymbol{Q}^T\beta, \frac{1}{N}\boldsymbol{I}\right) \qquad\qquad (Eq\ S3)$$

since $\boldsymbol{D} = \boldsymbol{Q}\boldsymbol{\Lambda}\boldsymbol{Q}^T$ and $\boldsymbol{Q}^T\boldsymbol{Q} = \boldsymbol{I}$. The sampling noise of $\hat{\eta}$ is now decorrelated with the covariance of $\frac{1}{N}\boldsymbol{I}$. Hence, the projection $\mathcal{P}$ removes correlations in both genotypes and summary statistics.

Interpretation of eigenvalues

Based on Eq S3, $\hat{\eta} \mid \beta$ approaches to $\boldsymbol{\Lambda}^{\frac{1}{2}}\boldsymbol{Q}^T\beta$ as the sample size goes to the infinity. Thus, we define true decorrelated effect η as:

$$\eta \coloneqq \boldsymbol{\Lambda}^{\frac{1}{2}}\boldsymbol{Q}^T\beta \qquad\qquad (Eq\ S4)$$

Let us assume that the distribution of β is symmetric at 0 and independent at each SNP. Then,

$$E[\eta_j] = E\left[\sqrt{\lambda_j}\, q_j^T\beta\right] = \sqrt{\lambda_j}\, q_j^T E[\beta] = 0$$

and

$$var[\eta_j] = E\left[\left(\sqrt{\lambda_j}\, q_j^T\beta\right)^2\right] - E[\eta_j]^2$$

$$= \lambda_j \sum_{s=1}^{m} q_{sj}^2\, E[\beta_s^2]$$

where the j-th eigenvector q_j is $(q_{1j} \quad \cdots \quad q_{mj})^T$. Therefore, the scale of η_j, namely, $var[\eta_j]$, is proportional to eigenvalue λ_j. Furthermore, in particular cases that all SNPs have the same variance of per-SNP effect sizes σ_g^2,

$$var[\eta_j] = \lambda_j \sigma_g^2$$

since $\sum_{s=1}^{m} q_{sj}^2 = 1$. In both non-infinitesimal and infinitesimal architecture, if the genetic effects are sampled i.i.d. for each SNP uniformly across the genome, $\sigma_g^2 = var(\beta_j) = h^2/M$. M is the total number of markers, and h^2 is the heritability of the phenotype.

Conditional mean effects under infinitesimal genetic architecture
Under infinitesimal genetic architecture, the conditional mean effect has been analytically derived in [1]:

$$E[\beta \mid \hat{\beta}] = \left(\frac{M}{Nh^2}I + D\right)^{-1} \hat{\beta} \qquad (Eq\ S5)$$

under the assumption that D is the LD matrix of full rank. Since

$$\left(\frac{M}{Nh^2}I + D\right) = Q\left(\frac{M}{Nh^2}I + \Lambda\right)Q^T$$

and

$$\left(\frac{M}{Nh^2}I + D\right)^{-1} = Q\left(\frac{M}{Nh^2}I + \Lambda\right)^{-1}Q^T$$

we can reformulate Eq S5 as follows:

$$E[\beta \mid \hat{\beta}] = Q\left(\frac{M}{Nh^2}I + \Lambda\right)^{-1}Q^T\hat{\beta} \qquad (Eq\ S6)$$

$$= Q\left(\frac{M}{Nh^2}I + \Lambda\right)^{-1}\Lambda^{\frac{1}{2}}\left(\Lambda^{-\frac{1}{2}}Q^T\hat{\beta}\right)$$

$$= Q\left(\frac{M}{Nh^2}I + \Lambda\right)^{-1}\Lambda^{\frac{1}{2}}\hat{\eta}$$

by the definition of $\hat{\eta}$ (Eq S2). Hence,

$$E[\eta \mid \hat{\eta}] = \Lambda^{\frac{1}{2}}Q^T E[\beta \mid \hat{\eta}] = \Lambda^{\frac{1}{2}}Q^T E[\beta \mid \hat{\beta}]$$

$$= \Lambda^{\frac{1}{2}}Q^T Q\left(\frac{M}{Nh^2}I + \Lambda\right)^{-1}\Lambda^{\frac{1}{2}}\hat{\eta}$$

$$= \left(\frac{M}{Nh^2} I + \Lambda \right)^{-1} \Lambda \, \hat{\eta} \qquad\qquad (Eq\ S7)$$

using the definition of η (Eq S4). For the j-th projection defined by λ_j and q_j, the conditional mean effect is given as the following:

$$E[\eta_j \mid \hat{\eta}_j] = \frac{\lambda_j}{\lambda_j + \dfrac{M}{Nh^2}} \hat{\eta}_j$$

Thus, in infinitesimal architecture, the conditional mean effect $E[\eta_j \mid \hat{\eta}_j]$ simplifies to $\omega \, \hat{\eta}_j$, where ω is the optimal shrinkage weight and depends only on eigenvalues as follow:

$$\omega = \frac{\lambda_j}{\lambda_j + \dfrac{M}{Nh^2}}$$

Back-conversion from the decorrelated space to the SNP space

Let $\hat{y}$ be an N'-dimensional vector of predicted phenotypes. We construct $\hat{y}$ by summing over all projected genotypes multiplied by conditional mean effects in the decorrelated space as follows:

$$\hat{y} = \sum_{l=1}^{L} X_l'^P E[\eta_l \mid \hat{\eta}_l]$$

where the conditional mean effect $E[\eta_l \mid \hat{\eta}_l]$ is obtained by non-parametric shrinkage. By the definition of $X_l'^P$ (Eq S2),

$$\hat{y} = \sum_{l=1}^{L} X_l' \left(\Lambda_l^{-\frac{1}{2}} Q_l^T \right)^T E[\eta_l \mid \hat{\eta}_l]$$

$$= \sum_{l=1}^{L} X_l' \left(Q_l \Lambda_l^{-\frac{1}{2}} E[\eta_l \mid \hat{\eta}_l] \right)$$

Note that X_l' is the genotype matrix in the original SNP space. Thus, $E[\eta_l \mid \hat{\eta}_l]$ can be converted back to the SNP space by the following transformation:

$$Q_l \Lambda_l^{-\frac{1}{2}} E[\eta_l \mid \hat{\eta}_l]$$

In the infinite sample size, $E[\eta_l \mid \hat{\eta}_l] = \hat{\eta}_l$. Hense, using Eq S1, we can recover

$$\hat{y} = \sum_{l=1}^{L} X_l' \left(\boldsymbol{Q}_l \boldsymbol{\Lambda}_l^{-\frac{1}{2}} \hat{\eta}_l \right) = \sum_{l=1}^{L} X_l' \left(\boldsymbol{Q}_l \boldsymbol{\Lambda}_l^{-\frac{1}{2}} \boldsymbol{\Lambda}_l^{-\frac{1}{2}} \boldsymbol{Q}_l^T \hat{\beta} \right) = \sum_{l=1}^{L} X_l' (\boldsymbol{D}^{-1} \hat{\beta}) = \sum_{l=1}^{L} X_l' \beta$$

as expected.

Rank deficiency of LD matrix

Even when the LD matrix $\boldsymbol{D}$ is not full rank, it is symmetric and non-negative semi-definite. In this case, spectral decomposition on $\boldsymbol{D}$ yields only r positive eigenvalues, where r is the rank of the matrix and $r < m$, and the rest of eigenvalues are 0. Without the loss of generality, we can reorder the eigenvalues and corresponding eigenvectors in such a way that only the first r eigenvalues are positive. We truncate the components corresponding to eigenvalues $r + 1, \dots, m$ and reduce the dimension to r. Specifically, the truncated matrices are defined as the following:

$$\boldsymbol{Q}' = (q_1 \quad \cdots \quad q_r)$$

$$\boldsymbol{\Lambda}' = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \cdots & 0 \\ 0 & 0 & \lambda_r \end{pmatrix}$$

where λ_j and q_j are the j-th eigenvalues and eigenvectors, respectively. Since $\boldsymbol{Q}'$ and $\boldsymbol{\Lambda}'$ satisfy the following:

$$\boldsymbol{D} = \boldsymbol{Q}' \boldsymbol{\Lambda}' \boldsymbol{Q}'^T$$

and

$$\boldsymbol{Q}'^T \boldsymbol{Q}' = \boldsymbol{I}_r$$

all results we derived in the previous sections hold with $\boldsymbol{Q}'$ and $\boldsymbol{\Lambda}'$ in place of $\boldsymbol{Q}$ and $\boldsymbol{\Lambda}$, respectively.

However, the analysis of infinitesimal model (Eqs S5-S7) requires further discussion since it is non-trivial to generalize to rank deficient $\boldsymbol{D}$. Note that Eq S5 was derived under the assumption that $\boldsymbol{D}$ has full rank [1], thus cannot be used directly for rank deficient $\boldsymbol{D}$ even though the matrix $\frac{M}{Nh^2} \boldsymbol{I} + \boldsymbol{D}$ is always invertible.

We re-derived the posterior mean effects from joint probability density function of $\hat{\eta}$ and η in reduced r-dimensional space. Now $E[\eta \mid \hat{\eta}]$ of Eq S7 becomes the following equation:

$$E[\eta \mid \hat{\eta}] = \left(\frac{M}{Nh^2} \boldsymbol{I}_r + \boldsymbol{\Lambda}' \right)^{-1} \boldsymbol{\Lambda}' \hat{\eta}$$

Therefore,

$$E[\beta \mid \hat{\beta}] = \boldsymbol{Q}' \boldsymbol{\Lambda}'^{-\frac{1}{2}} E[\eta \mid \hat{\eta}]$$

$$= Q' \Lambda'^{-\frac{1}{2}} \left(\frac{M}{Nh^2} I_r + \Lambda' \right)^{-1} \Lambda' \left(\Lambda'^{-\frac{1}{2}} Q'^T \hat{\beta} \right)$$

$$= Q' \left(\frac{M}{Nh^2} I_r + \Lambda' \right)^{-1} Q'^T \hat{\beta} \qquad (Eq\ S8)$$

Note that this result is not identical to the previous Eq S6, which was derived for full-rank D:

$$E[\beta \mid \hat{\beta}] = Q \left(\frac{M}{Nh^2} I + \Lambda \right)^{-1} Q^T \hat{\beta} \qquad (Eq\ S6)$$

In Eq S6, M/Nh^2 term remains for $q_{r+1}, \dots, q_m$ whereas it was truncated in Eq S8.

Empirical analysis of partitioning schemes based on simulations of unlinked markers

      To explore an effective partitioning strategy for non-parametric shrinkage, we compared the prediction accuracy of different partitioning approaches under simulated genetic architecture of unlinked markers. While a higher number of partitions is desirable to better interpolate the mean conditional effect curve, fully optimizing the partitioning cut-offs rapidly becomes impractical due to the combinatorial search space and limited training sample size. To explore this issue, we benchmarked fully optimized three-way partitioning, of which two cut-off values were optimized in the training cohort, against seven-way partitioning, of which cut-off points were evenly placed without further optimization. In our simulation, the seven-way partitioning performed equivalently as or significantly more accurately than the three-way partitioning (Fig S1, $P < 10^{-4}$ for the causal SNP fractions simulated at 1% and 3%).

      Note that the widely-used *p*-value thresholding approach is a special case of non-parametric shrinkage with the number of partition K being 2. When the effect sizes are measured relative to the standardized genotypes, thresholding on $|\hat{\beta}_j|$ is equivalent to thresholding on *p*-values of association. This is because $|\hat{\beta}_j|$ has the following monotonic relationship with *p*-values:

$$|\hat{\beta}_j| = \frac{1}{N} \Phi^{-1} \left(1 - \frac{1}{2} p_j \right)$$

where Φ is the standard normal cumulative distribution function and p_j is the *p*-value of association at SNP j. As expected, thresholding was always significantly inferior to finer partitioning (Fig S1, $P < 0.01$).

      For this analysis, we simulated a spike-and-slab genetic architecture under a liability threshold model. The total number of markers was 100,000, and 0.3, 1, or 3% of all markers were simulated to be causal SNPs with effect sizes randomly sampled from a normal distribution. SNPs were independent with each other without LD in this simulation. We assumed the heritability of 0.5 and disease prevalence of 5% and over-sampled cases relative to

controls to generate cohorts with case:control ratio of 1:1. For each simulated causal fraction, we generated 500 sets of three cohorts: discovery, training and validation with sample sizes of 20,000, 4,000 and 4,000, respectively. Discovery cohort data were accessed only as summary statistics. Prediction models were optimized in training cohort and evaluated in validation cohort. For the *p*-value thresholding, the optimal *p*-value cut-off for polygenic risk score was scanned from 10^{-7} to 1 at increments of 0.25 on the $\log_{10}$ scale. The cut-off yielding the highest Area Under the Curve (AUC) in the training cohort was selected for the final model tested in the validation cohort. For seven-way partitioning, we used fixed partition boundaries on the interval of $|\hat{\beta}|$ corresponding to every order of magnitudes in *p*-values of association. Specifically, the intervalof $|\hat{\beta}|$ was split at six *p*-values of association: 5×10^{-7}, 5×10^{-6}, …, and 5×10^{-2}. For three-way partitioning, all 15 combinations of the above six *p*-value cut-off values were examined to find the most optimal partition. In all three partitioning schemes, the shrinkage weight for the partition of smallest $|\hat{\beta}|$ was set to 0.

References

[1]   Vilhjalmsson BJ, Yang J, Finucane HK, Gusev A, Lindstrom S, Ripke S, et al. Modeling Linkage Disequilibrium Increases Accuracy of Polygenic Risk Scores. Am J Hum Genet. 2015 Oct 01;97(4):576-92. PubMed PMID: 26430803. Pubmed Central PMCID: 4596916.