Title

Novel cytonuclear combinations modify Arabidopsis seed physiology and vigor

Clément Boussardon[1,*], Marie-Laure Martin-Magniette[2,3,4], Béatrice Godin[1], Abdelilah Benamar[5], Benjamin Vittrant[2], Sylvie Citerne[1], Tristan Mary-Huard[2,6], David Macherel[5], Loïc Rajjou[1], Françoise Budar[1].

Author affiliations

[1] Institut Jean-Pierre Bourgin, INRA, AgroParisTech, CNRS, Université Paris-Saclay, 78000 Versailles, France

[2] UMR MIA-Paris, AgroParisTech, INRA, Université Paris-Saclay, 75005, Paris, France

[3] Institute of Plant Sciences Paris Saclay IPS2, CNRS, INRA, Université Paris-Sud, Université Evry, Université Paris-Saclay, Bâtiment 630, 91405 Orsay, France

[4] Institute of Plant Sciences Paris-Saclay IPS2, Paris Diderot, Sorbonne Paris-Cité, Bâtiment 630, 91405, Orsay, France

[5] IRHS, Université d'Angers, INRA, Agrocampus Ouest, UMR 1345, SFR 4207 QUASAV, Angers, France

[6] Génétique Quantitative et Evolution–Le Moulon, INRA, Université Paris-Sud, CNRS, AgroParisTech, Université Paris-Saclay, 91190 Gif-sur-Yvette, France

[*] ***Present address:*** Department of Plant Physiology, Umeå Plant Science Centre, Umeå University, S-90187 Umeå, Sweden

Authors emails: clement.boussardon@umu.se, marie_laure.martin-magniette@agroparistech.fr, beatrice.godin@inra.fr, Abdelilah.Benamar@Univ-Angers.Fr, benjamin.vittrant.1@ulaval.ca, sylvie.citerne@inra.fr, tristan.mary-huard@agroparistech.fr, David.Macherel@Univ-Angers.Fr, loic.rajjou@agroparistech.fr, francoise.budar@inra.fr

Corresponding author: Françoise Budar. Phone number: +33 130833015

Running title

Seed traits under cytonuclear control

Issue section

Growth and development

Title Novel cytonuclear combinations modify Arabidopsis seed physiology and vigor

Highlight

Natural variation in Arabidopsis organelles and cytonuclear interactions influence seed dormancy, longevity and germination performance. Enhanced seed vigor was obtained through the creation of novel cytonuclear combinations.

Abstract

The influence of intraspecific variation in cytoplasmic genomes and cytonuclear interactions on key seed traits that can impact adaptation and agriculture has not been thoroughly explored, so far. Here, dormancy, germination performance and longevity of seeds have been assessed in Arabidopsis plants with novel cytonuclear combinations that disrupt coadaptation between natural variants of nuclear and cytoplasmic genomes. Although all three traits were affected by cytonuclear reshuffling, the sensitivity of seed traits to cytoplasmic change was dependent on the nuclear background. Both deleterious and, more surprisingly, favorable effects of novel cytonuclear combinations (in comparison with the nuclear parent) were observed, suggesting suboptimal genetic combinations exist in natural populations for these traits. Significant changes on dormancy and germination performance due to specific cytonuclear interacting combinations mainly occurred in opposite directions, in accordance with the previously proposed 'dormancy continuum'. Consistently, reduced sensitivity to exogenous ABA and faster endogenous ABA decay during germination were observed in a novel cytonuclear combination that also exhibited enhanced longevity and better germination performance, compared to its natural nuclear parent. Cytoplasmic genomes, therefore, represent an additional resource of natural variation for breeding seed vigor traits.

Keywords

cytolines, cytonuclear co-adaptation, cytonuclear interaction, dormancy, seed longevity, germination, seed vigor.

Abbreviations

CDT, controlled deterioration treatment; Gmax, maximum germination percentage; PH, post-harvest; RH, relative humidity

Introduction

The evolutionary success of flowering plants is largely due to the invention of seeds, which

protect and scatter the next generation and provide it with resources upon germination.

Germination is a complex process determined by the intrinsic properties of seeds and

influenced by biotic and abiotic environment, aging-related damage, which can occur during

seed storage, and dormancy, which prevents germination under favorable conditions. In

nature, dormancy allows the seed to await for the favorable season for seedling success

(Finch-Savage and Footitt, 2017). Once dormancy is released, the capacity of seeds to

germinate in a wide range of environmental conditions contributes to adaptation, and

germination speed has a fitness impact in competitive situations (Dubois and Cheptou,

2012). Germination is also a trait of primary interest for agriculture, where seed vigor,

defined as the properties that ensure a fast and synchronized germination in field conditions

(Finch-Savage and Bassel, 2016), is crucial for the quality of seed lots. In contrast to its

adaptive role in natural conditions, dormancy is not a desirable trait in crops, as it leads to

low and unsynchronized seedling emergence rates in agricultural production (Shu *et al.*,

2015). In the laboratory, seed vigor is assessed through the speed and uniformity of

germination in favorable conditions, germination performance under stress conditions, and

the ability to survive storage (Finch-Savage *et al.*, 2010; Rajjou *et al.*, 2012). Seed longevity,

i.e. ability to survive and maintain germination performance despite physiological damage

occurring during storage, is an important component of seed vigor (Rajjou and Debeaujon,

2008; Sano *et al.*, 2016).

Arabidopsis thaliana (hereafter Arabidopsis) has been widely used to decipher seed biology

traits through mutant studies and genome-wide analyses (North *et al.*, 2010). Arabidopsis

natural accessions display genetic variation for seed dormancy, longevity and germination

vigor (Alonso-Blanco *et al.*, 2003; Clerkx *et al.*, 2004; Bentsink *et al.*, 2010; Vallejo *et al.*,

2010; Yuan *et al.*, 2016*a*). For example, the release of seed dormancy during post-harvest

(PH) dry storage occurred at variable speeds in 112 Arabidopsis accessions, with the time of

storage required to reach 50% of germination ranging from 3.5 to 264 days (Debieu *et al.*,

2013). In addition, Arabidopsis natural variation in seed dormancy and longevity has been

shown relevant for adaptation in ecological studies (Donohue *et al.*, 2005*a*; Huang *et al.*,

2010; Kronholm *et al.*, 2012; Debieu *et al.*, 2013; Postma *et al.*, 2015). A number of

quantitative trait loci (QTL) for dormancy, longevity, germination speed, and germination

tolerance to stresses have been reported (van Der Schaar *et al.*, 1997; Alonso-Blanco *et al.*,
2003; Clerkx *et al.*, 2004; Galpaz and Reymond, 2010; Bentsink *et al.*, 2010; Huang *et al.*,
2010; Joosen *et al.*, 2012; Nguyen *et al.*, 2012; Yuan *et al.*, 2016*b*). Further major advances in
our understanding of the genetic bases of natural variation in seed dormancy and
germination are being provided from the functional study of the genes underlying these loci
(Bentsink *et al.*, 2006; Shu *et al.*, 2015).

Comparatively, little is known about the role of cytoplasmic variation in the establishment of
seed traits important for adaptation or agriculture, despite the crucial role of mitochondria
and chloroplasts in seed quality and germination performance. Upon imbibition,
mitochondrial respiration resumes almost immediately to fuel the very active metabolism
which is required during imbibition and germination (Paszkiewicz *et al.*, 2017). Plastids are
the site of the early steps of ABA and GA syntheses. These two major hormones govern the
balance between dormancy and germination through their antagonist effects: ABA promotes
the induction and maintenance of seed dormancy, whereas GAs promote germination by
stimulating tegument and endosperm rupture and embryo cellular elongation (Finch-Savage
and Leubner-Metzger, 2006; Shu *et al.*, 2016). Both mitochondria and chloroplasts have
endosymbiotic origins and their co-evolution with the nucleus of the host cells has shaped
nuclear and organellar genomes, while organizing the compartmentalized metabolism of the
plant cell (Dyall *et al.*, 2004; Kutschera and Niklas, 2005). The tuning of mitochondrial and
chloroplast functions is under a complex and still elusive network of genetic and metabolic
regulations (Rurek, 2016; De Souza *et al.*, 2017). This is mainly ensured and controlled by
nuclear-encoded factors, some of which interact with organelle genes or their products.
Numerous genes encoding organellar proteins are expressed during germination and their
inactivation often affects the process (Yang *et al.*, 2011; Demarsy *et al.*, 2011; Savage *et al.*,
2013; Sew *et al.*, 2016). In addition, transcriptional regulation of specific mitochondrial
protein-encoding genes during seed germination has been largely documented (Howell *et*
al., 2008; Narsai *et al.*, 2011; Law *et al.*, 2012). Also, plastid gene expression is required for
proper seed development in Arabidopsis (Bryant *et al.*, 2011) and maize (Sosso *et al.*, 2012).

Based on the severe consequences for seed development and germination arising from
anomalies in the genetic and physiological activities of plastids and mitochondria, it could be
assumed that natural cytoplasmic variation affecting these traits is strongly limited, if not
suppressed, by natural selection. However, a few studies have reported results suggesting

an impact of natural variation in cytoplasmic genomes on seed physiological traits. A seminal

study on reciprocal Arabidopsis F1s and backcross generations showed that cytoplasm

variation impacted germination efficiency (Corey *et al.*, 1976). A recent report on seed

germination performance of reciprocal hybrids between *Arabidopsis lyrata* populations

suggested that a disruption of cytonuclear coadaptation contributed to the lower

germination of F1 and F2 hybrid generations (Hämälä *et al.*, 2017). Furthermore, in a study

using Arabidopsis cytolines, each combining the nuclear genome of a natural variant with

the cytoplasmic genomes of a different variant, we previously demonstrated the impact of

cytoplasmic natural variation and cytonuclear interactions on adaptive traits in the field,

including germination (Roux *et al.*, 2016). Here, using the same cytolines, we have

investigated further the effect of natural genetic variation within organelles on specific

physiological seed traits relevant to adaptation and agriculture, namely dormancy,

germination performance and longevity. We found that genetic variation in organelle

genomes could impact these traits, depending on the nuclear background. Furthermore, we

uncovered novel cytonuclear combinations that displayed higher performance than their

natural nuclear parent, suggesting that cytoplasmic variation could potentially be used in

breeding of seed traits. The improved germination performance and seed longevity of a

novel cytonuclear combination was associated with more efficient endogenous ABA

degradation and lower exogenous ABA sensitivity.

Materials and Methods

Plant material

Cytolines are genotypes combining the nuclear genome of one parent with the organelle genomes of another conspecific parent. Hereafter, a cytoline possessing the cytoplasm of accession "A" and the nucleus of accession "B" is designated [A]B. We used series of cytolines derived from eight natural accessions of Arabidopsis, namely: Blh-1, Bur-0, Ct-1, Cvi-0, Ita-0, Jea, Oy-0, and Sha, which were selected for their wide genetic diversity (Mckhann *et al.*, 2004; Moison *et al.*, 2010). Cytolines were obtained by recurrent paternal backcrosses of hybrids from the di-allele cross (Roux *et al.*, 2016). Seed stocks of the 64 genotypes (56 cytolines and 8 parental accessions) are available at the Versailles Arabidopsis Stock Center (http://publiclines.versailles.inra.fr/cytoline/index).

Unless specified, the seed stocks used were obtained in a growth chamber and previously used in the study of (Roux *et al.*, 2016). They are conserved in controlled conditions (4°C, 15-30% relative humidity, RH) in the Versailles Arabidopsis Stock Center. These seed lots had been stored for at least 18 months when used for the experiments described here.

Two additional seed productions were realized for dormancy studies in a growth chamber under the same conditions as in Roux *et al.* (2016) (light 16 h at 21°C, dark 8 h at 18°C), using two plants per genotype, each on either of the two available shelves. An additional seed production of the Sha and [Blh-1]Sha lines was realized in the greenhouse, seeds from four individuals were pooled for each genotype.

Seed dormancy assays

For the 64 genotypes, the depth of dormancy was measured on freshly harvested seeds then at 3, 6 and 9 months of after-ripening (designated 0PH, 3PH, 6PH and 9PH). Seed imbibition was conducted in Petri dishes (Ø 55 mm) on filter papers with three sheets of absorbent paper (Roundfilter paper circles, Ø 45 mm; Schleicher & Schuell, Dassel, Germany) covered by a black membrane filter (ME 25/31, Ø 45 mm; Schleicher & Schuell, Dassel, Germany) wetted with 1.3 mL of ultrapure water. Seeds were then incubated in a controlled culture room under continuous light (Philips TRM HOW/33 RS tubes, 70 $\mu mol.m^{-2}.s^{-1}$) at 15°C or 25°C. Germination (protrusion of the radicle after rupture of the testa and endosperm) was assessed daily during twelve days by visual examination.

117 Results were analyzed independently for each PH time and considering both germination
118 temperatures in order to better control the variability of germination success, except for
119 0PH whose only results at 15°C were considered because none of the genotypes germinated
120 at 25°C upon harvest. The [Sha]Cvi-0 and [Ct-1]Jea genotypes were excluded from the
121 analyzed dataset due to missing data for at least one biological replicate.
122
Germination performance tests in challenging conditions
The complete set of 64 genotypes was tested for germination performance on water and
NaCl, after stratification. Seeds were placed on a blue blotter papers (Anchor Paper
Company, SGB1924B) positioned on the germination medium (Phytoblend, 0.5% in water)
with or without NaCl, in square Petri dishes (120 x 120 x 17 mm). Because seed germination
tolerance to NaCl is variable from one accession to another (Galpaz and Reymond, 2010),
each nuclear series was tested on the NaCl concentration (Table S1) that resulted in
approximately 50% germination of the parental accession in a preliminary experiment. Each
dish contained four to eight genotypes and two or three replicates were realized.
Stratification treatment was applied at 4°C in the dark for four days. Then, dishes were
transferred into a phytotron system providing continuous light from three sides at 25°C
(Sanyo MLR-351H, Versatile environmental test chamber). Germination time was counted
from the transfer into the phytotron. For the following of germination kinetics, images were
taken at 5 to 6 time points between 24 and 72 hours of germination with a digital camera
(Nikon Coolpix-P510, Nikon, France). Germination was visually assessed using ImageJ
(http://rsbweb.nih.gov/ij/). The maximum germination percentage (Gmax) was considered
reached when no additional germination was observed after another 24 hours.
Following the same procedure, the Sha nuclear series was tested on mannitol (200 mM) or
KCl (100 mM). ABA (50 µM) sensitivity tests were identically realized on Sha, [Blh-1]Sha and
[Ita-0]Sha, except that germination was monitored visually daily during four days.
143
Controlled Deterioration Treatment (CDT)
The CDT was performed in order to mimic natural seed aging (Tesnier and Strookman-
Donkers, 2002; Rajjou *et al.*, 2008). These experiments were performed on the Sha and Ct-1
nuclear series, on seeds from the Versailles Arabidopsis Stock Center.

Briefly, dry mature seeds were first equilibrated at 75% relative humidity (20°C) during 3

days. After this step, day 0 controls were immediately dried back in a desiccator with

Silicagel for three days. Treatment was done by storing the seeds (at 75% RH, using sealed

boxes with saturated NaCl solution) for 10, 20, 30 and 40 days at 35°C. After the treatment,

seeds were dried back on Silicagel as above. Germination percentage was measured after

seed imbibition on water under continuous light at 23°C.

Measurement of Na and K ion contents in Arabidopsis seeds

A germination experiment was set as described above for germination performance, with or

without NaCl (100 mM), except that seeds were collected either immediately after

stratification or after 20 hours of germination. Three replicates were realized. Seeds were

rinsed three times with 50 mL of sterile water and dried at 100°C overnight. The dry weight

of each sample was measured and samples were digested in 2 mL of 70% HNO_3 for a total of

3h with temperature ramping from 80 to 120°C. Potassium and sodium contents were

determined with a Varian AA240FS atomic absorption spectrometer (Agilent Technologies,

Santa Clara, Ca, USA) and concentrations calculated by comparison with Na^+ and K^+

standards.

Measurement of ABA content in Arabidopsis seeds

ABA content was measured on dry seeds and on stratified seeds after 6 h of germination,

using the same experiment setting as described above for germination performance, with or

without NaCl (100 mM). Twenty mg of seeds were ground in a mortar with liquid nitrogen

and lyophilized with a freeze-drier. For each sample, 10 mg of freeze-dried powder were

extracted with 0.8 mL of acetone/water/acetic acid (80/19/1 v:v:v). ABA stable labelled

isotope used as internal standard was prepared as described in Le Roux *et al.* (2014). Two

nanograms of each standard were added to the sample. The extract was vigorously shaken

for 1 min, sonicated for 1 min at 25 Hz, shaken for 10 min at 4°C in a Thermomixer

(Eppendorf®), and then centrifuged (8,000 *g*, 4°C, 10 min). The supernatants were collected,

and the pellets were re-extracted twice with 0.4 mL of the same extraction solution, then

vigorously shaken (1 min) and sonicated (1 min; 25 Hz). After the centrifugations, the three

supernatants were pooled and dried (final volume of 1.6 mL). Each dry extract was dissolved

in 140 µL acetone, filtered, and analyzed using a Waters Acquity ultra performance liquid

chromatograph coupled to a Waters Xevo triple quadrupole mass spectrometer TQS (UPLC-

ESI-MS/MS). The compounds were separated on a reverse-phase column (Uptisphere C18

UP3HDO, 100*2.1 mm*3µm particle size; Interchim, France) using a flow rate of 0.4 mL.min⁻

^{1} and a binary gradient: (A) acetic acid 0.1 % in water (v/v) and (B) acetonitrile with 0.1 %

acetic acid. The following binary gradient was used (t, % A): (0 min, 98 %), (3 min, 70 %), (7.5

min, 50 %), (8.5 min, 5 %), (9.6 min, 0%), (13.2 min, 98 %), (15.7 min, 98 %). Mass

spectrometry was conducted in electrospray and Multiple Reaction Monitoring scanning

mode (MRM mode), in negative ion mode (see Le Roux *et al.* 2014 for monitored

transitions). Relevant instrumental parameters were set as follows: capillary 1.5 kV (negative

mode), source block and desolvation gas temperatures 130°C and 500°C, respectively.

Nitrogen was used to assist the cone and desolvation (150 L.h^{-1} and 800 L.h^{-1}, respectively),

argon was used as the collision gas at a flow of 0.18 mL.min^{-1}.

Statistical analyses

All analyses were performed with the R software. For all experiments, the selected models

and the contrast tests are detailed in Methods S1.

For all experiments measuring seed germination, germination success was analyzed using a

generalized linear model (McCullagh and Nelder, 1989) assuming a binomial distribution

with the dispersion parameter fixed at unity and a logit link function.

For each experiment, the model was selected as follows: (i) factors included in the model

were defined according to the experimental design; (ii) starting from the most complete

model, nested models were sequentially fitted and the best one according to BIC was

selected. Once the model selected, the contrasts relevant to the addressed question were

tested, and p-values were adjusted using the Bonferroni procedure to control the family-

wise error rate (FWER). A contrast was declared significant if its adjusted p-value was lower

than 0.05.

We performed a contrast test procedure to detect significant interacting cytoplasm x

nucleus combinations: each test concerned a pair of cytoplasms (C1 and C2) and a pair of

nuclei (Na and Nb), designed in the text as 'cytonuclear interacting combination'; a

significant p-value indicated that the effect of changing cytoplasm C1 to C2 in the Na nuclear

background differed from the effect of the same change in the Nb background, H0

{([C1]Na-[C2]Na) - ([C1]Nb-[C2]Nb) = 0}. The analysis of data from the complete set of 64

genotypes (8 parents and 56 cytolines) thus allows for the testing of 784 cytonuclear

interacting combinations (28 pairs of nuclei x 28 pairs of cytoplasms).

ABA content and Na^+/K^+ ion ratio were analyzed using linear models. The selection of the

model and tests for effects were conducted as described above.

Results

Effects of cytoplasmic variations and cytonuclear interactions on seed dormancy

Following on from the observation of differences in germination rate in field-grown Arabidopsis cytolines (Roux *et al.*, 2016), we sought to evaluate whether cytoplasmic variation and cytonuclear interactions specifically impacted seed primary dormancy and the kinetics of its release during after ripening storage. Germination of seeds of the complete set of 64 genotypes (8 natural accessions and 56 cytolines) was tested at harvest and three, six and nine months post-harvest (0PH, 3PH, 6PH or 9PH), without any stratification treatment, in two temperature conditions, 15°C and 25°C (Fig. 1 and S1). As expected from the wide variation for dormancy among the panel of parental accessions, the nuclear background had a strong effect on germination. Notably, no or low germination was obtained with seeds of the Ita-0 nuclear series in any conditions. For this reason, these genotypes were excluded from the subsequent analyses. Most seed lots germinated better at 15°C (Fig. 1) than at 25°C (Fig. S1), as previously reported for Col-0 freshly harvested seeds (Leymarie *et al.*, 2012), except those of the Bur-0 nuclear series, which reached almost 100% germination at 25°C at harvest, but displayed lower germination success at 15°C.

We tested whether germination of cytolines was modified compared to their natural nuclear parent. Table 1 shows that changing organelle genomes had effects on dormancy, but not to the same extent nor in the same direction in all nuclear backgrounds. When effects were observed, foreign cytoplasms tended to enhance dormancy in Ct-1, Jea, Oy-0 and Sha nuclear backgrounds (lower germination in cytolines compared to nuclear natural parent), whereas they rather lowered dormancy in Blh-1, Bur-0, and Cvi-0 backgrounds (higher germination in cytolines). Interestingly, the impact of a given cytoplasm differed according to the nucleus it was associated with, suggesting that dormancy was influenced by cytonuclear interactions. We thus tested for significant cytonuclear interacting combinations impacting dormancy, *i.e.* for differences between two given nuclei in the effect of a given change of cytoplasm, for all possible pairs of cytoplasms and all possible pairs of nuclei (see Material and Methods and Methods S1 for details). As some genotypes were excluded from the analysis due to missing data or total absence of germination, 441 combinations were tested for 0PH data, and 505 for data at other PH times. After Bonferroni corrections for multiple tests, we found 177 significant cytonuclear interacting combinations at 0PH, and 232, 186, and 204 at 3PH, 6PH and 9PH, respectively (Table S2). This indicated that the depth

of seed primary dormancy is modulated by genetic variation in interacting nuclear- and

organelle-encoded partners.

As seen in Table 1, observed effects of foreign cytoplasms on dormancy often persisted in

successive PH times, which was somewhat expected since an effect on the depth of

dormancy at harvest would still be detectable after some after-ripening, according to the

kinetics of dormancy release. Similarly, we observed persisting effects of cytonuclear

interacting combinations over several PH times (Table S2): among the 391 cytonuclear

interacting combinations that influenced germination at any PH time, 237 (60%) were

significant at two PH times or more and 47 (12%) at all four PH times and, in the vast

majority (between 86 % to 100 %) of the cases, they modified germination in the same

direction (Table 2), in accordance with a persistence of the observed effects during after-

ripening.

Interestingly, the rate of dormancy release during after-ripening also seemed to be modified

by foreign cytoplasms in some cases: as seen in Table 1, no difference compared to the

natural nuclear parent at 0PH could be followed by either better (*e.g.* [Jea]Cvi-0) or lower

(*e.g.* [Sha]Blh-1) germination percentage in later PH times. In addition, a modification in the

kinetics of dormancy release was indicated by a change in the contrast sign between PH

times in two cases: [Oy-0]Sha germinated less than Sha at 0PH but better at following PH

times; oppositely, [Bur-0]Cvi-0 germinated better than Cvi-0 at 0PH, but less at 9PH.

Similarly, we interpreted opposite signs of contrast for the effects of a cytonuclear

interacting combination significant at to PH times (Table 2) as indication that it altered the

kinetics of dormancy release.

Effects of cytoplasmic variations and cytonuclear interactions on germination performance

In order to assess whether a disruption of cytonuclear co-adaptation also affected the

germination performance of seeds after dormancy release, we measured the germination of

the complete set of genotypes under permissive and challenging conditions, specifically in

the presence of NaCl, at least 18 months after harvest and after stratification. We used salt

stress to enhance differences in germination performance that could be difficult to assess

after the stratification treatment. Because seed germination tolerance to NaCl is variable

from one accession to another (Galpaz and Reymond, 2010), each nuclear series was tested

on the NaCl concentration (Table S1) that resulted in 50% germination of the parental

accession in a preliminary experiment. Time plots are shown in Fig. S2. Despite the
stratification treatment, 100% of germination was not reached on water in Ct-1 and Cvi-0
nuclear series, although germination had reached the plateau (Gmax). Nevertheless, we
considered the Gmax as an indicator of germination performance and tested, in each
nuclear series, the germination performance of cytolines compared to the natural accession,
averaged on the two conditions (Fig. 2, see Methods S1 for details). The results clearly
showed that the sensitivity of germination to a cytoplasm change was very variable among
the nuclear backgrounds, Blh-1 being very sensitive whereas Ct-1 and Oy-0 were not. In
general, when affecting germination, foreign cytoplasms had a negative effect. However,
three cytolines ([Ct-1]Ita-0, [Ct-1]Jea and [Blh-1]Sha) did germinate better than their
respective nuclear parent. After Bonferroni correction for multiple tests, 202 out of 784
cytonuclear interacting combinations (26%, Table S2) significantly impacted germination
success in this experiment. Thus, as dormancy, germination performance was influenced by
interactions between organelle- and nuclear-encoded factors that vary among the panel of
parental accessions.
Among the 505 cytonuclear interacting combinations that were tested both for germination
performance and for dormancy, 92 were significant for both traits. This represented 80% of
the interacting combinations impacting germination performance (92/115), but only 23% of
those impacting dormancy for at least one PH (92/392). In addition, the cytonuclear
interacting combinations that influenced both dormancy and germination performance
impacted germination success more often in the same direction in both experiments than
would be randomly expected (Table 3, Chi^2 test pval <0.001), indicating a negative
correlation between dormancy and germination performance.
304
Contribution of individual accessions to cytonuclear interactions that impact germination
The above analyses, the impact on germination traits of a shuffling of genetic compartments
depended on the origin of the genomes. To better assess this variation, we analyzed the
contribution of individual accessions to cytonuclear interacting combinations affecting
dormancy or germination performance. Fig. 3 indicates, for each accession, the number of
significant cytonuclear interacting combinations that involved its cytoplasm or its nucleus
and that impacted dormancy depth, conservatively estimated as effects maintained through
all PH times (Fig. 3a) or germination performance after storage (Fig. 3b). Among the 47

cytonuclear interacting combinations that impacted dormancy depth, 28 involved the Ita-0

cytoplasm and 33 involved the Jea nucleus. Among the 202 significant interacting

combinations that affected germination performance after storage, 100 involved the Blh-1

cytoplasm and 76 the Sha nucleus. When considering the contribution of accessions to

cytonuclear interacting combinations that affect both dormancy and germination

performance, the Blh-1 and Ita-0 cytoplasms and the Sha nucleus clearly emerged (Fig.S3).

Effects of cytoplasm variation on seed longevity in Sha and Ct-1 nuclear backgrounds

Along with dormancy and germination performance, longevity is an important component of

seed vigor. Thus, we addressed whether this trait was also under the influence of

cytoplasmic variation. We selected the Sha and Ct-1 series because these two nuclear

backgrounds represented two extreme cases in their contribution to cytonuclear effects in

the previous experiments (Fig. 3 and S3). The longevity of seeds from these two cytoline

series was estimated by measuring their viability (germination percentage in standard

conditions) after 0, 10, 20, 30 or 40 days of controlled deterioration treatment (CDT) (Fig. 4).

We used the same seed lots as in the assays for germination performance after storage. In

this experiment, all genotypes of the Ct-1 series reached 100% germination. We concluded

that the partial seed imbibition followed by desiccation included in the CDT treatment, even

for the control, acted as a priming treatment releasing the residual dormancy observed in

the experiment on germination performance described above. The kinetics of viability loss

were different in the two nuclear backgrounds, revealing an expected nuclear effect on seed

longevity (Fig. 4). Interestingly, four cytolines appeared less sensitive to CDT than their

natural nuclear parent, whereas none was more sensitive (Table 4, see Methods S1 for

details). In addition, in both nuclear backgrounds, the selected statistical models included an

interaction between the cytoplasm and the length of CDT treatment, which suggested that

the loss of germination ability between two treatment durations was under the influence of

the cytoplasmic genomes. We could not test directly for significant cytonuclear interacting

combinations, because the two series were measured in different experiments and the

results analyzed independently. Nevertheless, in Blh-1/Sha and Ita-0/Sha cytoplasm pairs,

individual cytoplasms had opposite effects in the two nuclear backgrounds (Table S3),

indicating an influence of cytonuclear epistasis on germination after CDT. Strikingly, the Sha

cytoplasm performed better than the Blh-1 and the Ita-0 cytoplasms in the Ct-1 nuclear

background but worse when associated with its natural nuclear partner.

Physiological clues for germination performance of the [Blh-1]Sha line

A closer look at the 202 cytonuclear interacting combinations that influence germination

performance revealed that the [Blh-1]Sha combination was involved in 46 (23%, Table S2) of

them, indicating that this cytonuclear combination had a major effect on germination. In

addition, this genetic combination also had a strong impact on CDT tolerance. We thus

decided to get further insight into the behavior of [Blh-1]Sha seeds.

In order to test whether the peculiar behavior of the [Blh-1]Sha seeds was due to the

conditions of seed production, we tested germination performance of [Blh-1]Sha and Sha

seeds from two productions in different environments, growth chamber and greenhouse.

We observed higher germination performance of the cytoline compared to the natural

accession for both seed productions (Fig. S4). An environmental cause of the higher

germination performance of [Blh-1]Sha was therefore very unlikely. However, we also

observed that this positive effect did not persist in seed samples that had been stored

several months at room temperature in the laboratory.

We then tested whether the germination performance of [Blh-1]Sha was possibly

maintained in the presence of mannitol (200 mM) and KCl (100 mM), by analyzing the

germination efficiency of the Sha cytoline series (Fig. S5). The concentrations applied were

chosen to compare results to those of the NaCl treatment (100 mM) in the germination

performance experiment. The germination rate of the [Blh-1]Sha cytoline was consistently

higher than that of the other genotypes under both osmotic and ionic stresses, although

these stresses had a milder effect than the NaCl treatment on all genotypes, as previously

reported (Munns and Tester, 2008). The germination curve of [Blh-1]Sha on KCl grouped

with those of the other lines in the absence of KCl.

It was conceivable that the better germination performance of [Blh-1]Sha seeds would be

amplified by an increased tolerance to salt in germination under NaCl stress. Since Na^+/K^+

ratio is a key indicator of salt tolerance (Shabala and Cuin, 2008), potassium and sodium

contents were analyzed in seeds of [Blh-1]Sha, [Ita-0]Sha and Sha. The [Ita-0]Sha was used as

a control cytoline with germination performance similar to Sha (Fig. 2). Very little sodium

accumulated in seeds in the absence of NaCl. When germinated on NaCl, both [Blh-1]Sha

and [Ita-0]Sha seeds showed a lower Na^+/K^+ ratio than Sha seeds (Fig. 5). As the two

cytolines had different germination responses on NaCl compared to Sha, we considered it

unlikely that the Na^+/K^+ balance is the primary cause of the increased germination success of

the [Blh-1]Sha cytoline on NaCl. These results reinforced our hypothesis that the intrinsic

germination capacity of [Blh-1]Sha seeds was enhanced compared to its natural parent.

We then focused on ABA content and sensitivity to further explore physiological parameters

that could contribute to the enhanced germination performance of [Blh-1]Sha seeds. We

measured the endogenous ABA content of seeds from [Blh-1]Sha, [Ita-0]Sha, and Sha lines.

ABA contents were measured in both dry and stratified seeds placed under germination

conditions for six hours, in the presence or absence of NaCl (Fig. 6a). The experiment was

repeated after 10 further months of storage in controlled conditions (Versailles Stock

Centre), yielding similar results. The combined analysis of data from both experiments (see

Methods S1 for details) indicated that dry seeds of [Blh-1]Sha contained more ABA than

those of [Ita-0]Sha and Sha (p-values < 0.001), whereas they contained less ABA after six

hours of germination, either on water or NaCl (p-values < 0.01). The sensitivity of

germinating seeds to exogenous ABA (50 µM) was tested on the same seed stocks after the

second experiment (Fig. 6b). The [Blh-1]Sha seeds were less sensitive to ABA treatment than

Sha seeds. Taken together, the results suggest that both endogenous ABA metabolism and

ABA sensitivity are modified in [Blh-1]Sha seeds compared to Sha seeds.

Discussion

In this study, we examined seed physiological traits important for adaptation and of relevance to agriculture, in relation to cytoplasmic genetic variants and to the coadaptation between genomic compartments. Cytolines are original and valuable genetic resources to enlighten cytoplasmic and cytonuclear effects (Roux *et al.*, 2016). The analysis of a cytoline series from a given nuclear parent allows assessment of the consequences of a disruption of cytonuclear coadaptation in the considered nuclear background. The di-allele cross design used to produce the cytolines is suited to the detection of cytonuclear genetic interactions. When examining seed traits, this approach does not distinguish effects acting at the level of the mother plant from those acting at the level of the zygote(s). Nevertheless, as both the mother plant and progeny have the same genotype, it does not preclude the detection of cytoplasmic or cytonuclear effects, keeping in mind that they could act in the mother plant and/or in the progeny.

Dormancy was assessed by monitoring germination of freshly harvested seeds, or seeds after three, six or nine months of after-ripening, without stratification. Disruption of cytonuclear coadaptation had variable effects on dormancy depending on the nuclear background, both in the number of cytolines with modified dormancy and in the direction of the effect. When dormancy differed in a cytoline compared to its parent, it tended to be deeper when the parent was weakly dormant (*e.g.* Jea, Sha) and weaker in the deeply dormant Cvi-0 background. This suggests that coadapted variants of nuclear and organellar genes contributed to adaptive dormancy. However, this was not always verified: in the weakly dormant Bur-0, novel cytonuclear combinations enhanced germination. The Bur-0 nuclear series was remarkably different from the others regarding germination at the two tested temperature conditions: in contrast to the other nuclear series, the seeds of the Bur-0 nuclear series had no dormancy phenotype at 25°C (Fig. S1) but displayed dormancy at 15°C (Fig.1). A peculiar response to sowing temperature of Bur-0 seeds was also reported in a study examining a set of 73 accessions where they displayed almost 100% germination at 26°C and around 50% germination at 18°C (Schmuths *et al.*, 2006). Genetic variation for the rate of dormancy loss after-ripening was previously reported for Arabidopsis (Barua *et al.*, 2012). Here, by revealing that novel genetic combinations modify the kinetics of dormancy release, we provide evidence that this rate can be modulated by cytonuclear genetic interactions.

Germination performance was estimated through the germination percentage of seeds in
normal and challenging conditions. The effects of foreign cytoplasms tended to lower
germination performance, which is expected if optimal germinative success is assumed for
natural coadapted genotypes. However, some novel cytonuclear combinations had better
germination performance than their nuclear parent, suggesting that natural accessions do
not always have optimal germination performance, at least in laboratory conditions. A good
germination efficiency under challenging stress conditions was reported as a component of
seed vigor and related to germination speed (Finch-Savage and Bassel, 2016; Yuan *et al.*,
2016*a*). Although the limited number of time points precluded the precise fitting of
germination curves and reliable comparison of germination speeds, examination of the
germination time plots (Fig. S2) strongly suggested that cytolines with higher germination
performance also germinated quicker than their natural parent, whereas those performing
worse germinated slower. In addition, the germination percentage was influenced by
cytonuclear interactions independently of the presence of salt (no third-order level of
interaction), which was consistent with the idea that the detected cytonuclear effects affect
germination performance *per se* rather than salt tolerance. This was further verified in
[Blh-1]Sha and [Ita]Sha where a marker of salt tolerance (lower Na^+/K^+ ratio) was not linked
to germination performance, whereas the specific effect of the Blh-1 cytoplasm on ABA
decay rate was observed both on water and NaCl. We therefore conclude that our analysis
identified effects of cytoplasm variation and cytonuclear interactions on a component of
seed vigor. As longevity is also associated to seed vigor, we assessed germination after CDT
in Ct-1 and Sha nuclear series, which presented contrasted sensitivities to cytoplasmic
change for germination performance and dormancy. Strikingly, disruption of cytonuclear co-
adaptation had no or positive effect on seed longevity, at least for the genetic variation
tested. This was unexpected since lower seed longevity is generally indicative of low seed
vigor and an unfit genotype. It is interesting to note that [Sha]Ct-1 and [Blh-1]Sha displayed
both higher longevity and germination performance compared to their natural parents. It is
conceivable that both phenotypes revealed the same underlying beneficial effect on seed
vigor of the new cytonuclear combinations. However, the [Ita-0]Sha cytoline, which had a
higher seed longevity than Sha, did not germinated better. A negative correlation between
dormancy and seed longevity determined by co-located QTLs was observed in recombinant
inbred-line populations of Arabidopsis (Nguyen *et al.*, 2012). This negative correlation was

not verified here: we observed deeper dormancy at 0PH and higher tolerance to CDT in the

[Ita-0]Sha cytoline than in Sha. This suggests that the cytonuclear interaction effects on seed

longevity and dormancy in this cytoline are not driven by the loci reported in Nguyen's work

or, not exclusively, that this cytoplasm change has different impacts on the Sha allelic effects

on seed dormancy and longevity at these loci. It is also conceivable that the apparently

contradiction between our result and that from Nguyen *et al.* (2012) come from differences

in the conditions used to estimate seed longevity, *i.e.* survival after several years of dry

storage versus CDT for 10-40 days.

Some accessions contributed more than others to the cytonuclear interacting combinations

affecting germination, and individual accessions contributed differently to interactions that

affect dormancy depth and germination performance after storage. For example, the Jea

nucleus had the highest contribution to significant cytonuclear interacting combinations for

dormancy. This suggests that, when changing the organellar partners of cytonuclear

epistasis involved in dormancy tuning, Jea alleles lead to different consequences on the

phenotype than alleles from other parents.

We found that the majority of cytonuclear interacting combinations that modulated

germination performance also altered dormancy, with a negative correlation between

dormancy and germination performance in most cases (Table 3). This is consistent with the

idea that dormancy and germination speed are parts of a continuum that control

germination (Finch-Savage and Leubner-Metzger, 2006). We exclude that residual dormancy

of Ct-1 and Cvi-0 series induced a bias toward this conclusion because the contributions of

these nuclei to significant cytonuclear interacting combinations shared by germination

performance and dormancy experiments are among the fewest observed (Fig. S3). The

continuity of the biological process underlying dormancy and seed vigor was recently

brought back to light by a report that linked genetic variation in ABA sensitivity with seed

germination performance in *Brassica oleracea* (Awan *et al.*, 2018). ABA content was also

reported to be related to germination speed in *B. oleracea* (Morris *et al.*, 2016). In this

context, our results on the [Blh-1]Sha cytoline seem particularly relevant. At the moment, a

formal causal link cannot be made between the enhanced germination vigor of the cytoline

on one hand, and faster ABA decay or reduced ABA sensitivity on the other hand.

Nevertheless, these observations provide an interesting path for further research. In parallel

to deciphering the physiological consequences of novel genetic combinations that modulate

seed performance, genetic studies should also lead to the identification of gene variants that

underlie the effects reported here.

One remarkable and unexpected result of the present study is the enhanced seed behavior

of some cytolines compared to natural genotypes that were shaped by coevolution.

Interestingly, the cytoplasm of the drought adapted Kas-1 accession was reported to have a

strong and negative effect on water use efficiency in a QTL analysis (Mckay *et al.*, 2008), also

suggesting that the natural genotype was not optimal for this trait. It is conceivable that

trade-offs between adaptive characters have led to the maintenance, in natural variants of

Arabidopsis, of suboptimal genomic combinations for adaptive traits. Indeed, trade-offs

between traits under selection, including germination, have been previously shown to

underlie natural genetic variation in Arabidopsis (Donohue *et al.*, 2005*b*; Todesco *et al.*,

2010; Debieu *et al.*, 2013). In light of these results, we suggest that more attention is given

to cytoplasmic variation and its effect on seed vigor in the genetic resources of cultivated

species.

Supplementary data

Methods S1. Statistical models and contrast tests.

Fig. S1. Dormancy release during after ripening in cytoline series, germination at 25°C.

Fig. S2. Germination time curves of cytoline series on water and NaCl.

Fig. S3. Contribution of individual accessions to cytonuclear interacting combinations

impacting dormancy and germination performance.

Fig. S4. Germination performance of [Blh-1]Sha and Sha seeds from two environments.

Fig. S5. Germination time curves of the Sha cytoline series on mannitol and KCl.

Table S1. NaCl concentrations used in the germination performance experiment

Table S2. Results of tests for significant cytonuclear interacting combinations in dormancy

and germination performance experiments.

Table S3. Results of tests for significant effect of cytoplasms on germination after CDT.

Acknowledgments

We thank Simon Law, Annie Marion-Poll and Helen North for helpful comments on the

manuscript. We are grateful to Sébastien Thomine for his help in ion dosages. This work was

funded by the Plant Biology and the Genetics and Plant Breeding INRA Departments

("Cytolignées pilotes"-2010 and "Cytoressources"-2011) and by the ANR BIOADAPT program

(ANR-12_ADAP-0004). Institute Jean-Pierre Bourgin (IJPB) and Institute of Plant Sciences

Paris-Saclay (IPS2) benefit from support from the LABEX Saclay Plant Sciences-SPS (ANR-10-

LABX-0040-SPS).

References

Alonso-Blanco C, Bentsink L, Hanhart CJ, Blankestijn-de Vries H, Koornneef M. 2003. Analysis of natural allelic variation at seed dormancy loci of Arabidopsis thaliana. Genetics **164**, 711–729.

Awan S, Footitt S, Finch-Savage WE. 2018. Interaction of maternal environment and allelic differences in seed vigour genes determines seed performance in Brassica oleracea. The Plant Journal **94**, 1098–1108.

Barua D, Butler C, Tisdale TE, Donohue K. 2012. Natural variation in germination responses of Arabidopsis to seasonal cues and their associated physiological mechanisms. Annals of Botany **109**, 209–226.

Bentsink L, Hanson J, Hanhart CJ, *et al.* 2010. Natural variation for seed dormancy in Arabidopsis is regulated by additive genetic and molecular pathways. Proceedings of the National Academy of Sciences of the United States of America **107**, 4264–4269.

Bentsink L, Jowett J, Hanhart CJ, Koornneef M. 2006. Cloning of DOG1, a quantitative trait locus controlling seed dormancy in Arabidopsis. Proceedings of the National Academy of Sciences of the United States of America **103**, 17042–17047.

Bryant N, Lloyd J, Sweeney C, Myouga F, Meinke D. 2011. Identification of nuclear genes encoding chloroplast-localized proteins required for embryo development in Arabidopsis. PLANT PHYSIOLOGY **155**, 1678–1689.

Clerkx EJM, El-Lithy ME, Vierling E, Ruys GJ, Blankestijn-de Vries H, Groot SPC, Vreugdenhil D, Koornneef M. 2004. Analysis of natural allelic variation of Arabidopsis seed germination and seed longevity traits between the accessions Landsberg erecta and Shakdara, using a new recombinant inbred line population. PLANT PHYSIOLOGY **135**, 432–443.

Corey LA, Matzinger DF, Cockerham CC. 1976. Maternal and Reciprocal Effects on Seedling Characters in ARABIDOPSIS THALIANA (L.) Heynh. Genetics **82**, 677–683.

De Souza A, Wang J-Z, Dehesh K. 2017. Retrograde Signals: Integrators of Interorganellar Communication and Orchestrators of Plant Development. Annual Review of Plant Biology **68**, 85–108.

Debieu M, Tang C, Stich B, Sikosek T, Effgen S, Josephs E, Schmitt J, Nordborg M, Koornneef M, de Meaux J. 2013. Co-Variation between Seed Dormancy, Growth Rate and Flowering Time Changes with Latitude in Arabidopsis thaliana (JO Borevitz, Ed.). PLoS ONE **8**, e61075.

Demarsy E, Buhr F, Lambert E, Lerbs-Mache S. 2011. Characterization of the plastid-specific germination and seedling establishment transcriptional programme. Journal of Experimental Botany **63**, 925–939.

Donohue K, Dorn L, Griffith C, Kim E, Aguilera A, Polisetty CR, Schmitt J. 2005a. The evolutionary ecology of seed germination of Arabidopsis thaliana: variable natural selection on germination timing. Evolution **59**, 758–770.

Donohue K, Dorn L, Griffith C, Kim E, Aguilera A, Polisetty CR, Schmitt J. 2005b. Environmental and genetic influences on the germination of Arabidopsis thaliana in the field. Evolution **59**, 740–757.

Dubois J, Cheptou P-O. 2012. Competition/colonization syndrome mediated by early germination in non-dispersing achenes in the heteromorphic species Crepis sancta. Annals of Botany **110**, 1245–1251.

Dyall SD, Brown MT, Johnson PJ. 2004. Ancient invasions: from endosymbionts to organelles. Science (New York, NY) **304**, 253–257.

Finch-Savage WE, Bassel GW. 2016. Seed vigour and crop establishment: extending performance beyond adaptation. Journal of Experimental Botany **67**, 567–591.

Finch-Savage WE, Footitt S. 2017. Seed dormancy cycling and the regulation of dormancy mechanisms to time germination in variable field environments. Journal of Experimental Botany **68**, 843–856.

Finch-Savage WE, Leubner-Metzger G. 2006. Seed dormancy and the control of germination. The New phytologist **171**, 501–523.

Finch-Savage WE, Clay HA, Lynn JR, Morris K. 2010. Towards a genetic understanding of seed vigour in small-seeded crops using natural variation in Brassica oleracea. Plant Science **179**, 582–589.

Galpaz N, Reymond M. 2010. Natural variation in Arabidopsis thaliana revealed a genetic network controlling germination under salt stress. PLoS ONE **5**, e15198.

Hämälä T, Mattila TM, Leinonen PH, Kuittinen H, Savolainen O. 2017. Role of seed germination in adaptation and reproductive isolation in Arabidopsis lyrata. Molecular Ecology **26**, 3484–3496.

Howell KA, Narsai R, Carroll A, Ivanova A, Lohse M, Usadel B, Millar AH, Whelan J. 2008. Mapping Metabolic and Transcript Temporal Switches during Germination in Rice Highlights Specific Transcription Factors and the Role of RNA Instability in the Germination Process. PLANT PHYSIOLOGY **149**, 961–980.

Huang X, Schmitt J, Dorn L, Griffith C, Effgen S, Takao S, Koornneef M, Donohue K. 2010. The earliest stages of adaptation in an experimental plant population: strong selection on QTLS for seed dormancy. Molecular Ecology **19**, 1335–1351.

Joosen RVL, Arends D, Willems LAJ, Ligterink W, Jansen RC, Hilhorst HWM. 2012. Visualizing the genetic landscape of Arabidopsis seed performance. PLANT PHYSIOLOGY **158**, 570–589.

Kronholm I, Picó FX, Alonso-Blanco C, Goudet J, Meaux J de. 2012. Genetic basis of adaptation in Arabidopsis thaliana: local adaptation at the seed dormancy QTL DOG1. Evolution **66**, 2287–2302.

Kutschera U, Niklas KJ. 2005. Endosymbiosis, cell evolution, and speciation. Theory in biosciences = Theorie in den Biowissenschaften **124**, 1–24.

Law SR, Narsai R, Taylor NL, Delannoy E, Carrie C, Giraud E, Millar AH, Small I, Whelan J. 2012. Nucleotide and RNA metabolism prime translational initiation in the earliest events of

mitochondrial biogenesis during Arabidopsis germination. PLANT PHYSIOLOGY **158**, 1610–1627.

Le Roux C, Del Prete S, Boutet-Mercey S, Perreau F, Balagué C, Roby D, Fagard M, Gaudin V. 2014. The hnRNP-Q Protein LIF2 Participates in the Plant Immune Response (X Zhang, Ed.). PLoS ONE **9**, e99343.

Leymarie J, Vitkauskaité G, Hoang HH, Gendreau E, Chazoule V, Meimoun P, Corbineau F, El-Maarouf-Bouteau H, Bailly C. 2012. Role of reactive oxygen species in the regulation of Arabidopsis seed dormancy. Plant and Cell Physiology **53**, 96–106.

McCullagh P, Nelder JA. 1989. *Generalized Linear Models*. London: Chapman & Hall.

Mckay JK, Richards JH, Nemali KS, Sen S, Mitchell-Olds T, Boles S, Stahl EA, Wayne T, Juenger TE. 2008. Genetics of drought adaptation in Arabidopsis thaliana II. QTL analysis of a new mapping population, KAS-1 x TSU-1. Evolution **62**, 3014–3026.

Mckhann HI, Camilleri C, Berard A, Bataillon T, David JL, Reboud X, Le Corre V, Caloustian C, Gut IG, Brunel D. 2004. Nested core collections maximizing genetic diversity in Arabidopsis thaliana. The Plant Journal **38**, 193–202.

Moison M, Roux F, Quadrado M, Duval R, Ekovich M, Lê D-H, Verzaux M, Budar F. 2010. Cytoplasmic phylogeny and evidence of cyto-nuclear co-adaptation in Arabidopsis thaliana. The Plant Journal **63**, 728–738.

Morris K, Barker GC, Walley PG, Lynn JR, Finch-Savage WE. 2016. Trait to gene analysis reveals that allelic variation in three genes determines seed vigour. The New phytologist **212**, 964–976.

Munns R, Tester M. 2008. Mechanisms of Salinity Tolerance. Annual Review of Plant Biology **59**, 651–681.

Narsai R, Law SR, Carrie C, Xu L, Whelan J. 2011. In-depth temporal transcriptome profiling reveals a crucial developmental switch with roles for RNA processing and organelle metabolism that are essential for germination in Arabidopsis. PLANT PHYSIOLOGY **157**, 1342–1362.

Nguyen TP, Keizer P, van Eeuwijk F, Smeekens S, Bentsink L. 2012. Natural Variation for Seed Longevity and Seed Dormancy Are Negatively Correlated in Arabidopsis. PLANT PHYSIOLOGY **160**, 2083–2092.

North H, Baud S, Debeaujon I, *et al.* 2010. Arabidopsis seed secrets unravelled after a decade of genetic and omics-driven research. The Plant Journal **61**, 971–981.

Paszkiewicz G, Gualberto JM, Benamar A, Macherel D, Logan DC. 2017. Arabidopsis seed mitochondria are bioenergetically active immediately upon imbibition and specialize via biogenesis in preparation for autotrophic growth. The Plant Cell, tpc.00700.2016.

Postma FM, Lundemo S, Agren J. 2015. Seed dormancy cycling and mortality differ between two locally adapted populations of Arabidopsis thaliana. Annals of Botany, mcv171.

Rajjou L, Debeaujon I. 2008. Seed longevity: Survival and maintenance of high germination ability of dry seeds. C.R. Biologies **331**, 796–805.

Rajjou L, Duval M, Gallardo K, Catusse J, Bally J, Job C, Job D. 2012. Seed Germination and Vigor. Annual Review of Plant Biology **63**, 507–533.

Rajjou L, Lovigny Y, Groot SPC, Belghazi M, Job C, Job D. 2008. Proteome-wide characterization of seed aging in Arabidopsis: a comparison between artificial and natural aging protocols. PLANT PHYSIOLOGY **148**, 620–641.

Roux F, Mary-Huard T, Barillot E, Wenes E, Botran L, Durand S, Villoutreix R, Martin-Magniette M-L, Camilleri C, Budar F. 2016. Cytonuclear interactions affect adaptive traits of the annual plant Arabidopsis thaliana in the field. Proceedings of the National Academy of Sciences of the United States of America **113**, 3687–3692.

Rurek M. 2016. Participation of non-coding RNAs in plant organelle biogenesis. Acta biochimica Polonica **63**.

Sano N, Rajjou L, North HM, Debeaujon I, Marion-Poll A, Seo M. 2016. Staying Alive: Molecular Aspects of Seed Longevity. Plant and Cell Physiology **57**, 660–674.

Savage LJ, Imre KM, Hall DA, Last RL. 2013. Analysis of Essential Arabidopsis Nuclear Genes Encoding Plastid-Targeted Proteins (S Park, Ed.). PLoS ONE **8**, e73291.

Schmuths H, Bachmann K, Weber WE, Horres R, Hoffmann MH. 2006. Effects of preconditioning and temperature during germination of 73 natural accessions of Arabidopsis thaliana. Annals of Botany **97**, 623–634.

Sew YS, Ströher E, Fenske R, Millar AH. 2016. Loss of Mitochondrial Malate Dehydrogenase Activity Alters Seed Metabolism Impairing Seed Maturation and Post-Germination Growth in Arabidopsis. PLANT PHYSIOLOGY **171**, 849–863.

Shabala S, Cuin TA. 2008. Potassium transport and plant salt tolerance. Physiologia Plantarum **133**, 651–669.

Shu K, Liu X-D, Xie Q, He Z-H. 2016. Two Faces of One Seed: Hormonal Regulation of Dormancy and Germination. Molecular Plant **9**, 34–45.

Shu K, Meng YJ, Shuai HW, Liu WG, Du JB, Liu J, Yang WY. 2015. Dormancy and germination: How does the crop seed decide? Plant Biology **17**, 1104–1112.

Sosso D, Canut M, Gendrot G, Dedieu A, Chambrier P, Barkan A, Consonni G, Rogowsky PM. 2012. PPR8522 encodes a chloroplast-targeted pentatricopeptide repeat protein necessary for maize embryogenesis and vegetative development. Journal of Experimental Botany **63**, 5843–5857.

Tesnier K, Strookman-Donkers HM. 2002. A controlled deterioration test for Arabidopsis thaliana reveals genetic variation in seed quality. Seed Science and Technology **30**, 149–165.

Todesco M, Balasubramanian S, Hu TT, *et al.* 2010. Natural allelic variation underlying a major fitness trade-off in Arabidopsis thaliana. Nature **465**, 632–636.

Vallejo AJ, Yanovsky MJ, Botto JF. 2010. Germination variation in Arabidopsis thaliana accessions under moderate osmotic and salt stresses. Annals of Botany **106**, 833–842.

van Der Schaar W, Alonso-Blanco C, Léon-Kloosterziel KM, Jansen RC, van Ooijen JW, Koornneef M. 1997. QTL analysis of seed dormancy in Arabidopsis using recombinant inbred lines and MQM mapping. Heredity **79 (Pt 2)**, 190–200.

Yang L, Peng X, Sun M-X. 2011. AtNG1 encodes a protein that is required for seed germination. Plant Science **181**, 457–464.

Yuan W, Flowers JM, Sahraie DJ, Purugganan MD. 2016*a*. Cryptic Genetic Variation for Arabidopsis thaliana Seed Germination Speed in a Novel Salt Stress Environment. G3 (Bethesda, Md.).

Yuan W, Flowers JM, Sahraie DJ, Ehrenreich IM, Purugganan MD. 2016*b*. Extreme QTL mapping of germination speed in Arabidopsis thaliana. Molecular Ecology **25**, 4177–4196.

Table 1. Impact of foreign cytoplasms on germination after various post-harvest times in the different nuclear backgrounds.

Nuclear background[a]	cytoplasm[b]	0 PH		3 PH		6 PH		9 PH	
		Pval[c]	direction of effect[d]	Pval	direction of effect	Pval	direction of effect	Pval	direction of effect
Blh1	Bur-0	***	+	***	+	***	+	ns	
	Ct-1	ns		ns		ns		*	-
	Cvi-0	ns		ns		ns		ns	
	Ita-0	*	+	**	+	ns		ns	
	Jea	ns		ns		**	+	ns	
	Oy-0	ns		ns		ns		ns	
	Sha	ns		ns		***	-	***	-
Bur-0	Blh-1	ns		ns		ns		ns	
	Ct-1	**	+	***	+	ns		ns	
	Cvi-0	***	+	*	+	***	+	ns	
	Ita-0	**	+	***	+	ns		ns	
	Jea	*	+	***	+	***	+	***	+
	Oy-0	ns		ns		ns		ns	
	Sha	ns		***	+	ns		*	+
Ct-1	Blh-1	ns		ns		ns		ns	
	Bur-0	ns		ns		ns		ns	
	Cvi-0	*	-	ns		**	-	***	-
	Ita-0	ns		ns		ns		ns	
	Jea	ns		ns		ns		ns	
	Oy-0	ns		ns		ns		ns	
	Sha	ns		***	+	ns		ns	
Cvi-0	Blh-1	NA		ns		ns		**	-
	Bur-0	***	+	ns		ns		**	-
	Ct-1	***	+	***	+	***	+	ns	
	Ita-0	*	+	ns		***	+	ns	
	Jea	NA		ns		ns		***	+
	Oy-0	NA		ns		ns		ns	
Jea	Blh-1	***	-	**	-	***	-	***	-
	Bur-0	***	-	***	-	ns		ns	
	Cvi-0	ns		*	-	***	-	***	-
	Ita-0	***	-	***	-	***	-	***	-
	Oy-0	ns		ns		ns		ns	
	Sha	ns		ns		ns		ns	
Oy-0	Blh-1	ns		ns		ns		ns	
	Bur-0	**	-	ns		ns		ns	
	Ct-1	***	-	ns		ns		ns	
	Cvi-0	***	-	ns		ns		ns	
	Ita-0	***	+	ns		ns		ns	
	Jea	*	-	ns		ns		ns	
	Sha	ns		ns		ns		ns	
Sha	Blh-1	ns		ns		ns		*	+
	Bur-0	***	-	ns		***	-	**	-
	Ct-1	**	-	ns		ns		ns	
	Cvi-0	**	-	ns		ns		ns	
	Ita-0	***	-	ns		ns		ns	
	Jea	ns		ns		ns		ns	
	Oy-0	***	-	***	+	***	+	*	+

[a] The results for the Ita-0 nuclear background were not analyzed due to extremely low germination in all conditions (see Fig. 1 and Fig. S1).

[b] For each nuclear background, missing foreign cytoplasms correspond to genotypes excluded before analysis due to missing data or total absence of germination (NAs at 0PH).

[c] For each PH time, a difference in germination success between the genotype and its nuclear parent was tested (see Methods S1 for details). For 3PH, 6PH and 9PH experiments, the cytoplasm effects must be interpreted as averaged on germination temperatures. Pvalue : ***<0.001<**<0.01<*<0.05; ns, non significant.

[d] the direction of the effect is indicated by the sign of the contrast result. (+) indicates that the tested genotype had a higher germination than its natural nuclear donor. (-) indicates the opposite result.

Table 2. Comparison of directions of the effects of significant cytonuclear interacting combinations shared by two PH times.

comparison	number of shared cytonuclear interactions	number of identical direction of effects[a]	number of opposite directions of effects[b]
0PH vs 3PH	103	95	8
0PH vs 6PH	82	71	11
0PH vs 9PH	77	69	8
3PH vs 6PH	124	123	1
3PH vs 9PH	110	103	7
6PH vs 9PH	130	130	0

[a] Identical direction of effects corresponds to identical signs of contrast results in Table S2. It indicates that the considered cytonuclear interacting combination affected germination success in the same direction at the two PH times.

[b] Opposite directions of effects corresponds to opposite signs of the contrast results in Table S2. It indicates that the considered cytonuclear interacting combination affected germination success in opposite directions at the two PH times.

Table 3. Comparison of directions of effects of cytonuclear interacting combinations significant both in the dormancy and germination performance experiments.

comparison[a]	number of shared significant cytonuclear interacting combinations	number of identical direction of effects[b]	number of opposite directions of effects[c]
Gmax vs 0PH	49	36	13
Gmax vs 3PH	59	45	14
Gmax vs 6PH	37	20	17
Gmax vs 9PH	43	30	13

[a] Gmax stands for germination success in the experiment testing germination performance; xPH stands for germination success after x months PH in the dormancy experiment.

[b] Identical direction of the effects correspond to identical signs of the contrasts results in Table S2. Based on the writing of the contrasts, it indicates that the considered cytonuclear interacting combination affected germination success in the same direction in both experiments, thus dormancy and germination performance in opposite directions.

[c] On the same basis, opposite directions of the effects correspond to opposite signs of the contrasts results in Table S2. It indicates that the considered cytonuclear interacting combination affected germination success in opposite directions in both experiments, thus dormancy and germination performance in the same direction.

Table 4. Impact of foreign cytoplasms on germination after CDT in the Sha and Ct-1 nuclear backgrounds

cytoplasm[a]	Ct-1 nuclear background		Sha nuclear background	
	Pval[b]	direction of effect[c]	Pval	direction of effect
Blh-1	ns		***	+
Bur-0	*	+	ns	
Ct-1	/	/	ns	
Cvi-0	ns		ns	
Ita-0	ns		*	+
Jea	ns		ns	
Oy-0	ns		ns	
Sha	***	+	/	/

[a] Each foreign cytoplasm was tested for effect compared to the parental cytoplasm (see Methods S1 for details). The cytoplasm effects must be interpreted as averaged on {10, 20 ,30} days of CDT.

[b] Pvalue : ***<0.001<**<0.01<*<0.05; ns, non significant.

[c] (+) indicates that the tested genotype had a higher germination after CDT than its natural nuclear donor. (-) indicates that the tested genotype had a lower germination after CDT than its natural nuclear donor.

Figure legends

Fig. 1 Dormancy release during after ripening in cytoline series.

Each panel shows boxplots of the maximum germination percentage at 15°C for one cytonuclear combination 0, 3, 6 and 9 months after harvest (mph). Rows present genotypes sharing their nucleus and columns those sharing their cytoplasm. Natural accessions are framed in red. The [Sha]Cvi-0 cytoline could not be tested due to the small amount of seeds obtained by hand pollination on this male-sterile genotype.

Fig. 2 Germination performance of cytoline series.

Each panel shows boxplots of the maximum germination percentage (Gmax) of a nuclear series of cytolines, highlighting the natural parental accession in a red dotted frame. White, germination on water; blue, germination on NaCl. Significant differences of germination performance, averaged on the presence of salt, of cytolines compared to their nuclear natural parent are indicated with red stars. Pvalue: ***<0.001<**<0.01<*<0.05 (see Methods S1 for details).

Fig. 3 Contribution of individual accessions to cytonuclear interacting combinations impacting germination traits.

Cytonuclear interacting combinations are plotted according to the contribution of individual accessions. For each accession, the number of combinations involving its cytoplasm (orange) or its nucleus (blue) that significantly impacted dormancy depth at the four PH times (panel a) or germination performance after storage (panel b) are indicated upon the total number of tested cytonuclear interacting combinations (grey).

Fig. 4 Kinetics of loss of germination efficiency during artificial aging.

Controlled deterioration treatment (35°C, 75% relative humidity) was applied for different durations to seeds from cytolines of the Ct-1 (a) and Sha (b) nuclear series. The means of maximum germination percentage from three replicates were plotted. Vertical bars indicate standard deviations from the mean.

The colors of the graphs indicate the cytoplasms of the genotypes: black, Blh-1; blue, Bur-0; orange, Ct-1; green, Cvi-0; purple, Ita-0; brown, Jea; cyan, Oy-0; red, Sha.

Fig. 5 Ratios of Na+ and K+ ion contents in seeds from lines with different cytoplasms associated with the Sha nucleus

Ratios of Na+ and K+ ion contents just after stratification (white boxplots) or after 20h of germination (blue boxplots) measured in seeds on 100 mM NaCl. The Na/K ratios were different between the three genotypes in pairwise contrast tests. On the figure, only p-values for comparisons of the cytolines to Sha are shown, ***<0.001<**<0.01.

Fig. 6 ABA content and sensitivity in seeds from lines of the Sha nucleus series.

a. ABA content of seeds from [Blh-1]Sha, [Ita-0]Sha and Sha genotypes, at the dry stage (left panel), or after stratification followed by 6 hours of germination on water (middle panel) or on salt (right panel). Results of comparisons between the cytolines and Sha in the plotted experiment are shown, **<0.01<*<0.05; ns, not significant.

b. Germination of stratified seeds from [Blh-1]Sha, [Ita-0]Sha and Sha on water (white boxplots) or on 50 µM ABA (blue boxplots). Germination rate was monitored after 4 days. Results of the test for a different germination response to treatment of each cytoline compared to the parent (significant cytoplasm x treatment effect) are indicated; * pval <0.05; ns, not significant (see Methods S1 for details).

Fig. 1

Fig. 2

Fig. 3

a

b

Fig. 4

Fig. 5

Fig. 6

a

dry   H₂O   NaCl

b