1
2
A different state of mind: neural activity related to volitionally up- versus downregulating cortical excitability

Kathy Ruddy[1,2], Joshua Balsters[1,3], Dante Mantini[1,4], Quanying Liu[1,4], Pegah Kassraian-Fard[1], Nadja Enz[1], Ernest Mihelj[1], Bankim Chander[5], Surjo R. Soekadar[5,6], Nicole Wenderoth[1]*.

1. Neural Control of Movement Lab, ETH, Zürich, Switzerland.
2. Institute of Neuroscience, Trinity College Dublin, Ireland
3. Department of Psychology, Royal Holloway, University of London, UK
4. Movement Control and Neuroplasticity Research Group, KU Leuven, Leuven, Belgium
5. Applied Neurotechnology Laboratory, University of Tübingen, Tübingen, Germany
6. Clinical Neurotechnology Laboratory, Clinical Neurotechnology Laboratory, Neuroscience Research Center (NWFZ) & Department of Psychiatry and Psychotherapy, Charité – University Medicine Berlin, Berlin, Germany

*Corresponding Author

Neural Control of Movement Lab,
Department of Health Sciences and Technology,
ETH Zürich
Winterthurerstrasse 190
8057 Zürich
Switzerland

55 **Abstract**
56

To date there exists no reliable method to non-invasively upregulate or downregulate

the state of the resting motor system over a large dynamic range. Here we show that an

operant conditioning paradigm which provides neurofeedback of the size of motor

evoked potentials (MEPs) in response to transcranial magnetic stimulation (TMS),

enables participants to self-modulate their own brain state. Following training,

participants were able to robustly increase (by 83.8%) and decrease (by 30.6%) their

MEP amplitudes. This volitional up-versus down-regulation of corticomotor

excitability caused an increase of late-cortical disinhibition (LCD), a TMS derived

read-out of presynaptic $GABA_B$ disinhibition, which was accompanied by an increase

of gamma and a decrease of alpha oscillations in the trained hemisphere. This approach

paves the way for future investigations into how altered brain state influences motor

neurophysiology and recovery of function in a neurorehabilitation context.

Rhythmic oscillatory brain activity at rest is associated with high versus low

neuronal responsiveness, or 'excitability' of region [1,2]. Measuring these momentary

fluctuations of neural activity via electro- or magnetoencephalography (EEG/MEG)

over human primary motor cortex (M1), it has been demonstrated that frequency,

amplitude and phase of the ongoing oscillation cycle systematically modulate responses

evoked by transcranial magnetic stimulation (TMS) [3-8]. In particular, it has been shown

that corticomotor excitability is significantly higher when the power (amplitude) of

sensorimotor rhythms in the alpha band (8-14 Hz, also called the 'mu'-rhythm), or beta

band (15-30 Hz) are low, or when M1 is stimulated during the trough of the oscillatory

cycle of these rhythms [9]. This concept has inspired neurofeedback interventions

whereby, for example, stroke patients learn to volitionally desynchronize sensorimotor

rhythms with the goal of bringing the sensorimotor system into a more excitable state

as a precursor for enhanced neural plasticity and accelerated recovery [10-12].

Previous research has focussed on interactions between corticomotor

excitability and cortical dynamics at rest, but much less is known about whether it is

possible to *voluntarily* control the excitability of sensorimotor circuits while keeping

motor output and sensory feedback constant. In the case of stroke rehabilitation, this

mechanism may become particularly relevant as patients are unable to move or receive

sensory feedback from the paretic limb. Therefore, interventions that optimally harness

the residual ability to voluntarily and endogenously activate relevant brain circuits in

the days and weeks after the incident, may provide the crucial innervation necessary to

promote re-wiring for functional recovery [13].

It is well known that primates [14,15], and humans [10, 16-19] can gain volitional

control of neural activity by receiving neurofeedback via a brain-computer interface

(BCI). Here, we used a BCI-neurofeedback approach as an effective method for

training participants to both volitionally upregulate and downregulate corticomotor

excitability as reflected by the size of TMS-evoked motor potentials (MEPs), with the

aim to modulate their amplitudes over a much larger dynamic range than observed

during rest. Using this approach enabled us to investigate the neural mechanisms that

underlie volitional up- versus down-regulation of corticospinal excitability in the

motor system and the associated oscillatory signatures. By modulating one neural

marker, i.e. motor evoked potential amplitude, while measuring independent

modalities using EEG, or paired-pulse TMS, this approach allows us to causally relate

voluntary rather than incidental changes of corticomotor excitability to cortical

dynamics.

To achieve this goal we developed a BCI by stimulating the cortex with TMS and

providing neurofeedback of MEP amplitudes (Fig. 1). The feedback was designed such

that participants were rewarded for larger than average MEPs in one condition, and

smaller than average in another condition.

Figure 1. Outline of experimental setup. Each trial of neurofeedback training commenced with a

display of four circles (A), each representing the background EMG in one of the recorded hand muscles

(right FDI, ADM and OP, and left FDI). The circles were red if the root mean squared (rms) EMG at rest

was greater than 7 microvolts. It was essential that all four circles were green for at least 500ms before

the trial could proceed. When this condition was met a fixation cross appeared for a random period (in

order to prevent anticipation of the TMS pulse). During the fixation cross, it was still essential to keep

the background EMG below 7 microvolts in order for a TMS pulse to be delivered. (B) The peak-peak

amplitude of the motor evoked potential (MEP) evoked by the TMS was calculated in real-time and displayed immediately to the participant on screen in the form of a rectangular bar.

(C) Different feedback for UP training and DOWN training. In the UP training If the MEP was greater than the baseline mean, the rectangle was green, with a green tick, a dollar sign to indicate a small financial reward, a display of the current score, and a positive encouraging sound bite was heard. If the MEP did not meet the criterion amplitude, the bar was red, there was no dollar sign, and a negative sound bite was heard. (D) A custom 3D printed 'coil spacer' device was used to prevent direct contact of the TMS coil on the EEG electrodes and allow the pre-TMS EEG period to be recorded artefact free.

In a within-subject cross-over design, participants performed four training sessions with TMS-neurofeedback of MEP amplitudes in order to learn how to up- and downregulate their corticomotor excitability (two 'UP' sessions, two 'DOWN' sessions, order counterbalanced across participants). After the training we characterised the neural underpinnings of these two distinct activity states in detail by conducting a series of multimodal experiments using EEG and paired pulse TMS to profile the associated oscillatory and neurophysiological processes. As it has been proposed that dynamic modulation of neuronal activity is realized via synchronization of high frequency rhythms [20] which are tightly coupled to desynchronizing sensorimotor rhythms [21], we hypothesised that Gamma synchronisation (31-80Hz) and alpha (8-13Hz) /beta (14-30Hz) desynchronization play a critical role in actively determining the state of the motor cortex. Specifically, we hypothesised that following UP neurofeedback training, MEP amplitudes would be increased, and this volitional upregulation would be associated with increased gamma synchronisation and alpha/beta desynchronization. Concurrent reductions in TMS-derived measures of inhibition were also hypothesised during upregulation. The reverse pattern was predicted following DOWN neurofeedback training.

RESULTS

Bidirectional changes in corticospinal excitability were observed in the MEP neurofeedback group but not in a control group.

We first tested whether participants could learn to volitionally increase or decrease (bidirectional) corticomotor excitability when using a motor imagery strategy shaped by neurofeedback of MEP amplitudes. Across two training sessions, we found that

MEP amplitudes increased during UP training (Fig. 2A, orange symbols) and decreased

during DOWN training (Fig. 2A, blue symbols) relative to the baseline measurement

(BS), revealing a significant dissociation over time (*neurofeedback type* x *block*

number interaction during training session 1 [$F(4,115.9)=3.87$, $p=0.006$], session 2

[$F(4,125.0)=3.7$, $p=0.007$] and EEG session [$F(2,70)=6.9$, $p=0.002$], *F tests following*

mixed effects models, $n=15$; see supplementary Fig. 1 for additional analyses). Since

MEP amplitudes are a compound measure of excitability influenced by multiple neural

elements [22], including background muscle activity [23,24], we repeated this analysis using

the root mean squared (rms) background muscle activation (EMG) recorded in the

100ms prior to each TMS pulse. Importantly, this control analysis revealed no such

interactions on any of the sessions, suggesting that the observed modulation was not

driven by changes in peripheral activity of the target muscle, nor any of the additional

3 control muscles (OP, ADM, left FDI) (all p>0.18, see Supplementary Table 2).

Figure 2. MEP amplitudes during neurofeedback. Panel A depicts MEP amplitude in millivolts during

the two types of MEP neurofeedback. UP training is shown in orange and DOWN training in blue, across

all 10 training blocks. Filled triangles labelled 'BS' indicate the baseline measurement block that

occurred at the beginning of that particular session, prior to any neurofeedback. Dotted vertical lines

indicate the separation of the blocks into different 'sessions', which occurred on separate days. Panel B

shows the same data for the control group who received no veridical neurofeedback. Panel C shows the

UP-DOWN difference (in the normalised % change from baseline data) for each block in the

experimental group and the control group. Higher values represent greater deviations between the UP

and DOWN data points and therefore more modulation of MEP amplitude. Thus, these values are

significantly higher in the experimental group than in the control group. # symbols indicate blocks in

which the Cohen's *d* effect size for the difference between the experimental and control group was large-

very large (>0.8). All data are shown as mean ±SEM.

In order to isolate the effect of the neurofeedback, we included a control group who undertook the same protocol, using the same mental imagery strategies, but with feedback that was not contingent on the MEP amplitudes. This group exhibited no systematic changes of corticomotor excitability across training (Fig. 2B) and mixed effects models revealed no significant *neurofeedback type* x *block number* interactions on any of the separate testing sessions in the control group (all $p>0.06$, note that statistics approached significance for the second session because MEPs were randomly higher in the DOWN than in the UP condition; see Supplementary Fig. 1B for further details). Additionally, there were no significant differences in background EMG (All $p>0.09$, Supplementary Table 2). Next we compared the performance of the experimental and the control group, by normalizing MEP amplitudes to baseline (% change) and calculating the difference between UP and DOWN (Fig. 2C). The differences were substantial in the experimental group, who exhibited on average MEP amplitudes twice as large during UP than during DOWN, and differed significantly from the control group where systematic differences were virtually absent (effect of 'Group' [$F(1,25.6)=13.32$, $p=0.001$], *F tests following mixed effects*, *n*=28). The effect sizes (Cohen's *d*) of the between-group differences were small for the first two blocks (<0.5), but consistently increased during training (*d*= 1.27 for block 8), and remained high in the two blocks of the EEG session (*d*>0.97). As the control group were executing the same mental imagery strategies as the experimental group, this comparison demonstrates that veridical TMS neurofeedback was essential for gaining volitional control over corticomotor excitability.

Neurofeedback training effects are retained for at least 6 months

In a follow-up investigation approximately 6 months following initial neurofeedback training, we showed for a subset of the participants (n=11) that they had retained the ability to upregulate and downregulate their MEP amplitude with neurofeedback (Fig.3; significant effect of neurofeedback type (UP vs DOWN) in a retention block carried out with no top-up training (F(1,10)=6.64, p=0.028). Measurements of resting MEP amplitude taken 5 and 10 minutes following the retention block indicated no after-effects (all p>0.2) indicating that subjects could acutely control corticomotor excitability without long-lasting after-effects. Having verified that the ability to modulate brain states had been robustly retained, we next tested whether participants could sustain this performance even when feedback was removed. Performing a

feedback-free block, we found that MEP amplitude was significantly larger in the UP versus DOWN condition (F(1,10)=12.32, p=0.006), indicating that when participants have reached a sufficient level of training they have optimised their mental strategies and no longer require continuous feedback.

Figure 3. Retention, aftereffects and feedback-free measurements. Filled bars represent blocks of neurofeedback, and unfilled bars represent MEPs collected at rest. Panel A shows MEP amplitudes with their preceding resting baseline values subtracted. Values above 0 represent increases relative to baseline, and below 0 represent decreases. State-dependent neurofeedback training feedback effects were still evident in a retention block carried out approximately 6 months following the initial experiment. No aftereffects were observed on resting MEP amplitude 5 and 10 minutes later. In a separate block participants were capable of upregulating and downregulating MEP amplitudes with feedback removed (FB free). MEPs measured from the opposite hemisphere during neurofeedback exhibited a similar pattern of modulation.

In order to measure whether the feedback-induced changes in corticospinal excitability were specific to the hemisphere targeted by the neurofeedback, we used two TMS coils simultaneously and performed one block of 40 trials, where half of the TMS pules were applied to the left hemisphere (i.e. the usual feedback hemisphere), and the other half to the right (i.e. the opposite hemisphere). We found that the same pattern of upregulation and downregulation of MEP amplitudes was observed in the opposite hemisphere, an effect that approached significance (F(1,20)=4.032, p=0.07) but was much smaller than in the neurofeedback hemisphere particularly for the UP condition (UP d = 1.01 for neurofeedback hemi, d=0.27 for opp. hemi, DOWN d= 0.40 for neurofeedback hemi, d=0.35 for opp. hemi, Fig. 3).

Paired pulse TMS investigation of mechanisms

Finally, we investigated which excitatory/inhibitory circuits may have contributed to the changes in corticomotor excitability, using paired pulse TMS measures of three different neurophysiological processes: (i) Short-interval intracortical inhibition (SICI), believed to reflect postsynaptic GABA$_A$ inhibition [25]; (ii) Long interval intra-cortical inhibition (LICI), considered as a marker for postsynaptic GABA$_B$ inhibition; and (iii) late-cortical disinhibition (LCD), which is thought to measure presynaptic GABA$_B$ disinhibition, and manifests as a period in which MEP amplitude returns to and typically overshoots baseline levels, in a time window following LICI (~220ms after a suprathreshold conditioning TMS pulse) [26-28]. In the following analyses we determined the time point (baseline vs during NF) x neurofeedback type (UP, DOWN) interaction and applied FDR correction for multiple testing. Single pulse MEPs collected during these measurement blocks (25% of all trials) revealed significantly larger MEP amplitudes for the UP than the DOWN condition, replicating the findings of the main experiment (Fig. 4A; significant time point x neurofeedback type interaction: $F(1,27.67)=14.36$, $p_{FDR}=0.001$). Surprisingly, there were no significant differences in the magnitude of SICI (% of single pulse MEPs) between the resting baseline data and the SICI MEPs collected during neurofeedback, nor between the UP versus DOWN states (time point x neurofeedback type interaction [$F(1,28.31)=0.08$, $p_{FDR}=0.77$]). The same was true for LICI (time point x neurofeedback type interaction [$F(1,28.90)=0.02$, $p_{FDR}=0.88$]). Thus, circuits controlling postsynaptic inhibition did not seem to be differentially modulated by the UP versus DOWN state. However, for LCD there was a significant time point x neurofeedback type interaction ($F(1,28.35)=12.09$, $p_{FDR}=0.002$, Fig.4B).

Pairwise comparisons revealed that LCD was significantly elevated in the UP condition, when compared to the baseline measurement taken immediately before neurofeedback (Fig. 4B, right panel, MeanDiff=50.9%, df=28.35, p<0.001) and when compared to the equivalent data recorded in the DOWN condition (MeanDiff=56.1%,

297 df=28.76, p<0.001). For the DOWN condition LCD did not differ significantly from
298 baseline (p=0.45).
299

300
Figure 4. Investigation into mechanisms of MEP neurofeedback. The data show paired pulse TMS
measurements taken during neurofeedback blocks to probe distinct neurophysiological processes. In all
subsequent panels, unfilled bars represent baseline MEP amplitudes collected at rest prior to the block.
Panel A Shows that MEP amplitudes from the single pulses (from which neurofeedback was provided)
exhibited the same state-dependent modulation as observed previously. In Panel B MEP amplitudes are
expressed as a percentage of the single pulse MEPs. While expected levels of inhibition were observed
for both SICI and LICI paired pulses, there was no state-dependent modulation. LCD was, however,
significantly increased in the UP condition relative to baseline, and relative to the DOWN condition.
309
310
311
312 *Distinct oscillatory signatures for high versus low corticospinal excitability*
313
As part of the initial training study (see Figure 2A, 'Ses 3' for the behavioural results),
we investigated whether the two different activity states evoked differential cortical
dynamics extracted from EEG recordings which were acquired simultaneously while
TMS was being performed to provide neurofeedback of MEP amplitude. As distinct
functions have been ascribed to 8 different sub-frequency bands across the known range
of brain signals (0.1 - 80Hz), we now probed whether volitional changes in
corticospinal excitability of M1, drives neural activity measured in the delta (0.1-4Hz),
theta (5-7Hz), low alpha (8-10Hz), high alpha (11-13Hz), low beta (14-21Hz), high
beta (22-30Hz), low gamma (31-50Hz) and high gamma (51-80Hz) bands. Using the
portion of EEG data collected in the 1.5 seconds prior to each TMS pulse, we calculated
relative power in the UP and DOWN states for the eight frequency bands of interest.
Figure 5a-f (n=15) shows that UP- versus DOWN regulating corticomotor excitability
caused reduced band-limited power in the theta and alpha band (blue areas in Fig.5b-

d) while gamma power was clearly increased. (red areas in Fig.5e,f). For each participant, we extracted the information for the electrode closest to their individual motor hotspot (Figure 5a shows the different locations across participants) and calculated whether the UP-DOWN difference (Δ relative power %) deviated significantly from 0 (Fig. 5g). Wilcoxon signed rank tests revealed significantly higher power for the UP than DOWN condition in the Delta (p_{FDR}=0.024, d=0.754), Low Gamma (p_{FDR}=0.024, d=0.753) and High Gamma (p_{FDR}=0.016, d=0.712) band and significantly lower power for UP than DOWN in the theta (p_{FDR}=0.003, d=0.947), low alpha (p_{FDR}=0.004, d=0.805) and high alpha band (p_{FDR}=0.007, d=0.714). Although the feedback was lateralised to MEPs from the right limb (left hemisphere motor hotspot), we also quantified the same neural oscillations at the corresponding location in the opposite hemisphere. Here, only the theta rhythm showed significantly lower power for the UP than the DOWN state (p_{FDR}<0.001, d=1.071, see Supplementary Fig. 2).

Figure 5. Neural oscillations associated with the trained brain states. Panels b-f show topographical representations of the relative power (in % of whole spectrum) in the UP condition minus the DOWN condition, for 5 distinct frequency bands (Averaged group data, n=14, 3 other frequency bands shown in Supplementary Fig 3). Red colours indicate regions that demonstrated greater synchronisation in the UP condition. Blue colours indicate greater synchronization in the DOWN condition. The location of the electrode nearest to the TMS hotspot varied between participants but was always within the region indicated in a). Colours are scaled from blue-red by minimum-maximum (range shown to right of each plot). Panel g shows the same data (UP-DOWN) extracted for each participant's hotspot electrode. Values greater than 0 indicate larger amplitude oscillations in the UP condition, and lower than 0 indicate larger oscillations in the DOWN condition. Stars indicate significant deviations from 0 (Wilcoxon Signed Rank tests). Panel h shows group level data for regression analyses performed on MEP amplitudes with

relative power in each frequency band. This included all 120 trials (60 UP, 60 DOWN) collected during
the combined TMS-EEG recording session. The Y axis depicts the slope of the regression model. Stars
indicate significant deviations from 0 (0 would indicate no slope, Wilcoxon Signed Rank test). Individual
regression plots are shown for one representative participant in Supplementary Fig. 4.

Next, we tested whether the amplitude of neural oscillations recorded at the
hotspot at the time of each TMS pulse could predict the amplitude of the resulting
MEPs. For each participant, MEP amplitudes of the 120 trials (60 UP, 60 DOWN) were
entered as the outcome variable in a robust regression model with trial-by-trial relative
power values for each frequency band as predictor variables. Regression slopes (beta
values) for each participant were carried forward into a group level analysis (Fig. 5m),
and Wilcoxon signed rank tests were used to establish whether the slopes were
significantly different from 0 (a 0 slope would indicate no statistical relationship
between predictor and outcome variable). Lower amplitude oscillations in theta
(p_{FDR}=0.024, Fig.5h), low alpha (p_{FDR}<0.001) and high alpha (p_{FDR}=0.002) were
predictive of larger MEP amplitudes, and higher amplitude oscillations in low gamma
(p_{FDR}=0.020) and high gamma (p_{FDR}=0.020) were significant predictors of larger MEP
amplitudes. In a previous study, it was reported that a strong predictor of cortical
excitability was the low gamma : high alpha ratio [3]. We replicated this finding,
demonstrating that this ratio was a significant predictor of MEP amplitude (p_{FDR}=0.016)
with larger ratios predicting larger MEP amplitudes.

373
EEG data classification

We next tested whether the distinction between the two trained states was large enough
that the individual data trials could be successfully predicted as 'UP' state or 'DOWN'
state, using machine learning based solely on the EEG power values (relative power
data, scaled by 1/f transformation, in the 1.5s prior to TMS) of the 8 frequency bands
of interest. A linear support vector machine (SVM) was applied to the data of each
participant (60 UP 60 DOWN epochs). The SVM has been shown to be particularly
powerful on EEG data, which is noisy and contains many features that are correlated.
This approach additionally allowed us to perform feature selection, to quantify which
EEG features most heavily contributed to the distinction between the two states. Using
only data from the electrode closest to the hotspot (8 rhythms plus
LowGamma:HighAlpha ratio) the SVM was capable of classifying the brain states with
an average accuracy of 81.5% (±5.1%) based on 10-fold cross validation which differed

significantly (p=0.001, n=14) from a null model revealed by permutation testing

(accuracy null model: 49.0% ±13). Additionally, incorporating data from the same

rhythms recorded at the corresponding electrode in the opposite hemisphere increased

this accuracy to 85.1% (±4.6%) across participants (see Supplementary Table 1). Using

feature ranking based on Recursive Feature Elimination (RFE), taking the mode of the

top ranked features across participants revealed that the strongest contribution to the

high classification accuracy of the latter SVM was the High Gamma rhythm in the

hotspot electrode, followed by High Alpha at the hotspot, then the

LowGamma:HighAlpha ratio (for full ranking order see Supplementary Table 1).

398
399
400

.

Discussion

Here we aimed to uncover neural activity evoked by voluntarily facilitating or

suppressing excitability within sensorimotor circuits, while keeping motor output and

sensory feedback constant. We show that using a bidirectional TMS-neurofeedback

approach is critical to gain volitional control over MEP amplitudes, a skill that is

retained for at least 6 months without further training. This voluntary state-setting with

a large dynamic range is causally related to modulating pre-synaptic GABA$_B$ mediated

disinhibition and to a prominent increase of gamma power in sensorimotor cortex for

the UP state which was accompanied by a clear reduction of power in the theta, low

and high alpha bands.

Previous studies have shown that it is possible to gain voluntary control over

activity in the central nervous system if appropriate neurofeedback is embedded in a

reinforcement learning task, with food rewards for animals [14,15] and visually rewarding

stimuli for humans [19,29]. Here we confirm that this approach is also suitable for learning

how corticomotor excitability can be bidirectionally up- or down-regulated. Our

participants were initially familiarized with two motor imagery strategies which are

known to modulate corticospinal excitability in the required manner [29-32]. Learning,

however, indicated by progressively stronger dissociation between the UP and the

DOWN state, only took place when direct low-latency feedback regarding the MEP

amplitude was provided. After training, participants were able to modulate

corticomotor excitability across a large range so that MEP amplitudes were

approximately twice as large during the UP than the DOWN condition. UP training, in

particular, resulted in an 83.8% increase of MEP amplitudes from baseline, while

downregulation of MEP amplitude was possible [eg. see 29] but more difficult (30.6%

decrease from baseline). Once acquired, volitional control of corticomotor excitability

was retained for at least 6 months and could be performed even without online feedback

indicating true, long-term learning [33].

Once participants could control their corticomotor excitability, we uncovered the

electrophysiological underpinnings by applying measurements that were independent

of the feedback modality (single pulse TMS) and investigated whether there were

differences between the UP versus DOWN state. This approach cancelled out the

effects that were common to both mental strategies, isolating the mechanisms

underlying the MEP modulation. This revealed two key novel electrophysiological

findings, involving presynaptic $GABA_B$ disinhibition, and gamma oscillations.

Additionally, the pronounced changes of cortical physiology despite the absence of

EMG activity or changes in sensory input suggests that the increase vs decrease of

corticomotor excitability was -a least partly- of cortical origin rather than mediated by

spinal cord mechanisms.

The UP state was associated with a significant increase of LCD while other

measurements probing inhibitory M1 circuits failed to reveal differential effects for the

UP versus DOWN state. LCD is thought to represent a read-out of the presynaptic self-

inhibition of GABAergic neurons which is thought to be mediated by extrasynaptic

$GABA_B$ auto-receptors[34,35]. This mechanism is hypothesized to result in a net

facilitatory effect as observed during the UP condition in our study. Previously, LCD

was found to be elevated during motor imagery (MI), but this increase relative to rest

was observed irrespective of whether participants imagined voluntarily activating or

relaxing hand muscles [36]. However, this investigation was conducted in a single

session, and did not employ neurofeedback, so MEP modulation by these two

imagination conditions could be expected to be substantially smaller than observed in

our study, particularly for the voluntary relaxation condition which had a similar

excitability state as the rest condition. Thus, it is possible that the clear modulation of

LCD observed here only manifested after neurofeedback training, i.e. when the two

excitability states became clearly distinct. It is important to note here that in our

protocol, LCD was detected only during the UP state and not at rest. In our search

procedure to decide upon the optimum conditioning stimulus (CS) intensities, we

prioritized SICI and LICI, finding a CS intensity that elicited as close to 50% inhibition

of the test MEP as possible. We tested intensities between 106-114% RMT for LICI

(and above or below this if no appropriate inhibition was found), and applied these also

to LCD (such that the only difference between the LCD and LICI protocols was the

ISI). This may have simply been too low to consistently elicit LCD, which appears to

be more robustly evoked at higher intensities [27,37] or during mild contraction [38], and is

not observed in all individuals [37] . It may also be that LCD is more readily observed

using triple pulse procedures in which disinhibition can more directly be measured by

the reduction of SICI following a priming stimulus [35,37,39]. The increase in MEP

amplitude observed in the TMS-feedback induced UP state, may also be related to the

increase in I-wave recruitment that has been reported during LCD [26] and suggests that

further research into this effect is warranted, with parameters more specifically tailored

to investigate LCD.

We observed significant modulation of the alpha and gamma rhythms close to

M1 of the trained hemisphere. Focusing on data from the recording electrode closest to

each individual's hotspot revealed a significant association between low alpha and high

gamma power for the UP versus DOWN state. Trial-by-trial modulation of these

rhythms correlated significantly with MEP amplitude, and a support vector machine

(SVM) classifying the two states based on EEG data ranked the high gamma and high

alpha band as the two top features characterizing the distinction. Our observation of

reciprocal changes in the alpha and gamma band are in line with previous studies using

transcranial as well as intracranial recording methods [1]. The 'pulsed inhibition' theory

suggests that repeated bursts of inhibitory alpha activity serve to temporarily silence

gamma oscillations [1]. Thus, these two rhythms are seen to exhibit a reciprocal

relationship, whereby when alpha is high, gamma is low. In periods of high alpha,

gamma may still burst periodically, but only at the troughs of the oscillation cycle,

meaning that the gamma 'duty cycle' (window for neural processing) is short, and only

brief messages can be sent. By contrast, in periods of low alpha power, the gamma duty

cycle is longer, and more extensive neuronal processing and inter-regional

communication may occur. Our finding of increased gamma activity is also consistent

with previous animal literature, showing that the pharmacological removal of $GABA_B$-

mediated inhibition (by receptor blockage) in rats results in increased gamma

oscillations [40] which have been shown to be largest in M1's layer V [41].

Gamma has often been considered difficult to detect using scalp electrodes

because it is highly localised [42] and may also reflect non-cortical sources when recorded

with EEG [43,44]. However, it is tempting to speculate that, in our experiment, gamma

activity was strongly synchronized as a consequence of the neurofeedback training,

where participants learned to substantially facilitate corticomotor excitability while

keeping EMG activity constant, such that EMG amplitude differed only minimally

between the UP and DOWN conditions. This suggestion is in line with previous

neurofeedback studies that provided direct feedback of gamma activity, showing that

gamma power could be upregulated to a substantial amount which even exceeded

power values observed during movement execution [15,45]. By keeping the visual feedback

for the two conditions identical, we ensured that differences in eye movements between

the UP and DOWN states were minor. As we were particularly interested in gamma

oscillations, we additionally performed all EEG recordings in an electromagnetically

shielded room, using a gel-based electrode system to maximize signal to noise ratio.

Previous studies have taken a correlational approach to investigating the

relationships between brain rhythms and corticomotor excitability. These have shown

that low alpha [4,46] or beta power [47] as well as high gamma power [3] during natural

fluctuations at rest are associated with larger MEP amplitudes. We confirm and extend

these results by introducing causality to this relationship for the first time, showing that

experimentally driving excitability into two distinct states causes specific patterns of

neural dynamics in the volitionally controlled cortical area.

While changes in alpha and gamma were specific to the hemisphere from which

feedback was provided, theta showed a bilateral pattern of modulation, being higher in

the DOWN than the UP state in motor areas in both hemispheres. While mid-frontal

theta activity has been linked to error monitoring [48] the role of lateralized theta activity

close to the sensorimotor hotspot electrode and its symmetric counterpart is less clear.

Slower rhythms exert effects over larger distances, and are thought to be involved in

long-range communication [42]. A similar pattern of upregulation and downregulation

was observed in the homologous muscle in the opposite limb, albeit weaker and not

statistically significant. This is likely a reflection of the extensive transcallosal

structural connectivity and functional coupling of homologous regions of the cortical

motor network [49-51]. It is tempting to speculate that the bilateral theta activity observed

in the current study served to regulate the inhibition/facilitation of functional coupling

or 'spillover' of activation from motor areas in the target hemisphere to their

homologous counterparts.

Surprisingly we did not observe differential modulation of the Beta band, which

is the predominant oscillatory frequency in sensorimotor cortical regions [52,53]. It

typically desynchronizes (together with alpha) during motor execution and motor

imagery [54-57] and has been associated with corticomotor excitability at rest [3]. As our

results represent the direct contrast between the UP and DOWN states, the lack of Beta

involvement may firstly be due to the fact that both conditions involved a mental

strategy targeted at the sensorimotor system and, secondly, that no temporal structure

was imposed so that we could not perform analyses which are, for example, time-locked

to the potential onset of these mental strategies. However, our data further confirm that

the two 'inhibitory' rhythms alpha and beta might serve different functions in selecting

and activating the appropriate sensorimotor representations [58].

Here we present an innovative approach to voluntarily and bidirectionally

change the state of the motor cortex, by directly targeting MEP amplitudes in a

neurofeedback paradigm. This method provided a unique opportunity to reveal the

oscillatory and neurochemical underpinnings of the two distinct trained brain states,

using concurrent TMS EEG measurements, and mechanistic follow-up investigations

using paired-pulse TMS. The results comparing UP and DOWN states indicate that

voluntary upregulation of corticomotor excitability causes increased presynaptic

GABA$_B$-mediated disinhibition, elevated neural oscillations in the gamma frequency
range, and reduced alpha and theta rhythms.

This paves the way for new technologies that allow the user to regulate aspects
of their own brain function in order to reach desired states that are, for example,
associated with enhanced motor performance. In the context of stroke rehabilitation,
training volitional modulation of corticomotor excitability may hold promise as a
rehabilitative therapy early after stroke, i.e. when patients are deprived of rehabilitation
training because they are unable to execute overt movements with the impaired upper
limb. As it is known that LCD is recruited during actual movement [28,59,60], the elevated
LCD we observed in the UP condition may reflect that the neurofeedback had engaged
similar mechanisms to those involved in movement execution, using only voluntary
endogenous processes. Furthermore, as pathological hyperexcitability of the non-
damaged hemisphere has been hypothesized to limit recovery in some patients [61], the
TMS-neurofeedback protocol can be individually tailored either to upregulate the
damaged hemisphere, down-regulate the intact hemisphere, or a combination of both,
depending on the patient's specific needs.

579 **Materials and Methods**
580
Participants

Fifteen healthy volunteers (age 23± 3.14 s.d, 7 female) participated in the
experimental group. An additional thirteen participants (age 25± 3.06 s.d, 3 female)
formed a control group. All participants were right handed according to the Edinburgh
Handedness Inventory [62], and gave informed consent to procedures. The experiments
were approved by the Kantonale Ethikkommission Zürich, and were conducted in
accordance with the Declaration of Helsinki (1964).

TMS-based neurofeedback

Participants undertook five sessions of TMS-based neurofeedback, on separate

days. The first four days comprised of neurofeedback training, and on the fifth day

neurofeedback was performed with simultaneous electroencephalography (EEG)

recording to investigate state specific neural dynamics. On two of the training days

neurofeedback was adjusted so that a rewarding visual stimulus was displayed when

MEPs were larger than baseline (the 'UP' condition) and on the other two days, the

rewarding stimulus was displayed when MEPs were smaller than baseline (the

'DOWN' condition). On each of the training days, 4 separate blocks of neurofeedback

were preformed, each comprising of 30 individual MEP feedback trials (total 120 trials

per day). The format of individual trials and feedback is described in more detail below.

Baseline corticospinal excitability was measured on each day prior to training (20

MEPs) and post-measurements were taken during the rest periods between each of the

4 blocks (12 MEPs per measurement).

Subjects sat in a comfortable chair with both arms and legs resting in a neutral

position supported by foam pillows. Surface electromyography (EMG, Trigno

Wireless; Delsys) was recorded from right First Dorsal Interosseous (FDI), Abductor

Digiti Minimi (ADM), Opponens Pollicis (OP), and left FDI. EMG data were sampled

at 2000Hz (National Instruments, Austin, Texas), amplified and stored on a PC for off-

line analysis.

TMS was performed with a figure-of-eight coil (internal coil diameter 50mm)

connected to a Magstim 200 stimulator (Magstim, Whitland, UK). The coil was held

on the left hemisphere over the 'hotspot' of the right FDI at the location with the largest

and most consistent MEPs, and with the optimal orientation for evoking a descending

volley in the corticospinal tract (approximately 45 degrees from the sagittal plane in

order to induce posterior-anterior current flow). Once the hotspot was established, the

lowest stimulation intensity at which MEPs with peak-to-peak amplitude of

approximately 50μV were evoked in at least 5 of 10 consecutive trials was taken as

Resting Motor Threshold (RMT).

The stimulation intensity used to evoke MEPs during the experiment was chosen

using the following procedure in order to obtain a baseline MEP amplitude that was

50% of the participant's maximum. A recruitment curve [eg. 63] was performed at the

beginning of the first experimental session, whereby 6 TMS pulses were applied at 10

different intensities relative to RMT (90%, 100%, 110%, 120%, 130%, 140%, 150%,

160%, 180%, 190%) in a randomized order. MEP amplitude at each intensity was

plotted to determine the point on the curve at which plateau occurs and the MEPs do

not continue to increase. Maximal MEP amplitude was recorded, and the intensity

required to evoke 50% of this amplitude was used for all subsequent testing. With this

approach, there was scope for MEP amplitude to both increase and decrease to similar

extents from this 'intermediate' value. Post-hoc analyses revealed that this procedure

resulted in an average stimulation intensity corresponding to 130% RMT. Immediately

following this procedure and prior to the first block of neurofeedback, 20 MEPs were

collected at rest at the chosen intensity to determine 'baseline' corticospinal

excitability. The mean MEP amplitude at baseline was recorded and used during

neurofeedback to establish the criterion amplitude that determined whether participants

received either positive or negative feedback.

Format of neurofeedback

Neurofeedback was performed using custom written MATLAB software.

Participants kept eyes open with attention directed to a monitor in front of them. They

were instructed to relax their limbs and avoid tensing any muscles throughout the

experiment. In order to ensure that MEP amplitude could not be influenced by

background muscle activation, the root mean square (rms) of the EMG signal for each

muscle for the previous 100ms of data was calculated and displayed in real-time on

screen at the beginning of each trial in the form of four coloured 'traffic lights',

representing each muscle (Fig. 1A). If the background EMG in a muscle exceeded 7μV,

the corresponding light turned red. Participants were instructed that a trial could not

begin unless all four lights were green (all muscles relaxed below 7μV) for at least a

continuous 500ms period. When a trial commenced the traffic lights disappeared, but

background EMG continued to be monitored and the trial was automatically paused if

any muscle exceeded the threshold. At the beginning of each trial a fixation cross

appeared in the center of the screen. After a variable period of time (between 5.5 - 8.5

seconds, or longer if muscle activation delayed the trial) a TMS pulse was fired. The

MEP amplitude for the target muscle (right FDI) was immediately measured and

displayed to the participant on screen within 500ms. The display consisted of a vertical

bar indicating MEP amplitude relative to a horizontal line in the middle of the screen representing the mean recorded at baseline (Fig. 1B). In 'UP' sessions if the MEP was larger than the criterion amplitude, the bar was shown as green with a tick beside it, a positive soundbyte was heard, and a number adjacent to a dollar sign incremented to indicate that a small financial reward had been gained. If the MEP was smaller than the criterion amplitude, the bar was red with a cross beside it, a negative soundbyte was heard, and no financial reward was shown. The reverse was true in the 'DOWN' sessions (Fig. 1C). The feedback remained on screen for 4 seconds, before being replaced by the traffic lights display preceding the next trial. Participants were instructed to attend to the feedback and that the goal was to increase (or decrease) the size of the MEP represented by the bar. Prior to the experiment participants read an instruction sheet explaining the procedures above and providing recommended mental strategies that were reported in previous literature in which corticospinal excitability was downregulated [29] and upregulated [31] by motor imagery (Specific task instructions are provided in Supplementary Material). Initially the criterion amplitude corresponded to the baseline MEP measure. After each block of 30 MEPs, performance was quantified and the task difficulty was adjusted if necessary. If the success rate was >70% difficulty was increased by raising (or lowering in the DOWN condition) the criterion MEP amplitude that needed to be reached by 10% in order for the positive reward to be presented. If performance was > 90%, this was adjusted by 20%.

EEG session

On the fifth day neurofeedback was provided during simultaneous EEG recording. The participant's TMS hotspot was determined and marked on the scalp prior to EEG capping. EEG signals were recorded using a 64 channel gel-based TMS-compatible cap (Electrical Geodesics Inc., Oregon, USA), and the channel closest to the TMS hotspot was noted. EEG data were amplified and sampled at 1000hz. In order to minimize artefacts associated with the direct contact of the TMS coil resting on the electrodes of the EEG cap, we designed and 3D-printed a custom plastic 'coil spacer' device [64], which has four wide legs positioned to provide a platform distributing the weight of the TMS coil, so that it hovers over the electrodes without contact (Fig. 1D). This allowed quality recordings to be obtained even from the channel of interest closest to the participant's 'hotspot'. The participants RMT was established while wearing the

EEG cap with TMS coil spacer, and the same % above threshold that was used for all

previous sessions was applied for neurofeedback. Impedances were monitored

throughout and maintained below 50kΩ.

Baseline corticospinal excitability was measured in the same fashion as for the

first four sessions, followed by two blocks of neurofeedback (UP or DOWN,

counterbalanced) with brief (12 MEP) post measurements following each. After the

final post measurement, a 15 minute rest break was scheduled for the participant.

Following this, the procedure was repeated and baseline excitability was measured

again, followed by two blocks of either UP or DOWN neurofeedback (whichever was

not performed in the first half of the session). At the end of this session participants

were debriefed.

Control group

Participants were blinded as to whether they were allocated to the experimental

or control group. The control group experienced identical conditions to the

experimental group, with the exception that direct neurofeedback was not provided.

The visual feedback bar demonstrating MEP amplitude was always the same height

(reaching the 'mean' horizontal line). 'Positive' feedback/rewards were presented in

the same proportion as in the experimental group (66% of all trials - calculated upon

completion of experimental group), but at a fixed and predicable rate in order to prevent

the development of illusory correlations. Participants were instructed to attend to the

visual feedback on screen, and that rewards would occur at a fixed rate. Aside from

this, they were otherwise given identical instructions as the participants in the

experimental group- i.e. the same recommended mental strategies were provided on

control 'UP' and 'DOWN' blocks.

717
718   **Data processing and analysis**

719
720   *MEP data*

721
EMG data from all four hand muscles were band-pass filtered (30–800 Hz),

separately for the portion of data containing the 100ms of 'pre-TMS' background EMG,

and for the portion of EMG containing the MEP, in order to prevent smearing of the

MEP into the background EMG data chunk. The root mean squared (rms) of the

background EMG was calculated, and peak-peak MEP amplitude was measured.

Trimming (removal) of the maximum and minimum MEP in each block was performed

in order to screen out extreme values. MEP amplitude is known to be modulated by

EMG background activation [23,24]. Therefore, the rms pre-stimulus EMG recordings

were used to assess the presence of unwanted background EMG activity in the period

110 to 10ms preceding the magnetic pulse. MEPs preceded by background EMG higher

than 0.01mV were excluded. For each subject and over all trials we calculated the mean

and standard deviation of the background EMG. MEPs that occurred when the

background EMG value exceeded 2.5 standard deviations above the mean, and MEPs

with a peak-to-peak amplitude which exceeded $Q3 + 1.5 \times (Q3 - Q1)$ were removed

from further analysis, with Q1 denoting the first quartile and Q3 the third quartile

computed over the whole set of trials for each subject.

Inferential statistics were computed using Mixed Effects Models in SPSS (Version

16.0, SPSS Inc. Chicago, US), as they account for covariances between related data

samples in repeated measures designs, and have greater flexibility for modeling effects

over time than traditional ANOVA approaches [65]. Fitting of the mixed effects models

employed restricted maximum likelihood estimation (REML) and a compound

symmetry covariance matrix. Model fit indices (Akaike Information Criterion and

Schwarz Bayesian Criterion) were considered prior to choosing the covariance matrix

and model type. Fixed effects were *neurofeedback type* (UP or DOWN) and *block*

number (1-10). The influence of each of the fixed effects on the model was estimated

using *F* tests. In all models *subject* was designated as a random effect with random

intercepts.

The criterion alpha value was set to 0.05 for all inferential tests. In cases where

multiple comparisons were performed, *p* values were false discovery rate (FDR)

corrected.

EEG data

Signals from all 64 channels were first epoched to extract only the data on each

trial from the 4 seconds before the TMS pulse. This was to remove the substantial

artefacts that arise during the magnetic pulses, prior to conducting any filtering or further processing. These separate chunks of unpolluted data were then concatenated into one continuous epoch, and highpass filtered at 1Hz, prior to conducting an independent components analysis (ICA). Independent components were visualized and those containing artefacts arising from eye movements, facial EMG, cardiac signals, bad channels or other non-brain activity related signals were removed.

The cleaned data were average-referenced, and re-epoched into chunks of data containing only the 1.5s on each trial prior to the TMS pulse (ie. to capture the ongoing oscillatory activity at the instance in which the TMS occurred, while the fixation cross was on screen and the 'traffic lights' had disappeared).

A power spectrum was computed (Welch periodogram/FFT) for each single epoch and the mean power (and relative power) in each of the relevant bandwidths were extracted (delta (0.1-4Hz), theta (5-7Hz), low alpha (8-10Hz), high alpha (11-13Hz), low beta (14-21Hz), high beta (22-30Hz), low gamma (31-50Hz) and high gamma (51-80Hz). Power values were computed separately for UP and DOWN trials, and non-parametric Wilcoxon signed rank tests (with FDR correction) were used to compare neural oscillations in these two states.

We also analysed whether trial-by-trial variation of EEG data was associated with trial-by-trial variation of MEP amplitudes. Therefore, relative power in each bandwidth for each epoch was entered into a multiple regression model with MEP amplitudes measured in the muscle from which neurofeedback was provided (right FDI). The beta (slope) values resulting from each regression model for each participant were forwarded into a group-level analysis.

Classification of distinct brain states

Individual epochs of EEG data (60 UP 60 DOWN) were classified by a linear support vector machine (SVM, 10-fold cross validation), to test separately for each participant whether the epochs could be successfully predicted as 'UP' state or 'DOWN' state based solely on the power values (scaled by using 1/f transformed relative power) of the 4 frequency bands of interest. The SVM was chosen as it is known to perform particularly well in BCI settings using EEG data which is noisy and has features that are correlated. In order to validate the results the same procedure was repeated with randomly permuted labels, and this null model was statistically compared to the model

794 with true labels (C=1). Feature selection was conducted using feature ranking based on

795 Recursive Feature Elimination [66].

796
797
798 ***Follow-up experiment 6 months later***
799
A sub-set of 11 participants from the experimental group returned approximately 6

months later to participate in a follow-up experiment probing retention and mechanisms

underlying the two distinct states. This was conducted over a further 4 days of testing.

On one day, retention, aftereffects, and excitability in the opposite 'untrained'

hemisphere were tested for the 'UP' condition. On another, neurophysiolocial

mechanisms were probed using paired pulse TMS. These two days were repeated for

the 'DOWN' condition, and the order of these sessions was counterbalanced. We

additionally tested whether trained participants were able to upregulate and

downregulate when feedback was temporarily removed.

Retention testing & aftereffects measurement

After a 6-month break and no top-up training, participants were tested with one block

of TMS-neurofeedback (20 MEPs) in order to assess retention of learning. All other

procedures were identical to those carried out in the main experiment.

Following this block, 12 MEPs were collected at rest after 5 and 10 minutes.

Excitability in the opposite hemisphere

During one block, two TMS coils were used, placed over the right and left motor

hotspots (as described previously). This block contained 40 trials, 20 of which were

normal TMS neurofeedback trials. The other 20 were trials where TMS was applied to

the opposite hemisphere, rather than to the hemisphere that was the target for

neurofeedback. No feedback was given in these trials. The presentation of left and right

hemisphere TMS pulses was randomized.

Feedback- free measurements

We additionally tested whether trained participants were able to upregulate and

downregulate when feedback was temporarily removed. In this feedback-free block,

the timing of trials and participant instructions were identical to normal neurofeedback

blocks, but in place of the usual feedback bar showing MEP amplitude, the white

fixation cross simply turned red during this period. The onset of trials was still

contingent on muscles being completely relaxed, and the traffic lights display still

preceded every trial.

Paired pulse TMS measurements

On separate days (one 'UP' one 'DOWN) from the measurements described above, we

performed three additional blocks of TMS neurofeedback (24 trials per block x 3 = 72

total trials), in which just 25% of trials were standard single pulse TMS-neurofeedback

trials, with the usual feedback. The remaining trials contained paired pulses in place of

the usual single pulse TMS. For all paired pulse measurements, the test stimulus

intensity was identical to that which had been chosen for the TMS neurofeedback (ie.

that produced MEPs that were 50% of the maximum on the recruitment curve). On 25%

of trials Short Interval Intracortical Inhibition (SICI) was measured. This was with a

conditioning stimulus intensity that was chosen using a personalized search procedure

testing intensities ranging from 50%-90% RMT, to achieve as close to 50% inhibition

as possible, and an inter-stimulus interval of 1.97ms [67]. The reduction in the size of the

test MEP is believed to represent postsynaptic $GABA_A$ inhibition [25]. On 25% of trials

Long Interval Intracortical Inhibition (LICI) was measured. This was with two

suprathreshold pulses, with the conditioning stimulus intensity chosen using a search

procedure between 106-114% RMT, and an inter-stimulus interval of 100ms [27]. This is

believed to reflect postsynaptic $GABA_B$ inhibition [68]. On the remaining 25% of trials,

Late Cortical Disinhibition (LCD) was tested. This was with the exact same pulse

intensities as used for LICI, but with a 220ms inter-stimulus interval [27], and is thought

to measure presynaptic $GABA_B$ disinhibition [26-28]. The order of presentation of paired

pulses and single pulses was randomized.

Baseline measurements were taken at rest with each of these three paired-pulse TMS

protocols, prior to the beginning of neurofeedback blocks (20 of each type of paired

pulse measurement, and 20 single pulse MEPs).

1

REFERENCES

1.   Jensen, O. & Mazaheri, A. Shaping Functional Architecture by
868   Oscillatory Alpha Activity: Gating by Inhibition. *Frontiers in human*
869   *neuroscience* **4,** (2010).
870   2.   Jensen, O. *et al.* Using Brain–Computer Interfaces and Brain-State
871   Dependent Stimulation as Tools in Cognitive Neuroscience. *Front*
872   *Psychol* **2,** (2011).
873   3.   Zarkowski, P., Shin, C. J., Dang, T., Russo, J. & Avery, D. EEG and the
874   Variance of Motor Evoked Potential Amplitude. *Clinical EEG and*
875   *Neuroscience* **37** *(3)* 247-251 (2006).
876   4.   Sauseng, P., Klimesch, W., Gerloff, C. & Hummel, F. C. Spontaneous
877   locally restricted EEG alpha activity determines cortical excitability in
878   the motor cortex. *Neuropsychologia* **47,** 284–288 (2009).
879   5.   Schaworonkow, N., Triesch, J., Ziemann, U. & Zrenner, C. EEG-
880   triggered TMS reveals stronger brain state-dependent modulation of
881   motor evoked potentials at weaker stimulation intensities. *bioRxiv*
882   251363 (2018). doi:10.1101/251363
883   6.   Kelly, S. P., Gomez-Ramirez, M. & Foxe, J. J. The strength of
884   anticipatory spatial biasing predicts target discrimination at attended
885   locations: a high-density EEG study. *Eur. J. Neurosci.* **30,** 2224–2234
886   (2009).
887   7.   Mazaheri, A., Nieuwenhuis, I. L. C., van Dijk, H. & Jensen, O.
888   Prestimulus alpha and mu activity predicts failure to inhibit motor
889   responses. *Hum Brain Mapp* **30,** 1791–1800 (2009).
890   8.   Schubert, R., Blankenburg, F., Lemm, S., Villringer, A. & Curio, G. Now
891   you feel it--now you don't: ERP correlates of somatosensory awareness.
892   *Psychophysiology* **43,** 31–40 (2006).
893   9.   Zaehle, T., Rach, S. & Herrmann, C. S. Transcranial alternating current
894   stimulation enhances individual alpha activity in human EEG. *PLoS ONE*
895   **5,** e13766 (2010).
896   10.   Buch, E. *et al.* Think to move: a neuromagnetic brain-computer interface
897   (BCI) system for chronic stroke. *Stroke* **39,** 910–917 (2008).
898   11.   Caria, A. *et al.* Chronic stroke recovery after combined BCI training and
899   physiotherapy: A case report. *Psychophysiology* **48,** 578–582 (2010).
900   12.   Soekadar, S. R., Birbaumer, N., Slutzky, M. W. & Cohen, L. G. Brain–
901   machine interfaces in neurorehabilitation of stroke. *Neurobiology of*
902   *Disease* **83,** 172–179 (2015).
903   13.   Maulden, S. A., Gassaway, J., Horn, S. D., Smout, R. J. & DeJong, G.
904   Timing of Initiation of Rehabilitation After Stroke. *Arch Phys Med*
905   *Rehabil* **86,** 34–40 (2005).
906   14.   Fetz, E. E. Volitional Control of Cortical Oscillations and Synchrony.
907   *Neuron* **77,** 216–218 (2013).

15.   Engelhard, B., Ozeri, N., Israel, Z., Bergman, H. & Vaadia, E. Inducing
Gamma Oscillations and Precise Spike Synchrony by Operant
Conditioning via Brain-Machine Interface. *Neuron* (2013).

16.   Mellinger, J. *et al.* An MEG-based brain-computer interface (BCI).
Neuroimage **36,** 581–593 (2007).

17.   Soekadar, S. R., Witkowski, M., Birbaumer, N. & Cohen, L. G.
Enhancing Hebbian Learning to Control Brain Oscillatory Activity.
Cerebral Cortex **83**, 172-179 (2014). doi:10.1093/cercor/bhu043

18.   Sitaram, R. *et al.* Closed-loop brain training: the science of
neurofeedback. *Nat Rev Neurosci* **18,** 86–100 (2017).

19.   Thompson, A. K., Chen, X. Y. & Wolpaw, J. R. Acquisition of a Simple
Motor Skill: Task-Dependent Adaptation Plus Long-Term Change in the
Human Soleus H-Reflex. *J. Neurosci.* **29,** 5784–5792 (2009).

20.   Fries, P. A mechanism for cognitive dynamics: neuronal communication
through neuronal coherence. *Trends Cogn. Sci. (Regul. Ed.)* **9,** 474–480
(2005).

21.   Grosse-Wentrup, M., Schölkopf, B. & Hill, J. Causal influence of gamma
oscillations on the sensorimotor rhythm. *Neuroimage* **56,** 837–842
(2011).

22.   Carson, R. G., Ruddy, K. L. & McNickle, E. *Progress in Motor Control;*
Theories and Translations (eds Laczko, J. & Latash, M.L) ISBN 978-3-
319-47312-3 (Springer International Publishing AG).

23.   Hess, C. W., Mills, K. R. & Murray, N. M. Magnetic stimulation of the
human brain: facilitation of motor responses by voluntary contraction of
ipsilateral and contralateral muscles with additional observations on an
amputee. *Neurosci Lett* **71,** 235–240 (1986).

24.   Devanne, H., Lavoie, B. A. & Capaday, C. Input-output properties and
gain changes in the human corticospinal pathway. *Exp Brain Res* **114,**
329–338 (1997).

25.   Fisher, R. J., Nakamura, Y., Bestmann, S., Rothwell, J. C. & Bostock, H.
Two phases of intracortical inhibition revealed by transcranial magnetic
threshold tracking. *Exp Brain Res* **143,** 240–248 (2002).

26.   Cash, R. F. H., Ziemann, U. & Thickbroom, G. W. Inhibitory and
disinhibitory effects on I-wave facilitation in motor cortex. *Journal of*
Neurophysiology **105,** 100–106 (2011).

27.   Cash, R. F. H., Ziemann, U., Murray, K. & Thickbroom, G. W. Late
cortical disinhibition in human motor cortex: a triple-pulse transcranial
magnetic stimulation study. *Journal of Neurophysiology* **103,** 511–518
(2010).

28.   Caux-Dedeystère, A., Derambure, P. & Devanne, H. Late cortical
disinhibition in relaxed versus active hand muscles. *Neuroscience* **298,**
52–62 (2015).

29.   Majid, D. S. A., Lewis, C. & Aron, A. R. Training voluntary motor
suppression with real-time feedback of motor evoked potentials. *Journal*
of Neurophysiology **113,** 3446–3452 (2015).

30.   Stinear, C. M., Byblow, W. D., Steyvers, M., Levin, O. & Swinnen, S. P.
Kinesthetic, but not visual, motor imagery modulates corticomotor
excitability. *Exp Brain Res* **168,** 157–164 (2005).

31.   Izumi, S. *et al.* Facilitatory effect of thinking about movement on motor-
evoked potentials to transcranial magnetic stimulation of the brain.

959 *American Journal of Physical Medicine & Rehabilitation* **74,** 207–213 (1995).

32. Fadiga, L. *et al.* Corticospinal excitability is specifically modulated by 961 962 motor imagery: a magnetic stimulation study. *Neuropsychologia* **37,** 147–158 (1998).

33. Magill, R. *Motor Learning and Control: Concepts and Applications*. 964 (McGraw-Hill Education, 2010).

34. Mott, D. D. & Lewis, D. V. Facilitation of the induction of long-term 966 potentiation by GABAB receptors. *Science* **252,** 1718–1720 (1991).

35. Sanger, T. D., Garg, R. R., Chen, R. Interactions between two different 968 inhibitory systems in the human motor cortex. *J Physiol* 530, 307-317 969 (2001). doi:10.1111/j.1469-7793.2001.0307l.x

36. Chong, B. W. X. & Stinear, C. M. Modulation of motor cortex inhibition 971 during motor imagery. *Journal of Neurophysiology* **117,** 1776–1784 972 (2017).

37. Cash, R. F. H., Murakami, T., Chen, R., Thickbroom, G. W. & Ziemann, 974 U. Augmenting Plasticity Induction in Human Motor Cortex by 975 Disinhibition Stimulation. *Cereb. Cortex* **26,** 58–69 (2016).

38. Caux Dedeystère, A. *et al.* Task-dependent changes in late inhibitory and 977 disinhibitory actions within the primary motor cortex in humans. 978 *European Journal of Neuroscience* **39,** 1485–1490 (2014).

39. Ni, Z., Gunraj, C. & Chen, R. Short interval intracortical inhibition and 980 facilitation during the silent period in human. *The Journal of Physiology* 981 **583,** 971–982 (2007).

40. Leung, L. S. & Shen, B. GABAB receptor blockade enhances theta and 983 gamma rhythms in the hippocampus of behaving rats. *Hippocampus* **17,** 984 281–291 (2007).

41. Johnson, N. W. *et al.* Phase-amplitude coupled persistent theta and 986 gamma oscillations in rat primary motor cortex in vitro. 987 *Neuropharmacology* **119,** 141–156 (2017).

42. Buzsaki, G. *Rhythms of the Brain*. (2006) ISBN-13-978-0-19-530106-9 989 (Oxford University Press, USA).

43. Yuval-Greenberg, S., Tomer, O., Keren, A. S., Nelken, I. & Deouell, L. 991 Y. Transient induced gamma-band response in EEG as a manifestation of 992 miniature saccades. *Neuron* **58,** 429–441 (2008).

44. Whitham, E. M. *et al.* Scalp electrical recording during paralysis: 994 Quantitative evidence that EEG frequencies above 20 Hz are 995 contaminated by EMG. *Clinical Neurophysiology* **118,** 1877–1888 996 (2007).

45. Miller, K. J. *et al.* Cortical activity during motor execution, motor 998 imagery, and imagery-based online feedback. *Proceedings of the* 999 *National Academy of Sciences* **107,** 4430–4435 (2010).

46. Zrenner, C., Desideri, D., Belardinelli, P. & Ziemann, U. Real-time EEG-1001 defined excitability states determine efficacy of TMS-induced plasticity 1002 in human motor cortex. *Brain Stimul* **11,** 374–389 (2018).

47. Mäki, H. & Ilmoniemi, R. J. EEG oscillations and magnetically evoked 1004 motor potentials reflect motor system excitability in overlapping neuronal 1005 populations. *Clinical Neurophysiology* **121,** 492–501 (2010).

48. Pezzetta, R., Nicolardi, V., Tidoni, E. & Aglioti, S. M. Error, rather than 1007 its probability, elicits specific electrocortical signatures: a combined

EEG-immersive virtual reality study of action observation. *Journal of Neurophysiology* jn.00130.2018 (2018). doi:10.1152/jn.00130.2018

49. Ruddy, K. L., Leemans, A. & Carson, R. G. Transcallosal connectivity of the human cortical motor network. *Brain Struct Funct* 222, 1243-1252 (2017). doi:DOI: 10.1007/s00429-016-1274-1

50. Ruddy, K. L., Leemans, A., Woolley, D. G., Wenderoth, N. & Carson, R. G. Structural and Functional Cortical Connectivity Mediating Cross Education of Motor Function. *Journal of Neuroscience* **37,** 2555–2564 (2017).

51. Ruddy, K. L. Directionality of interhemispheric communication. *Brain Structure and Function*. DOI 10.1107/s00429-017-1557-1 (2017).

52. Baker, S. N., Olivier, E. & Lemon, R. N. Coherent oscillations in monkey motor cortex and hand muscle EMG show task-dependent modulation. *Journal of Physiology* **501**(1), 225-241 (1997).

53. Murthy, V. N. & Fetz, E. E. Oscillatory activity in sensorimotor cortex of awake monkeys: synchronization of local field potentials and relation to behavior. *Journal of Neurophysiology* **76,** 3949–3967 (1996).

54. Ramos-Murguialday, A. & Birbaumer, N. Brain oscillatory signatures of motor tasks. *Journal of Neurophysiology* **113,** 3663–3682 (2015).

55. Pfurtscheller, G. & Neuper, C. Motor imagery activates primary sensorimotor area in humans. *Neurosci Lett* **239,** 65–68 (1997).

56. Pfurtscheller, G. & Berghold, A. Patterns of cortical activation during planning of voluntary movement. *Electroencephalogr Clin Neurophysiol* **72,** 250–258 (1989).

57. Alegre, M. *et al.* Alpha and beta oscillatory activity during a sequence of two movements. *Clinical Neurophysiology* **115,** 124–130 (2004).

58. Brinkman, L., Stolk, A., Dijkerman, H. C., de Lange, F. P. & Toni, I. Distinct Roles for Alpha- and Beta-Band Oscillations during Mental Simulation of Goal-Directed Actions. *Journal of Neuroscience* **34,** 14783–14792 (2014).

59. Hammond, G. & Vallence, A.-M. Modulation of long-interval intracortical inhibition and the silent period by voluntary contraction. *Brain Res* **1158,** 63–70 (2007).

60. Opie, G. M., Ridding, M. C. & Semmler, J. G. Task-related changes in intracortical inhibition assessed with paired- and triple-pulse transcranial magnetic stimulation. *Journal of Neurophysiology* **113,** 1470–1479 (2015).

61. Murase, N., Duque, J., Mazzocchio, R. & Cohen, L. G. Influence of interhemispheric interactions on motor function in chronic stroke. *Ann. Neurol.* **55,** 400–409 (2004).

62. Oldfield, R. C. The assessment and analysis of handedness: The Edinburgh Inventory. *Neuropsychologia* **9,** 97–113 (1971).

63. Carson, R. G. *et al.* Characterizing changes in the excitability of corticospinal projections to proximal muscles of the upper limb. *Brain Stimul* **6,** 760–768 (2013).

64. Ruddy, K. L. *et al.* Improving the quality of combined EEG-TMS neural recordings: Introducing the coil spacer. *Journal of Neuroscience Methods* **294,** 34–39 (2018).

65. Gueorguieva, R. & Krystal, J. H. Move over ANOVA: progress in analyzing repeated-measures data and its reflection in papers published in

the Archives of General Psychiatry. *Arch. Gen. Psychiatry* **61,** 310–317
(2004).
66.   Guyon, I., Weston, J., Barnhill, S. & Vapnik, V. Gene Selection for
Cancer Classification using Support Vector Machines. *Machine Learning*
46, 389–422 (2002).
67.   Peurala, S. H., Müller-Dahlhaus, J. F. M., Arai, N. & Ziemann, U.
Interference of short-interval intracortical inhibition (SICI) and short-
interval intracortical facilitation (SICF). *Clinical Neurophysiology* **119,**
2291–2297 (2008).
68.   McDonnell, M. N., Orekhov, Y. & Ziemann, U. The role of $GABA_B$
receptors in intracortical inhibition in the human motor cortex. *Exp Brain*
Res **173,** 86–93

1070
1071

1072 **Acknowledgements**

1073 This work was supported by grant number 320030_175616 from the Swiss National
1074 Science Foundation. KLR is supported by a research fellowship from the Irish
1075 Research Council, GOIPD/2017/798. We thank Dr. Robin Cash for helpful comments
1076 in preparing this manuscript

1077
1078 **Author contributions**

1079 KR and NW conceived the experiment. KR and NE performed the measurements.
1080 KR, EM and BC analyzed the data. KR, SRS, DM, JB and NW edited the manuscript.
1081 All authors contributed to the scientific discussion and manuscript revisions.

1082

1083 **Competing interests:** The authors declare no competing financial interests

1084

1085

1086

1087 **FIGURE LEGENDS**
1088

Figure 1. Outline of experimental setup. Each trial of neurofeedback training

commenced with a display of four circles (A), each representing the background EMG

in one of the recorded hand muscles (right FDI, ADM and OP, and left FDI). The circles

were red if the root mean squared (rms) EMG at rest was greater than 7 microvolts. It

was essential that all four circles were green for at least 500ms before the trial could

proceed. When this condition was met a fixation cross appeared for a random period

(in order to prevent anticipation of the TMS pulse). During the fixation cross, it was

still essential to keep the background EMG below 7 microvolts in order for a TMS pulse to be delivered. (B) The peak-peak amplitude of the motor evoked potential (MEP) evoked by the TMS was calculated in real-time and displayed immediately to the participant on screen in the form of a rectangular bar.

(C) Different feedback for UP training and DOWN training. In the UP training If the MEP was greater than the baseline mean, the rectangle was green, with a green tick, a dollar sign to indicate a small financial reward, a display of the current score, and a positive encouraging sound bite was heard. If the MEP did not meet the criterion amplitude, the bar was red, there was no dollar sign, and a negative sound bite was heard. (D) A custom 3D printed 'coil spacer' device was used to prevent direct contact of the TMS coil on the EEG electrodes and allow the pre-TMS EEG period to be recorded artefact free.

Figure 2. MEP amplitudes during neurofeedback. Panel A depicts MEP amplitude in millivolts during the two types of MEP neurofeedback. UP training is shown in orange and DOWN training in blue, across all 10 training blocks. Filled triangles labelled 'BS' indicate the baseline measurement block that occurred at the beginning of that particular session, prior to any neurofeedback. Dotted vertical lines indicate the separation of the blocks into different 'sessions', which occurred on separate days. Panel B shows the same data for the control group who received no veridical neurofeedback. Panel C shows the UP-DOWN difference (in the normalised % change from baseline data) for each block in the experimental group and the control group. Higher values represent greater deviations between the UP and DOWN data points and therefore more modulation of MEP amplitude. Thus, these values are significantly higher in the experimental group than in the control group. # symbols indicate blocks in which the Cohen's d effect size for the difference between the experimental and control group was large-very large (>0.8). All data are shown as mean ±SEM.

Figure 3. Retention, aftereffects and feedback-free measurements. Filled bars represent blocks of neurofeedback, and unfilled bars represent MEPs collected at rest. Panel A shows MEP amplitudes with their preceding resting baseline values subtracted. Values above 0 represent increases relative to baseline, and below 0 represent decreases. State-dependent neurofeedback training feedback effects were still evident in a retention block carried out approximately 6 months following the initial

1130   experiment. No aftereffects were observed on resting MEP amplitude 5 and 10 minutes

1131   later. In a separate block participants were capable of upregulating and downregulating

1132   MEP amplitudes with feedback removed (FB free). MEPs measured from the opposite

1133   hemisphere during neurofeedback exhibited a similar pattern of modulation.

1134

Figure 4. Investigation into mechanisms of MEP neurofeedback. The data show

paired pulse TMS measurements taken during neurofeedback blocks to probe distinct

neurophysiological processes. In all subsequent panels, unfilled bars represent baseline

MEP amplitudes collected at rest prior to the block. Panel A Shows that MEP

amplitudes from the single pulses (from which neurofeedback was provided) exhibited

the same state-dependent modulation as observed previously. In Panel B MEP

amplitudes are expressed as a percentage of the single pulse MEPs. While expected

levels of inhibition were observed for both SICI and LICI paired pulses, there was no

state-dependent modulation. LCD was, however, significantly increased in the UP

condition relative to baseline, and relative to the DOWN condition.

1145

1146

Figure 5. Neural oscillations associated with the trained brain states. Panels b-f

show topographical representations of the relative power (in % of whole spectrum) in

the UP condition minus the DOWN condition, for 5 distinct frequency bands (Averaged

group data, n=14, 3 other frequency bands shown in Supplementary Fig 3). Red colours

indicate regions that demonstrated greater synchronisation in the UP condition. Blue

colours indicate greater synchronization in the DOWN condition. The location of the

electrode nearest to the TMS hotspot varied between participants but was always within

the region indicated in a). Colours are scaled from blue-red by minimum-maximum

(range shown to right of each plot). Panel g shows the same data (UP-DOWN) extracted

for each participant's hotspot electrode. Values greater than 0 indicate larger amplitude

oscillations in the UP condition, and lower than 0 indicate larger oscillations in the

DOWN condition. Stars indicate significant deviations from 0 (Wilcoxon Signed Rank

tests). Panel h shows group level data for regression analyses performed on MEP

amplitudes with relative power in each frequency band. This included all 120 trials (60

UP, 60 DOWN) collected during the combined TMS-EEG recording session. The Y

axis depicts the slope of the regression model. Stars indicate significant deviations from

0 (0 would indicate no slope, Wilcoxon Signed Rank test). Individual regression plots

are shown for one representative participant in Supplementary Fig. 4.

1165

A

B

Neurofeedback of
MEP size

Transcranial Magnetic
Stimulation   (TMS)

$ 2/4

Motor evoked potential
(MEP)

C

UP Training   DOWN Training

$ 2/4 $ 2/4   SUCCESS

FAILURE

D

UP
DOWN

Change from baseline (mV)

Retention
*

+5 mins

+10 mins

FB free
*

Opp Hemi