A different state of mind: neural activity related to volitionally up- versus downregulating cortical excitability

Kathy Ruddy[1,2], Joshua Balsters[1,3], Dante Mantini[1,4], Quanying Liu[1,4], Pegah Kassraian-Fard[1], Nadja Enz[1]., Ernest Mihelj[1]., Bankim Chander[5], Surjo Soekadar[5], Nicole Wenderoth[1].

1. Neural Control of Movement Lab, ETH, Zürich, Switzerland.
2. Institute of Neuroscience, Trinity College Dublin, Ireland
3. Royal Holloway, University of London, UK
4. Movement Control and Neuroplasticity Research Group, KU Leuven, Leuven, Belgium
5. University of Tübingen, Germany

Abstract

To date there exists no reliable method to non-invasively upregulate or downregulate the state of the resting motor system over a large dynamic range. Here we show that an operant conditioning paradigm which provides neurofeedback of the size of motor evoked potentials (MEPs) in response to transcranial magnetic stimulation (TMS), enables participants to self-modulate their own brain state. Following training, participants were able to robustly increase (by 83.8%) and decrease (by 30.6%) their MEP amplitudes. This volitional up- versus downregulation of corticomotor excitability caused an increase of late-cortical disinhibition (LCD), a read-out of presynaptic $GABA_B$ disinhibition which was accompanied by an increase of gamma and a decrease of alpha oscillations in the trained hemisphere. This approach paves the way for future investigations into how altered brain state influences motor neurophysiology and recovery of function in a neurorehabilitation context.

Introduction

Rhythmic oscillatory brain activity at rest is associated with high versus low neuronal responsiveness, or 'excitability' of a region [1,2]. Measuring these momentary fluctuations of neural activity via electro- or magnetoencephalography (EEG/MEG) over human primary motor cortex (M1), it has been demonstrated that frequency, amplitude and phase of the ongoing oscillation cycle systematically modulate responses evoked by transcranial magnetic stimulation (TMS) [3-5] [6-8]. In particular, it has been shown that corticomotor excitability is significantly higher when the power (amplitude) of sensorimotor rhythms in the alpha band (8-14 Hz, also called the 'mu'-rhythm), or beta band (15-30 Hz) are low, or when M1 is stimulated during the trough of the oscillatory cycle of these rhythms [9]. This concept has inspired neurofeedback interventions whereby, for example, stroke patients learn to volitionally desynchronize sensorimotor rhythms with the goal of bringing the

sensorimotor system into a more excitable state as a precursor for enhanced neural

plasticity and accelerated recovery [10-12].

Previous research has focussed on interactions between corticomotor

excitability and cortical dynamics at rest, but much less is known about whether it is

possible to *voluntarily* control the excitability of sensorimotor circuits while keeping

motor output and sensory feedback constant. In the case of stroke rehabilitation, this

mechanism may become particularly relevant as patients are unable to move or

receive sensory feedback from the paretic limb. Therefore, interventions that

optimally harness the residual ability to voluntarily and endogenously activate

relevant brain circuits in the days and weeks early after the incident, may provide the

crucial innervation necessary to promote re-wiring for functional recovery [13].

It is well known that primates [14,15], and humans [10,eg. 16-19] can gain volitional

control of neural activity by receiving neurofeedback via a brain-computer interface

(BCI). Here, we used a BCI-neurofeedback approach as an effective method for

training participants to both volitionally upregulate and downregulate corticomotor

excitability as reflected by the size of TMS-evoked motor potentials (MEPs), with the

aim to modulate their amplitudes over a much larger dynamic range than observed

during rest. Using this approach enabled us to investigate the neural mechanisms that

underlie volitional up- versus down-regulation of corticospinal excitability in the

motor system and the associated oscillatory signatures. By modulating one neural

marker, i.e. motor evoked potential amplitude, while measuring independent

modalities using EEG, or paired-pulse TMS, this approach allows us to causally relate

voluntary rather than incidental changes of corticomotor excitability to cortical

dynamics.

To achieve this goal we developed a BCI by stimulating the cortex with TMS

and providing neurofeedback of MEP amplitudes (Fig. 1). The feedback was designed

such that participants were rewarded for larger than average MEPs in one condition,

and smaller than average in another condition.

In a within-subject cross-over design, participants performed four training

sessions with TMS-neurofeedback of MEP amplitudes in order to learn how to up-

and downregulate their corticomotor excitability (two 'UP' sessions, two 'DOWN'

sessions, order counterbalanced across participants). After the training we
characterised the neural underpinnings of these two distinct activity states in detail by
conducting a series of multimodal experiments using EEG and paired pulse TMS to
profile the associated oscillatory and neurophysiological processes. As it has been
proposed that dynamic modulation of neuronal activity is realized via synchronization
of high frequency rhythms [20] which are tightly coupled to desynchronizing
sensorimotor rhythms [21], we hypothesised that Gamma synchronisation (31-80Hz)
and alpha (8-13Hz) /beta (14-30Hz) desynchronization play a critical role in actively
determining the state of the motor cortex.

131
132 **RESULTS**
133
134 *Bidirectional changes in corticospinal excitability were observed in the MEP*
135 *neurofeedback group but not in a control group.*
136
We first tested whether participants could learn to volitionally increase or decrease
corticomotor excitability when using a motor imagery strategy shaped by
neurofeedback of MEP amplitudes. Across two training sessions, we found that MEP
amplitudes increased during UP training (Fig. 2A, orange symbols) and decreased
during DOWN training (Fig. 2A, blue symbols) relative to the baseline measurement
(BS), revealing a significant dissociation over time (*neurofeedback type* x *block*
number interaction during training session 1 [$F(4,115.9)=3.87$, *p*=0.006], session 2
[$F(4,125.0)=3.7$, *p*=0.007] and EEG session [$F(2,70)=6.9$, *p*=0.002], *F tests following*
mixed effects models, *n*=15; see supplementary Fig. 1 for additional analyses). Since
MEP amplitudes are a compound measure of excitability influenced by multiple
neural elements [22], including background muscle activity [23,24], we repeated this
analysis using the root mean squared (rms) background muscle activation (EMG)
recorded in the 100ms prior to each TMS pulse. Importantly, this control analysis
revealed no such interactions on any of the sessions, suggesting that the observed
modulation was not driven by changes in activity of the target muscle, nor any of the
additional 3 control muscles (OP, ADM, left FDI) (all p>0.18, see Supplementary
Table 2).

In order to isolate the effect of the neurofeedback, we included a control group who
undertook the same protocol, using the same mental imagery strategies, but with

feedback that was not contingent on the MEP amplitudes. This group exhibited no
systematic changes of corticomotor excitability across training (Fig. 2B) and mixed
effects models revealed no significant *neurofeedback type* x *block number*
interactions on any of the separate testing sessions in the control group (all $p>0.06$,
note that statistics approached significance for the second session because MEPs were
randomly higher in the DOWN than in the UP condition ; see Supplementary Fig. 1B
for further details). Additionally, there were no significant differences in background
EMG (All p>0.09, Supplementary Table 2). Next we compared the performance of
the experimental and the control group, by normalizing MEP amplitudes to baseline
(% change) and calculating the difference between UP and DOWN (Fig. 2C). The
differences were substantial in the experimental group, who exhibited on average
MEP amplitudes twice as large during UP than during DOWN, and differed
significantly from the control group where systematic differences were virtually
absent (effect of 'Group' [$F(1,25.6)=13.32$, $p=0.001$], *F tests following mixed effects*,
n=28). The effect sizes (Cohen's *d*) of the between-group differences were small for
the first two blocks (<0.5), but consistently increased during training (*d*= 1.27 for
block 8), and remained high in the two blocks of the EEG session (*d*>0.97). As the
control group were executing the same mental imagery strategies as the experimental
group, this comparison demonstrates that veridical TMS neurofeedback was essential
for gaining volitional control over corticomotor excitability.

177
178 *Neurofeedback training effects are retained for at least 6 months*
179
In a follow-up investigation approximately 6 months following initial neurofeedback
training, we showed for a subset of the participants (n=11) that they had retained the
ability to upregulate and downregulate their MEP amplitude with neurofeedback
(Fig.3; significant effect of neurofeedback type (UP vs DOWN) in a retention block
carried out with no top-up training (F(1,10)=6.64, p=0.028). Measurements of resting
MEP amplitude taken 5 and 10 minutes following the retention block indicated no
after-effects (all p>0.2) indicating that subjects could acutely control corticomotor
excitability without long-lasting after-effects. Having verified that the ability to
modulate brain states had been robustly retained, we next tested whether participants
could sustain this performance even when feedback was removed. Performing a
feedback-free block, we found that MEP amplitude was significantly larger in the UP
versus DOWN condition (F(1,10)=12.32, p=0.006), indicating that when participants

have reached a sufficient level of training they have optimised their mental strategies

and no longer require continuous feedback.

In order to measure whether the feedback-induced changes in corticospinal

excitability were specific to the hemisphere targeted by the neurofeedback, we used

two TMS coils simultaneously and performed one block of 40 trials, where half of the

TMS pules were applied to the left hemisphere (i.e. the usual feedback hemisphere),

and the other half to the right (i.e. the opposite hemisphere). We found that the same

pattern of upregulation and downregulation of MEP amplitudes was observed in the

opposite hemisphere, an effect that approached significance (F(1,20)=4.032, p=0.07)

but was much smaller than in the neurofeedback hemisphere particularly for the UP

condition (UP d = 1.01 for neurofeedback hemi, d=0.27 for opp. hemi, DOWN d=

0.40 for neurofeedback hemi, d=0.35 for opp. hemi, Fig. 3).

Paired pulse TMS investigation of mechanisms

Finally, we investigated which excitatory/inhibitory circuits may have contributed to

the changes in corticomotor excitability, using paired pulse TMS measures of three

different neurophysiological processes: (i) Short-interval intracortical inhibition

(SICI), believed to reflect postsynaptic $GABA_A$ inhibition [25]; (ii) Long interval intra-

cortical inhibition (LICI), considered as a marker for postsynaptic $GABA_B$ inhibition;

and (iii) late-cortical disinhibition (LCD), which is thought to measure presynaptic

$GABA_B$ disinhibition, and manifests as a period in which MEP amplitude returns to

and typically overshoots baseline levels, in a time window following LICI (~220ms

after a suprathreshold conditioning TMS pulse) [26-28]. In the following analyses we

determined the time point (baseline vs during NF) x neurofeedback type (UP,

DOWN) interaction and applied FDR correction for multiple testing. Single pulse

MEPs collected during these measurement blocks (25% of all trials) revealed

significantly larger MEP amplitudes for the UP than the DOWN condition, replicating

the findings of the main experiment (Fig. 4A; significant time point x neurofeedback

type interaction: F(1,27.67)=14.36, p_{FDR}=0.001). Surprisingly, there were no

significant differences in the magnitude of SICI (% of single pulse MEPs) between

the resting baseline data and the SICI MEPs collected during neurofeedback, nor

between the UP versus DOWN states (time point x neurofeedback type interaction
[F(1,28.31)=0.08, p_{FDR}=0.77]). The same was true for LICI (time point x
neurofeedback type interaction [F(1,28.90)=0.02, p_{FDR}=0.88]). Thus, circuits
controlling postsynaptic inhibition did not seem to be differentially modulated by the
UP versus DOWN state. However, for LCD there was a significant time point x
neurofeedback type interaction (F(1,28.35)=12.09, p_{FDR}=0.002, Fig.4B).

Pairwise comparisons revealed that LCD was significantly elevated in the UP
condition, when compared to the baseline measurement taken immediately before
neurofeedback (Fig. 4B, right panel, MeanDiff=50.9%, df=28.35, p<0.001) and when
compared to the equivalent data recorded in the DOWN condition (MeanDiff=56.1%,
df=28.76, p<0.001). For the DOWN condition LCD did not differ significantly from
baseline (p=0.45).

239
240
241 *Distinct oscillatory signatures for high versus low corticospinal excitability*
242
As part of the initial training study (see Figure 2A, 'Ses 3' for the behavioural
results), we investigated whether the two different activity states evoked differential
cortical dynamics extracted from EEG recordings which were acquired
simultaneously while TMS was being performed to provide neurofeedback of MEP
amplitude. As distinct functions have been ascribed to 8 different sub-frequency
bands across the known range of brain signals (0.1 - 80Hz), we now probed whether
volitional changes in corticospinal excitability of M1, drives neural activity measured
in the delta (0.1-4Hz), theta (5-7Hz), low alpha (8-10Hz), high alpha (11-13Hz), low
beta (14-21Hz), high beta (22-30Hz), low gamma (31-50Hz) and high gamma (51-
80Hz) bands. Using the portion of EEG data collected in the 1.5 seconds prior to each
TMS pulse, we calculated relative power in the UP and DOWN states for the eight
frequency bands of interest. Figure 5a-f (n=15) shows that UP- versus DOWN
regulating corticomotor excitability caused reduced band-limited power in the theta
and alpha band (blue areas in Fig.5b-d) while gamma power was clearly increased.
(red areas in Fig.5e,f). For each participant, we extracted the information for the
electrode closest to their individual motor hotspot (Figure 5a shows the different
locations across participants) and calculated whether the UP-DOWN difference (Δ
relative power %) deviated significantly from 0 (Fig. 5g). Wilcoxon signed rank tests
revealed significantly higher power for the UP than DOWN condition in the Delta

(p=0.024, d=0.754), Low Gamma (p_{FDR}=0.024, d=0.753) and High Gamma

(p_{FDR}=0.016, d=0.712) band and significantly lower power for UP than DOWN in the

theta (p_{FDR}=0.003, d=0.947), low alpha (p_{FDR}=0.004, d=0.805) and high alpha band

(p_{FDR}=0.007, d=0.714). Although the feedback was lateralised to MEPs from the right

limb (left hemisphere motor hotspot), we also quantified the same neural oscillations

at the corresponding location in the opposite hemisphere. Here, only the theta rhythm

showed significantly lower power for the UP than the DOWN state (p_{FDR}<0.001,

d=1.071, see Supplementary Fig. 2).

Next, we tested whether the amplitude of neural oscillations recorded at the

hotspot at the time of each TMS pulse could predict the amplitude of the resulting

MEPs. For each participant, MEP amplitudes of the 120 trials (60 UP, 60 DOWN)

were entered as the outcome variable in a robust regression model with trial-by-trial

relative power values for each frequency band as predictor variables. Regression

slopes (beta values) for each participant were carried forward into a group level

analysis (Fig. 5m), and Wilcoxon signed rank tests were used to establish whether the

slopes were significantly different from 0 (a 0 slope would indicate no statistical

relationship between predictor and outcome variable). Lower amplitude oscillations in

theta (p_{FDR}=0.024, Fig.5h), low alpha (p_{FDR}<0.001) and high alpha (p_{FDR}=0.002) were

predictive of larger MEP amplitudes, and higher amplitude oscillations in low gamma

(p_{FDR}=0.020) and high gamma (p_{FDR}=0.020) were significant predictors of larger

MEP amplitudes. In a previous study, it was reported that a strong predictor of

cortical excitability was the low gamma : high alpha ratio [3]. We replicated this

finding, demonstrating that this ratio was a significant predictor of MEP amplitude

(p_{FDR}=0.016) with larger ratios predicting larger MEP amplitudes.

EEG data classification

We next tested whether the distinction between the two trained states was large

enough that the individual data trials could be successfully predicted as 'UP' state or

'DOWN' state, using machine learning based solely on the EEG power values

(relative power data, scaled by 1/f transformation) of the 8 frequency bands of

interest. A linear support vector machine (SVM) was applied to the data of each

participant (60 UP 60 DOWN epochs). The SVM has been shown to be particularly

powerful on EEG data, which is noisy and contains many features that are correlated. This approach additionally allowed us to perform feature selection, to quantify which EEG features most heavily contributed to the distinction between the two states. Using only data from the electrode closest to the hotspot (8 rhythms plus LowGamma:HighAlpha ratio) the SVM was capable of classifying the brain states with an average accuracy of 81.5% (±5.1%) based on 10-fold cross validation which differed significantly (p=0.001, n=14) from a null model revealed by permutation testing (accuracy null model: 49.0% ±13). Additionally, incorporating data from the same rhythms recorded at the corresponding electrode in the opposite hemisphere increased this accuracy to 85.1% (±4.6%) across participants (see Supplementary Table 1). Using feature ranking based on Recursive Feature Elimination (RFE), taking the mode of the top ranked features across participants revealed that the strongest contribution to the high classification accuracy of the latter SVM was the High Gamma rhythm in the hotspot electrode, followed by High Alpha at the hotspot, then the LowGamma:HighAlpha ratio (for full ranking order see Supplementary Table 1).

.

Discussion

Here we aimed to uncover neural activity evoked by voluntarily facilitating or suppressing excitability within sensorimotor circuits, while keeping motor output and sensory feedback constant. We show that using a bidirectional TMS-neurofeedback approach is critical to gain volitional control over MEP amplitudes, a skill that is retained for at least 6 months without further training. This voluntary state-setting with a large dynamic range is causally related to modulating pre-synaptic $GABA_B$ mediated disinhibition and to a prominent increase of gamma power in sensorimotor cortex for the UP state which was accompanied by a clear reduction of power in the theta, low and high alpha bands.

Volitional control of corticomotor excitability

Previous studies have shown that it is possible to gain voluntary control over activity in the central nervous system if appropriate neurofeedback is embedded in a

reinforcement learning task, with food rewards for animals [14,15] and visually
rewarding stimuli for humans [19,29]. Here we confirm that this approach is also suitable
for learning how corticomotor excitability can be bidirectionally up- or down-
regulated. Our participants were initially familiarized with two motor imagery
strategies which are known to modulate corticospinal excitability in the required
manner [29-32]. Learning, however, indicated by progressively stronger dissociation
between the UP and the DOWN state, only took place when direct low-latency
feedback regarding the MEP amplitude was provided. After training, participants
were able to modulate corticomotor excitability across a large range so that MEP
amplitudes were approximately twice as large during the UP than the DOWN
condition. UP training, in particular, resulted in an 83.8% increase of MEP amplitudes
from baseline, while downregulation of MEP amplitude was possible [eg. see 29] but
more difficult (30.6% decrease from baseline). Once acquired, volitional control of
corticomotor excitability was retained for at least 6 months and could be performed
even without online feedback indicating true, long-term learning [33].

Once participants could control their corticomotor excitability, we uncovered
the electrophysiological underpinnings by applying measurements that were
independent of the feedback modality (single pulse TMS) and investigated whether
there were differences between the UP versus DOWN state. This approach cancelled
out the effects that were common to both mental strategies, isolating the mechanisms
underlying the MEP modulation. This revealed two key novel electrophysiological
findings, involving presynaptic GABA$_B$ disinhibition, and gamma oscillations.
Additionally, the pronounced changes of cortical physiology despite the absence of
EMG activity or changes in sensory input suggests that the increase vs decrease of
corticomotor excitability was -a least partly- of cortical origin rather than mediated by
spinal cord mechanisms.

Different excitability states cause modulation of GABA$_B$ mediated inhibitory
circuits

The UP state was associated with a significant increase of LCD while other
measurements probing inhibitory M1 circuits failed to reveal differential effects for

the UP versus DOWN state. LCD is thought to represent a read-out of the presynaptic self-inhibition of GABAergic neurons which is thought to be mediated by extrasynaptic $GABA_B$ auto-receptors [34,35]. This mechanism is hypothesized to result in a net facilitatory effect as observed during the UP condition in our study. Previously, LCD was found to be elevated during motor imagery (MI), but this increase relative to rest was observed irrespective of whether participants imagined voluntarily activating or relaxing hand muscles [36]. However, this investigation was conducted in a single session, and did not employ neurofeedback, so MEP modulation by these two imagination conditions could be expected to be substantially smaller than observed in our study, particularly for the voluntary relaxation condition which had a similar excitability state as the rest condition. Thus, it is possible that the clear modulation of LCD observed here only manifested after neurofeedback training, i.e. when the two excitability states became clearly distinct. It is important to note here that group level results indicated no LCD at rest, and in fact it was only evident during the UP state. While LCD is elicited more readily during contraction [28], some studies have reported LCD at rest [27,36], whereas others have only reported occasional or non-significant facilitation occurring beyond 200ms after the suprathreshold conditioning stimulus, ie. in the period immediately following LICI (Valls-Sole et al, 1992). In our search procedure (to decide upon the optimum conditioning stimulus (CS) intensities), we prioritized SICI and LICI, finding a CS intensity that elicited as close to 50% inhibition of the test MEP as possible. We tested intensities between 106-114% RMT for LICI (and above or below this if no appropriate inhibition was found), and applied these parameters also to LCD (such that the only difference between the LCD and LICI protocols was the ISI). This may simply have been too low to elicit strong LCD at rest. Other studies have reported no LCD at 110% RMT, neither at rest [27] nor with contraction [37]. It is nonetheless interesting that the lack of LCD at rest in the current study was overshadowed by the strong facilitation observed while in the TMS-feedback induced UP state, indicating that future indepth investigation into this effect with a larger range of conditioning stimulus intensities may be warranted.

Different excitability states cause distinct neural dynamics in motor cortex

We observed significant modulation of the alpha and gamma rhythms close to

M1 of the trained hemisphere. Focusing on data from the recording electrode closest

to each individual's hotspot revealed a significant association between low alpha and

high gamma power for the UP versus DOWN state. Trial-by-trial modulation of these

rhythms correlated significantly with MEP amplitude, and a support vector machine

(SVM) classifying the two states based on EEG data ranked the high gamma and high

alpha band as the two top features characterizing the distinction. Our observation of

reciprocal changes in the alpha and gamma band are in line with previous studies

using transcranial as well as intracranial recording methods [1]. The 'pulsed inhibition'

theory suggests that repeated bursts of inhibitory alpha activity serve to temporarily

silence gamma oscillations [1]. Thus, these two rhythms are seen to exhibit a reciprocal

relationship, whereby when alpha is high, gamma is low. In periods of high alpha,

gamma may still burst periodically, but only at the troughs of the oscillation cycle,

meaning that the gamma 'duty cycle' (window for neural processing) is short, and

only brief messages can be sent. By contrast, in periods of low alpha power, the

gamma duty cycle is longer, and more extensive neuronal processing and inter-

regional communication may occur. Our finding of increased gamma activity is also

consistent with previous animal literature, showing that the pharmacological removal

of $GABA_B$-mediated inhibition (by receptor blockage) in rats results in increased

gamma oscillations [38] which have been shown to be largest in M1's layer V [39].

Gamma has often been considered difficult to detect using scalp electrodes

because it is highly localised [40] and may also reflect non-cortical sources when

recorded with EEG [41,42]. However, it is tempting to speculate that, in our experiment,

gamma activity was strongly synchronized as a consequence of the neurofeedback

training, where participants learned to substantially facilitate corticomotor excitability

while keeping EMG activity constant, such that EMG amplitude differed only

minimally between the UP and DOWN conditions. This suggestion is in line with

previous neurofeedback studies that provided direct feedback of gamma activity,

showing that gamma power could be upregulated to a substantial amount which even

exceeded power values observed during movement execution [15,43]. By keeping the

visual feedback for the two conditions identical, we ensured that differences in eye movements between the UP and DOWN states were minor. As we were particularly interested in gamma oscillations, we additionally performed all EEG recordings in an electromagnetically shielded room, using a gel-based electrode system to maximize signal to noise ratio.

Previous studies have taken a correlational approach to investigating the relationships between brain rhythms and corticomotor excitability. These have shown that low alpha [4,44] or beta power [45] as well as high gamma power [3] during natural fluctuations at rest are associated with larger MEP amplitudes. We confirm and extend these results by introducing causality to this relationship for the first time, showing that experimentally driving excitability into two distinct states causes specific patterns of neural dynamics in the volitionally controlled cortical area.

While changes in alpha and gamma were specific to the hemisphere from which feedback was provided, theta showed a bilateral pattern of modulation, being higher in the DOWN than the UP state in motor areas in both hemispheres. While mid-frontal theta activity has been linked to error monitoring [46] the role of lateralized theta activity close to the sensorimotor hotspot electrode and its symmetric counterpart is less clear. Slower rhythms exert effects over larger distances, and are thought to be involved in long-range communication [40]. A similar pattern of upregulation and downregulation was observed in the homologous muscle in the opposite limb, albeit weaker and not statistically significant. This is likely a reflection of the extensive transcallosal structural connectivity and functional coupling of homologous regions of the cortical motor network [47-49]. It is tempting to speculate that the bilateral theta activity observed in the current study served to regulate the inhibition/facilitation of functional coupling or 'spillover' of activation from motor areas in the target hemisphere to their homologous counterparts.

Surprisingly we did not observe differential modulation of the Beta band, which is the predominant oscillatory frequency in sensorimotor cortical regions [50,51]. It typically desynchronizes (together with alpha) during motor execution and motor

imagery [52-55] and has been associated with corticomotor excitability at rest [3]. As our

results represent the direct contrast between the UP and DOWN states, the lack of

Beta involvement may firstly be due to the fact that both conditions involved a mental

strategy targeted at the sensorimotor system and, secondly, that no temporal structure

was imposed so that we could not perform analyses which are, for example, time-

locked to the potential onset of these mental strategies. However, our data further

confirm that the two 'inhibitory' rhythms alpha and beta might serve different

functions in selecting and activating the appropriate sensorimotor representations [56].

Conclusion and future applications

Here we present an innovative approach to voluntarily and bidirectionally change the

state of the motor cortex, by directly targeting MEP amplitudes in a neurofeedback

paradigm. This method provided a unique opportunity to reveal the oscillatory and

neurochemical underpinnings of the two distinct trained brain states, using concurrent

TMS EEG measurements, and mechanistic follow-up investigations using paired-

pulse TMS. The results comparing UP and DOWN states indicate that voluntary

upregulation of corticomotor excitability causes increased presynaptic $GABA_B$-

mediated disinhibition, elevated neural oscillations in the gamma frequency range,

and reduced alpha and theta rhythms.

This paves the way for new technologies that allow the user to regulate aspects

of their own brain function in order to reach desired states that are, for example,

associated with enhanced motor performance. In the context of stroke rehabilitation,

training volitional modulation of corticomotor excitability may hold promise as a

rehabilitative therapy early after stroke, i.e. when patients are deprived of

rehabilitation training because they are unable to execute overt movements with the

impaired upper limb. As it is known that LCD is recruited during actual movement

[28,57,58], the elevated LCD we observed in the UP condition may reflect that the

neurofeedback had engaged similar mechanisms to those involved in movement

execution, using only voluntary endogenous processes. Furthermore, as pathological

hyperexcitability of the non-damaged hemisphere has been hypothesized to limit

recovery in some patients [59], the TMS-neurofeedback protocol can be individually

tailored either to upregulate the damaged hemisphere, down-regulate the intact
hemisphere, or a combination of both, depending on the patient's specific needs.
500
501
502
503
504 **FIGURE LEGENDS**
505
Figure 1. Outline of experimental setup. Each trial of neurofeedback training
commenced with a display of four circles (A), each representing the background
EMG in one of the recorded hand muscles (right FDI, ADM and OP, and left FDI).
The circles were red if the root mean squared (rms) EMG at rest was greater than 7
microvolts. It was essential that all four circles were green for at least 500ms before
the trial could proceed. When this condition was met a fixation cross appeared for a
random period (in order to prevent anticipation of the TMS pulse). During the fixation
cross, it was still essential to keep the background EMG below 7 microvolts in order
for a TMS pulse to be delivered. (B) The peak-peak amplitude of the motor evoked
potential (MEP) evoked by the TMS was calculated in real-time and displayed
immediately to the participant on screen in the form of a rectangular bar.
(C) Different feedback for UP training and DOWN training. In the UP training If the
MEP was greater than the baseline mean, the rectangle was green, with a green tick, a
dollar sign to indicate a small financial reward, a display of the current score, and a
positive encouraging sound bite was heard. If the MEP did not meet the criterion
amplitude, the bar was red, there was no dollar sign, and a negative sound bite was
heard. (D) A custom 3D printed 'coil spacer' device was used to prevent direct
contact of the TMS coil on the EEG electrodes and allow the pre-TMS EEG period to
be recorded artefact free.
525
Figure 2. MEP amplitudes during neurofeedback. Panel A depicts MEP amplitude
in millivolts during the two types of MEP neurofeedback. UP training is shown in
orange and DOWN training in blue, across all 10 training blocks. Filled triangles
labelled 'BS' indicate the baseline measurement block that occurred at the beginning
of that particular session, prior to any neurofeedback. Dotted vertical lines indicate
the separation of the blocks into different 'sessions', which occurred on separate days.
Panel B shows the same data for the control group who received no veridical

neurofeedback. Panel C shows the UP-DOWN difference (in the normalised % change from baseline data) for each block in the experimental group and the control group. Higher values represent greater deviations between the UP and DOWN data points and therefore more modulation of MEP amplitude. Thus, these values are significantly higher in the experimental group than in the control group. # symbols indicate blocks in which the Cohen's d effect size for the difference between the experimental and control group was large-very large (>0.8). All data are shown as mean ±SEM.

Figure 3. Retention, aftereffects and feedback-free measurements. Filled bars represent blocks of neurofeedback, and unfilled bars represent MEPs collected at rest. Panel A shows MEP amplitudes with their preceding resting baseline values subtracted. Values above 0 represent increases relative to baseline, and below 0 represent decreases. State-dependent neurofeedback training feedback effects were still evident in a retention block carried out approximately 6 months following the initial experiment. No aftereffects were observed on resting MEP amplitude 5 and 10 minutes later. In a separate block participants were capable of upregulating and downregulating MEP amplitudes with feedback removed (FB free). MEPs measured from the opposite hemisphere during neurofeedback exhibited a similar pattern of modulation.

Figure 4. Investigation into mechanisms of MEP neurofeedback. The data show paired pulse TMS measurements taken during neurofeedback blocks to probe distinct neurophysiological processes. In all subsequent panels, unfilled bars represent baseline MEP amplitudes collected at rest prior to the block. Panel A Shows that MEP amplitudes from the single pulses (from which neurofeedback was provided) exhibited the same state-dependent modulation as observed previously. In Panel B MEP amplitudes are expressed as a percentage of the single pulse MEPs. While expected levels of inhibition were observed for both SICI and LICI paired pulses, there was no state-dependent modulation. LCD was, however, significantly increased in the UP condition relative to baseline, and relative to the DOWN condition.

Figure 5. Neural oscillations associated with the trained brain states. Panels b-f
show topographical representations of the relative power (in % of whole spectrum) in
the UP condition minus the DOWN condition, for 5 distinct frequency bands
(Averaged group data, n=14, 3 other frequency bands shown in Supplementary Fig 3).
Red colours indicate regions that demonstrated greater synchronisation in the UP
condition. Blue colours indicate greater synchronization in the DOWN condition. The
location of the electrode nearest to the TMS hotspot varied between participants but
was always within the region indicated in a). Colours are scaled from blue-red by
minimum-maximum (range shown to right of each plot). Panel g shows the same data
(UP-DOWN) extracted for each participant's hotspot electrode. Values greater than 0
indicate larger amplitude oscillations in the UP condition, and lower than 0 indicate
larger oscillations in the DOWN condition. Stars indicate significant deviations from
0 (Wilcoxon Signed Rank tests). Panel h shows group level data for regression
analyses performed on MEP amplitudes with relative power in each frequency band.
This included all 120 trials (60 UP, 60 DOWN) collected during the combined TMS-
EEG recording session. The Y axis depicts the slope of the regression model. Stars
indicate significant deviations from 0 (0 would indicate no slope, Wilcoxon Signed
Rank test). Individual regression plots are shown for one representative participant in
Supplementary Fig. 4.

585
586
587
588

589 **Materials and Methods**
590
Participants

Fifteen healthy volunteers (age 23± 3.14 s.d, 7 female) participated in the
experimental group. An additional thirteen participants (age 25± 3.06 s.d, 3 female)
formed a control group. All participants were right handed according to the Edinburgh
Handedness Inventory [60], and gave informed consent to procedures. The experiments
were approved by the Kantonale Ethikkommission Zürich, and were conducted in
accordance with the Declaration of Helsinki (1964).

TMS-based neurofeedback

Participants undertook five sessions of TMS-based neurofeedback, on separate

days. The first four days comprised of neurofeedback training, and on the fifth day

neurofeedback was performed with simultaneous electroencephalography (EEG)

recording to investigate state specific neural dynamics. On two of the training days

neurofeedback was adjusted so that a rewarding visual stimulus was displayed when

MEPs were larger than baseline (the 'UP' condition) and on the other two days, the

rewarding stimulus was displayed when MEPs were smaller than baseline (the

'DOWN' condition). On each of the training days, 4 separate blocks of neurofeedback

were preformed, each comprising of 30 individual MEP feedback trials (total 120

trials per day). The format of individual trials and feedback is described in more detail

below. Baseline corticospinal excitability was measured on each day prior to training

(20 MEPs) and post-measurements were taken during the rest periods between each

of the 4 blocks (12 MEPs per measurement).

Subjects sat in a comfortable chair with both arms and legs resting in a neutral

position supported by foam pillows. Surface electromyography (EMG, Trigno

Wireless; Delsys) was recorded from right First Dorsal Interosseous (FDI), Abductor

Digiti Minimi (ADM), Opponens Pollicis (OP), and left FDI. EMG data were

sampled at 2000Hz (National Instruments, Austin, Texas), amplified and stored on a

PC for off-line analysis.

TMS was performed with a figure-of-eight coil (internal coil diameter 50mm)

connected to a Magstim 200 stimulator (Magstim, Whitland, UK). The coil was held

on the left hemisphere over the 'hotspot' of the right FDI at the location with the

largest and most consistent MEPs, and with the optimal orientation for evoking a

descending volley in the corticospinal tract (approximately 45 degrees from the

sagittal plane in order to induce posterior-anterior current flow). Once the hotspot was

established, the lowest stimulation intensity at which MEPs with peak-to-peak

amplitude of approximately 50μV were evoked in at least 5 of 10 consecutive trials

was taken as Resting Motor Threshold (RMT).

The stimulation intensity used to evoke MEPs during the experiment was

chosen using the following procedure in order to obtain a baseline MEP amplitude

that was 50% of the participant's maximum. A recruitment curve [eg. 61] was performed

at the beginning of the first experimental session, whereby 6 TMS pulses were

applied at 10 different intensities relative to RMT (90%, 100%, 110%, 120%, 130%,

140%, 150%, 160%, 180%, 190%) in a randomized order. MEP amplitude at each

intensity was plotted to determine the point on the curve at which plateau occurs and

the MEPs do not continue to increase. Maximal MEP amplitude was recorded, and the

intensity required to evoke 50% of this amplitude was used for all subsequent testing.

With this approach, there was scope for MEP amplitude to both increase and decrease

to similar extents from this 'intermediate' value. Post-hoc analyses revealed that this

procedure resulted in an average stimulation intensity corresponding to 130% RMT.

Immediately following this procedure and prior to the first block of neurofeedback, 20

MEPs were collected at rest at the chosen intensity to determine 'baseline'

corticospinal excitability. The mean MEP amplitude at baseline was recorded and

used during neurofeedback to establish the criterion amplitude that determined

whether participants received either positive or negative feedback.

Format of neurofeedback

Neurofeedback was performed using custom written MATLAB software.

Participants kept eyes open with attention directed to a monitor in front of them. They

were instructed to relax their limbs and avoid tensing any muscles throughout the

experiment. In order to ensure that MEP amplitude could not be influenced by

background muscle activation, the root mean square (rms) of the EMG signal for each

muscle for the previous 100ms of data was calculated and displayed in real-time on

screen at the beginning of each trial in the form of four coloured 'traffic lights',

representing each muscle (Fig. 1A). If the background EMG in a muscle exceeded

7µV, the corresponding light turned red. Participants were instructed that a trial could

not begin unless all four lights were green (all muscles relaxed below 7µV) for at least

a continuous 500ms period. When a trial commenced the traffic lights disappeared,

but background EMG continued to be monitored and the trial was automatically

paused if any muscle exceeded the threshold. At the beginning of each trial a fixation

cross appeared in the center of the screen. After a variable period of time (between 5.5

- 8.5 seconds, or longer if muscle activation delayed the trial) a TMS pulse was fired.

The MEP amplitude for the target muscle (right FDI) was immediately measured and

displayed to the participant on screen within 500ms. The display consisted of a

vertical bar indicating MEP amplitude relative to a horizontal line in the middle of the

screen representing the mean recorded at baseline (Fig. 1B). In 'UP' sessions if the

MEP was larger than the criterion amplitude, the bar was shown as green with a tick

beside it, a positive soundbyte was heard, and a number adjacent to a dollar sign

incremented to indicate that a small financial reward had been gained. If the MEP was

smaller than the criterion amplitude, the bar was red with a cross beside it, a negative

soundbyte was heard, and no financial reward was shown. The reverse was true in the

'DOWN' sessions (Fig. 1C). The feedback remained on screen for 4 seconds, before

being replaced by the traffic lights display preceding the next trial. Participants were

instructed to attend to the feedback and that the goal was to increase (or decrease) the

size of the MEP represented by the bar. Prior to the experiment participants read an

instruction sheet explaining the procedures above and providing recommended mental

strategies that were reported in previous literature in which corticospinal excitability

was downregulated [29] and upregulated [31] by motor imagery (Specific task instructions

are provided in Supplementary Material). Initially the criterion amplitude

corresponded to the baseline MEP measure. After each block of 30 MEPs,

performance was quantified and the task difficulty was adjusted if necessary. If the

success rate was >70% difficulty was increased by raising (or lowering in the DOWN

condition) the criterion MEP amplitude that needed to be reached by 10% in order for

the positive reward to be presented. If performance was > 90%, this was adjusted by

20%.

EEG session

On the fifth day neurofeedback was provided during simultaneous EEG

recording. The participant's TMS hotspot was determined and marked on the scalp

prior to EEG capping. EEG signals were recorded using a 64 channel gel-based TMS-

compatible cap (Electrical Geodesics Inc., Oregon, USA), and the channel closest to

the TMS hotspot was noted. EEG data were amplified and sampled at 1000hz. In

order to minimize artefacts associated with the direct contact of the TMS coil resting

on the electrodes of the EEG cap, we designed and 3D-printed a custom plastic 'coil

spacer' device [62], which has four wide legs positioned to provide a platform
distributing the weight of the TMS coil, so that it hovers over the electrodes without
contact (Fig. 1D). This allowed quality recordings to be obtained even from the
channel of interest closest to the participant's 'hotspot'. The participants RMT was
established while wearing the EEG cap with TMS coil spacer, and the same % above
threshold that was used for all previous sessions was applied for neurofeedback.
Impedances were monitored throughout and maintained below 50kΩ.

Baseline corticospinal excitability was measured in the same fashion as for the
first four sessions, followed by two blocks of neurofeedback (UP or DOWN,
counterbalanced) with brief (12 MEP) post measurements following each. After the
final post measurement, a 15 minute rest break was scheduled for the participant.
Following this, the procedure was repeated and baseline excitability was measured
again, followed by two blocks of either UP or DOWN neurofeedback (whichever was
not performed in the first half of the session). At the end of this session participants
were debriefed.

Control group

Participants were blinded as to whether they were allocated to the experimental
or control group. The control group experienced identical conditions to the
experimental group, with the exception that direct neurofeedback was not provided.
The visual feedback bar demonstrating MEP amplitude was always the same height
(reaching the 'mean' horizontal line). 'Positive' feedback/rewards were presented in
the same proportion as in the experimental group (66% of all trials - calculated upon
completion of experimental group), but at a fixed and predicable rate in order to
prevent the development of illusory correlations. Participants were instructed to attend
to the visual feedback on screen, and that rewards would occur at a fixed rate. Aside
from this, they were otherwise given identical instructions as the participants in the
experimental group- i.e. the same recommended mental strategies were provided on
control 'UP' and 'DOWN' blocks.

728
729   ***Data processing and analysis***
730
MEP data

EMG data from all four hand muscles were band-pass   filtered (30–800 Hz),

separately for the portion of data containing the 100ms of 'pre-TMS' background

EMG, and for the portion of EMG containing the MEP, in order to prevent smearing

of the MEP into the background EMG data chunk. The root mean squared (rms) of

the background EMG was calculated, and peak-peak MEP amplitude was measured.

Trimming (removal) of the maximum and minimum MEP in each block was

performed in order to screen out extreme values. MEP amplitude is known to be

modulated by EMG background activation [23,24]. Therefore, the rms pre-stimulus EMG

recordings were used to assess the presence of unwanted background EMG activity in

the period 110 to 10ms preceding the magnetic pulse. MEPs preceded by background

EMG higher than 0.01mV were excluded. For each subject and over all trials we

calculated the mean and standard deviation of the background EMG. MEPs that

occurred when the background EMG value exceeded 2.5 standard deviations above

the mean, and MEPs with a peak-to-peak amplitude which exceeded Q3 + 1.5 x (Q3 -

Q1) were removed from further analysis, with Q1 denoting the first quartile and Q3

the third quartile computed over the whole set of trials for each subject.

Inferential statistics were computed using Mixed Effects Models in SPSS

(Version 16.0, SPSS Inc. Chicago, US), as they account for covariances between

related data samples in repeated measures designs, and have greater flexibility for

modeling effects over time than traditional ANOVA approaches [63]. Fitting of the

mixed effects models employed restricted maximum likelihood estimation (REML)

and a compound symmetry covariance matrix. Model fit indices (Akaike Information

Criterion and Schwarz Bayesian Criterion) were considered prior to choosing the

covariance matrix and model type. Fixed effects were *neurofeedback type* (UP or

DOWN) and *block number* (1-10). The influence of each of the fixed effects on the

model was estimated using *F* tests. In all models *subject* was designated as a random

effect with random intercepts.

The criterion alpha value was set to 0.05 for all inferential tests. In cases where

multiple comparisons were performed, *p* values were false discovery rate (FDR)

corrected.

 EEG data

Signals from all 64 channels were first epoched to extract only the data on each

trial from the 4 seconds before the TMS pulse. This was to remove the substantial

artefacts that arise during the magnetic pulses, prior to conducting any filtering or

further processing. These separate chunks of unpolluted data were then concatenated

into one continuous epoch, and highpass filtered at 1Hz, prior to conducting an

independent components analysis (ICA). Independent components were visualized

and those containing artefacts arising from eye movements, facial EMG, cardiac

signals, bad channels or other non-brain activity related signals were removed.

The cleaned data were average-referenced, and re-epoched into chunks of data

containing only the 1.5s on each trial prior to the TMS pulse (ie. to capture the

ongoing oscillatory activity at the instance in which the TMS occurred, while the

fixation cross was on screen and the 'traffic lights' had disappeared).

A power spectrum was computed for each single epoch and the mean power (and

relative power) in each of the relevant bandwidths were extracted (delta (0.1-4Hz),

theta (5-7Hz), low alpha (8-10Hz), high alpha (11-13Hz), low beta (14-21Hz), high

beta (22-30Hz), low gamma (31-50Hz) and high gamma (51-80Hz). Power values

were computed separately for UP and DOWN trials, and non-parametric Wilcoxon

signed rank tests (with FDR correction) were used to compare neural oscillations in

these two states.

We also analysed whether trial-by-trial variation of EEG data was associated with

trial-by-trial variation of MEP amplitudes. Therefore, relative power in each

bandwidth for each epoch was entered into a multiple regression model with MEP

amplitudes measured in the muscle from which neurofeedback was provided (right

FDI). The beta (slope) values resulting from each regression model for each

participant were forwarded into a group-level analysis.

794
795 *Classification of distinct brain states*
796
797 Individual epochs of EEG data (60 UP 60 DOWN) were classified by a linear support

798 vector machine (SVM, 10-fold cross validation), to test separately for each participant

799 whether the epochs could be successfully predicted as 'UP' state or 'DOWN' state

800 based solely on the power values (scaled by using 1/f transformed relative power) of

the 4 frequency bands of interest. The SVM was chosen as it is known to perform particularly well in BCI settings using EEG data which is noisy and has features that are correlated. In order to validate the results the same procedure was repeated with randomly permuted labels, and this null model was statistically compared to the model with true labels (C=1). Feature selection was conducted using feature ranking based on Recursive Feature Elimination [64].

Follow-up experiment 6 months later

A sub-set of 11 participants from the experimental group returned approximately 6 months later to participate in a follow-up experiment probing retention and mechanisms underlying the two distinct states. This was conducted over a further 4 days of testing. On one day, retention, aftereffects, and excitability in the opposite 'untrained' hemisphere were tested for the 'UP' condition. On another, neurophysiolocial mechanisms were probed using paired pulse TMS. These two days were repeated for the 'DOWN' condition, and the order of these sessions was counterbalanced. We additionally tested whether trained participants were able to upregulate and downregulate when feedback was temporarily removed.

Retention testing & aftereffects measurement

After a 6-month break and no top-up training, participants were tested with one block of TMS-neurofeedback (20 MEPs) in order to assess retention of learning. All other procedures were identical to those carried out in the main experiment.

Following this block, 12 MEPs were collected at rest after 5 and 10 minutes.

Excitability in the opposite hemisphere

During one block, two TMS coils were used, placed over the right and left motor hotspots (as described previously). This block contained 40 trials, 20 of which were normal TMS neurofeedback trials. The other 20 were trials where TMS was applied to the opposite hemisphere, rather than to the hemisphere that was the target for neurofeedback. No feedback was given in these trials. The presentation of left and right hemisphere TMS pulses was randomized.

Feedback- free measurements

We additionally tested whether trained participants were able to upregulate and
downregulate when feedback was temporarily removed. In this feedback-free block,
the timing of trials and participant instructions were identical to normal
neurofeedback blocks, but in place of the usual feedback bar showing MEP
amplitude, the white fixation cross simply turned red during this period. The onset of
trials was still contingent on muscles being completely relaxed, and the traffic lights
display still preceded every trial.

Paired pulse TMS measurements

On separate days (one 'UP' one 'DOWN) from the measurements described above,
we performed three additional blocks of TMS neurofeedback (24 trials per block x 3
= 72 total trials), in which just 25% of trials were standard single pulse TMS-
neurofeedback trials, with the usual feedback. The remaining trials contained paired
pulses in place of the usual single pulse TMS. For all paired pulse measurements, the
test stimulus intensity was identical to that which had been chosen for the TMS
neurofeedback (ie. that produced MEPs that were 50% of the maximum on the
recruitment curve). On 25% of trials Short Interval Intracortical Inhibition (SICI) was
measured. This was with a conditioning stimulus intensity that was chosen using a
personalized search procedure testing intensities ranging from 50%-90% RMT, to
achieve as close to 50% inhibition as possible, and an inter-stimulus interval of
1.97ms [65]. The reduction in the size of the test MEP is believed to represent
postsynaptic $GABA_A$ inhibition [25]. On 25% of trials Long Interval Intracortical
Inhibition (LICI) was measured. This was with two suprathreshold pulses, with the
conditioning stimulus intensity chosen using a search procedure between 106-114%
RMT, and an inter-stimulus interval of 100ms [27]. This is believed to reflect
postsynaptic $GABA_B$ inhibition [66]. On the remaining 25% of trials, Late Cortical
Disinhibition (LCD) was tested. This was with the exact same pulse intensities as
used for LICI, but with a 220ms inter-stimulus interval [27], and is thought to measure
presynaptic $GABA_B$ disinhibition [26-28]. The order of presentation of paired pulses and
single pulses was randomized.

Baseline measurements were taken at rest with each of these three paired-pulse TMS

protocols, prior to the beginning of neurofeedback blocks (20 of each type of paired

pulse measurement, and 20 single pulse MEPs).

1

REFERENCES

1. Jensen, O. & Mazaheri, A. Shaping Functional Architecture by Oscillatory
Alpha Activity: Gating by Inhibition. *Frontiers in human neuroscience* **4,**
(2010).
2. Jensen, O. *et al.* Using Brain–Computer Interfaces and Brain-State
Dependent Stimulation as Tools in Cognitive Neuroscience. *Front Psychol* **2,**
(2011).
3. Zarkowski, P., Shin, C. J., Dang, T., Russo, J. & Avery, D. EEG and the
Variance of Motor Evoked Potential Amplitude. *Clinical EEG and …* (2006).
4. Sauseng, P., Klimesch, W., Gerloff, C. & Hummel, F. C. Spontaneous locally
restricted EEG alpha activity determines cortical excitability in the motor
cortex. *Neuropsychologia* **47,** 284–288 (2009).
5. Schaworonkow, N., Triesch, J., Ziemann, U. & Zrenner, C. EEG-triggered
TMS reveals stronger brain state-dependent modulation of motor evoked
potentials at weaker stimulation intensities. *bioRxiv* 251363 (2018).
doi:10.1101/251363
6. Kelly, S. P., Gomez-Ramirez, M. & Foxe, J. J. The strength of anticipatory
spatial biasing predicts target discrimination at attended locations: a high-
density EEG study. *Eur. J. Neurosci.* **30,** 2224–2234 (2009).
7. Mazaheri, A., Nieuwenhuis, I. L. C., van Dijk, H. & Jensen, O. Prestimulus
alpha and mu activity predicts failure to inhibit motor responses. *Hum Brain*
Mapp **30,** 1791–1800 (2009).
8. Schubert, R., Blankenburg, F., Lemm, S., Villringer, A. & Curio, G. Now you
feel it--now you don't: ERP correlates of somatosensory awareness.
Psychophysiology **43,** 31–40 (2006).
9. Zaehle, T., Rach, S. & Herrmann, C. S. Transcranial alternating current
stimulation enhances individual alpha activity in human EEG. *PLoS ONE* **5,**
e13766 (2010).
10. Buch, E. *et al.* Think to move: a neuromagnetic brain-computer interface
(BCI) system for chronic stroke. *Stroke* **39,** 910–917 (2008).
11. Caria, A. *et al.* Chronic stroke recovery after combined BCI training and
physiotherapy: A case report. *Psychophysiology* **48,** 578–582 (2010).
12. Machado, S. *et al.* EEG-based brain-computer interfaces: an overview of
basic concepts and clinical applications in neurorehabilitation. *Reviews in the*
Neurosciences **21,** 451–468 (2010).
13. Maulden, S. A., Gassaway, J., Horn, S. D., Smout, R. J. & DeJong, G.
Timing of Initiation of Rehabilitation After Stroke. *Arch Phys Med Rehabil*
86, 34–40 (2005).

14. Fetz, E. E. Volitional Control of Cortical Oscillations and Synchrony. *Neuron*
77, 216–218 (2013).

15. Engelhard, B., Ozeri, N., Israel, Z., Bergman, H. & Vaadia, E. Inducing
Gamma Oscillations and Precise Spike Synchrony by Operant Conditioning
via Brain-Machine Interface. *Neuron* (2013).

16. Mellinger, J. *et al.* An MEG-based brain-computer interface (BCI).
Neuroimage **36,** 581–593 (2007).

17. Soekadar, S. R., Witkowski, M., Birbaumer, N. & Cohen, L. G. Enhancing
Hebbian Learning to Control Brain Oscillatory Activity. *Cerebral Cortex*
(2014). doi:10.1093/cercor/bhu043

18. Sitaram, R. *et al.* Closed-loop brain training: the science of neurofeedback.
Nat Rev Neurosci **18,** 86–100 (2017).

19. Thompson, A. K., Chen, X. Y. & Wolpaw, J. R. Acquisition of a Simple
Motor Skill: Task-Dependent Adaptation Plus Long-Term Change in the
Human Soleus H-Reflex. *J. Neurosci.* **29,** 5784–5792 (2009).

20. Fries, P. A mechanism for cognitive dynamics: neuronal communication
through neuronal coherence. *Trends Cogn. Sci. (Regul. Ed.)* **9,** 474–480
(2005).

21. Grosse-Wentrup, M., Schölkopf, B. & Hill, J. Causal influence of gamma
oscillations on the sensorimotor rhythm. *Neuroimage* **56,** 837–842 (2011).

22. Carson, R. G., Ruddy, K. L. & McNickle, E. *What do TMS evoked motor*
potentials tell us about motor learning? (2016).

23. Hess, C. W., Mills, K. R. & Murray, N. M. Magnetic stimulation of the
human brain: facilitation of motor responses by voluntary contraction of
ipsilateral and contralateral muscles with additional observations on an
amputee. *Neurosci Lett* **71,** 235–240 (1986).

24. Devanne, H., Lavoie, B. A. & Capaday, C. Input-output properties and gain
changes in the human corticospinal pathway. *Exp Brain Res* **114,** 329–338
(1997).

25. Fisher, R. J., Nakamura, Y., Bestmann, S., Rothwell, J. C. & Bostock, H.
Two phases of intracortical inhibition revealed by transcranial magnetic
threshold tracking. *Exp Brain Res* **143,** 240–248 (2002).

26. Cash, R. F. H., Ziemann, U. & Thickbroom, G. W. Inhibitory and
disinhibitory effects on I-wave facilitation in motor cortex. *Journal of*
Neurophysiology **105,** 100–106 (2011).

27. Cash, R. F. H., Ziemann, U., Murray, K. & Thickbroom, G. W. Late cortical
disinhibition in human motor cortex: a triple-pulse transcranial magnetic
stimulation study. *Journal of Neurophysiology* **103,** 511–518 (2010).

28. Caux-Dedeystère, A., Derambure, P. & Devanne, H. Late cortical
disinhibition in relaxed versus active hand muscles. *Neuroscience* **298,** 52–62
(2015).

29. Majid, D. S. A., Lewis, C. & Aron, A. R. Training voluntary motor
suppression with real-time feedback of motor evoked potentials. *Journal of*
Neurophysiology **113,** 3446–3452 (2015).

30. Stinear, C. M., Byblow, W. D., Steyvers, M., Levin, O. & Swinnen, S. P.
Kinesthetic, but not visual, motor imagery modulates corticomotor
excitability. *Exp Brain Res* **168,** 157–164 (2005).

31. Izumi, S. *et al.* Facilitatory effect of thinking about movement on motor-
evoked potentials to transcranial magnetic stimulation of the brain. *American*
Journal of Physical Medicine & Rehabilitation **74,** 207–213 (1995).

32. Fadiga, L. *et al.* Corticospinal excitability is specifically modulated by motor
imagery: a magnetic stimulation study. *Neuropsychologia* **37,** 147–158
(1998).
33. Magill, R. *Motor Learning and Control: Concepts and Applications.*
(McGraw-Hill Education, 2010).
34. Mott, D. D. & Lewis, D. V. Facilitation of the induction of long-term
potentiation by GABAB receptors. *Science* **252,** 1718–1720 (1991).
35. Sanger, T. D., Garg, R. R. & Chen, R. Interactions between two different
inhibitory systems in the human motor cortex. *The Journal of Physiology*
530, 307–317 (2004).
36. Chong, B. W. X. & Stinear, C. M. Modulation of motor cortex inhibition
during motor imagery. *Journal of Neurophysiology* **117,** 1776–1784 (2017).
37. Claus, D., Weis, M., Jahnke, U., Plewe, A. & Brunhölzl, C. Corticospinal
conduction studied with magnetic double stimulation in the intact human. *J.*
Neurol. Sci. **111,** 180–188 (1992).
38. Leung, L. S. & Shen, B. GABAB receptor blockade enhances theta and
gamma rhythms in the hippocampus of behaving rats. *Hippocampus* **17,** 281–
291 (2007).
39. Johnson, N. W. *et al.* Phase-amplitude coupled persistent theta and gamma
oscillations in rat primary motor cortex in vitro. *Neuropharmacology* **119,**
141–156 (2017).
40. Buzsaki, G. *Rhythms of the Brain.* (Oxford University Press, USA, 2006).
41. Yuval-Greenberg, S., Tomer, O., Keren, A. S., Nelken, I. & Deouell, L. Y.
Transient induced gamma-band response in EEG as a manifestation of
miniature saccades. *Neuron* **58,** 429–441 (2008).
42. Whitham, E. M. *et al.* Scalp electrical recording during paralysis:
Quantitative evidence that EEG frequencies above 20 Hz are contaminated by
EMG. *CLINICAL NEUROPHYSIOLOGY* **118,** 1877–1888 (2007).
43. Miller, K. J. *et al.* Cortical activity during motor execution, motor imagery,
and imagery-based online feedback. *Proceedings of the National Academy of*
Sciences **107,** 4430–4435 (2010).
44. Zrenner, C., Desideri, D., Belardinelli, P. & Ziemann, U. Real-time EEG-
defined excitability states determine efficacy of TMS-induced plasticity in
human motor cortex. *Brain Stimul* **11,** 374–389 (2018).
45. Mäki, H. & Ilmoniemi, R. J. EEG oscillations and magnetically evoked motor
potentials reflect motor system excitability in overlapping neuronal
populations. *CLINICAL NEUROPHYSIOLOGY* **121,** 492–501 (2010).
46. Pezzetta, R., Nicolardi, V., Tidoni, E. & Aglioti, S. M. Error, rather than its
probability, elicits specific electrocortical signatures: a combined EEG-
immersive virtual reality study of action observation. *Journal of*
Neurophysiology jn.00130.2018 (2018). doi:10.1152/jn.00130.2018
47. Ruddy, K. L., Leemans, A. & Carson, R. G. Transcallosal connectivity of the
human cortical motor network. *Brain Struct Funct* (2016). doi:DOI:
10.1007/s00429-016-1274-1
48. Ruddy, K. L., Leemans, A., Woolley, D. G., Wenderoth, N. & Carson, R. G.
Structural and Functional Cortical Connectivity Mediating Cross Education
of Motor Function. *Journal of Neuroscience* **37,** 2555–2564 (2017).
49. Ruddy, K. L.2017. Directionality of interhemispheric communication.
Springer
1013

1014   50.   Baker, S. N., Olivier, E. & Lemon, R. N. Coherent oscillations in monkey
1015   motor cortex and hand muscle EMG show task-dependent modulation.
1016   (1997).
1017   51.   Murthy, V. N. & Fetz, E. E. Oscillatory activity in sensorimotor cortex of
1018   awake monkeys: synchronization of local field potentials and relation to
1019   behavior. *Journal of Neurophysiology* **76,** 3949–3967 (1996).
1020   52.   Ramos-Murguialday, A. & Birbaumer, N. Brain oscillatory signatures of
1021   motor tasks. *Journal of Neurophysiology* **113,** 3663–3682 (2015).
1022   53.   Pfurtscheller, G. & Neuper, C. Motor imagery activates primary sensorimotor
1023   area in humans. *Neurosci Lett* **239,** 65–68 (1997).
1024   54.   Pfurtscheller, G. & Berghold, A. Patterns of cortical activation during
1025   planning of voluntary movement. *Electroencephalogr Clin Neurophysiol* **72,**
1026   250–258 (1989).
1027   55.   Alegre, M. *et al.* Alpha and beta oscillatory activity during a sequence of two
1028   movements. *CLINICAL NEUROPHYSIOLOGY* **115,** 124–130 (2004).
1029   56.   Brinkman, L., Stolk, A., Dijkerman, H. C., de Lange, F. P. & Toni, I. Distinct
1030   Roles for Alpha- and Beta-Band Oscillations during Mental Simulation of
1031   Goal-Directed Actions. *Journal of Neuroscience* **34,** 14783–14792 (2014).
1032   57.   Hammond, G. & Vallence, A.-M. Modulation of long-interval intracortical
1033   inhibition and the silent period by voluntary contraction. *Brain Res* **1158,** 63–
1034   70 (2007).
1035   58.   Opie, G. M., Ridding, M. C. & Semmler, J. G. Task-related changes in
1036   intracortical inhibition assessed with paired- and triple-pulse transcranial
1037   magnetic stimulation. *Journal of Neurophysiology* **113,** 1470–1479 (2015).
1038   59.   Murase, N., Duque, J., Mazzocchio, R. & Cohen, L. G. Influence of
1039   interhemispheric interactions on motor function in chronic stroke. *Ann.*
1040   *Neurol.* **55,** 400–409 (2004).
1041   60.   Oldfield, R. C. The assessment and analysis of handedness: The Edinburgh
1042   Inventory. *Neuropsychologia* **9,** 97–113 (1971).
1043   61.   Carson, R. G. *et al.* Characterizing changes in the excitability of corticospinal
1044   projections to proximal muscles of the upper limb. *Brain Stimul* **6,** 760–768
1045   (2013).
1046   62.   Ruddy, K. L. *et al.* Improving the quality of combined EEG-TMS neural
1047   recordings: Introducing the coil spacer. *Journal of Neuroscience Methods*
1048   **294,** 34–39 (2018).
1049   63.   Gueorguieva, R. & Krystal, J. H. Move over ANOVA: progress in analyzing
1050   repeated-measures data and its reflection in papers published in the Archives
1051   of General Psychiatry. *Arch. Gen. Psychiatry* **61,** 310–317 (2004).
1052   64.   Guyon, I., Weston, J., Barnhill, S. & Vapnik, V. Gene Selection for Cancer
1053   Classification using Support Vector Machines. *Machine Learning* **46,** 389–
1054   422 (2002).
1055   65.   Peurala, S. H., Müller-Dahlhaus, J. F. M., Arai, N. & Ziemann, U.
1056   Interference of short-interval intracortical inhibition (SICI) and short-interval
1057   intracortical facilitation (SICF). *CLINICAL NEUROPHYSIOLOGY* **119,**
1058   2291–2297 (2008).
1059   66.   McDonnell, M. N., Orekhov, Y. & Ziemann, U. The role of
1060 $GABA_B$ receptors in intracortical inhibition in the
1061   human motor cortex. *Exp Brain Res* **173,** 86–93
1062

A

B

Neurofeedback of MEP size

Transcranial Magnetic Stimulation (TMS)

$ 2/4

Motor evoked potential (MEP)

C

UP Training

DOWN Training

$ 2/4

$ 2/4

SUCCESS

FAILURE

D

UP
DOWN

Retention

+5 mins

+10 mins

FB free

*

Opp Hemi

Change from baseline (mV)

1.5

1.0

0.5

0.0

-0.5

*

*