Genome-wide associations reveal human-mouse genetic convergence and

modifiers of myogenesis, *CPNE1* and *STC2*

Authors

Ana I. Hernandez Cordero[1], Natalia M. Gonzales[2], Clarissa C. Parker[3], Greta Sokoloff[4], David

J. Vandenbergh[5], Riyan Cheng[6], Mark Abney[2], Andrew Skol[7], Alex Douglas[8], Abraham A.

Palmer[9,10], Jennifer S. Gregory[1], and Arimantas Lionikas[1*]

Affiliations:

1. School of Medicine, Medical Sciences and Nutrition, College of Life Sciences and Medicine,

University of Aberdeen, Aberdeen, UK

2. Department of Human Genetics, University of Chicago, Chicago, IL 60637, USA

3. Department of Psychology and Program in Neuroscience, Middlebury College, Middlebury,

VT 05753, USA

4. Department of Psychological and Brain Sciences, The 25 University of Iowa, Iowa City, IA

52242, USA

5. Department of Biobehavioral Health, Penn State Institute for the Neurosciences, &

Molecular, Cellular & Integrative Sciences Program. The Pennsylvania State University,

University Park, PA 16802, USA

6. Department of Health Sciences, University of California San Diego, La Jolla, CA 92093,

USA

7. Department of Medicine, University of Chicago, Chicago, IL 60637, USA

8. Institute of Biological and Environmental Sciences, University of Aberdeen, Aberdeen,

AB24 3FX, UK

9. Department of Psychiatry, University of California San Diego, La Jolla, CA 92093, USA

10. Institute for Genomic Medicine, University of California San Diego, La Jolla, CA 92093,

26   USA

28   *Corresponding Author: Dr. Arimantas Lionikas, e-mail: a.lionikas@abdn.ac.uk

Abstract

Muscle bulk in adult healthy humans is highly variable even after accounting for height, age and sex. Low muscle mass, due to fewer and/or smaller constituent muscle fibers, would exacerbate the impact of muscle loss occurring in aging or disease. Genetic variability substantially influences muscle mass differences, but causative genes remain largely unknown. In a genome-wide association study (GWAS) on appendicular lean mass (ALM) in a population of 85,750 middle-age (38-49 years) individuals from the UK Biobank (UKB) we found 182 loci associated with ALM ($P<5x10^{-8}$). We replicated associations for 78% of these loci ($P<5x10^{-8}$) with ALM in a population of 181,862 elderly (60-74 years) individuals from UKB. We also conducted a GWAS on hindlimb skeletal muscle mass of 1,867 mice from an advanced intercross between two inbred strains (LG/J and SM/J) which identified 23 quantitative trait loci. 38 positional candidates distributed across 5 loci overlapped between the two species. *In vitro* studies of positional candidates confirmed *CPNE1* and *STC2* as modifiers of myogenesis. Collectively, these findings shed light on the genetics of muscle mass variability in humans and identify targets for the development of interventions for treatment of muscle loss. The overlapping results between humans and the mouse model GWAS point to shared genetic mechanisms across species.

Introduction

Skeletal muscle plays key roles in locomotion, respiration, thermoregulation, maintenance of glucose homeostasis and protection of bones and viscera. The loss of muscle due to aging, known as sarcopenia, affects mobility and can lead to frailty and deterioration of quality of life[1]. The risk of disability is 1.5 to 4.6 times higher in the sarcopenic elderly than in the age-matched individuals with normal muscle mass[2]. However, lean mass, a non-invasive proxy for muscle mass, differs by more than two-fold between healthy adult individuals of same sex, age and height[3]. Therefore, we hypothesize that differential accretion of muscle mass by adulthood may influence the risk of sarcopenia and frailty later in life.

Genetic factors contribute substantially to the variability in lean mass in humans, with heritability estimates of 40 – 80 %[4]. A continuous distribution of the trait and data obtained from animal models[5-7] indicates a polygenic causality. However, thus far, genome-wide association studies (GWAS) have implicated fewer than a dozen genes, explaining a small fraction of this heritability[8; 9]. Limited sample size in early studies[10-14] and the effects of confounders such as subject age[8], size of the skeleton and lean mass components (non-fat organs and tissues, heterogeneity of muscle fibers) have hindered detection of genes. The UK Biobank is a resource of demographic, phenotypic and genotypic data collected on ~500,000 individuals[15]. It includes the arm and leg lean mass, body composition and morphometric information, providing a model for improving our understanding of the genetic basis for variability in muscle mass. Skeletal muscle mass, however, changes over the course of an individual's lifespan. It reaches a peak around mid-twenties and remains largely stable through mid-forties, before succumbing to gradual decline, which accelerates after about 70 years of age[16]. There is a substantial degree of individual variability in the slope of muscle change across both the increasing and decreasing phases of the lifespan trajectory[17]. Both the trajectory itself and the slope of individual variability may impede identification of genes.

The indirect estimates of lean mass impose limitations because muscle mass is not an exclusive contributor to this variable. Furthermore, the cellular basis of variability in muscle mass (i.e. if it is caused by the differences in the number of constituent muscle fibers, their size, or both)

remains poorly understood. Using the laboratory mouse circumvents a number of those limitations.

The mouse shares approximately 90% of the genome with humans[18], and dissection permits

analyses of traits that are difficult to study directly in humans, such as the mass of individual

muscles [6; 7; 19] and whole-muscle fiber characteristics[20; 21]. The phenotypic differences between the

LG/J and SM/J mouse strains make them particularly attractive for complex trait analyses[22-24]. LG/J

mice were selected for large body size[25], while SM/J mice were selected for small body size[26]. The

second filial generation (F_2) of intercross derived from the LG/J and SM/J strains (LGSM)[6; 27] and

an advanced intercross line (AIL) of the LGSM (LGSM AIL), developed using a breeding strategy

proposed by Darvasi and Soller[28], led to multiple quantitative trait loci (QTLs) for hindlimb muscle

mass[6; 27]. However, these QTLs still encompass tens or even hundreds of genes and require

further prioritizing. We hypothesized that the detection power of a modest sample size of the

LGSM AIL and the superior resolution of a human cohort will facilitate identification of the

quantitative trait genes (QTGs) underlying muscle QTLs.

The aim of this study was to identify the genomic loci and the underlying genes for

variability in skeletal muscle mass and to assess their effects in the elderly. We addressed this in

three stages: (1) we conducted a GWAS in a human cohort of middle-aged individuals from the UK

Biobank and tested the effect of the identified set of loci in an elderly cohort; (2) we conducted a

GWAS on hindlimb muscle mass in a population of LGSM AIL mice. (3) In the final stage, we

nominated candidate genes by comparing mouse and human loci and validated the myogenic role

of selected candidates *in vitro*.

Methods

Stage one: Genome mapping in human populations

UK Biobank cohort

The population in this study consisted of 316,589 adult individuals of 37 to 74 years of age (project

ID: 26746). We drew this cohort from the UK Biobank (UKB) project[15]; all participants recruited

were identified from the UK National Health Service (NHS) records and attended a baseline visit

assessment between 2006 and 2010. During the assessment, participants gave written consent,

answered a questionnaire, and were interviewed about their health and lifestyle. Blood samples

and anthropometric measurements were collected from each participant. Assessments were

conducted at 22 facilities in Scotland, England, and Wales.

We divided the sample into middle-aged and elderly cohorts. The middle-aged cohort

consisted of 99,065 adults ranging from 38 to 49 years of age; based on previous studies we

assumed that these individuals were not affected by sarcopenia[29]. We excluded 3,520 participants

that were reported to be ill with cancer, pregnant, or had undergone a leg amputation procedure,

as well as individuals with discordant genetic sex and self-reported sex records. In addition, we

excluded non-white Europeans (self-reported) from the analyses (n = 9,599) and individuals

without imputed genotypes. We retained a total of 85,750 adult individuals (46,353 females and

39,397 males) for further analyses.

The elderly cohort consisted of 217,524 adults ranging from 60 to 74 years of age. We

selected this cohort to test if the effect of the genetic variants identified on middle-aged individuals

could also influenced phenotypes later in life. We excluded 35,662 individuals based on the same

criteria used for the middle-aged cohort. After exclusions, the elderly cohort included 181,862

individuals of 60 to 73 years of age (94,229 females and 87,633 males; Table S1).

UK biobank traits

We used the data for standing height (UKB field ID: 50), sitting height (UKB field ID: 20015), whole

body fat (UKB field ID: 23100), arm lean mass (UKB field ID: 23121 and 23125), and leg lean mass

(UKB field ID: 23113 and 23117) measured as part of the UK Biobank project. Body composition

measurements were taken using bioelectric impedance (this was preferred to the DXA scan data

because of the substantially larger number of phenotyped individuals). We calculated leg length by

subtracting sitting height from standing height (all measurements were recorded in cm). Because

lean mass in the limbs primarily consists of skeletal muscle tissue, we used ALM as a proxy for

muscle mass. We calculated ALM as the sum of the muscle mass of two arms and two legs. We

checked that all traits were normally distributed by examining the QQ-plot and histogram of

residuals from a simple linear model that included sex as a covariate. Residuals were normally

distributed, and we did not transform any of the traits.

130   UK biobank genotypes

We obtained genotype data for all participants from the UKB v3 genotypes release[30], which

includes genotype calls from the Affymetrix UK BiLEVE Axiom array and the Affymetrix UK

Biobank Axiom array, and imputed genotypes from the UK10K and 1000 Genomes Phase 3

reference panels[31]. We kept all imputed genotype data (21,375,087 genetic variants (SNPs, Indels

and structural variants)) with MAF > 0.001 and imputation quality > 0.30. The software (BOLT-LMM

v2.3.4) [32] we used to perform GWAS was developed for large data sets (i.e.: UK Biobank cohort)

and it was only tested for human cohorts, which have different LD patterns from animals; BOLT-

LMM uses a linear mixed model, which have been shown to successfully control for confounding

due to population structure or cryptic relatedness in individuals (related and unrelated) from the UK

biobank[33-36]. For these reasons, we used BOLT-LMM v2.3.4 for the analyses of human data only.

142   Appendicular lean mass GWAS

We used BOLT-LMM (v2.3.4)[37] to perform a GWAS for ALM in the middle-aged cohort. The linear

mixed model (LMM) approach implemented in BOLT-LMM is capable of analyzing large data sets

while also accounting for cryptic relatedness between individuals. Specifically, BOLT-LMM

calibrates the association statistics using a linkage disequilibrium (LD) score regression

approach[38]; this allowed us to evaluate the impact of confounding factors on the GWAS test

statistics[38] and calibrate them accordingly. In the absence of confounding factors, P values should

not be inflated, and the LD score regression intercept should be equal to 1[38]. The LD Score

regression intercept in this study was 1.051 ± 0.007, suggesting minimal inflation of P values due

to linkage between markers. After calibrating the test statistics, the mean χ^2 of the ALM GWAS

was 1.29 and lambda (λ_{GC}) or genomic control inflation factor was 1.20 (Figure S1), which

indicated polygenicity of the trait as described by Bulik-Sullivan and colleagues[38].

We also assessed population structure by running principal component analysis on the

genotype calls. We included sex, leg length, whole body fat, and the first four principal components

as fixed effects in the LMM used for the ALM GWAS. Sex was included to account for differences

in muscle mass caused by higher testosterone levels in males[39]. Testosterone is a potent

stimulator of muscle growth and if systematically varied in males, it can also influence muscle

mass (e.g. as a result of hypogonadism[40]). However, if there was a common genetic basis for such

variability it could be captured in the association analysis. It needs to be noted that inclusion of sex

as a covariate would not permit capturing sex-by-locus interactions. Identification of sex specific

loci, albeit of interest, was not attempted due to complexity posed by the number of genetic

markers and the sample size. An outcome of a GWAS would also depend of the complexity of

mechanisms affecting the phenotype and adjustments included in a model[8; 41]. Leg length and

whole body fat were included because they are biologically related to muscle mass: longer bones

result in longer muscles, while fat shares part of its developmental origin with skeletal muscle

tissue[42]. Furthermore, each of these traits is correlated with muscle mass. An association was

considered statistically significant if its $P < 5 \times 10^{-8}$ ($\alpha = 0.05$). This threshold is the standard for

GWAS of complex traits[43; 44].

We obtained variance components and SNP heritability estimates of ALM from the middle-

aged cohort using BOLT-REML[37]. The BOLT-REML method robustly estimates the variance of

genotyped SNPs and fixed effects on the LMM. As described by Loh et al. [45], BOLT-REML

partitions the SNP heritability across common alleles; hence, the additive variance is calculated as

the cumulative variance of genotyped SNPs.

Phenotypic variance explained by ALM loci

We defined ALM genomic loci using the web-based platform Functional Mapping and Annotation of

Genome-Wide Association Studies (FUMA GWAS[46]). A key feature of this tool is the identification

of genomic regions and independent genomic signals based on the provided summary statistics of

a GWAS depending on LD structure; this process is automated using pairwise LD of SNPs in the

reference panel (1000 genomes project phase 3 EUR [47]) previously calculated by PLINK[48]. We

provided to FUMA GWAS the summary statistic of our GWAS on ALM with the following

parameters: 250kb window (maximum distance between LD blocks), $r^2 > 0.6$ (minimum r^2 for

determining LD with independent genome-wide significant SNPs used to determine the limits of

significant genomic loci), MAF > 0.001 (minimum minor allele frequency to be included in the

annotation), $P < 5 \times 10^{-8}$ (threshold of significantly associated variants). We refer to the identified

regions and the independent signals as loci throughout the text.

We used the top variant (based on the outcome from FUMA[46]) of each locus identified to

estimate the proportion of phenotypic variance explained by each locus. We estimated phenotype

residuals using a model that included the fixed effects and principal components described above.

We then regressed the residuals on the genotype of the top SNP in a linear model. We estimated

the coefficients of determination and reported them as the proportion of phenotypic variance

explained by each locus.

Genetic effects in the elderly cohort

We tested the combined effect of all 182 genome-wide significant ALM loci identified in the middle-

aged cohort in the elderly cohort using the top SNP at each locus. We used PLINK2[48] to extract

genotype dosages for each variant identified in the middle-aged GWAS in the elderly cohort. We

then estimated a 'genetic lean mass score' for each individual using the following procedure. First,

we estimated the contribution of each variant on the phenotype as a product of the SNP effect size

obtained from BOLT-LMM (β, calculated based on the reference allele) and the genotype dosage.

Second, we calculated the 'lean mass score' for each individual as the sum of the products for all

selected variants. We ranked the resulting distribution of lean mass scores in ascending order and

partitioned it into five quantiles. We used ALM without any adjustment (raw ALM) because

estimates of effects size already accounted for sex, whole body fat and leg length differences.

However, since the raw ALM did not meet the assumption of normality, we used a Kruskal-Wallis

test (non-parametrical) to evaluate the difference in the median of the phenotypes between the

quantiles, and a Wilcoxon test (non-parametrical) for pairwise comparisons between quantiles. We

conducted five replicates of a negative control test that consisted of randomly selecting a subset (n

~ 185) of non-significant SNPs in the middle-aged cohort and generating 'lean mass score' as

described above for the elderly cohort; this set of SNPs had MAF > 0.001.

We also aimed to replicate the individual variants effects on the ALM of the elderly cohort.

We checked normality of ALM in the elderly cohort as described for the middle-aged cohort. We

tested a subset of genetic variants (n=17,914,406) selected based on their MAF > 0.001 and

imputation quality > 0.3, and we used the same LMM, fixed covariates, and genome-wide

significance threshold ($P < 5 \times 10^{-8}$) as described for the middle-aged cohort. We conducted a

Fisher's exact test to evaluate if overlapping loci between the middle-aged and elderly cohorts

were significantly different from random. The null hypothesis was rejected at $P < 0.05$ (two-tailed).

Genomic regions tagged by loci

We used the 'biomaRT' package in R[49; 50] to retrieve gene and regulatory element annotations at

the genomic position of each statistically significant SNP ($P < 5 \times 10^{-8}$) and Polyphen 2[51] and

SIFT[52; 53] to predict the functional consequences of each SNP. We retrieved additional information

about the positional candidate genes and their expression levels from Ensembl[54] (release 94 -

October 2018) and the Genotype Tissue Expression Project (GTEx) portal[55] (See Web

Resources).

Stage two: LGSM AIL mouse cohort and GWAS

To maximize QTL detection power, we combined three cohorts of LGSM AIL mice for the second

stage of this study (n = 1,867). The LGSM AIL was initiated by Dr. James Cheverud at Washington

University in St. Louis [56]. Cohort 1 included 490 mice (253 males and 237 females) from LGSM

filial generation 54 (F_{34}). Phenotype data was collected from these mice between 80-102 days of

age. Cohort 2 consisted of 506 male mice (~ 84 days of age) from filial generation 50-54 (F_{50-54}).

Cohort 3 includes 887 mice (447 males and 440 females) from filial generation 50-56 (F_{50-56});

between 64 - 111 days of age. Mice were housed at room temperature (70 - 72°F) at 12:12 h light-

dark cycle, with 1-4 same-sex animals per cage and with *ad libitum* access to standard lab chow

and water.

Mouse traits and genotypes

We collected muscle phenotypes after the animals were sacrificed and frozen. We dissected four

muscles and one long bone (tibia or femur) from each mouse at the Pennsylvania State University

(n = 584) and the University of Aberdeen (n = 1,283). Each of the four muscles exhibits a different

proportion of muscle fiber types and often revealed muscle specific QTLs[6; 7; 19; 27].The dissection

procedure involved defrosting the carcasses and removing the muscles (TA, EDL, gastrocnemius

and soleus) and tibia from the hind limbs under a dissection microscope. We weighed the muscles

to 0.1-mg precision on a Pioneer balance (Pioneer, Ohaus) and measured long bone length of the

hindlimb (mm) using an electronic digital calliper (Powerfix, Profi). We quantile normalized all

LGSM AIL traits before mapping QTLs.

Cohort 1 was genotyped using a custom SNP genotyping array[57]. These SNPs (n=2,965)

were evenly distributed along the autosomes (*Mus musculus* genome assembly MGSCv36

(mm8)). The median distance between adjacent SNPs was 446 Kb, and the maximum was 18 Mb.

Large gaps are due to regions identical by descent between the LG/J and SM/J founders[58]. Cohort

2 was genotyped at 75,746 SNPs (73,301 on the autosomes and 2,386 on X and Y) using the

MEGA Mouse Universal Genotyping Array (MegaMUGA; *Mus musculus*) genome assembly

MGSCv37 (mm9)); after removing SNP markers that are not polymorphic between the LG/J and

SM/J strains we retained 7,168 autosomal SNPs for subsequent analyses. The median distance

between adjacent SNPs was 126.9 Kb and the maximum distance was 15 Mb for all chromosomes

except for chromosomes 8, 10, and 14, which had distances of 19, 16, and 16 Mb, respectively.

We used a conversion tool in Ensembl to convert SNP positions from mm8 and mm9 to *Mus*

musculus genome assembly GRCm38 (mm10). Cohort 3 genotypes were obtained from Gonzales

and colleagues[59]. These genotypes were generated using genotyping by sequencing. This

approach has been recently used and described in detail[19]. Only autosomal SNPs known to be

polymorphic in the LG/J and SM/J founder strains (n=523,027; mm10, build 38) were retained for

subsequent analyses. We combined the genotype data from Cohorts 1-3 using PLINK v.1.9 and

imputed missing genotypes using BEAGLE v.4.1[60]. For these steps, we used a reference panel

obtained from whole genome sequencing data of the LG/J and SM/J strains[58]. Dosage estimates

(expected allele counts) were extracted from the output and used for the GWAS; these estimates

captured the degree of uncertainty from the imputation procedure. To ensure the quality of the

genotype data, we excluded SNP genotypes with MAF < 0.20 (because it is an AIL, almost all

SNPs have MAF > 0.2) and dosage R^2 < 0.70 (dosage R^2 corresponds the estimated squared

correlation between the allele dosage and the "true allele dosage" from the genetic marker, and is

used as a measure of imputation quality). After applying these filters, we retained 434,249 SNPs.

Mouse association analyses

Population structure can potentially lead to a rise of false positive associations[61; 62]. The LMM

approach is used to map QTLs while dealing with confounding effects due to relatedness[57; 63; 64].

We used the LMM method implemented in the software GEMMA (genome-wide efficient mixed-

model association)[65] to analyze the mouse phenotypes. In our LMM model we included the

genotypes, a set of fixed effects described later in this section, and a polygenic effect to deal with

population structure.

The polygenic effect is a random vector which was derived from a multivariate normal

distribution with mean zero and a $n \times n$ covariance matrix $\sigma^2 \lambda K$; where n is the number of samples.

The relatedness matrix K was defined by the genotypes. The two parameters, σ^2 and λ, were

estimated from the data by GEMMA; they represent the polygenic and residual variance

components of the phenotypic variance, respectively.

Relatedness matrix and proximal contamination

We used the genotype data to estimate the relatedness matrix K, which was part of the covariance matrix. Although genotype-based and pedigree-based K matrices yield very similar results[66; 67], we have shown that in general, genotype-based estimates are more accurate[66; 68-70]. We constructed the relatedness matrix as $K = XX'/p$, where X is the genotype matrix of entries x_{ij} and $n \times p$ dimensions, p is the number of SNPs.

The relatedness matrix K was estimated taking into account the potential problem of proximal contamination[67], which involves loss of power due to including genetic markers in multiple components of the LMM equation. Furthermore, because of LD, markers in close proximity to the genetic marker that is being tested can also lead to deflation of the *P* values[59; 68]. To avoid this problem, the K matrix was estimated by excluding from the calculations the SNPs within the chromosome that was analyzed (this approach is termed leave one chromosome out (LOCO)), therefore, K matrix was slightly different for each chromosome.

Genetic and fixed effects

We did not include non-additive effects in the LMMs used for GWAS in the LGSM AIL. Our previous studies[6] suggest that musculoskeletal traits in this population are mostly influenced by additive loci, and by ignoring dominance effects we avoid introducing an additional degree of freedom, hence potentially avoiding a decrease of power to detect QTLs.

To analyze the muscle weights of the combined data, we used four fixed effects in the LMM: sex, dissector of the samples, age, and long bone length of the hindlimb. We selected these variables after using a linear model to estimate their effect on the four muscles; only statistically significant effects were included (P < 0.01). Sex and dissector were included as binary variables; whereas age and long bone were included as continuous variables. Including long bone length of the hindlimb allowed us to capture genetic effects associated with variation in muscle weight *per se* (as opposed to genetic effects on bone length)[19]. In other words, failing to include long bone as a covariate would yield QTLs that are more likely to be genetic contributors to general growth of the

skeleton instead of specifically muscle. We used two bones for the long bone variable, for cohort 1

femur, and for cohorts 2 and 3 tibia. Based on personal communication with Dr. Cheverud, the

femur and tibia bones were found to be positively and highly correlated ($r > 0.8$) in LGSM AIL

(F34). We did not include generation ($r = 1$) and bone type of each cohort ($r = 1$) as fixed effects

since the dissector variable functioned as a proxy for these two variables. Body weight was not

used as a fixed effect because muscle weight accounts for a considerable amount of the body

weight.

SNP heritability

To estimate the SNP heritability or proportion of phenotypic variance explained by all genotypes,

we used the $n \times n$ realized relatedness matrix K, which was constructed using all the available

genotypes. We extracted the SNP heritability from the QTL mapping outputs of the LGSM AIL

cohort described before; GEMMA provides an estimate of the heritability and its standard error[65].

The SNPs available to estimate the heritability do not capture all genetic causal variants, hence the

SNP heritability underestimates the true narrow sense heritability.

Threshold of significance and QTLs intervals

The *P* values estimated from the likelihood ratio test statistic performed by GEMMA were

transformed to $-\log_{10} P$ values. We calculated a threshold to evaluate whether or not a given SNP

significantly contributes to a QTL. We estimated the distribution of minimum *P* values under the

null hypothesis and selected the threshold of significance to be $100(1 - \alpha)^{\text{th}}$ percentile of this

distribution, with $\alpha = 0.05$. In order to estimate this distribution, we randomly permuted phenotypes

1,000 times, as described previously [6; 7; 19; 71]. We did not include the relatedness matrix in the

permutation tests due to computational restrictions, and because, past studies have found that

relatedness does not have a major effect on the permutation test[6; 7].

339   We estimated QTL intervals in three steps. 1) We used Manhattan plots to identify the top

340   SNP within each statistically significant region (SNP with highest $-\log_{10} P$ values), which we refer to

as the peak QTL position. 2) We transformed *P* values from each analysis to LOD scores (base-10

logarithm of the likelihood ratio). 3) We applied the LOD interval function implemented in the r/qtl

package[72] to the regions tagged by each peak SNP, and obtained the QTL start and end positions

based on the 1.5 LOD score interval. 1.5 LOD intervals are commonly used to approximate the ~

95% confidence interval of mouse QTLs[5; 73]. The 1.5 LOD interval estimation is comparable to the

95% CI in the case of a dense marker map[74]; hence, its coverage depends on the location of the

peak QTL marker relative to the adjacent genotyped markers. We estimated the direction of the

QTL effect by calculating the phenotypic mean of each allele based on the peak SNP of each QTL.

We adjusted the phenotypic means and standard errors by fitting the fixed effects used in the

association analyses in a linear model.

We explored the QTL intervals to identify genes that potentially affect hindlimb muscle

mass. We retrieved the genomic location of all genes located within the intervals using the BioMart

database through the 'biomaRT' package implemented in R[49; 50].

Meta-analysis in the LGSM AIL mice

Although we adjusted our GWAS analyses on the LGSM AIL mice for confounding effects, it was

possible that uncontrolled factors could have affected the phenotypes. Therefore, we conducted an

additional meta-analysis on the three LGSM AIL cohorts. We first analyzed each cohort separately

using the same approach as for the combined data, except the dissector variable was not used as

a covariate because it was largely confounded with the cohort. We extracted *P* values, estimated

SNP effects and standard errors at each scanning locus. We considered two popular meta-

analysis approaches: the inverse variance-weighted average and the weighted sum of z-scores[75-

77]. For weighted sum of z-scores, we tried two weighing schemes, i.e., the sample size and

squared root of the sample size, and found the results were very similar, and were slightly better

than that of the inverse variance-weighted average. Therefore, we chose to report the result of the

weighted sum of z-scores with the squared root of the sample size being the weight as suggested

in meta-analysis literature[78]. The test statistic (Z) for each SNP was constructed as follows:

$$Z = \frac{z_1 w_1 + z_2 w_2 + z_3 w_3}{\sqrt{w_1^2 + w_2^2 + w_3^2}}$$

368 where z_i is the z-score that was obtained by transforming the likelihood ratio test P value and w_i is

369 the squared root of the sample size in cohort i=1 to 3. We compared this result with the combined

370 GWAS of the LGSM AIL. This statistical analysis was performed in R[79].

Overlap of mouse and human results

373 The significantly associated muscle QTLs (mice) and lean mass loci (humans) were compared by

374 exploring the genomic regions and genes tagged in each analysis. We used a Fisher's exact test to

375 evaluate whether the number of overlapping loci from the human and mouse analyses exceeded

376 the number expected by chance; the null hypothesis was rejected at $P < 0.05$.

Gene validation using siRNA in C2C12 myoblasts

To validate efficiency of siRNA-mediated gene knockdown, the C2C12 cells were lysed and RNA

isolated using RNeasy mini kit (QIAGEN) following manufacturers recommendations.

Concentration was assessed using NanoDrop (Thermo Scientific) spectrophotometer and ~1.5 μg

of RNA was applied to 1.5% agarose gel to validate its integrity. The cDNA was synthesized using

random primers (Invitrogen) and SuperScript II reverse transcriptase (Invitrogen). Quantitative

PCR for expression of the target *Cpne1*, *Sbf2* and *Stc2* and the reference *Actb* was carried out in

duplicates on LightCycler 480 II (Roche) using SYBR green Master mix (Roche), 10 ng cDNA and

0.5 μM forward and reverse primers (Table S2). Quantification of gene expression was performed

using the comparative Ct method[80].

C2C12 myoblasts, validated for differentiation, were seeded on 8-chamber slides (Lab-Tek

II), batch 1, and 13 mm diameter Thermanox Plastic coverslips (Thermo Fisher Scientific), batch 2,

at 100 cells/mm^2 in high glucose growth medium (D5671, Sigma), containing 10% foetal calf serum

and 2% glutamine. Next day the cells were washed with phosphate-buffered saline (PBS) and

transferred to differentiation medium (D5671, Sigma) supplemented with 10 nM siRNA and

Lipofectamine RNAiMAX (Invitrogen) as per manufacturer protocol. We used the following siRNAs (Life Technologies): negative control #1, s113938 and 93494 (*Cpne1*), 151885 and 151886 (*Stc2*), s115441 and s115442 (*Sbf2*). The treatment achieved expression knockdown by 55-70%. The differentiation medium with 10nM siRNA and Lipofectamine RNAiMAX were replaced once, after 3 days of incubation. After 6 days of incubation, cells were fixed in 4% paraformaldehyde (PFA). We examined 8 cultures for *Stc2* and 12 for the remaining genes (equally divided between the two siRNAs) and negative control that were generated in two batches on separate occasions.

Cells were washed in PBS, fixed in 4% PFA for 15 min, PBS washed again and permeabilized for 6 min with 0.5% Triton X-100 in PBS. The cells were then blocked for 30 min in blocking buffer (10% foetal calf serum in PBS) and incubated overnight at 4 ⁰C with primary anti-myosin heavy chains antibody (Monoclonal Anti-Myosin skeletal, Fast, Clone My-32, Mouse Ascities Fluid, M4276, Sigma-Aldrich) diluted (1:400) in PBS. After three washes in 0.025% Tween-20 in PBS at room temperature, secondary donkey anti-mouse IgG H&L antibody (ab150109, abcam) conjugated to fluorescent dye (Alexa Fluor 488) in PBS (1:400) were applied and incubated for 90 minutes. Following three washes in 0.025% Tween-20 in PBS cells were incubated in 300 nM DAPI in PBS for 15 min. After that cells were covered by coverslip using Mowiol 4-88 (Sigma-Aldrich), sealed with nail polish, and stored at 4 °C in the dark.

Slides were scanned using Axioscan Z1 slide scanner (Zeiss) using X20 magnification. The entire 0.7 cm^2 chamber of a slide or a coverslip was imaged using the wavelength spectrum band of 353-465 nm and 493-517 nm and exposure time 4 ms and 100 ms for DAPI and Alexa Fluor, respectively, at 50% Colibri 7 UV-free LED light source intensity. Alexa Fluor and DAPI channel images of a rectangular area free of artefacts and covering at 14-91% of a chamber of batch 1 and 70% of a coverslip of batch 2 were exported separately for analyses with Fiji[81]. Note that the rectangle area of the majority of batch 1 samples (88%), covered more than 40% of the cell culture. A sensitivity analysis testing the exclusion of small coverage images (14-31%) from the statistical analyses described below, showed results comparable to the analysis of all samples; therefore, we reported significance values (*P* values) corresponding to the statistical analysis of all samples.

Three indices characterizing the effect of treatment on myogenesis were quantified in an

unbiased, automated analysis of the entire exported area: 1) percentage of fluorescent area in the

Alexa Fluor channel, reflecting the level of myosin expression, and 2) the longest-shortest-path

reflecting the length and number of myotubes (Figure S2). The longest-shortest-path analysis was

carried out using the Analyze Skeleton plugin[82] and the shortest path calculation function[83]

implemented in Fiji[81]. We carried out the image analyses on a Linux computer and we allocated

190 GB of RAM for these analyses. The myotube threshold was set at 103.97 μm for batch 1 and

191.63 μm for batch 2, i.e. the mean (batch 1: 54.34 μm, batch 2:100.95 μm) plus 3 standard

deviations (batch 1: SD = 16.54 μm, batch 2: SD = 30.23 μm) of the length of mononucleated and

myosin expressing myocytes (n=35) measured in the negative control #1 cells. The myotube

length variable did not follow normality, therefore quantile normalization was applied to the

variable. All statistical analyses were adjusted for the image area of each sample and batch of

cells, by fitting a linear model on the three indices investigated; all subsequent statistical analyses

were conducted on the residuals, which met the assumptions of normality and homoscedasticity of

residuals. The effect of gene knockdown on these indices was assessed using ANOVA. After, a t-

test was carried out to evaluate the mean differences between the control group and the gene

knockdown groups. In addition, we evaluated the myosin expressing area (as percentage of the

total) present within each knockdown versus control groups using ANOVA.

Data availability

The human data used for this study can be obtained upon application to the UK biobank project[15].

Results

Over 180 genomic loci associated with appendicular lean mass in humans

The appendicular lean mass (ALM) ranged from 12.2 to 41.6 kg and 15.3 to 54.5 kg in healthy

middle age females and males, respectively (Table 1). SNP heritability estimates indicated that

36% of phenotypic variability was due to genetic factors. The GWAS analysis results presented in

Figure 1 revealed 6,693 autosomal variants (MAF > 0.001) associated ($P < 5 \times 10^{-8}$) with ALM

(Table S3). The associated variants tagged 331 genes and 753 regulatory elements. We used the

Functional Mapping and Annotation of Genome-Wide Association Studies (FUMA GWAS[46]) to

define genomic regions containing the associated variants, and we identified 77 of them that were

on average 0.40 Mb long and contained 182 independent signals (Table S4). We refer to the

identified regions and the independent signals as "loci" throughout the text. The 182 loci identified

indicate that ALM is influenced by multiple genetic elements. The LD score intercept that we

estimated during this ALM GWAS (1.05 ± 0.007 (mean $\pm$ SE)) provides further evidence

suggesting polygenicity. Cumulative effects of these loci explained 24% of SNP heritability.

78% of the same loci affect appendicular lean mass in older adults

As expected due to the aging effect on skeletal muscle, the ALM in the cohort of elderly declined

by 4 and 8% in comparison to the middle-age cohort of females and males, respectively ($P < 2\times10^{-16}$

). A more prominent decline in males is consistent with earlier reports[84]. We then used a 'genetic

lean mass score' (see Methods for details) to test if the identified 182 loci contributed to ALM

variability in the elderly population. The genetic lean mass score had a statistically significant

overall effect ($\chi^2 = 376.13$, df = 4, $P = 3.99\times10^{-80}$) on ALM variability in the elderly population

(Figure 2). On average, individuals with the highest genetic lean mass score had 0.73 kg, or 3.2%,

more ALM compared to those with the lowest scores (Figure 2). Negative controls showed no

statistically significant effects (Table S5 and Figure S3).

We also asked if the variants identified in the middle-aged cohort were associated with ALM

in the elderly. A GWAS in the elderly cohort replicated 5,291 variants based on their P value ($P < 5$

$\times 10^{-8}$) and allelic effect (*beta*); moreover, the replicated variants tagged 78% of the ALM loci of the

middle-aged cohort (two tailed Fisher test P value $< 2.2 \times 10^{-16}$). Overall, the set of genomic loci in

the elderly cohort appeared similar to that of the middle-aged adults, with the exception of an

approximately 5 Mb region on chromosome 5 (Figure S4). This region showed a very strong

association with the ALM variability in older adults (lowest P value = 4.50×10^{-56}, *beta* = 0.12 ±

0.01 kg), and had a modest albeit significant association with the ALM of middle-aged individuals

(lowest P value = 3.30×10^{-11}) with an effect size of 0.07 ± 0.01 kg.

23 QTLs contribute to muscle weight variability in LG/J and SM/J strain-derived advanced intercross lines

23 QTLs contribute to muscle weight variability in LG/J and SM/J strain-

derived advanced intercross lines

We examined the weight of four hindlimb muscles of the LGSM AIL (F_{34} and F_{50}-F_{56}): tibialis

anterior (TA), extensor digitorium longus (EDL), gastrocnemius and soleus. The LGSM AIL

muscles showed extensive individual variability (Table 2); furthermore, the SNP heritabilities of the

TA, EDL, gastrocnemius and soleus muscles were 0.39, 0.42, 0.31 and 0.30, respectively (Table

2). The genome mapping of LGSM AIL muscles yielded 23 QTLs ($P < 6.45 \times 10^{-06}$). The TA, EDL

and gastrocnemius QTLs explained more than the 50% of the SNP heritability of each trait (Table

S6). The soleus muscle phenotypic variability explained by QTLs was 23% of its SNP heritability.

Three QTLs were shared between the four hindlimb muscles (chromosome 7, 11 and 13; (Figure

3); the QTL on chromosome 13 resulted in the strongest association (EDL $P = 2.95 \times 10^{-21}$), with

its peak position at 104,435,003 bp, and the percentage of phenotypic variance explained by this

locus was 5.2%; the SM/J allele conferred increased muscle mass (Figure 3). Furthermore, six

QTLs were shared between two or three hindlimb muscles, while fourteen identified QTLs were

only associated with one specific hindlimb muscle (Figure 3).

The mapping resolution was comparable to that attained in the previous study in the LGSM

AIL cohort[59]. On average, mouse QTLs spanned 2.80 Mb (based on the 1.5 LOD interval) and

encompassed 2,267 known genes (Table S7). The median number of genes per QTL was 55;

more than half of the mouse QTLs contained a modest number of genes; however, 7 QTLs

contained more than 100 genes each, and a single QTL located on chromosome 7 as many as 644

genes (Table S6). Although all mouse QTLs identified in the LGSM AIL contained SNPs, at least

seven QTLs covered long genomic regions characterized as identical by descent between the LG/J

and SM/J strains[58]. We also analyzed the LGSM AIL using a meta-analysis approach and identified

14 QTLs that on average were 3.78 Mb long. The majority of the QTLs from the meta-analysis, 12

out of the 14, overlapped with the findings of the mega-analysis (Figure S5). The meta-analysis

results are shown in Table S8.

Interspecies overlap between appendicular lean mass loci and muscle weight

QTLs

The ALM mainly consists of the skeletal muscle of the extremities; however, other tissues also

contribute. To test the hypothesis that ALM-associated genetic variants primarily affect skeletal

muscle mass, we overlaid human ALM finding with the mouse where skeletal muscle was weighed

directly. Specifically, we overlaid the captured genomic regions restricted by the significant SNPs

used in the GWAS of each species. The mouse QTL regions were notably larger, partially due to

the median distance between adjacent genetic markers of 126.9 Kb. Our analysis identified five

syntenic regions associated with ALM in humans and hindlimb muscle mass in mice (Table 3). We

used Fisher's exact test to discover that the number of overlapping regions significantly ($P =$

0.0019; Table S9) exceeded that which could be expected by chance. This analysis permitted us to

shorten the list of positional candidates. Assuming the same causative entity for an overlapping

mouse and human locus, these five loci harbour 38 homologous genes (Table 3). Encouragingly,

four of these five genomic loci tagged by rs148833559, rs9469775, rs4837613 and rs57153895

SNPs (Table S3) were replicated in the ALM of the elderly cohort (Table S10).

Modifiers of *in vitro* myogenesis

We used siRNA-mediated gene knockdown in C2C12 cells to test if candidate genes affected

myogenic differentiation. The *STC2*[85], *CPNE1*[5] and *SBF2*[86] were prioritized for this assay because

they were highlighted by both mouse and human GWAS. We assessed indices of myogenic

differentiation (the number and length of the myotubes, and expression of myosin) of C2C12 cells.

In total, 34,989 myotubes were identified and measured in 44 cell cultures (see Methods for

details). The gene knockdown had a significant effect on myotube length, with *Cpne1 (P = 0.001,*

95% confidence interval = 0.019-0.068, effect size = 0.024) and *Stc2 (P = 0.015, 95% confidence*

interval= 0.007-0.066, effect size = 0.017) showing an increase in length compared to the control

cells (Figure 4). There was no significant difference for the *Sbf2* gene. The pattern of the effect on

myosin expressing area was similar to that of myotube length but was not statistically significant (*P*

= 0.21). The number of myotubes was also unaffected.

Discussion

The key findings of the present report are as follows: 1) we identified a set of over 180 loci

associated with ALM, a substantial expansion in comparison to previous human studies. 2) There

is a substantial overlap of the genetic effects between middle-aged and elderly subjects. 3)

Integration of mouse and human GWAS indicates that skeletal muscle is the primary component

affected by the ALM loci, facilitates prioritization of candidate genes, and helps prediction of their

effect on cellular mechanisms underlying muscle mass variation. 4) *In vitro* studies validated two

genes, *CPNE1* and *STC2*, as modifiers of muscle mass in humans.

We estimated SNP heritability for ALM to be of 0.36, which is lower than heritability

estimates previously reported (0.44) [8], this difference could be due to different fixed effects used to

estimate variance components. In total, we mapped 182 loci that collectively explain 24% of the

SNP heritability of ALM. The most recent report, a meta-analysis of 47 independent cohorts

(dbGAP), comparable in sample size but ranging in subjects aged 18 to 100 years, reported five

significant associations with lean body mass[8]. Even fewer associations were detected in the

earlier, small sample size studies[10; 12-14; 87]. However, our results indicate that ALM is a highly

polygenic trait in humans. We hypothesize that multiple factors contributed to the improved locus

detection in the present GWAS. We restricted subjects' age to a narrow range, 38 to 49 years,

minimizing the effects of the developmental and aging-related processes on phenotypic variance.

The skeletal muscle is a dynamic tissue reaching its peak mass by late 20s, then a trend of decline

emerges after 40s and accelerates about two decades later[1]. An estimated 30-50% decline in

muscle mass can be expected between 40 and 80 years of age[88]. These developmental and

aging-related changes are not linear in progression and therefore would hamper detection of loci

even if accounted for in a linear model. In addition, unlike Zillikens and colleagues[8], the data set

we used was systematically collected as described by the UK Biobank project[15] and we only

employed bioelectric impedance measurements of lean mass. Furthermore, we used a LMM to test

the effects of > 21 million variants (MAF > 0.001), and our analysis was adjusted for a different set

of fixed effects than in previous research[8; 10; 12; 14]. Our analysis captured three loci identified by

Zillikens and colleagues[8], containing the *VCAM, ADAMTSL3* and *FTO* genes, suggesting that their

effects are not influenced by age. We hypothesize that a combination of a homogeneous age

group, the optimized genomic coverage and the method used to conduct this association analysis

contributed to improved detection of loci in the present study.

The analyses presented here shed light into the complex genetic mechanisms behind the

appendicular muscle mass of humans. In the past, concern was expressed about the

reproducibility of association analyses of complex traits; however, an increasing number of human

GWAS have shown that their findings are remarkably reproducible[89]. The present study provides

further support for the reliability of association studies, demonstrating replication of 78% of ALM

loci in the elderly cohort. Furthermore, we show that the genetic profile characterized by depletion

of ALM-increasing alleles leads to a lower ALM in elderly individuals (Figure 2). Hence, it is

conceivable that genetic architecture predisposing individuals to lower muscle mass may lead to

elevated risk of sarcopenia[1].

Combining two experimental models, mouse and human, facilitated prioritization of

candidate genes for functional validation and indicated that skeletal muscle is the primary

component of lean tissues affected by the identified loci. Furthermore, the mouse model revealed

that genetic effects may not all be uniform across skeletal muscle tissue, instead some of the

effects can be muscle type- or muscle-specific. To establish the association between the QTGs of

the identified loci and the muscular phenotype, we focused on the overlapping human and mouse

results. Integration of results from these two species permitted circumvention of the limitations

imposed by the individual models. While human GWAS often identify loci containing single genes,

it is often unclear which tissue is most relevant to the phenotype. Although mouse QTLs often

contain multiple positional candidate genes, mice can be used as experimental models to identify

loci specifically associated with skeletal muscle. In this study, we used a mouse model to show that

the association with hindlimb skeletal muscle mass was specifically related to differences in the

cross-sectional area of the constituent muscle fibers, rather than to the number of fibers in the

muscle. This is because between the two founders of the LGSM AIL, the LG/J strain compared to

the SM/J strain shows over 50% larger cross-sectional area of muscle fibers, but no difference in

the number of fibers in soleus muscle[21]. Hence, it is conceivable that the QTGs of the majority of

the overlapping loci affected hindlimb muscle mass specifically via the hypertrophy of muscle

fibers. Such prioritization between the two cellular mechanisms of muscle mass variability is

important because genes specifically influencing cross-sectional area of muscle fibers can be

targeted pharmacologically to prevent and reverse atrophy of muscle fibers in aging muscle[90]. In

humans, the bone, muscle and skin tissues contribute to lean mass determined by bioelectric

impedance. Approximately 1-2 mm thick skin[91] constitutes a rather minor component of lean mass

compared to the size of human extremities. The long bones determine axial dimensions of a limb

but we adjusted for that to minimize bone effect on variation of lean mass. Whereas the magnetic

resonance imaging (MRI) assessed muscle mass accounts for ~38% of body weight in humans,

and the MRI data strongly and positively correlates with the estimates of bioelectric impedance[92].

This, collectively with the overlap of the ALM loci and mouse muscle QTLs, provides a strong

support for the notion that skeletal muscle is the primary tissue affected by the ALM loci.

For the functional validation we prioritized three candidate genes (*STC2*, *SBF2* and

CPNE1) implicated by both human and mouse analyses. *STC2* had the largest effect size on the

ALM (beta = 0.88 ± 0.13 kg; Table S3) with the minor allele (A) of a missense SNP (rs148833559

(A/C) associated with the increase in ALM. Prediction tools (SIFT[53], PolyPhen[51], CADD[93], and

REVEL[94]) suggested a detrimental consequence of rs148833559 on STC2 structure. *SBF2* has

been linked to Charcot-Marie-Tooth hereditary motor and sensory neuropathy[86], and is expressed

in skeletal muscle and associated with a cis-eQTL[55]. Although little is known about *CPNE1,* it is an

intriguing candidate because the minor allele of the missense variant (rs12481228) is predicted by

SIFT[53] and PolyPhen[52] to be detrimental on the structure of CPNE1. That allele was associated

with increased ALM in the middle-aged cohort, and a frameshift variant (rs147019139) leading to

premature stop codon was also associated with an increase in ALM in the elderly cohort (Table

S10). Furthermore, in a previous GWAS using outbred (CFW) mice[5], *Cpne1* was implicated in

hindlimb muscle mass. To validate these QTGs for their effects on skeletal muscle, we tested the

siRNA-mediated knockdown effect on myogenesis *in vitro*. A knockdown of two genes, *CPNE1* and

STC2, increased the length of the myotubes. Although it is not completely understood how

changes in the indices of *in vitro* myogenesis correlate with the fiber hypertrophy and/or

hyperplasia *in vivo*, our findings implicate an upregulation of myogenic differentiation. We interpret

this *in vitro* observation as being consistent with the allelic effect of the two loci identified in human

GWAS. *CPNE1* encodes for Copine 1, a soluble calcium-dependent membrane-binding protein[95]

expressed in skeletal muscle[24]. *STC2* encodes Stanniocalcin 2, a homodimeric glycoprotein

hormone abundantly expressed in skeletal[55] and cardiac muscle[96], and involved in regulation of

IGF1 through interaction with pregnancy-associated plasma protein-A[97]. A suppressive role of

STC2 is consistent with reduced muscle mass in the STC2 overexpressing mice[85]. Hence, our

analyses and recent reports provide support for *CPNE1* and *STC2* as suppressors of muscle mass

development and/or maintenance in humans.

In conclusion, the present study integrated human and mouse GWAS and used *in vitro*

validation to further interrogate a subset of the genes implicated in both species. Our results

revealed over 180 genomic loci contributing to ALM in middle-aged humans. The effects of the

majority of these loci persist in the elderly human population. Integration of human and mouse data

also highlighted candidate genes affecting skeletal muscle mass in mammals. Two genes, *CPNE1*

and *STC2* were confirmed to be modifiers of *in vitro* myogenesis.

Supplemental Data

Supplemental Data include 5 figures, 10 tables and 1 macro script.

Declaration of interest

The authors declare no competing interests.

Acknowledgements

The authors would like to acknowledge Dr David A. Blizard for his role in the development of the ideas that led to this study and feedback on the manuscript, Professor Helen Macdonald for valuable advice on study design, Dr Leslie R. Noble for help with the UK Biobank data, and Dr Joseph P. Gyekis for help genotyping cohort 2 mice. The authors would like to acknowledge funding from the University of Aberdeen for the Maxwell computer cluster, the Elphinstone and IMS studentship for AIHC; a Schweppe Foundation Career Development Award (AAP), and the NIH (NIAMS (AL: R01AR056280) and NIDA (AAP:R01DA021336, AAP:R21DA024845, AAP:T32MH020065, NMG:F31DA03635803), NIGMS (NMG:T32GM007197), NHGRI (MA:R01 HG002899)).

Web Resources

Functional Mapping and Annotation of Genome-Wide Association Studies (FUMA GWAS)[46]. URL: http://fuma.ctglab.nl/

Ensembl[54]. URL: https://www.ensembl.org

Gene Tissue Expression Project (GTEx) portal[36]. URL: https://gtexportal.org/home/

References

1. Lauretani, F., Russo, C.R., Bandinelli, S., Bartali, B., Cavazzini, C., Di Iorio, A., Corsi, A.M., Rantanen, T., Guralnik, J.M., and Ferrucci, L. (2003). Age-associated changes in skeletal muscles and their effect on mobility: an operational diagnosis of sarcopenia. J Appl Physiol 95, 1851-1860.
2. Janssen, I., Shepard, D.S., Katzmarzyk, P.T., and Roubenoff, R. (2004). The healthcare costs of sarcopenia in the United States. Journal of the American Geriatrics Society 52, 80-85.
3. Kim, J., Wang, Z., Heymsfield, S.B., Baumgartner, R.N., and Gallagher, D. (2002). Total-body skeletal muscle mass: estimation by a new dual-energy X-ray absorptiometry method. Am J Clin Nutr 76, 378-383.
4. Livshits, G., Gao, F., Malkin, I., Needhamsen, M., Xia, Y., Yuan, W., Bell, C.G., Ward, K., Liu, Y., and Wang, J. (2016). Contribution of Heritability and Epigenetic Factors to Skeletal Muscle Mass Variation in United Kingdom Twins. J Clin Endocrinol Metab 101, 2450-2459.
5. Nicod, J., Davies, R.W., Cai, N., Hassett, C., Goodstadt, L., Cosgrove, C., Yee, B.K., Lionikaite, V., McIntyre, R.E., Remme, C.A., et al. (2016). Genome-wide association of multiple complex traits in outbred mice by ultra-low-coverage sequencing. Nature Genetics 48, 912-918.
6. Hernandez Cordero, A.I., Carbonetto, P., Riboni Verri, G., Gregory, J.S., Vandenbergh, D.J., J, P.G., Blizard, D.A., and Lionikas, A. (2018). Replication and discovery of musculoskeletal QTLs in LG/J and SM/J advanced intercross lines. Physiological reports 6.
7. Carbonetto, P., Cheng, R., Gyekis, J.P., Parker, C.C., Blizard, D.A., Palmer, A.A., and Lionikas, A. (2014). Discovery and refinement of muscle weight QTLs in B6 x D2 advanced intercross mice. Physiol Genomics 46, 571-582.
8. Zillikens, M.C., Demissie, S., Hsu, Y.-H., Yerges-Armstrong, L.M., Chou, W.-C., Stolk, L., Livshits, G., Broer, L., Johnson, T., Koller, D.L., et al. (2017). Large meta-analysis of genome-wide association studies identifies five loci for lean body mass. Nat Commun 8, 80.
9. Urano, T., and Inoue, S. (2015). Recent genetic discoveries in osteoporosis, sarcopenia and obesity. Endocrine journal 62, 475-484.
10. Hai, R., Pei, Y.-F., Shen, H., Zhang, L., Liu, X.-G., Lin, Y., Ran, S., Pan, F., Tan, L.-J., and Lei, S.-F. (2012). Genome-wide association study of copy number variation identified gremlin1 as a candidate gene for lean body mass. J Hum Genet 57, 33.
11. Liu, X.G., Tan, L.J., Lei, S.F., Liu, Y.J., Shen, H., Wang, L., Yan, H., Guo, Y.F., Xiong, D.H., Chen, X.D., et al. (2009). Genome-wide association and replication studies identified TRHR as an important gene for lean body mass. Am J Hum Genet 84, 418-423.
12. Guo, Y.-F., Zhang, L.-S., Liu, Y.-J., Hu, H.-G., Li, J., Tian, Q., Yu, P., Zhang, F., Yang, T.-L., and Guo, Y. (2013). Suggestion of GLYAT gene underlying variation of bone size and body lean mass as revealed by a bivariate genome-wide association study. Hum Genet 132, 189-199.
13. Ran, S., Liu, Y.-J., Zhang, L., Pei, Y., Yang, T.-L., Hai, R., Han, Y.-Y., Lin, Y., Tian, Q., and Deng, H.-W. (2014). Genome-wide association study identified copy number variants important for appendicular lean mass. PloS ONE 9, e89776.
14. Urano, T., Shiraki, M., Sasaki, N., Ouchi, Y., and Inoue, S. (2014). Large-scale analysis reveals a functional single-nucleotide polymorphism in the 5′-flanking region of PRDM16 gene associated with lean body mass. Aging cell 13, 739-743.
15. Sudlow, C., Gallacher, J., Allen, N., Beral, V., Burton, P., Danesh, J., Downey, P., Elliott, P., Green, J., Landray, M., et al. (2015). UK biobank: an open access resource for identifying the causes of a wide range of complex diseases of middle and old age. PLoS medicine 12, e1001779.
16. Lynch, N.A., Metter, E.J., Lindle, R.S., Fozard, J.L., Tobin, J.D., Roy, T.A., Fleg, J.L., and Hurley, B.F. (1999). Muscle quality. I. Age-associated differences between arm and leg muscle groups. J Appl Physiol (1985) 86, 188-194.
17. Kallman, D.A., Plato, C.C., and Tobin, J.D. (1990). The role of muscle loss in the age-related decline of grip strength: cross-sectional and longitudinal perspectives. J Gerontol 45, M82-88.

18. Guigo, R., Dermitzakis, E.T., Agarwal, P., Ponting, C.P., Parra, G., Reymond, A., Abril, J.F.,
Keibler, E., Lyle, R., Ucla, C., et al. (2003). Comparison of mouse and human genomes
followed by experimental verification yields an estimated 1,019 additional genes.
Proceedings of the National Academy of Sciences of the United States of America 100,
1140-1145.
19. Parker, C.C., Gopalakrishnan, S., Carbonetto, P., Gonzales, N.M., Leung, E., Park, Y.J.,
Aryee, E., Davis, J., Blizard, D.A., Ackert-Bicknell, C.L., et al. (2016). Genome-wide
association study of behavioral, physiological and gene expression traits in outbred CFW
mice. Nat Genet 48, 919-926.
20. Carroll, A., Cheng, R., Collie-Duguid, E., Meharg, C., Scholz, M.E., Fiering, S.N., Fields, J.L.,
Palmer, A.A., and Lionikas, A. (2017). Fine mapping of genes determining extrafusal fiber
properties in murine soleus muscle. Physiological genomics 49, 141-150.
21. Carroll, A.M., Palmer, A.A., and Lionikas, A. (2012). QTL analysis of type I and type IIA fibers
in soleus muscle in a cross between LG/J and SM/J mouse strains. Frontiers in genetics 2,
99.
22. Partridge, C.G., Fawcett, G.L., Wang, B., Semenkovich, C.F., and Cheverud, J.M. (2014). The
effect of dietary fat intake on hepatic gene expression in LG/J AND SM/J mice. BMC
genomics 15, 1.
23. Cheverud, J.M., Vaughn, T.T., Pletscher, L.S., Peripato, A.C., Adams, E.S., Erikson, C.F., and
King-Ellison, K.J. (2001). Genetic architecture of adiposity in the cross of LG/J and SM/J
inbred mice. Mammalian genome : official journal of the International Mammalian Genome
Society 12, 3-12.
24. Lionikas, A., Meharg, C., Derry, J., Ratkevicius, A., Carroll, A., Vandenbergh, D., and Blizard,
D.A. (2012). Resolving candidate genes of mouse skeletal muscle QTL via RNA-Seq and
expression network analyses. BMC Genomics 13.
25. Goodale, H. (1938). A study of the inheritance of body weight in the albino mouse by selection.
Journal of Heredity 29, 101-112.
26. MacArthur, J.W. (1944). Genetics of Body Size and Related Characters. I. Selecting Small and
Large Races of the Laboratory Mouse. The American Naturalist 78, 142-157.
27. Lionikas, A., Cheng, R., Lim, J.E., Palmer, A.A., and Blizard, D.A. (2010). Fine-mapping of
muscle weight QTL in LG/J and SM/J intercrosses. Physiological genomics 42, 33-38.
28. Darvasi, A., and Soller, M. (1995). Advanced intercross lines, an experimental population for
fine genetic mapping. Genetics 141, 1199-1207.
29. Jackson, A.S., Janssen, I., Sui, X., Church, T.S., and Blair, S.N. (2012). Longitudinal changes
in body composition associated with healthy ageing: men, aged 20-96 years. The British
journal of nutrition 107, 1085-1091.
30. Bycroft, C., Freeman, C., Petkova, D., Band, G., Elliott, L.T., Sharp, K., Motyer, A., Vukcevic,
D., Delaneau, O., O'Connell, J., et al. (2018). The UK Biobank resource with deep
phenotyping and genomic data. Nature 562, 203-209.
31. Howie, B., Marchini, J., and Stephens, M. (2011). Genotype imputation with thousands of
genomes. G3 (Bethesda, Md) 1, 457-470.
32. Wood, A.R., Esko, T., Yang, J., Vedantam, S., Pers, T.H., Gustafsson, S., Chu, A.Y., Estrada,
K., Luan, J., Kutalik, Z., et al. (2014). Defining the role of common variation in the genomic
and biological architecture of adult human height. Nat Genet 46, 1173-1186.
33. Loh, P.-R., Kichaev, G., Gazal, S., Schoech, A.P., and Price, A.L. (2018). Mixed-model
association for biobank-scale datasets. Nature Genetics 50, 906-908.
34. Shrine, N., Guyatt, A.L., Erzurumluoglu, A.M., Jackson, V.E., Hobbs, B.D., Melbourne, C.A.,
Batini, C., Fawcett, K.A., Song, K., Sakornsakolpat, P., et al. (2019). New genetic signals
for lung function highlight pathways and chronic obstructive pulmonary disease
associations across multiple ancestries. Nat Genet 51, 481-493.
35. Lloyd-Jones, L.R., Robinson, M.R., Yang, J., and Visscher, P.M. (2018). Transformation of
Summary Statistics from Linear Mixed Model Association on All-or-None Traits to Odds
Ratio. Genetics 208, 1397-1408.
36. Pulit, S.L., Stoneman, C., Morris, A.P., Wood, A.R., Glastonbury, C.A., Tyrrell, J., Yengo, L.,
Ferreira, T., Marouli, E., Ji, Y., et al. (2019). Meta-analysis of genome-wide association
studies for body fat distribution in 694 649 individuals of European ancestry. Hum Mol
Genet 28, 166-174.

37. Loh, P.-R., Tucker, G., Bulik-Sullivan, B.K., Vilhjalmsson, B.J., Finucane, H.K., Salem, R.M.,
Chasman, D.I., Ridker, P.M., Neale, B.M., Berger, B., et al. (2015). Efficient Bayesian
mixed-model analysis increases association power in large cohorts. Nat Genet 47, 284-
290.
38. Bulik-Sullivan, B.K., Loh, P.R., Finucane, H.K., Ripke, S., Yang, J., Patterson, N., Daly, M.J.,
Price, A.L., and Neale, B.M. (2015). LD Score regression distinguishes confounding from
polygenicity in genome-wide association studies. Nat Genet 47, 291-295.
39. Egner, I.M., Bruusgaard, J.C., Eftestol, E., and Gundersen, K. (2013). A cellular memory
mechanism aids overload hypertrophy in muscle long after an episodic exposure to
anabolic steroids. J Physiol 591, 6221-6230.
40. Bhasin, S., Storer, T.W., Berman, N., Yarasheski, K.E., Clevenger, B., Phillips, J., Lee, W.P.,
Bunnell, T.J., and Casaburi, R. (1997). Testosterone replacement increases fat-free mass
and muscle size in hypogonadal men. The Journal of clinical endocrinology and metabolism
82, 407-413.
41. Karasik, D., Zillikens, M.C., Hsu, Y.H., Aghdassi, A., Akesson, K., Amin, N., Barroso, I.,
Bennett, D.A., Bertram, L., Bochud, M., et al. (2019). Disentangling the genetics of lean
mass. The American journal of clinical nutrition 109, 276-287.
42. Sanchez-Gurmaches, J., and Guertin, D.A. (2014). Adipocytes arise from multiple lineages that
are heterogeneously and dynamically distributed. Nat Commun 5, 4099.
43. The International HapMap Consortium. (2005). A haplotype map of the human genome. Nature
437, 1299-1320.
44. Pe'er, I., Yelensky, R., Altshuler, D., and Daly, M.J. (2008). Estimation of the multiple testing
burden for genomewide association studies of nearly all common variants. Genet Epidemiol
32, 381-385.
45. Loh, P.-R., Bhatia, G., Gusev, A., Finucane, H.K., Bulik-Sullivan, B.K., Pollack, S.J., de Candia,
T.R., Lee, S.H., Wray, N.R., and Kendler, K.S. (2015). Contrasting genetic architectures of
schizophrenia and other complex diseases using fast variance components analysis. Nat
Genet 47, 1385.
46. Watanabe, K., Taskesen, E., van Bochoven, A., and Posthuma, D. (2017). Functional mapping
and annotation of genetic associations with FUMA. Nature Communications 8, 1826.
47. Auton, A., Brooks, L.D., Durbin, R.M., Garrison, E.P., Kang, H.M., Korbel, J.O., Marchini, J.L.,
McCarthy, S., McVean, G.A., and Abecasis, G.R. (2015). A global reference for human
genetic variation. Nature 526, 68-74.
48. Chang, C.C., Chow, C.C., Tellier, L.C., Vattikuti, S., Purcell, S.M., and Lee, J.J. (2015).
Second-generation PLINK: rising to the challenge of larger and richer datasets.
Gigascience 4, 7.
49. Durinck, S., Moreau, Y., Kasprzyk, A., Davis, S., De Moor, B., Brazma, A., and Huber, W.
(2005). BioMart and Bioconductor: a powerful link between biological databases and
microarray data analysis. Bioinformatics (Oxford, England) 21, 3439-3440.
50. Durinck, S., Spellman, P.T., Birney, E., and Huber, W. (2009). Mapping identifiers for the
integration of genomic datasets with the R/Bioconductor package biomaRt. Nature
protocols 4, 1184-1191.
51. Adzhubei, I.A., Schmidt, S., Peshkin, L., Ramensky, V.E., Gerasimova, A., Bork, P.,
Kondrashov, A.S., and Sunyaev, S.R. (2010). A method and server for predicting damaging
missense mutations. Nat Methods 7, 248-249.
52. Flanagan, S.E., Patch, A.M., and Ellard, S. (2010). Using SIFT and PolyPhen to predict loss-of-
function and gain-of-function mutations. Genetic testing and molecular biomarkers 14, 533-
537.
53. Ng, P.C., and Henikoff, S. (2003). SIFT: Predicting amino acid changes that affect protein
function. Nucleic Acids Res 31, 3812-3814.
54. Frankish, A., Vullo, A., Zadissa, A., Yates, A., Thormann, A., Parker, A., Gall, A., Moore, B.,
Walts, B., Aken, B.L., et al. (2017). Ensembl 2018. Nucleic Acids Research 46, D754-D761.
55. Carithers, L.J., Ardlie, K., Barcus, M., Branton, P.A., Britton, A., Buia, S.A., Compton, C.C.,
DeLuca, D.S., Peter-Demchok, J., Gelfand, E.T., et al. (2015). A Novel Approach to High-
Quality Postmortem Tissue Procurement: The GTEx Project. Biopreservation and
biobanking 13, 311-319.

56. Cheverud, J.M., Routman, E.J., Duarte, F., van Swinderen, B., Cothran, K., and Perel, C.
(1996). Quantitative trait loci for murine growth. Genetics 142, 1305-1319.
57. Cheng, R., Lim, J.E., Samocha, K.E., Sokoloff, G., Abney, M., Skol, A.D., and Palmer, A.A.
(2010). Genome-wide association studies and the problem of relatedness among advanced
intercross lines and other highly recombinant populations. Genetics 185, 1033-1044.
58. Nikolskiy, I., Conrad, D.F., Chun, S., Fay, J.C., Cheverud, J.M., and Lawson, H.A. (2015).
Using whole-genome sequences of the LG/J and SM/J inbred mouse strains to prioritize
quantitative trait genes and nucleotides. BMC genomics 16, 415.
59. Gonzales, N.M., Seo, J., Hernandez Cordero, A.I., St. Pierre, C.L., Gregory, J.S., Distler, M.G.,
Abney, M., Canzar, S., Lionikas, A., and Palmer, A.A. (2018). Genome wide association
analysis in a mouse advanced intercross line. Nature Communications 9, 5162.
60. Browning, Brian L., and Browning, Sharon R. (2016). Genotype Imputation with Millions of
Reference Samples. The American Journal of Human Genetics 98, 116-126.
61. Astle, W., and Balding, D.J. (2009). Population structure and cryptic relatedness in genetic
association studies. Statistical Science, 451-471.
62. Price, A.L., Zaitlen, N.A., Reich, D., and Patterson, N. (2010). New approaches to population
stratification in genome-wide association studies. Nature Reviews Genetics 11, 459-463.
63. Yang, J., Zaitlen, N.A., Goddard, M.E., Visscher, P.M., and Price, A.L. (2014). Advantages and
pitfalls in the application of mixed-model association methods. Nature genetics 46, 100-
106.
64. Yu, J., Pressoir, G., Briggs, W.H., Bi, I.V., Yamasaki, M., Doebley, J.F., McMullen, M.D., Gaut,
B.S., Nielsen, D.M., and Holland, J.B. (2006). A unified mixed-model method for
association mapping that accounts for multiple levels of relatedness. Nature genetics 38,
203-208.
65. Zhou, X., and Stephens, M. (2012). Genome-wide efficient mixed-model analysis for
association studies. Nature Genetics 44, 821-824.
66. Parker, C.C., Carbonetto, P., Sokoloff, G., Park, Y.J., Abney, M., and Palmer, A.A. (2014).
High-resolution genetic mapping of complex traits from a combined analysis of F2 and
advanced intercross mice. Genetics 198, 103-116.
67. Cheng, R., and Palmer, A.A. (2013). A simulation study of permutation, bootstrap, and gene
dropping for assessing statistical significance in the case of unequal relatedness. Genetics
193, 1015-1018.
68. Cheng, R., Parker, C.C., Abney, M., and Palmer, A.A. (2013). Practical considerations
regarding the use of genotype and pedigree data to model relatedness in the context of
genome-wide association studies. G3 (Bethesda, Md) 3, 1861-1867.
69. Speed, D., and Balding, D.J. (2015). Relatedness in the post-genomic era: is it still useful?
Nature Reviews Genetics 16, 33-44.
70. Weir, B.S., Anderson, A.D., and Hepler, A.B. (2006). Genetic relatedness analysis: modern
data and new challenges. Nat Rev Genet 7, 771-780.
71. Churchill, G.A., and Doerge, R.W. (1994). Empirical threshold values for quantitative trit
mapping. Genetics 138, 963-971.
72. Broman, K.W., Wu, H., Sen, S., and Churchill, G.A. (2003). R/qtl: QTL mapping in experimental
crosses. Bioinformatics (Oxford, England) 19, 889-890.
73. Manichaikul, A., Dupuis, J., Sen, Ś., and Broman, K.W. (2006). Poor performance of bootstrap
confidence intervals for the location of a quantitative trait locus. Genetics 174, 481-489.
74. Dupuis, J., and Siegmund, D. (1999). Statistical methods for mapping quantitative trait loci from
a dense set of markers. Genetics 151, 373-386.
75. Whitlock, M.C. (2005). Combining probability from independent tests: the weighted Z-method is
superior to Fisher's approach. Journal of evolutionary biology 18, 1368-1373.
76. Lee, C.H., Cook, S., Lee, J.S., and Han, B. (2016). Comparison of Two Meta-Analysis
Methods: Inverse-Variance-Weighted Average and Weighted Sum of Z-Scores. Genomics
Inform 14, 173-180.
77. Borenstein, M., Hedges, L.V., Higgins, J.P., and Rothstein, H.R. (2011). Introduction to meta-
analysis.(John Wiley & Sons).
78. Zaykin, D.V. (2011). Optimally weighted Z-test is a powerful method for combining probabilities
in meta-analysis. Journal of evolutionary biology 24, 1836-1841.

79. R Core Team. (2019). R: a language and environment for statistical computing [online]. R
Foundation for Statistical Computing, Vienna, Austria. In. (
80. Pfaffl, M.W. (2001). A new mathematical model for relative quantification in real-time RT–PCR.
Nucleic acids research 29, e45-e45.
81. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,
S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for
biological-image analysis. Nature Methods 9, 676.
82. Arganda-Carreras, I., Fernández-González, R., Muñoz-Barrutia, A., and Ortiz-De-Solorzano, C.
(2010). 3D reconstruction of histological sections: application to mammary gland tissue.
Microscopy research and technique 73, 1019-1029.
83. Polder, G., Hovens, H., and Zweers, A. (2010). Measuring shoot length of submerged aquatic
plants using graph analysis. In Proceedings of the ImageJ User and Developer Conference
2010, Mondorf-les-Bains, Luxembourg, 27-29 October 2010. pp 172-177.
84. Yamada, M., Moriguch, Y., Mitani, T., Aoyama, T., and Arai, H. (2014). Age-dependent
changes in skeletal muscle mass and visceral fat area in Japanese adults from 40 to 79
years-of-age. Geriatrics & gerontology international 14 Suppl 1, 8-14.
85. Gagliardi, A.D., Kuo, E.Y., Raulic, S., Wagner, G.F., and DiMattia, G.E. (2005). Human
stanniocalcin-2 exhibits potent growth-suppressive properties in transgenic mice
independently of growth hormone and IGFs. American journal of physiology Endocrinology
and metabolism 288, E92-105.
86. Schneider, B.P., Lai, D., Shen, F., Jiang, G., Radovich, M., Li, L., Gardner, L., Miller, K.D.,
O'Neill, A., Sparano, J.A., et al. (2016). Charcot-Marie-Tooth gene, SBF2, associated with
taxane-induced peripheral neuropathy in African Americans. Oncotarget 7, 82244-82253.
87. Liu, X.G., Tan, L.J., Lei, S.F., Liu, Y.J., Shen, H., Wang, L., Yan, H., Guo, Y.F., Xiong, D.H.,
Chen, X.D., et al. (2009). Genome-wide association and replication studies identified TRHR
as an important gene for lean body mass. American journal of human genetics 84, 418-423.
88. Faulkner, J.A., Larkin, L.M., Claflin, D.R., and Brooks, S.V. (2007). Age-related changes in the
structure and function of skeletal muscles. Clinical and experimental pharmacology &
physiology 34, 1091-1096.
89. Marigorta, U.M., Rodríguez, J.A., Gibson, G., and Navarro, A. (2018). Replicability and
Prediction: Lessons and Challenges from GWAS. Trends in Genetics.
90. Brocca, L., McPhee, J.S., Longa, E., Canepari, M., Seynnes, O., De Vito, G., Pellegrino, M.A.,
Narici, M., and Bottinelli, R. (2017). Structure and function of human muscle fibres and
muscle proteome in physically active older men. The Journal of physiology 595, 4823-4844.
91. Arda, O., Goksugur, N., and Tuzun, Y. (2014). Basic histological structure and functions of
facial skin. Clinics in dermatology 32, 3-13.
92. Janssen, I., Heymsfield, S.B., Wang, Z.M., and Ross, R. (2000). Skeletal muscle mass and
distribution in 468 men and women aged 18-88 yr. Journal of applied physiology (Bethesda,
Md : 1985) 89, 81-88.
93. Rentzsch, P., Witten, D., Cooper, G.M., Shendure, J., and Kircher, M. (2019). CADD:
predicting the deleteriousness of variants throughout the human genome. Nucleic Acids
Res 47, D886-d894.
94. Ioannidis, N.M., Rothstein, J.H., Pejaver, V., Middha, S., McDonnell, S.K., Baheti, S., Musolf,
A., Li, Q., Holzinger, E., Karyadi, D., et al. (2016). REVEL: An Ensemble Method for
Predicting the Pathogenicity of Rare Missense Variants. American journal of human
genetics 99, 877-885.
95. Tomsig, J.L., and Creutz, C.E. (2002). Copines: a ubiquitous family of Ca2+-dependent
phospholipid-binding proteins. Cellular and Molecular Life Sciences CMLS 59, 1467-1477.
96. Ishibashi, K., Miyamoto, K., Taketani, Y., Morita, K., Takeda, E., Sasaki, S., and Imai, M.
(1998). Molecular cloning of a second human stanniocalcin homologue (STC2). Biochem
Biophys Res Commun 250, 252-258.
97. Jepsen, M.R., Kloverpris, S., Mikkelsen, J.H., Pedersen, J.H., Fuchtbauer, E.M., Laursen, L.S.,
and Oxvig, C. (2015). Stanniocalcin-2 inhibits mammalian growth by proteolytic inhibition of
the insulin-like growth factor axis. J Biol Chem 290, 3430-3439.

Figure legends

Figure 1. Map of genome associations with the appendicular lean mass (ALM) of humans.

Genome-wide association study (GWAS) on the ALM of middle-aged adults from the UK Biobank.

Significance level is presented on the vertical axis, while the chromosomal position of each genetic

marker is shown on the horizontal axis. Red line across the plot represents the genome-wide

threshold of significance ($P < 5 \times 10^{-8}$). This plot shows the association of variants with MAF >

0.001.

Figure 2. Genetic lean mass score affects the appendicular lean mass (ALM) in elderly humans.

The plot shows the ALM (kg) of the elderly cohort on the vertical axis. The elderly cohort was

ranked by genetic lean mass score and clustered in five quantiles (Q1 to Q5) (horizontal axis). The

average genetic lean mass score ($\pm$ standard error) of each quantile is shown in parenthesis below

the horizontal axis. The overall quantile effect of the genetic lean mass score on ALM was tested

with Kruskal-Wallis test and the resulting P value is presented on the top of horizontal line above

the bars. The ALM median differences between the groups were tested using a Wilcoxon test; the

significance level of each comparison is presented above the horizontal lines with a Holm adjusted

P value.

Figure 3. Muscle weight QTLs identified in mice of the LGSM AIL and density plot of the

genotypes. The circle plot (A) shows from the outer to the inner ring the GWAS of the TA, EDL,

gastrocnemius and soleus muscle weights. Chromosomal position of each SNP is shown in the

outer black circle of the plot; chromosome names are shown outside as "Chr". Dots within each

chromosome space represent the association ($-\log_{10} P\ value$) of each SNP tested. Dotted blue

lines represent the genome-wide threshold ($P < 6.45 \times 10^{-06}$) of significance, and red dots above

the genome-wide threshold are significantly associated SNPs. (B) Plots of the allelic effect of the

Skmw34, *Skmw55* and *Skmw46* QTLs on the EDL muscle mass. These QTLs were identified for

the four muscles investigated. Vertical axis represents the residual muscle mass adjusted for sex,

age, dissector and long bone length of the hindlimb, and the horizontal axis shows the genotypes

(LG/J homozygote, heterozygote and SM/J homozygote). Below the horizontal axis, the number of

individuals with a given genotype is provided. The violin shapes within the plot area represent the

distribution of individuals with the genotypes. Box whiskers represent minimum and maximum

values, distance between a whisker and the top or bottom of the box contains 25% of the

distribution, the box captures 50% of the distribution, and the bold horizontal line represents the

median. Pairwise comparison *P* value (t-test) is shown above horizontal lines at the top of the

plots.

Figure 4. Gene knockdown effect on C2C12 myotube length.

This figure shows the gene knockdown effect of the *Cpne1*, *Sbf2* and *Stc2* genes on myotube

length. The overall effect of the gene knockdown on myotube length was tested using ANOVA and

the resulting *P* value was 0.00017 ($F_{3,\ 34985}$ =6.63). Vertical axis represents the myotube length

(quantile normalised) residuals (adjusted for area analyzed and batch of cells), and the horizontal

axis shows control and knockdown gene groups. Boxes represent the distribution of the myotube

length for each group. Box whiskers represent minimum and maximum values within 1.5-fold

interquartile range above the 75th percentile and below the 25th percentile; the box captures 50% of

the distribution, and the bold horizontal line represents the median value of the myotube length

normalized residuals distribution for each knockdown group. Each red dot represents a single cell

culture sample for each knockdown group. Statistically significant t–test *P* values between control

and knockdown genes are presented above horizontal lines. Effects without a statistically

significant difference between the control and gene knockdown are presented as "ns". *Cpne1* and

Stc2 knockdown groups were not different from each other (*P > 0.05*). *Sbf2* knockdown differed

from *Cpne1 (P = 0.002)* and *Stc2 (P = 0.043).*

Tables

Table 1. Summary of the middle-aged cohort

Trait	N		MIN	MAX	AVERAGE	SD	SNP heritability ± SE
Age (years)	Females =	46,353	39.67	49.00	44.98	2.43	n/a
	Males =	39,397	38.83	49.00	44.89	2.46	
ALM (kg)	Females =	46,307	12.20	41.60	20.05	2.57	0.36 ± 0.003
	Males =	39,353	15.30	54.50	30.17	3.95	
Arm lean mass (kg)	Females =	46,314	1.00	5.10	2.29	0.31	0.32 ± 0.003
	Males =	39,362	1.60	7.10	3.85	0.57	
Leg lean mass (kg)	Females =	46,323	4.60	16.60	7.77	0.98	0.36 ± 0.003
	Males =	39,373	6.20	20.00	11.31	1.42	
WBF (kg)	Females =	46,308	5.00	109.80	25.54	10.63	0.33 ± 0.006
	Males =	39,171	5.00	88.50	21.12	8.29	
Leg (cm)	Females =	46,302	43.00	113.00	76.57	4.27	0.59 ± 0.010
	Males =	39,353	40.00	122.00	83.87	4.68	

Column description from left to right: 1) Trait, 2) Number of records, 3) Minimum value within the distribution of each trait, 4) Maximum value within the distribution of each trait, 5) Average value of each trait, 6) Standard deviation, 7) SNP heritability of the ALM across sex. All summary statistic values were calculated for each sex group. ALM: appendicular lean mass. WBF: whole body fat. n/a: no applicable.

Table 2. Summary of the LGSM AIL muscle traits

Trait	N		MIN	MAX	AVERAGE	SD	SNP heritability ± SE
Tibialis anterior (mg)	Females =	675	26.60	57.20	42.22	5.34	0.39 ± 0.03
	Males =	1,186	31.60	70.80	50.11	6.73	
Extensor digitorum	Females =	675	4.60	10.40	7.52	0.94	0.42 ± 0.03
longus (mg)	Males =	1,184	5.90	13.30	9.31	1.30	
Gastrocnemius (mg)	Females =	675	64.00	133.00	93.15	10.68	0.31 ± 0.03
	Males =	1,187	70.20	174.90	119.32	16.32	
Soleus (mg)	Females =	671	3.20	10.30	6.34	1.18	0.30 ± 0.03
	Males =	1,187	4.00	13.50	7.78	1.64	

997   Column description from left to right: 1) Trait, 2) Number of records, 3) Minimum value within the

998   distribution of each trait, 4) Maximum value within the distribution of each trait, 5) Average or mean

999   value of each trait distribution, 6) Standard deviation of the mean, 7) SNP heritability for each trait

1000   across sex. Summary statistic values were calculated for each sex group.

 Table 3. Syntenic regions between human and mouse QTLs and positional candidate genes

Human locus peak pos	Mouse QTL peak pos (syntenic to human)	Elderly cohort *P*	*Gene symbol*	Human gene name	Differential expression in mouse Soleus
5:64555615	13:104435003	NS	ADAMTS6	ADAM metallopeptidase with thrombospondin type 1 motif 6	0.44
5:172755066	11:31680504	5.6×10^{-11}	STC2	stanniocalcin 2	0.97
6:34335091	17:34968724	7.1×10^{-09}	HLA-B	major histocompatibility complex, class I, B	0.77
			HLA-DQB1	major histocompatibility complex, class II, DQ beta 1	0.76
			BTNL2	butyrophilin like 2	n/a
			TSBP1	testis expressed basic protein 1	n/a
			PBX2	PBX homeobox 2	n/a
			ATF6B	activating transcription factor 6 beta	0.74
			TNXB	tenascin XB	0.63
			C4B	complement C4B (Chido blood group)	0.37
			STK19	serine/threonine kinase 19	0.43
			SKIV2L	Ski2 like RNA helicase	0.83
			NELFE	negative elongation factor complex member E	n/a
			AL645922.1	novel complement component 2 (C2) and complement factor B (CFB) protein	0.99
			C2	complement C2	0.43
			EHMT2	euchromatic histone lysine methyltransferase 2	0.78
			SLC44A4	solute carrier family 44 member 4	n/a
			NEU1	neuraminidase 1	0.15
			HSPA1L	heat shock protein family A (Hsp70) member 1 like	0.04
			LSM2	LSM2 homolog, U6 small nuclear RNA and mRNA degradation associated	0.93
			VARS	valyl-tRNA synthetase	0.95
			VWA7	von Willebrand factor A domain containing 7	n/a
			MSH5	mutS homolog 5	n/a
			CLIC1	chloride intracellular channel 1	0.48
			AL662899.1	novel transcript	n/a
			ABHD16A	abhydrolase domain containing 16A	0.95
			AL662899.2	novel protein	n/a
			CSNK2B	casein kinase 2 beta	0.43
			GPANK1	G-patch domain and ankyrin repeats 1	0.02
			APOM	apolipoprotein M	n/a
			BAG6	BCL2 associated athanogene 6	0.86
			PRRC2A	proline rich coiled-coil 2A	0.5

			ATP6V1G2	ATPase H+ transporting V1 subunit G2	0.88
			DDX39B	DExD-box helicase 39B	0.46
9:119309525	4:65416188	1.2×10^{-08}	ASTN2	astrotactin 2	0.01
11:10322720	7:109218379	6.6×10^{-17}	SBF2	SET binding factor 2	0.76
			ADM	adrenomedullin	n/a
			AMPD3	adenosine monophosphate deaminase 3	0.06

1002   Column description from left to right: 1) ALM Human locus peak position as "chromosome: base

1003   pair position", 2) LGSM QTL peak position as "chromosome: base pair position" (syntenic to

1004   human), 3) Elderly cohort *P* value (NS – not significant), 4) Human gene symbol, 5) Human gene

1005   name, 6) Adjusted *P* value of differential expression between the soleus muscle of the LG/J and

1006   SM/J mouse strains[24].

1007