1
12   A novel virulence phenotype rapidly assesses *Candida* fungal pathogenesis in healthy
13   and immunocompromised *Caenorhabditis elegans* hosts
14
15
16

17   Dorian J. Feistel, Rema Elmostafa, Nancy Nguyen, McKenna Penley, Levi Morran, and
18   Meleah A. Hickman[*]

25   Running title: Fungal infection reduces multiple measures of host fitness
26

27   Department of Biology, Emory University
28   *corresponding author

29
30
31

Abstract

The yeast *Candida albicans* is an opportunistic pathogen of humans, meaning that despite commensal interactions with its host, it can transition to a harmful pathogen. While *C. albicans* is the predominant species isolated in the human mycobiome and implicated in fungal infection, infections due to non-albicans *Candida* species are rapidly rising. Studying the factors that contribute to virulence is often challenging and frequently depends on many contexts including host immune status and pathogen genetic background. Here, we utilize the nematode *Caenorhabditis elegans* as a perspicuous and efficient model host system to study fungal infections of *Candida* pathogens. We find that in addition to reducing lifetime host survival, exposure to *C. albicans* results in delayed reproduction, which significantly reduced lineage growth over multiple generations. Furthermore, we assessed fungal pathogen virulence in *C. elegans* hosts compromised for innate immune function and detected increased early mortality, reduced brood sizes and delayed reproduction relative to infected healthy hosts. Importantly, by assessing virulence in both healthy and immunocompromised host backgrounds we reveal the pathogen potential in non-*albicans Candida* species. We find that *C. tropicalis* is moderately virulent in both host backgrounds whereas *C. dublineinsis* and *C. parapsilosis* show little to no virulence in healthy hosts but significantly elevated virulence in immunocompromised hosts. Taken together, we present a novel lineage growth assay to measure reduction in host fitness associated with fungal infection and demonstrate significant interactions between pathogen and host immune function that contribute to virulence.

Introduction

Candida species are commensals of the human gastrointestinal microbiota and various other niches in the human body [1]. Despite their commensal existence in many humans, *Candida* species also account for the most fungal infections in the world and are the fourth most prevalent cause of all nosocomial blood stream infections [2,3]. The severity of fungal infection is often dependent on the immune status of the host, with superficial infections occurring in healthy individuals and bloodstream infections in immunocompromised hosts [4-8]. The majority of these infections are associated with

Candida albicans, however other non-*albicans Candida* species are increasingly

becoming implicated in fungal infections [9,10].

Several host models have been employed to better understand fungal infection

and disease progression. These experimental systems include mouse, zebrafish,

nematode, wax moth and human *ex vivo* models have been specifically developed to

study *Candida* infections [11]. Most often, virulence phenotypes are limited to survival

outcomes or tissue-specific fungal burden [12], the latter of which requires animal

sacrifice and cannot address the long-term consequence of fungal disease.

Furthermore, animal mortality and sacrifice, particularly in vertebrate models, constrain

the number of infected individuals to study and thus reduce the quantitative robustness

of these experiments. Nonetheless, these studies provide valuable insight into *Candida*

infection and disease and have uncovered important roles for innate immune function of

hosts [13-15], specific *C. albicans* genes regulating virulence [16,17] and differences

between genetic backgrounds within and between *Candida* species [18].

Other major limitations for most studies of *Candida* infection virulence include the

time-intensive nature of tracking virulence phenotypes in vertebrate systems such as

mice, the genetic intractability of zebrafish and wax moth larvae and the context-

independent nature of *ex vivo* models [4]. As such, most studies focus specifically on *C.*

albicans virulence while many non-albicans *Candida* species are evaluated to a lesser

degree. The nematode *Caenorhabditis elegans* has been developed as a model for

Candida infection to overcome these limitations [19,20] and to screen compounds for

antifungal activity [21]. *C. elegans* has proven useful for studying host-microbe

interactions [22] because many fungal and bacterial pathogens that cause illness in

humans also cause disease in *C. elegans* [23]. Infecting *C. elegans* is relatively simple

and often done by replacing or incorporating its standard laboratory food *E. coli* with a

desired pathogen which colonizes the gut and causes disease [23]. Furthermore, the

innate immune system in *C. elegans* includes the *SEK-1* gene which encodes a MAP

Kinase Kinase (MAPKK) that is homologous to the MKK3/6 and MKK4 family of

mammalian MAPKKs and activates the *C. elegans* p38 MAP kinase ortholog [24]. This

pathway has been suggested to be an ancient and conserved component of *C. elegans'*

immune response to pathogens [25]. As such, the *sek-1* mutation increases

susceptibility to microbial colonization [26], including infection with many *Candida*

species [27] and is necessary to induce the appropriate antifungal immune defense [28].

In addition to utilizing innate immune system mutants, most studies measure host

mortality in temperature sensitive sterile mutants [17,21,29,30] to more easily track

mortality in founder populations since *C. elegans* has a rapid lifecycle and large brood

sizes [31,32]. The effects on host reproduction are often overlooked in favor of

examination of mortality rates when assessing a pathogen's virulence [33]. Yet, it is

important to remember that virulence can be broadly measured as any reduction in host

fitness resulting from interactions between a pathogen and its host [34-36]. In this work,

we assessed virulence of *Candida* species in *C. elegans* hosts using a novel measure

of host fitness that incorporated both host survival and fecundity. We found that fungal

pathogens reduced host fitness by delaying reproduction, resulting in long-term

consequences for population growth in both healthy and immunocompromised host

backgrounds. Using this novel measure, we characterized virulence phenotypes for

three non-*albicans Candida* species; *C. dubliniensis*, *C. tropicalis*, and *C. parapsilosis* in

both healthy and immunocompromised *C. elegans* hosts. Our studies demonstrate that

differences in virulence can be identified between pathogen species and that

pathogenic potential is often revealed when host immune function is diminished.

Methods

Strains and media

For this study, the fungal pathogens *C. albicans* (SC5314 [37]), *C. dubliniensis*

(Wu284 [38]), *C. tropicalis* (ATCC 22109), and *C. parapsilosis* (ATCC 22109) strains

were used. *C. elegans* N2 Bristol (WT) and a *sek-1* mutant derivative [24] were used to

test host survival, fecundity and population growth. *C. elegans* populations were

maintained at 20°C on 100mm petri dishes with 25 mL of lite nematode growth medium

(NGM, US Biological) with *E. coli* OP50 as a food source. Nematodes were transferred

to a newly seeded *E. coli* plate every 3-4 days. For survival, fecundity and population

growth assays, NGM was supplemented with 0.08g/L uridine, 0.08g/L histidine, and

0.04g/L adenine to facilitate growth of auxotrophic *C. albicans* strains and 0.2g/L

streptomycin sulfate to inhibit *E. coli* overgrowth so fungal strains could proliferate.

Seeding NGM plates for survival, fecundity, and population growth assays

      Candida strains and *E. coli* OP50 strains were inoculated in 3 mL of YPD or 5 mL of LB, respectively, and cultured at 30°C for 1-2 days. *Candida* culture densities were measured with a spectrophotometer and diluted to a final volume of 3.0 OD_{600} per mL (~6 x 10^7 cells per ml). *E. coli* cultures were pelleted and washed twice with 1 mL of ddH_2O. The supernatant was removed and the pellet was centrifuged for 60 sec at maximum to remove any excess liquid. The pellet was weighed and suspended with sterilized water to a final volume of 200 mg/mL. A mix of 6.25 μL *E. coli*, 1.25 μL *Candida* was brought to a final volume of 50 μL with ddH_2O. The entire 50 μL was spotted onto the center of a 35mm supplemented-NGM Lite agar plate and incubated at room temperature overnight before addition of eggs or transferring nematode. *E. coli* OP50 was used as a control at the same concentration described above.

Egg preparation and synchronization for survival, fecundity and population growth assays

      For survival, fecundity and population growth assays, approximately 100 nematodes at the L3/L4 stage were transferred to a 100mm NGM plate seeded with *E. coli* OP50 and maintained at 20°C for 2-3 days prior to the start of an experiment. On the first day of an experiment, these NGM plates were washed with M9 buffer and contents (live nematodes and eggs) transferred to 15 mL conical tube and pelleted by centrifugation (2 min at 1200 rpm). The pellet was re-suspended in a 1:4 bleach (5.25%) solution and transferred to a micro-centrifuge tube. The suspension was mixed via inversion for 90-120 sec and subsequently centrifuged (30 sec at 1500 rpm). The pellet was washed with 1 mL M9 buffer three consecutive times to remove excess bleach solution and brought to a final suspension with 500 μL M9 buffer. To determine the concentration of eggs, 10 μL was placed on a concaved slide and eggs counted and the egg suspension was diluted with M9 to a final concentration of 10 eggs/μL. All assays were treated equally on the first day (Day 0) by adding roughly 100 eggs to a treatment or *E. coli* plate (described above).

Survival assays

This experimental procedure is a modified version of [39]. Briefly, 72h (Day 3)

after adding ~50-75 nematode eggs to a plate, 40 adult nematodes were randomly

selected and transferred to newly seeded plates with the same concentration of food as

described above and incubated at 20°C. Every other day nematodes were transferred to

freshly seeded plates until all nematode populations went extinct. The number of living,

dead, and censored worms were scored daily. Each survival assay was replicated at

least three independent times.

Fecundity and lineage expansion assays

For both fecundity and lineage expansion assays, 48h (Day 2) after adding

nematode eggs to a plate, a single L3/L4 reproductively immature hermaphroditic

nematode was randomly selected and transferred (6-10 independent biological

replicates per treatment per block) to a newly seeded 35mm petri plate containing either

a treatment of *C. albicans* and *E. coli* or *E. coli* alone (described above) containing one-

fifth the volume of food and incubated at 20°C. *C. elegans* were transferred to freshly

seeded plates in 24 hour time intervals from day two until the end of the experiment.

Eggs remained undisturbed on the plate and were incubated at 20°C for an additional

24 h to provide enough time to hatch, at which the number of viable progeny per day

was scored. Nematodes that died during the assay were scored dead at the time of

transfer. Nematodes that crawled off the plate or were otherwise unaccountable for

were considered censored and excluded from the analysis.

Lineage Expansion Assay

48 hours after adding nematode eggs to a plate (Day 2), a single L3/L4

reproductively immature hermaphroditic nematode was randomly selected and

transferred (6 biological replicates per treatment) to a freshly seeded 100mm treatment

or *E. coli* plate (described above) containing a 6-fold increase in food. On Day 7, each

plate was washed with M9 buffer until the majority of nematodes were displaced and

subsequently transferred to 15mL conical tubes. Tubes were placed at 4°C for 1h to

allow the nematodes to settle at the bottom. All tubes were concentrated to a final

volume of 10mL. Six-20uL samples were taken from each population and counted.

Results

Exposure to C. albicans reduces multiple measures of C. elegans fitness

Pathogens affect host fitness by decreasing lifespan and/or reducing fecundity.

To determine how *C. albicans* impacts *C. elegans* host fitness, we first measured

nematode lifespan in the presence or absence of *C. albicans*. We observed a significant

reduction in survival when *C. elegans* was reared on *C. albicans* compared to when it

was reared in its absence (Figure 1, p < 0.0001 Log-rank test), consistent with

previously published results [39]. 50% mortality was reached in eight days in nematode

populations exposed to *C. albicans* compared to 16 days in unexposed nematode

populations, indicating that *C. elegans* lifespan is substantially decreased by exposure

to *C. albicans*.

Figure 1: Reduced nematode survival with *C. albicans* infection. Survival curves for *C. elegans* populations that are either uninfected (exposed to just an *E. coli* food source, black) or when infected with *C. albicans* SC5314 (red). Mean data values are plotted with SEM error bars, number of worms analyzed (n) for each treatment is indicated in the legend. **** indicates p < 0.0001, Log-rank (Mantel-Cox) test.

While overall survival was reduced, nematode death was rarely observed earlier

than 5 days post-infection. Importantly, adult nematodes produce the majority of their

offspring before *C. albicans* infection reduces rates of survival. Given this information

and that fecundity is a key component of fitness from an evolutionary perspective, we

investigated whether *C. albicans* negatively impacts *C. elegans* reproduction. To test

this, we counted the total number of viable progeny (i.e. brood size) produced in the first

seven days of reproductive maturity from individual nematodes that were either

unexposed or exposed to either heat-killed or live *C. albicans*. As shown in Figure 2A,

exposure to *C. albicans* reduced the average number of viable progeny (267±6)

compared to the unexposed control (286±4; *p* < 0.05, one-way ANOVA) by ~7%. While

statistically significant, the cost of *C. albicans* toward host brood size is modest.

Furthermore, the brood size of nematodes exposed to heat-killed *C. albicans* (HK) was

not detectably different from uninfected or live *C. albicans* treatments (Figure 2A).

Figure 2. Live *C. albicans* reduces nematode fecundity. A) Box and whiskers plot of average brood sizes for *C. elegans* (N2) exposure to *E. coli* food source alone (uninfected, black), heat-killed *C. albicans* (HK, grey), or live *C. albicans* (red). Boxes indicate the 25-75th quartiles with median indicated. Error bars are the normalized range of the data and circles indicate outliers. **B)** Number of viable *C. elegans* progeny produced per day in uninfected, heat-killed and live *C. albicans* treatments. Data represent the mean and error bars indicate SEM. **C)** The fraction of *C. elegans* progeny produced after Day 3 during the 'Late Reproductive' window. Data represent the mean and error bars indicate SEM. **D)** Box and whiskers plot of average population size (representing the number of F1 and F2 progeny) produced from a single founder *C. elegans*. Boxes indicate the 25-75th quartiles with median indicated. Error bars are the normalized range of the data and circles indicate outliers. The number (n) of experimental samples analyzed is indicated for each treatment. * p < 0.05, **** p < 0.0001; one-way ANOVA with Tukey's multiple comparison test.

Intriguingly, we identified a significant delay in reproductive timing when we

measured the number of progeny produced every 24 hrs during the host reproductive

window. We observed a large reduction in reproduction on Day 3 in nematodes

exposed to *C. albicans* (102±4) relative to unexposed hosts (174±4) (Figure 2B; *p* <

0.001 two-tailed t-test) and increased progeny produced on Days 4-6 in nematodes

exposed to *C. albicans* (Figure 2B). We calculated the total fraction of reproduction

occurring in this 'late' window (Days 4-6) and identified a significant increase in the

progeny produced late in *C. elegans* exposed to *C. albicans* compared to unexposed

controls (Figure 2C; *p*=0.0205 two-tailed t-test). Taken together, our data indicate that

C. albicans severely delays and reduces reproduction, in addition to impacting overall

survival in *C. elegans.*

To investigate whether the reproductive delay observed in nematodes exposed

to *C. albicans* had any long-term consequences, we measured the total progeny

produced from a single founder worm over multiple generations through lineage

expansion assays. From these experiments, we find that a single nematode can

produce ~28,500 F1 and F2 progeny within seven days (Figure 2D). Lineage expansion

is significantly impacted in nematodes exposed to live *C. albicans*, with an average

population size of ~18,000 F1 and F2 progeny, a reduction of ~35% compared to

uninfected and heat-killed control treatments (Figure 2D, $p < 0.0001$ one-way ANOVA).

We conclude that delays in reproduction resulting from exposure to *C. albicans* severely

impacts *C. elegans* evolutionary fitness, as we detect a dramatic reduction in population

size in a timeframe that is unlikely to be severely impacted by mortality (Figure 1) nor

the modest reduction in total brood size (Figure 2A).

Immunocompromised hosts are susceptible to fungal infection

C. albicans is an opportunistic pathogen meaning that it only causes disease

under certain conditions and host contexts. A leading host-related risk factor for invasive

candidiasis is compromised immune function [40]. Given our novel results regarding

fecundity in a healthy *C. elegans* host, we were curious how exposure to *C. albicans*

impacts *C. elegans* hosts with compromised immune function. To address this point, we

utilized a *C. elegans* strain background carrying a mutation in *SEK*-1, a well-conserved

MAP kinase involved in the innate immune signaling cascade [24,26] and antifungal

response [28]. We detected substantial mortality in *sek-1* immunocompromised hosts

during the course of our seven-day fecundity assays, which we did not observe with

healthy wild-type hosts (data not shown). Furthermore, exposure to *C. albicans* resulted

in the largest amount of early mortality in immunocompromised hosts compared to heat-

killed *C. albicans* treatments and unexposed controls (Figure 3A). In addition to

increased early mortality, *C. albicans* exposure reduced average brood size by nearly

50% (84±6) in immunocompromised hosts compared to uninfected (157±5) and heat-

killed (127±13) controls (Figure 3B, p < 0.0001, one-way ANOVA). It is important to note

that in all treatments (unexposed, heat-killed, and live *C. albicans*),

immunocompromised hosts produced significantly less progeny than healthy hosts

(Table S1), but even more so when exposed to live *C. albicans*.

Figure 3. Live *C. albicans* severely impacts host fitness in immunocompromised nematodes. A) Percent mortality observed on or by Day3 in immunocompromised (*sek-1*) hosts exposed to *E. coli* food source alone (unin-fected, black), heat-killed *C. albicans* (HK, grey), or live *C. albicans* (red). **B)** Box and whiskers plot of average brood sizes of *sek-1 C. elegans* treated with the same conditions as (A). Boxes indicate the 25-75th quartiles with median indicated. Error bars are the normalized range of the data and circles indicate outliers. **C)** The fraction of *sek-1 C. elegans* progeny produced after Day3 in the 'Late Reproductive' window. Data represent the mean and error bars indicate SEM. D) Box and whiskers plot of average population size (representing the number of F1 and F2 progeny) produced from a single founder *sek-1 C. elegans*. Boxes indicate the 25-75th quartiles with median indicated. Error bars are the normalized range of the data and circles indicate outliers. The number (n) of experimental samples analyzed is indicated for each treatment. * p < 0.05, ** p < 0.01, *** p <0.001, **** p < 0.0001; one-way ANOVA with Tukey's multiple comparison test.

One explanation for the overall reduction in average brood size is the increased

early mortality we observe in immunocompromised hosts. To address this possibility,

we re- analyzed brood size only from hosts that survived past Day3 and found that while
this did slightly increase average brood size across all treatments, it was not significant
between analyses (p=0.2356 two-tailed t-test) and exposure to *C. albicans* still
significantly reduced average brood size (94±6) relative to uninfected (163±5) and heat-
killed controls (141±9) (Figure S1, *p* < 0.0001 one-way ANOVA). For hosts that survived
past three days, we also detected a significant delay in reproductive timing upon
exposure to *C. albicans*, with nearly 45% of all reproduction occurring in this late
window (Figure 3C), similar to what we observe in healthy hosts (Table S1).
We predicted that lineage expansion in immunocompromised hosts exposed to
C. albicans would be more dramatically impacted relative to wild-type nematodes
exposed to *C. albicans*, given the reproductive delays but also early mortality, and
overall reduced brood sizes observed. Indeed, we determined that the average
population size was reduced by ~45% in immunocompromised hosts exposed to *C.*
albicans compared to uninfected or heat-killed controls (Figure 3D). Furthermore, this
reduction in population size is significantly greater in immunocompromised hosts
compared to healthy hosts (Figure S2, p = 0.035 two-tailed t-test). Interestingly, we also
detected a significant reduction in population size in immunocompromised hosts
exposed to heat-killed *C. albicans* relative to uninfected immunocompromised hosts.
This reduction in population size in heat-killed treatments may result from the increased
early mortality in immunocompromised nematodes. Taken together, our results indicate
that immunocompromised hosts are more susceptible to fungal infection than healthy
wild-type hosts.
283
Host immune status reveals pathogenic potential in other Candida species
While *C. albicans* is the predominant fungal agent in invasive candidiasis, non-
albicans Candida species (NACs) are estimated to account for 35-65% of
candidaemia [41]. Here, we measured the impact on host fitness of three additional
Candida species: *C. dubliniensis*, *C. tropicalis* and *C. parapsilosis* analyzing both
healthy and immunocompromised *C. elegans* hosts. By evaluating NACs in both healthy
and immunocompromised hosts, we can begin to assess their pathogenic potential, and
thus extend the utility of *C. elegans* as a model host system for detecting virulence in

Candida species. For healthy hosts infected with *C. dubliniensis* and *C. tropicalis,* we

detected modest levels of early mortality, and we detected no early mortality in healthy

hosts infected with *C. parapsolsis*. However, there was increased early mortality in

immunocompromised hosts exposed to all three *Candida* species, with the highest

mortality observed for *C. parapsilosis* (Figure 4A). In regards to early mortality, we

determined that *C. parapsilosis* has high virulence in immunocompromised hosts, but

none in healthy hosts.

Figure 4. Non-*albicans Candida* species differentially impact host fitness depending on host immune status. A) Percent mortality observed on or by Day3 in N2 (healthy, black) or *sek-1* ('immunocompromised,' grey) *C. elegans* hosts exposed to *C. dubliniensis, C. tropicalis* and *C. parapsilosis*. **B)** Average brood sizes of N2 and *sek-1 C. elegans* hosts exposed to *C. dubliniensis, C. tropicalis* and *C. parapsilosis*. Data represent the mean and error bars indicate SEM. **C)** The fraction of N2 and *sek-1 C. elegans* progeny produced after Day 3 in the 'Late Reproductive' window when exposed to *C. dubliniensis, C. tropicalis* and *C. parapsilosis*. Data represent the mean and error bars indicate SEM. Refer to Table S1 for experimental sample sizes (n). * p < 0.05, ** p < 0.01, *** p <0.001, **** p < 0.0001; one-way ANOVA with Tukey's multiple comparison test.

We also detected different pathogen virulence levels depending on host immune

function for average brood size (Figure 4B) and reproductive delays (Figure 4C). In

healthy hosts, *C. dubliniensis* had no significant impact on average brood size (276±17)

or reproductive timing (21% late) compared to uninfected controls (286±5 brood size

and 21% late). However, *C. dubliniensis* was much more virulent in

immunocompromised hosts, where we observed a 40% reduction in total brood size

(94±17, Figure 4B) and increased reproductive delays compared to uninfected hosts

(Figure 4C). A similar pattern is observed for *C. parapsilosis*, in which healthy hosts

were not significantly impacted in regards to reproductive timing (17% late, Figure 4B)

and only modest reductions in average brood size (224±9) were observed compared to

uninfected healthy hosts. However, *C. parapsilosis* infection greatly reduced average

brood size (38±13) and delayed reproductive timing (72% late) in immunocompromised

hosts.

Not all *Candida* species exhibited elevated virulence in immunocompromised

hosts. For example, *C. tropicalis* reduced average brood size by about 37% in healthy

hosts (180±25) and 34% in immunocompromised hosts (105±16) (Figure 4B) compared

to uninfected treatments. Furthermore, we saw similar significant increases in

reproductive delays in healthy (32% late) and immunocompromised (34%) hosts (Figure

4C). If we compare these three *Candida* species to each other, we see significant

differences for both average brood size (Figure 4B, *p*=0.0012 and *p*=0.015 comparing

healthy and immunocompromised hosts, respectively, one-way ANOVA) and

reproductive timing in immunocompromised hosts (Figure 4C, p = 0.0029 one-way

ANOVA), but not in healthy hosts (p = 0.0573 one-way ANOVA). In healthy hosts, *C.*

tropicalis significantly reduced average brood size more than *C. dubliniensis,* but in

immunocompromised hosts, no significant difference is detected. In

immunocompromised hosts, *C. parapsilosis* displayed significantly more virulence in

regards to both average brood size and reproductive timing compared to *C. tropicalis*

and *C. dubliniensis*. Given the virulence phenotypes of these yeasts, our results

suggest that fungal pathogenicity depends not only on the species of pathogen but also

on the immune status of the host.

Discussion

Here we utilized a *C. elegans* experimental host system to identify novel

measures of host fitness associated with fungal infection. By tracking the number of

progeny produced per day, we can quickly and quantitatively assess three aspects of

host fitness: early mortality, total viable offspring produced and reproductive timing. We

show that exposure to *Candia albicans* in healthy *C. elegans* hosts not only reduces

survival (Figure 1), it modestly reduces total progeny produced, and dramatically delays

reproductive timing (Figure 2). This reproductive delay has long-term consequences for

lineage expansion, as single founder nematodes exposed to *C. albicans* reduces its

population growth by ~30% (Figure 2D). We found similar delays in reproductive timing

in immunocompromised hosts exposed to *C. albicans*, although immunocompromised
hosts are more susceptible to infection and have higher incidence of early mortality and
smaller brood sizes compared to healthy hosts (Figure 3). Importantly, the delayed
reproduction phenotype in infected *C. elegans* is easily ascertained, highly quantitative
and reproducible and can be used as a screening method to detect relatively small
differences in virulence across *Candida* strains or other fungal species. Furthermore, by
utilizing different host contexts, this assay revealed the pathogenic potential of other
important, yet understudied, *Candida* species (Figure 4), including *C. parapsilosis*,
which has dramatically higher levels of virulence in immunocompromised hosts
compared to healthy hosts.
The reductions in host fitness associated with *C. albicans* and other *Candida*
species indicate that we have established a bona fide model of fungal infection.
Importantly, these reductions in host fitness cannot be attributed to *Candida* species
being a sub-optimal food source or that the hosts are starving because we deliver the
pathogen with *E. coli*, the standard food source for *C. elegans*. Further, we do not
detect any host larval development into dauer, an alternative lifestage in response to
starvation and overpopulation [32]. While pathogen avoidance is a common defense
strategy for *C. elegans* [42], we censored any worms that have crawled off the plate or
have disappeared during the course of our experiment and removed them from our data
analysis. Furthermore, the reductions in host fitness depend on ingesting live *C.*
albicans, as heat-killed treatments do not cause significant reductions in total brood
size, reproductive timing, or lineage expansion (Figure 2, 'HK'). Microscopic analysis
reveals that fungal cells inhabit the gut of *C. elegans* ([39] and data not shown) and we
can extract viable host-associated fungal cells (data not shown). Taken together, these
data support a model of fungal infection in the host *C. elegans*.
The host reproductive delay we observed in *C. elegans* upon fungal infection is a
robust, highly quantitative measure of virulence that makes it amenable to screen a
variety of host and pathogen genetic backgrounds that has been challenging in
mammalian and insect models. *C. elegans* is easily maintained in the lab, has a short
lifecycle that generates a large number of progeny, and has been a fundamental model
genetic organism [32]. Previous work has shown that upon exposure to *C. albicans, C.*

elegans have a transcriptional response consistent with that of infection [28]. In this

work, we utilized a mutant *C. elegans* strain in which *SEK-1*, an important MAPK in

innate immunity and whose homologs include the MKK3/6 and MKK4 family of

mammalian MAPKKs, to determine the severity of fungal infection in

immunocompromised hosts. We demonstrate that deficiencies in innate immune

function result in hosts with high susceptibility to fungal infection and in the future can

extend this analysis for many other host backgrounds that are mutant for immune

function.

Previous studies using *C. elegans* as a host for fungal infection have described

additional phenotypes beyond reduced survival, including a deformed anal region (DAR)

[39,43]. It is possible that DAR may be contributing to the delayed reproductive

phenotype we observe in our experiments, however, the frequency of DAR phenotypes

was low and we did not observe any detectable differences between hosts exposed to

C. albicans and hosts unexposed (data not shown), nor did we observe dramatic

reductions in total number of viable offspring produced between these two groups of

hosts (Figure 2A), indicating that exhibiting a DAR phenotype does not necessarily

impede egg laying. While this deformity is a result of a local defense reaction of the

worm due to extracellular signal-regulated kinase activation of the innate immunity

MAKK cascade, it is a distinct marker from delayed reproduction for infection [44].

Wildtype reproductive timing utilizes the highly conserved DAF-2 DAF-16 insulin-like

signaling pathway [23], but has also been implicated in innate immunity and bacterial

infection [45] and is potentially disrupted in *C. elegans* hosts infected with *Candida*.

Impacts on host reproduction have often been overlooked in studies regarding

Candida virulence, yet host reproduction is an important evolutionary measure of an

organism's fitness. Not only can we utilize these fecundity assays to assess the roles of

host immune function on fungal pathogenesis, we also readily screen diverse pathogen

strain backgrounds and species. A large-scale, international 10-year study identified 31

species of *Candida* associated with clinical samples. While *C. albicans* is still the most

prevalent, its isolation from clinical isolates is decreasing with corresponding increases

isolation of *C. glabrata, C. tropicalis* and *C. parapsilosis* [46]. Despite its close

evolutionary relationship with *C. albicans*, *C. dubliniensis* does not seem to be highly

pathogenic [47]. However, other non-*albicans Candida* species have been increasingly
implicated in fungal infections of humans [48]. For example, *C. tropicalis* and *C.*
parapsilosis are common fungal species isolated in patients with candidemia (7 - 48%
and 11 - 27% respectively and depending on geographic region), candiduria (8 – 44%
and 0.5 – 11% respectively, and oral candidosis (5 – 13% and 7 – 15%, respectively)
([48] and references therein). Despite the increasing incidence of non-*albicans Candida*
infections [46], experimental studies using these pathogens remains limited. Here, we
analyzed the virulence phenotypes of three non-albicans *Candida* species, *C.*
dubliniensis, *C. tropicalis* and *C. parapsilosis* in both healthy and immunocompromised
host backgrounds. In healthy *C. elegans* hosts, *C. dubliniensis* is the least virulent; with
no significant differences in brood size or reproductive timing compared to uninfected
hosts. *C. parapsilosis* is moderately virulent in healthy hosts; there is no reproductive
delay, but reduced brood sizes are detected. For both *C. dubliniensis* and *C. tropicalis*,
virulence is more severe in immunocompromised hosts. In contrast, *C. tropicalis* has
similar degrees of virulence between healthy and immunocompromised host
backgrounds. Together, our results regarding virulence in NACs are consistent with
published case reports where there is higher incidence of isolating *C. tropicalis* in fungal
infections affecting patients with healthy immune function (oral candidosis and
candiduria) and infections affecting patients with compromised immune function
(candidemia), whereas we see significant *C. parapsilosis* virulence only in
immunocompromised hosts. Therefore, the *C. elegans* fecundity assays we developed
in this study can reveal the pathogenic potential by assessing virulence in multiple host
genetic backgrounds and across pathogens.
426
427   Literature Cited
428

1. Underhill, D. M.; Iliev, I. D. The mycobiota: interactions between commensal fungi
and the host immune system. *Nature reviews. Immunology* **2014**, *14*, 405–416.
2. Pappas, P. G.; Lionakis, M. S.; Arendrup, M. C.; Ostrosky-Zeichner, L.; Kullberg, B. J.
Invasive candidiasis. *Nat Rev Dis Primers* **2018**, *4*, 18026.

3. Edmond, M. B.; Wallace, S. E.; McClish, D. K.; Pfaller, M. A.; Jones, R. N.; Wenzel,
R. P. Nosocomial bloodstream infections in United States hospitals: a three-year
analysis. *Clin Infect Dis* **1999**, *29*, 239–244.
4. MacCallum, D. M. Hosting infection: experimental models to assay Candida
virulence. *Int J Microbiol* **2012**, *2012*, 363764–12.
5. Pfaller, M. A. Nosocomial candidiasis: emerging species, reservoirs, and modes of
transmission. *Clin Infect Dis* **1996**, *22 Suppl 2*, S89–94.
6. Coleman, D. C.; Sullivan, D. J.; Bennett, D. E.; Moran, G. P.; Barry, H. J.; Shanley, D.
B. Candidiasis: the emergence of a novel species, Candida dubliniensis. *AIDS* **1997**,
11, 557–567.
7. Kontoyiannis, D. P.; Vaziri, I.; Hanna, H. A.; Boktour, M.; Thornby, J.; Hachem, R.;
Bodey, G. P.; Raad, I. I. Risk Factors for Candida tropicalis fungemia in patients with
cancer. *Clin Infect Dis* **2001**, *33*, 1676–1681.
8. Clark, T. A.; Slavinski, S. A.; Morgan, J.; Lott, T.; Arthington-Skaggs, B. A.; Brandt, M.
E.; Webb, R. M.; Currier, M.; Flowers, R. H.; Fridkin, S. K.; Hajjeh, R. A.
Epidemiologic and molecular characterization of an outbreak of Candida parapsilosis
bloodstream infections in a community hospital. *J. Clin. Microbiol.* **2004**, *42*, 4468–
4472.
9. Pfaller, M. A.; Diekema, D. J.; Jones, R. N.; Sader, H. S.; Fluit, A. C.; Hollis, R. J.;
Messer, S. A.; SENTRY Participant Group International surveillance of bloodstream
infections due to Candida species: frequency of occurrence and in vitro
susceptibilities to fluconazole, ravuconazole, and voriconazole of isolates collected
from 1997 through 1999 in the SENTRY antimicrobial surveillance program. *J. Clin.
Microbiol.* **2001**, *39*, 3254–3259.
10. Hajjeh, R. A.; Sofair, A. N.; Harrison, L. H.; Lyon, G. M.; Arthington-Skaggs, B. A.;
Mirza, S. A.; Phelan, M.; Morgan, J.; Lee-Yang, W.; Ciblak, M. A.; Benjamin, L. E.;
Sanza, L. T.; Huie, S.; Yeo, S. F.; Brandt, M. E.; Warnock, D. W. Incidence of
bloodstream infections due to Candida species and in vitro susceptibilities of isolates
collected from 1998 to 2000 in a population-based active surveillance program. *J.
Clin. Microbiol.* **2004**, *42*, 1519–1527.

11. Harwood, C. G.; Rao, R. P. Host pathogen relations: exploring animal models for
fungal pathogens. *Pathogens* **2014**, *3*, 549–562.

12. Conti, H. R.; Huppler, A. R.; Whibley, N.; Gaffen, S. L. Animal models for
candidiasis. *Curr Protoc Immunol* **2014**, *105*, 19.6.1–17.

13. Verma, A.; Gaffen, S. L.; Swidergall, M. Innate Immunity to Mucosal Candida
Infections. *J Fungi (Basel)* **2017**, *3*, 60.

14. Naglik, J. R.; König, A.; Hube, B.; Gaffen, S. L. Candida albicans-epithelial
interactions and induction of mucosal innate immunity. *Curr Opin Microbiol* **2017**, *40*,
104–112.

15. Richardson, J. P.; Moyes, D. L.; Ho, J.; Naglik, J. R. Candida innate immunity at the
mucosa. *Seminars in Cell & Developmental Biology* **2018**.

16. Lu, Y.; Su, C.; Solis, N. V.; Filler, S. G.; Liu, H. Synergistic regulation of hyphal
elongation by hypoxia, CO_2, and nutrient conditions controls the virulence of
Candida albicans. *Cell Host Microbe* **2013**, *14*, 499–509.

17. Pukkila-Worley, R.; Peleg, A. Y.; Tampakakis, E.; Mylonakis, E. Candida albicans
hyphal formation and virulence assessed using a Caenorhabditis elegans infection
model. *Eukaryotic Cell* **2009**, *8*, 1750–1758.

18. Hirakawa, M. P.; Martinez, D. A.; Sakthikumar, S.; Anderson, M. Z.; Berlin, A.; Gujja,
S.; Zeng, Q.; Zisson, E.; Wang, J. M.; Greenberg, J. M.; Berman, J.; Bennett, R. J.;
Cuomo, C. A. Genetic and phenotypic intra-species variation in Candida albicans.
Genome Res **2015**, *25*, 413–425.

19. Powell, J. R.; Ausubel, F. M. Models of Caenorhabditis elegans infection by
bacterial and fungal pathogens. *Methods Mol Biol* **2008**, *415*, 403–427.

20. Aballay, A.; Ausubel, F. M. Caenorhabditis elegans as a host for the study of host-
pathogen interactions. *Curr Opin Microbiol* **2002**, *5*, 97–101.

21. Breger, J.; Fuchs, B. B.; Aperis, G.; Moy, T. I.; Ausubel, F. M.; Mylonakis, E.
Antifungal chemical compounds identified using a C. elegans pathogenicity assay.
PLoS Pathog **2007**, *3*, e18.

22. Sifri, C. D.; Begun, J.; Ausubel, F. M. The worm has turned--microbial virulence
modeled in Caenorhabditis elegans. *Trends Microbiol.* **2005**, *13*, 119–127.

23. Irazoqui, J. E.; Urbach, J. M.; Ausubel, F. M. Evolution of host innate defence:

insights from Caenorhabditis elegans and primitive invertebrates. *Nature reviews.*

Immunology **2010**, *10*, 47–58.

24. Kim, D. H.; Feinbaum, R.; Alloing, G.; Emerson, F. E.; Garsin, D. A.; Inoue, H.;

Tanaka-Hino, M.; Hisamoto, N.; Matsumoto, K.; Tan, M.-W.; Ausubel, F. M. A

conserved p38 MAP kinase pathway in Caenorhabditis elegans innate immunity.

Science **2002**, *297*, 623–626.

25. Ermolaeva, M. A.; Schumacher, B. Insights from the worm: The C. elegans model

for innate immunity. *Seminars in Immunology* **2014**, *26*, 303–309.

26. Youngman, M. J.; Rogers, Z. N.; Kim, D. H. A Decline in p38 MAPK Signaling

Underlies Immunosenescence in Caenorhabditis elegans. *PLOS Genet* **2011**, *7*,

e1002082.

27. Ortega-Riveros, M.; De-la-Pinta, I.; Marcos-Arias, C.; Ezpeleta, G.; Quindós, G.;

Eraso, E. Usefulness of the Non-conventional Caenorhabditis elegans Model to

Assess Candida Virulence. *Mycopathologia* **2017**, *182*, 785–795.

28. Pukkila-Worley, R.; Ausubel, F. M.; Mylonakis, E. Candida albicans Infection of

Caenorhabditis elegans Induces Antifungal Immune Defenses. *PLoS Pathog* **2011**,

7, e1002074–13.

29. Moy, T. I.; Ball, A. R.; Anklesaria, Z.; Casadei, G.; Lewis, K.; Ausubel, F. M.

Identification of novel antimicrobials using a live-animal infection model. *Proc Natl*

Acad Sci USA **2006**, *103*, 10414–10419.

30. Peleg, A. Y.; Tampakakis, E.; Fuchs, B. B.; Eliopoulos, G. M.; Moellering, R. C.;

Mylonakis, E. Prokaryote-eukaryote interactions identified by using Caenorhabditis

elegans. *Proc Natl Acad Sci U S A* **2008**, *105*, 14585–14590.

31. Muschiol, D.; Schroeder, F.; Traunspurger, W. Life cycle and population growth rate

of Caenorhabditis elegans studied by a new method. *BMC Ecol.* **2009**, *9*, 14.

32. Corsi, A. K.; Wightman, B.; Chalfie, M. A Transparent Window into Biology: A Primer

on Caenorhabditis elegans. *Genetics* **2015**, *200*, 387–407.

33. Abbate, J. L.; Kada, S.; Lion, S. Beyond Mortality: Sterility As a Neglected

Component of Parasite Virulence. *PLoS Pathog* **2015**, *11*, e1005229.

34. Read, A. F. The evolution of virulence. *Trends Microbiol.* **1994**, *2*, 73–76.

35. Lipsitch, M.; Siller, S.; Nowak, M. A. THE EVOLUTION OF VIRULENCE IN
525   PATHOGENS WITH VERTICAL AND HORIZONTAL TRANSMISSION. *Evolution*
526   **1996**, *50*, 1729–1741.

527 36. Cressler, C. E.; McLEOD, D. V.; Rozins, C.; van den Hoogen, J.; Day, T. The
528   adaptive evolution of virulence: a review of theoretical predictions and empirical
529   tests. *Parasitology* **2016**, *143*, 915–930.

530 37. Gillum, A. M.; Tsay, E. Y.; Kirsch, D. R. Isolation of the *Candida albicans* gene for
531   orotidine-5'-phosphate decarboxylase by complementation of *S. cerevisiae ura3* and
532   *E. coli pyrF* mutations. *Mol Gen Genet* **1984**, *198*, 179–182.

533 38. Morschhäuser, J.; Ruhnke, M.; Michel, S.; Hacker, J. Identification of CARE-2-
534   negative Candida albicans isolates as Candida dubliniensis. *Mycoses* **1999**, *42*, 29–
535   32.

536 39. Jain, C.; Pastor, K.; Gonzalez, A. Y.; Lorenz, M. C.; Rao, R. P. The role of Candida
537   albicans AP-1 protein against host derived ROS in in vivo models of infection.
538   *Virulence* **2013**, *4*, 67–76.

539 40. Yapar, N. Epidemiology and risk factors for invasive candidiasis. *Ther Clin Risk*
540   *Manag* **2014**, *10*, 95–105.

541 41. Krcméry, V.; Barnes, A. J. Non-albicans Candida spp. causing fungaemia:
542   pathogenicity and antifungal resistance. *J. Hosp. Infect.* **2002**, *50*, 243–260.

543 42. Meisel, J. D.; Kim, D. H. Behavioral avoidance of pathogenic bacteria by
544   Caenorhabditis elegans. *Trends Immunol.* **2014**, *35*, 465–470.

545 43. Jain, C.; Yun, M.; Politz, S. M.; Rao, R. P. A pathogenesis assay using
546   Saccharomyces cerevisiae and Caenorhabditis elegans reveals novel roles for yeast
547   AP-1, Yap1, and host dual oxidase BLI-3 in fungal pathogenesis. *Eukaryotic Cell*
548   **2009**, *8*, 1218–1227.

549 44. Gravato-Nobre, M. J.; Nicholas, H. R.; Nijland, R.; O'Rourke, D.; Whittington, D. E.;
550   Yook, K. J.; Hodgkin, J. Multiple genes affect sensitivity of Caenorhabditis elegans to
551   the bacterial pathogen Microbacterium nematophilum. *Genetics* **2005**, *171*, 1033–
552   1045.

45. Garsin, D. A.; Villanueva, J. M.; Begun, J.; Kim, D. H.; Sifri, C. D.; Calderwood, S.
B.; Ruvkun, G.; Ausubel, F. M. Long-lived C. elegans daf-2 mutants are resistant to
bacterial pathogens. *Science* **2003**, *300*, 1921–1921.
46. Pfaller, M. A.; Diekema, D. J.; Gibbs, D. L.; Newell, V. A.; Ellis, D.; Tullio, V.;
Rodloff, A.; Fu, W.; Ling, T. A.; Global Antifungal Surveillance Group Results from
the ARTEMIS DISK Global Antifungal Surveillance Study, 1997 to 2007: a 10.5-year
analysis of susceptibilities of Candida Species to fluconazole and voriconazole as
determined by CLSI standardized disk diffusion. *J. Clin. Microbiol.* **2010**, *48*, 1366–
1377.
47. Vilela, M. M. S.; Kamei, K.; Sano, A.; Tanaka, R.; Uno, J.; Takahashi, I.; Ito, J.;
Yarita, K.; Miyaji, M. Pathogenicity and virulence of Candida dubliniensis:
comparison with C. albicans. *Med Mycol* **2002**, *40*, 249–257.
48. Silva, S.; Negri, M.; Henriques, M.; Oliveira, R.; Williams, D. W.; Azeredo, J.
Candida glabrata, Candida parapsilosis and Candida tropicalis: biology,
epidemiology, pathogenicity and antifungal resistance. *FEMS Microbiol. Rev.* **2012**,
36, 288–305.
569