1

11 A novel virulence phenotype rapidly assesses *Candida* fungal pathogenesis in healthy

12 and immunocompromised *Caenorhabditis elegans* hosts

16 Dorian J. Feistel, Rema Elmostafa, Nancy Nguyen, McKenna Penley, Levi Morran, and

17 Meleah A. Hickman[*]

24 Running title: Fungal infection reduces multiple measures of host fitness

26 Department of Biology, Emory University

27 *corresponding author

Abstract

The yeast *Candida albicans* is an opportunistic pathogen of humans, meaning that
despite commensal interactions with its host, it can transition to a harmful pathogen.
While *C. albicans* is the predominant species isolated in the human mycobiome and
implicated in fungal infection, infections due to non-albicans *Candida* species are rapidly
rising. Studying the factors that contribute to virulence is often challenging and
frequently depends on many contexts including host immune status and pathogen
genetic background. Here, we utilize the nematode *Caenorhabditis elegans* as a
perspicuous and efficient model host system to study fungal infections of *Candida*
pathogens. We find that in addition to reducing lifetime host survival, exposure to *C.
albicans* results in delayed reproduction, which significantly reduced lineage growth
over multiple generations. Furthermore, we assessed fungal pathogen virulence in *C.
elegans* hosts compromised for innate immune function and detected increased early
mortality, reduced brood sizes and delayed reproduction relative to infected healthy
hosts. Importantly, by assessing virulence in both healthy and immunocompromised
host backgrounds we reveal the pathogen potential in non-*albicans Candida* species.
Taken together, we present a novel lineage growth assay to measure reduction in host
fitness associated with fungal infection and demonstrate significant interactions between
pathogen and host immune function that contribute to virulence.

Importance

Opportunistic pathogens are commensals capable of causing disease and are serious
threats to human health. It is critical to understand the mechanisms and host contexts
under which opportunistic pathogens become virulent. In this work, we present a novel
assay to quickly and quantitatively measure pathogen virulence in healthy and
immunocompromised hosts. We found that *Candida* species, the most prominent fungal
opportunistic pathogens of humans, decrease host fitness by reducing survival and
impacting host reproduction. Most importantly, by measuring virulence in hosts that
have intact or compromised immune function, we can reveal the pathogenic potential of
opportunistic fungal pathogens.

Introduction

Candida species are commensals of the human gastrointestinal microbiota and various other niches in the human body [1]. Despite their commensal existence in many humans, Candida species also account for the most fungal infections in the world and are the fourth most prevalent cause of all nosocomial blood stream infections [2,3]. The severity of fungal infection is often dependent on the immune status of the host, with superficial infections occurring in healthy individuals and bloodstream infections in immunocompromised hosts [4-8]. The majority of these infections are associated with Candida albicans, however other non-albicans Candida species are increasingly becoming implicated in fungal infections [9,10].

Several host models have been employed to better understand fungal infection and disease progression. These experimental systems include mouse, zebrafish, nematode, wax moth and human ex vivo models have been specifically developed to study Candida infections [11]. Most often, virulence phenotypes are limited to survival outcomes or tissue-specific fungal burden [12], the latter of which requires animal sacrifice and cannot address the long-term consequence of fungal disease. Furthermore, animal mortality and sacrifice, particularly in vertebrate models, constrain the number of infected individuals to study and thus reduce the quantitative robustness of these experiments. Nonetheless, these studies provide valuable insight into Candida infection and disease and have uncovered important roles for innate immune function of hosts [13-15], specific C. albicans genes regulating virulence [16,17] and differences between genetic backgrounds within and between Candida species [18].

Other major limitations for most studies of Candida infection virulence include the time-intensive nature of tracking virulence phenotypes in vertebrate systems such as mice, the genetic intractability of zebrafish and wax moth larvae and the context-independent nature of ex vivo models [4]. As such, most studies focus specifically on C. albicans virulence while many non-albicans Candida species are evaluated to a lesser degree. The nematode Caenorhabditis elegans has been developed as a model for Candida infection to overcome these limitations [19,20] and to screen compounds for antifungal activity [21]. C. elegans has proven useful for studying host-microbe interactions [22] because many fungal and bacterial pathogens that cause illness in

humans also cause disease in *C. elegans* [23]. Infecting *C. elegans* is relatively simple

and often done by replacing or incorporating its standard laboratory food *E. coli* with a

desired pathogen which colonizes the gut and causes disease [23]. Furthermore, the

innate immune system in *C. elegans* includes the *SEK-1* gene which encodes a MAP

Kinase Kinase (MAPKK) that is homologous to the MKK3/6 and MKK4 family of

mammalian MAPKKs and activates the *C. elegans* p38 MAP kinase ortholog [24]. This

pathway has been suggested to be an ancient and conserved component of *C. elegans'*

immune response to pathogens [25]. As such, the *sek-1* mutation increases

susceptibility to microbial colonization [26], including infection with many *Candida*

species [27] and is necessary to induce the appropriate antifungal immune defense [28].

In addition to utilizing innate immune system mutants, most studies measure host

mortality in temperature sensitive sterile mutants [17,21,29,30] to more easily track

mortality in founder populations since *C. elegans* has a rapid lifecycle and large brood

sizes [31,32]. The effects on host reproduction are often overlooked in favor of

examination of mortality rates when assessing a pathogen's virulence [33]. Yet, it is

important to remember that virulence can be broadly measured as any reduction in host

fitness resulting from interactions between a pathogen and its host [34-36]. In this work,

we assessed virulence of *Candida* species in *C. elegans* hosts using a novel measure

of host fitness that incorporated both host survival and fecundity. We found that fungal

pathogens reduced host fitness by delaying reproduction, resulting in long-term

consequences for population growth in both healthy and immunocompromised host

backgrounds. Using this novel measure, we characterized virulence phenotypes for

three non-*albicans Candida* species; *C. dubliniensis*, *C. tropicalis*, and *C. parapsilosis* in

both healthy and immunocompromised *C. elegans* hosts. Our studies demonstrate that

differences in virulence can be identified between pathogen species and that

pathogenic potential is often revealed when host immune function is diminished.

Methods

Strains and media

For this study, the fungal pathogens *C. albicans* (SC5314 [37]), *C. dubliniensis*

(Wu284 [38]), *C. tropicalis* (ATCC 22109), and *C. parapsilosis* (ATCC 22109) strains

were used. *C. elegans* N2 Bristol (WT) and a *sek-1* mutant derivative [24] were used to

test host survival, fecundity and population growth. *C. elegans* populations were

maintained at 20°C on 100mm petri dishes with 25 mL of lite nematode growth medium

(NGM, US Biological) with *E. coli* OP50 as a food source. Nematodes were transferred

to a newly seeded *E. coli* plate every 3-4 days. For survival, fecundity and population

growth assays, NGM was supplemented with 0.08 g/L uridine, 0.08 g/L histidine, and

0.04 g/L adenine to facilitate growth of auxotrophic *C. albicans* strains and 0.2 g/L

streptomycin sulfate to inhibit *E. coli* overgrowth so fungal strains could proliferate.

Seeding NGM plates for survival, fecundity, and population growth assays

Candida strains and *E. coli* OP50 strains were inoculated in 3 mL of YPD or 5 mL

of LB, respectively, and cultured at 30°C for 1-2 days. *Candida* culture densities were

measured with a spectrophotometer and diluted to a final volume of 3.0 OD_{600} per mL

(~6 x 10^7 cells per ml). *E. coli* cultures were pelleted and washed twice with 1 mL of

ddH_2O. The supernatant was removed and the pellet was centrifuged for 60 sec at

maximum to remove any excess liquid. The pellet was weighed and suspended with

sterilized water to a final volume of 200 mg/mL. A mix of 6.25 μL *E. coli*, 1.25 μL

Candida was brought to a final volume of 50 μL with ddH_2O. The entire 50 μL was

spotted onto the center of a 35mm supplemented-NGM Lite agar plate and incubated at

room temperature overnight before addition of eggs or transferring nematode. *E. coli*

OP50 was used as a control at the same concentration described above.

Egg preparation and synchronization for survival, fecundity and population growth

assays

For survival, fecundity and population growth assays, approximately 100

nematodes at the L3/L4 stage were transferred to a 100mm NGM plate seeded with *E.*

coli OP50 and maintained at 20°C for 2-3 days prior to the start of an experiment. On

the first day of an experiment, these NGM plates were washed with M9 buffer and

contents (live nematodes and eggs) transferred to 15 mL conical tube and pelleted by

centrifugation (2 min at 1200 rpm). The pellet was re-suspended in a 1:4 bleach (5.25%)

solution and transferred to a micro-centrifuge tube. The suspension was mixed via

inversion for 90-120 sec and subsequently centrifuged (30 sec at 1500 rpm). The pellet

was washed with 1 mL M9 buffer three consecutive times to remove excess bleach

solution and brought to a final suspension with 500 μL M9 buffer. To determine the

concentration of eggs, 10 μL was placed on a concaved slide and eggs counted and the

egg suspension was diluted with M9 to a final concentration of 10 eggs/μL. All assays

were treated equally on the first day (Day 0) by adding roughly 100 eggs to a treatment

or *E. coli* plate (described above).

Survival assays

This experimental procedure is a modified version of [39]. Briefly, 72 h after

adding ~50-75 nematode eggs to a plate, 40 adult nematodes were randomly selected

and transferred to newly seeded plates with the same concentration of food as

described above and incubated at 20°C. Every other day nematodes were transferred to

freshly seeded plates until all nematode replicate populations went extinct. The number

of living, dead, and censored worms were scored daily. Each survival assay was

performed at least three independent times.

Fecundity assays

For both fecundity and lineage expansion assays, 48 h after adding nematode

eggs to a plate, a single L4 reproductively immature hermaphroditic nematode was

randomly selected and transferred (6-10 independent biological replicates per treatment

per block) to a newly seeded 35 mm petri plate containing either a treatment of *C.*

albicans and *E. coli* or *E. coli* alone (described above) and incubated at 20°C.

Nematodes were transferred to freshly seeded plates every 24 h for seven consecutive

days. Eggs remained undisturbed on the plate and were incubated at 20°C for an

additional 24 h to provide enough time to hatch, at which point the number of viable

progeny per day was scored. Nematodes that died during the assay were scored as

dead at the time of transfer. Nematodes that crawled off the plate or were otherwise

unaccountable for were considered censored and excluded from the analysis.

Lineage Expansion Assay

A single L4 reproductively immature hermaphroditic nematode was randomly

selected and transferred (6 biological replicates per treatment) to a freshly seeded 100

mm treatment or *E. coli* plate (described above) containing a 6-fold increase in food.

Five days later, each plate was washed with M9 buffer until the majority of nematodes

were displaced and subsequently transferred to 15mL conical tubes. Tubes were placed

at 4°C for 1h to allow the nematodes to settle at the bottom. All tubes were

concentrated to a final volume of 10mL. Six, 20 μL samples were taken from each

population and counted.

Statistical analyses

All statistical analysis were performed with GraphPad Prism. Data sets were

tested for normality using the D'Agostino & Pearson omnibus normality test. For

comparisons across groups, one-way ANOVAs and post-hoc Tukey multiple

comparison tests or Kruskal-Wallis and Dunn's multiple comparison tests were

performed.

Results

Exposure to C. albicans reduces multiple measures of C. elegans fitness

Pathogens affect host fitness by decreasing lifespan and/or reducing fecundity.

To determine how *C. albicans* impacts *C. elegans* host fitness, we first measured

nematode lifespan in the presence or absence of *C. albicans*. We observed a significant

reduction in survival when *C. elegans* was reared on *C. albicans* compared to when it

was reared in its absence (Figure 1, $p < 0.0001$ Log-rank test), consistent with

previously published results [39]. 50% mortality was reached in eight days in nematode

populations exposed to *C. albicans* compared to 16 days in unexposed nematode

populations, indicating that *C. elegans* lifespan is substantially decreased by exposure

to *C. albicans*.

While overall survival was reduced, nematode death was rarely observed earlier

than 5 days post-infection. Importantly, adult nematodes produce the majority of their

offspring before *C. albicans* infection reduces rates of survival. Given this information

and that fecundity is a key component of fitness from an evolutionary perspective, we

investigated whether *C. albicans* negatively impacts *C. elegans* reproduction. To test

this, we counted the total number of viable progeny (i.e. brood size) produced in the first

seven days of reproductive maturity from individual nematodes that were either

unexposed or exposed to either heat-killed or live *C. albicans*. As shown in Figure 2A,

exposure to *C. albicans* reduced the average number of viable progeny (253±7)

compared to the unexposed control (285±4; $p < 0.0002$, Kruskal-Wallis test) by ~11%.

While statistically significant, the cost of *C. albicans* toward host brood size is modest.

Furthermore, the brood size of nematodes exposed to heat-killed *C. albicans* (HK) was

not detectably different from uninfected or live *C. albicans* treatments (Figure 2A).

Intriguingly, we identified a significant delay in reproductive timing when we

measured the number of progeny produced every 24 h during the host reproductive

window. We observed a substantial reduction in reproduction on Day 3 in nematodes

exposed to *C. albicans* (103±4) relative to unexposed hosts (187±3) or hosts associated

with heat-killed *C. albicans* (Figure 2B; $p < 0.0001$ Kruskal-Wallis test) and increased

progeny produced on Days 4-6 in nematodes exposed to *C. albicans* (Figure 2B). We

calculated the total fraction of reproduction occurring in this 'late' window (Days 4-6) and

identified a significant increase in the progeny produced late in *C. elegans* exposed to

C. albicans compared to unexposed or heat-killed controls (Figure 2C; $p=0.0001$,

Kruskal-Wallis test). Taken together, our data indicate that *C. albicans* severely delays

and reduces reproduction in addition to impacting overall survival in *C. elegans.*

To investigate whether this reproductive delay observed in nematodes exposed

to *C. albicans* had any long-term consequences, we took advantage that *C. elegans* are

hermaphrodites and measured the total population produced from a single founder

worm over multiple generations through lineage expansion assays. We find that the

median population produced within 7 days from a single uninfected nematode is

~30,250 F1 and F2 progeny (Figure 2D). Lineage expansion is significantly impacted in

nematodes exposed to live *C. albicans*, with a median population size of ~20,208 F1

and F2 progeny, a reduction of ~33% compared to uninfected and heat-killed control

treatments (Figure 2D, $p < 0.0001$ one-way ANOVA). We conclude that the delay in

reproduction, coupled with the reduced brood size resulting from exposure to *C.*

albicans severely impacts *C. elegans* evolutionary fitness.

Immunocompromised hosts are susceptible to fungal infection

A leading host-related risk factor for invasive candidiasis is compromised

immune function [40]. Given our novel results regarding fecundity in healthy *C. elegans*

hosts, we were curious how exposure to *C. albicans* impacts nematode hosts with

compromised immune function. To address this point, we utilized a *C. elegans* strain

carrying a mutation in *SEK*-1, a well-conserved MAP kinase involved in the innate

immune signaling cascade [24,26] and antifungal response [28]. Unlike healthy hosts,

immunocompromised hosts show high rates of early (within 7 days) mortality in both

uninfected and *C. albicans* treatments (Figure S1). *C. albicans* exposure reduced

average brood size by nearly 50% (78±6) in immunocompromised hosts compared to

uninfected (151±5) and heat-killed (127±13) controls (Figure 3A, $p < 0.0001$, Kruskal-

Wallis test). It is important to note that in all treatments (unexposed, heat-killed, and live

C. albicans), immunocompromised hosts produced significantly less progeny than

healthy hosts (Table S1), but the largest reductions occur in immunocompromised hosts

are exposed to live *C. albicans*.

One explanation for the overall reduction in average brood size is the increased

early mortality we observe in immunocompromised hosts. To address this possibility,

we re-analyzed brood size only from hosts that survived past Day 3 and found that while

this did slightly increase average brood size across all treatments, it was not significant

between analyses (p=0.2356 two-tailed t-test) and exposure to *C. albicans* still

significantly reduced average brood size (94±6) relative to uninfected (163±5) and heat-

killed controls (141±9) (Figure S2, *p* < 0.0001 Kruskal-Wallis test). For hosts that

survived past three days, we also detected a significant delay in reproductive timing

upon exposure to *C. albicans*, with nearly 45% of all reproduction occurring in this late

window (Figure 3C), similar to what we observe in healthy hosts (Table S1).

We predicted that lineage expansion in immunocompromised hosts exposed to

C. albicans would be more dramatically impacted relative to wild-type nematodes

exposed to *C. albicans*, given the reproductive delays coupled with the overall reduced

brood sizes and increased mortality observed. Indeed, we determined that the median

population size of *C. albicans* exposed nematodes was ~11,000 compared to ~17,000

and 19,500 in heat-killed and uninfected controls, respectively (Figure 3C). The overall

reduction in population size resulting from *C. albicans* exposure is ~43% in

immunocompromised hosts and is significantly greater than the ~32% reduction in

population size in healthy hosts (Figure 4 and Table S1). Taken together, our results

indicate that immunocompromised hosts are more susceptible to fungal infection than

healthy wild-type hosts.

Host immune status reveals pathogenic potential in other Candida species

While *C. albicans* is the predominant fungal agent in invasive candidiasis, non-

albicans Candida species (NACs) are estimated to account for 35-65% of

candidaemia [41]. By evaluating NACs in both healthy and immunocompromised hosts,

we can begin to assess their pathogenic potential, and thus extend the utility of *C.*

elegans as a model host system for detecting virulence in *Candida* species. Here, we

leveraged our simple, yet robust, lineage expansion assays to measure the impact of

three additional *Candida* species: *C. dubliniensis*, *C. tropicalis* and *C. parapsilosis* on

host fitness for healthy and immunocompromised *C. elegans*. All three non-*albicans*

Candida species significantly reduced the population size of both healthy and

immunocompromised hosts compared to uninfected controls (black asterisks, Figure S3

and Table S1), and sometimes were also significantly different than population sizes of

hosts exposed to *C. albicans* (red asterisks, Figure S3).

We next wanted to determine if there was a significant interaction between

Candida species and host immune function. To test this, we evaluated the population

size of infected nematodes relative to uninfected for healthy (Figure 4; black bars) and

immunocompromised (Figure 4; green bars) host backgrounds, since

immunocompromised hosts have significantly smaller populations even when

uninfected (Figure 2 and S4). There are significant differences among *Candida* species

and nematode host backgrounds ($p < 0.001$, two-way ANOVA; $p < 0.001$ between

pathogens, hosts, and their interaction). In healthy hosts, *C. albicans* and *C. tropicalis*

are the most virulent, with 32-35% reductions in total population size, whereas *C.*

parapsilosis and *C. dublineinsis* result in more modest population reductions (17-20%).

However, in immunocompromised hosts, *C. parapsilosis* is one of the most virulent, with

nearly a 67% reduction in population size. *C. albicans* and *C. tropicalis* are also highly

virulent in immunocompromised hosts and all three of these species more severely

impact immunocompromised hosts compared to healthy hosts. In contrast, *C. dublineinsis* is the least virulent, with no significant differences observed between the

relative population sizes of healthy and immunocompromised hosts (Figure 4). Given

the virulence phenotypes of these different yeasts, our results suggest that fungal

virulence depends not only on the species of pathogen but also on the immune status of

the host.

Discussion

Here we utilized a *C. elegans* experimental host system to identify novel

measures of host fitness associated with fungal infection. By tracking the number of

progeny produced per day, we can quickly and quantitatively assess three aspects of

host fitness: early mortality, total viable offspring produced and reproductive timing. We

show that exposure to *Candia albicans* in healthy *C. elegans* hosts not only reduces

survival (Figure 1), it modestly reduces total progeny produced, and dramatically delays

reproductive timing (Figure 2). This reproductive delay has long-term consequences for

lineage expansion, as single founder nematodes exposed to *C. albicans* reduces its

population growth by ~30% (Figure 2D). We found similar delays in reproductive timing

in immunocompromised hosts exposed to *C. albicans*, although immunocompromised

hosts are more susceptible to infection and have higher incidence of early mortality and

smaller brood sizes compared to healthy hosts (Figure 3). Importantly, the delayed

reproduction phenotype in infected *C. elegans* is easily ascertained, highly quantitative

and reproducible and can be used as a screening method to detect relatively small

differences in virulence across *Candida* strains or other fungal species. Furthermore, by

utilizing different host contexts, this assay revealed the pathogenic potential of other

important, yet understudied, *Candida* species (Figure 4), including *C. parapsilosis*,

which has dramatically higher levels of virulence in immunocompromised hosts

compared to healthy hosts.

The reductions in host fitness associated with *C. albicans* and other *Candida*

species indicate that we have established a bona fide model of fungal infection.

Importantly, these reductions in host fitness cannot be attributed to *Candida* species

being a sub-optimal food source or that the hosts are starving because we deliver the

pathogen with *E. coli*, the standard food source for *C. elegans*. Further, we do not

detect any host larval development into dauer, an alternative lifestage in response to

starvation and overpopulation [32]. While pathogen avoidance is a common defense

strategy for *C. elegans* [42], we censored any worms that have crawled off the plate or

have disappeared during the course of our experiment and removed them from our data

analysis. Furthermore, the reductions in host fitness depend on ingesting live *C.*

albicans, as heat-killed treatments do not cause significant reductions in total brood

size, reproductive timing, or lineage expansion (Figure 2, 'HK'). Microscopic analysis

reveals that fungal cells inhabit the gut of *C. elegans* ([39] and data not shown) and we

can extract viable host-associated fungal cells (data not shown). Taken together, these

data support a model of fungal infection in the host *C. elegans*.

The host reproductive delay we observed in *C. elegans* upon fungal infection is a

robust, highly quantitative measure of virulence that makes it amenable to screen a

variety of host and pathogen genetic backgrounds that has been challenging in

mammalian and insect models. *C. elegans* is easily maintained in the lab, has a short

lifecycle that generates a large number of progeny, and has been a fundamental model

genetic organism [32]. Previous work has shown that upon exposure to *C. albicans, C.*

elegans have a transcriptional response consistent with that of infection [28]. In this

work, we utilized a mutant *C. elegans* strain in which *SEK-1*, an important MAPK in

innate immunity and whose homologs include the MKK3/6 and MKK4 family of

mammalian MAPKKs, to determine the severity of fungal infection in

immunocompromised hosts. We demonstrate that deficiencies in innate immune

function result in hosts with high susceptibility to fungal infection and in the future can

extend this analysis for many other host backgrounds that are mutant for immune

function.

Previous studies using *C. elegans* as a host for fungal infection have described

additional phenotypes beyond reduced survival, including a deformed anal region (DAR)

[39,43]. It is possible that DAR may be contributing to the delayed reproductive

phenotype we observe in our experiments, however, the frequency of DAR phenotypes

was low and we did not observe any detectable differences between hosts exposed to

C. albicans and hosts unexposed (data not shown), nor did we observe dramatic

reductions in total number of viable offspring produced between these two groups of

hosts (Figure 2A. While this deformity is a result of a local defense reaction of the worm

due to extracellular signal-regulated kinase activation of the innate immunity MAKK

cascade, it is a distinct marker from delayed reproduction for infection [44]. Wildtype

reproductive timing utilizes the highly conserved DAF-2 DAF-16 insulin-like signaling

pathway [23], but has also been implicated in innate immunity and bacterial infection

[45] and is potentially disrupted in *C. elegans* hosts infected with *Candida*.

Impacts on host reproduction have often been overlooked in studies regarding

Candida virulence, yet host reproduction is an important evolutionary measure of an

organism's fitness. Not only can we utilize these fecundity assays to assess the roles of

host immune function on fungal pathogenesis, we also readily screen diverse pathogen

strain backgrounds and species. A large-scale, international 10-year study identified 31

species of *Candida* associated with clinical samples. While *C. albicans* is still the most

prevalent, its isolation from clinical isolates is decreasing with corresponding increases

isolation of *C. glabrata, C. tropicalis* and *C. parapsilosis* [46]. Despite its close

evolutionary relationship with *C. albicans*, *C. dubliniensis* does not seem to be highly

pathogenic [47]. However, other non-*albicans Candida* species have been increasingly

implicated in fungal infections of humans [48]. For example, *C. tropicalis* and *C.*

parapsilosis are common fungal species isolated in patients with candidemia (7 - 48%

and 11 - 27% respectively and depending on geographic region), candiduria (8 – 44%

and 0.5 – 11% respectively, and oral candidosis (5 – 13% and 7 – 15%, respectively)

([48] and references therein). Despite the increasing incidence of non-*albicans Candida*

infections [46], experimental studies using these pathogens remains limited. Here, we

analyzed the virulence phenotypes of three non-albicans *Candida* species, *C.*

dubliniensis, *C. tropicalis* and *C. parapsilosis* in both healthy and immunocompromised

host backgrounds. In healthy *C. elegans* hosts, *C. parapsilosis* is the least virulent

compared to *C. albicans* and *C. tropicalis* and *C. dublineinsis* has intermediate level of

virulence (Figure 4). For all species except for *C. dubliniensis,* virulence is more severe

in immunocompromised hosts and is particularly striking in *C. parapsilosis*, which is the

most virulent in hosts with compromised immune function. Therefore, the *C. elegans*

fecundity assays we developed in this study can rapidly reveal pathogenic potential by assessing virulence in multiple host genetic backgrounds and across pathogens in a highly quantitative and robust manner.

Literature Cited

1. Underhill, D. M.; Iliev, I. D. The mycobiota: interactions between commensal fungi and the host immune system. *Nature reviews. Immunology* **2014**, *14*, 405–416.

2. Pappas, P. G.; Lionakis, M. S.; Arendrup, M. C.; Ostrosky-Zeichner, L.; Kullberg, B. J. Invasive candidiasis. *Nat Rev Dis Primers* **2018**, *4*, 18026.

3. Edmond, M. B.; Wallace, S. E.; McClish, D. K.; Pfaller, M. A.; Jones, R. N.; Wenzel, R. P. Nosocomial bloodstream infections in United States hospitals: a three-year analysis. *Clin Infect Dis* **1999**, *29*, 239–244.

4. MacCallum, D. M. Hosting infection: experimental models to assay Candida virulence. *Int J Microbiol* **2012**, *2012*, 363764–12.

5. Pfaller, M. A. Nosocomial candidiasis: emerging species, reservoirs, and modes of transmission. *Clin Infect Dis* **1996**, *22 Suppl 2*, S89–94.

6. Coleman, D. C.; Sullivan, D. J.; Bennett, D. E.; Moran, G. P.; Barry, H. J.; Shanley, D. B. Candidiasis: the emergence of a novel species, Candida dubliniensis. *AIDS* **1997**, *11*, 557–567.

7. Kontoyiannis, D. P.; Vaziri, I.; Hanna, H. A.; Boktour, M.; Thornby, J.; Hachem, R.; Bodey, G. P.; Raad, I. I. Risk Factors for Candida tropicalis fungemia in patients with cancer. *Clin Infect Dis* **2001**, *33*, 1676–1681.

8. Clark, T. A.; Slavinski, S. A.; Morgan, J.; Lott, T.; Arthington-Skaggs, B. A.; Brandt, M. E.; Webb, R. M.; Currier, M.; Flowers, R. H.; Fridkin, S. K.; Hajjeh, R. A. Epidemiologic and molecular characterization of an outbreak of Candida parapsilosis bloodstream infections in a community hospital. *J. Clin. Microbiol.* **2004**, *42*, 4468–4472.

9. Pfaller, M. A.; Diekema, D. J.; Jones, R. N.; Sader, H. S.; Fluit, A. C.; Hollis, R. J.; Messer, S. A.; SENTRY Participant Group International surveillance of bloodstream infections due to Candida species: frequency of occurrence and in vitro

susceptibilities to fluconazole, ravuconazole, and voriconazole of isolates collected

from 1997 through 1999 in the SENTRY antimicrobial surveillance program. *J. Clin.*

Microbiol. **2001**, *39*, 3254–3259.

10. Hajjeh, R. A.; Sofair, A. N.; Harrison, L. H.; Lyon, G. M.; Arthington-Skaggs, B. A.;

Mirza, S. A.; Phelan, M.; Morgan, J.; Lee-Yang, W.; Ciblak, M. A.; Benjamin, L. E.;

Sanza, L. T.; Huie, S.; Yeo, S. F.; Brandt, M. E.; Warnock, D. W. Incidence of

bloodstream infections due to Candida species and in vitro susceptibilities of isolates

collected from 1998 to 2000 in a population-based active surveillance program. *J.*

Clin. Microbiol. **2004**, *42*, 1519–1527.

11. Harwood, C. G.; Rao, R. P. Host pathogen relations: exploring animal models for

fungal pathogens. *Pathogens* **2014**, *3*, 549–562.

12. Conti, H. R.; Huppler, A. R.; Whibley, N.; Gaffen, S. L. Animal models for

candidiasis. *Curr Protoc Immunol* **2014**, *105*, 19.6.1–17.

13. Verma, A.; Gaffen, S. L.; Swidergall, M. Innate Immunity to Mucosal Candida

Infections. *J Fungi (Basel)* **2017**, *3*, 60.

14. Naglik, J. R.; König, A.; Hube, B.; Gaffen, S. L. Candida albicans-epithelial

interactions and induction of mucosal innate immunity. *Curr Opin Microbiol* **2017**, *40*,

104–112.

15. Richardson, J. P.; Moyes, D. L.; Ho, J.; Naglik, J. R. Candida innate immunity at the

mucosa. *Seminars in Cell & Developmental Biology* **2018**.

16. Lu, Y.; Su, C.; Solis, N. V.; Filler, S. G.; Liu, H. Synergistic regulation of hyphal

elongation by hypoxia, CO(2), and nutrient conditions controls the virulence of

Candida albicans. *Cell Host Microbe* **2013**, *14*, 499–509.

17. Pukkila-Worley, R.; Peleg, A. Y.; Tampakakis, E.; Mylonakis, E. Candida albicans

hyphal formation and virulence assessed using a Caenorhabditis elegans infection

model. *Eukaryotic Cell* **2009**, *8*, 1750–1758.

18. Hirakawa, M. P.; Martinez, D. A.; Sakthikumar, S.; Anderson, M. Z.; Berlin, A.; Gujja,

S.; Zeng, Q.; Zisson, E.; Wang, J. M.; Greenberg, J. M.; Berman, J.; Bennett, R. J.;

Cuomo, C. A. Genetic and phenotypic intra-species variation in Candida albicans.

Genome Res **2015**, *25*, 413–425.

19. Powell, J. R.; Ausubel, F. M. Models of Caenorhabditis elegans infection by
bacterial and fungal pathogens. *Methods Mol Biol* **2008**, *415*, 403–427.

20. Aballay, A.; Ausubel, F. M. Caenorhabditis elegans as a host for the study of host-
pathogen interactions. *Curr Opin Microbiol* **2002**, *5*, 97–101.

21. Breger, J.; Fuchs, B. B.; Aperis, G.; Moy, T. I.; Ausubel, F. M.; Mylonakis, E.
Antifungal chemical compounds identified using a C. elegans pathogenicity assay.
PLoS Pathog **2007**, *3*, e18.

22. Sifri, C. D.; Begun, J.; Ausubel, F. M. The worm has turned--microbial virulence
modeled in Caenorhabditis elegans. *Trends Microbiol.* **2005**, *13*, 119–127.

23. Irazoqui, J. E.; Urbach, J. M.; Ausubel, F. M. Evolution of host innate defence:
insights from Caenorhabditis elegans and primitive invertebrates. *Nature reviews.*
Immunology **2010**, *10*, 47–58.

24. Kim, D. H.; Feinbaum, R.; Alloing, G.; Emerson, F. E.; Garsin, D. A.; Inoue, H.;
Tanaka-Hino, M.; Hisamoto, N.; Matsumoto, K.; Tan, M.-W.; Ausubel, F. M. A
conserved p38 MAP kinase pathway in Caenorhabditis elegans innate immunity.
Science **2002**, *297*, 623–626.

25. Ermolaeva, M. A.; Schumacher, B. Insights from the worm: The C. elegans model
for innate immunity. *Seminars in Immunology* **2014**, *26*, 303–309.

26. Youngman, M. J.; Rogers, Z. N.; Kim, D. H. A Decline in p38 MAPK Signaling
Underlies Immunosenescence in Caenorhabditis elegans. *PLOS Genet* **2011**, *7*,
e1002082.

27. Ortega-Riveros, M.; De-la-Pinta, I.; Marcos-Arias, C.; Ezpeleta, G.; Quindós, G.;
Eraso, E. Usefulness of the Non-conventional Caenorhabditis elegans Model to
Assess Candida Virulence. *Mycopathologia* **2017**, *182*, 785–795.

28. Pukkila-Worley, R.; Ausubel, F. M.; Mylonakis, E. Candida albicans Infection of
Caenorhabditis elegans Induces Antifungal Immune Defenses. *PLoS Pathog* **2011**,
7, e1002074–13.

29. Moy, T. I.; Ball, A. R.; Anklesaria, Z.; Casadei, G.; Lewis, K.; Ausubel, F. M.
Identification of novel antimicrobials using a live-animal infection model. *Proc Natl*
Acad Sci USA **2006**, *103*, 10414–10419.

30. Peleg, A. Y.; Tampakakis, E.; Fuchs, B. B.; Eliopoulos, G. M.; Moellering, R. C.;
Mylonakis, E. Prokaryote-eukaryote interactions identified by using Caenorhabditis
elegans. *Proc Natl Acad Sci U S A* **2008**, *105*, 14585–14590.

31. Muschiol, D.; Schroeder, F.; Traunspurger, W. Life cycle and population growth rate
of Caenorhabditis elegans studied by a new method. *BMC Ecol.* **2009**, *9*, 14.

32. Corsi, A. K.; Wightman, B.; Chalfie, M. A Transparent Window into Biology: A Primer
on Caenorhabditis elegans. *Genetics* **2015**, *200*, 387–407.

33. Abbate, J. L.; Kada, S.; Lion, S. Beyond Mortality: Sterility As a Neglected
Component of Parasite Virulence. *PLoS Pathog* **2015**, *11*, e1005229.

34. Read, A. F. The evolution of virulence. *Trends Microbiol.* **1994**, *2*, 73–76.

35. Lipsitch, M.; Siller, S.; Nowak, M. A. The evolution of virulence in pathogens with
vertical and horizontal transmission. *Evolution* **1996**, *50*, 1729–1741.

36. Cressler, C. E.; McLEOD, D. V.; Rozins, C.; van den Hoogen, J.; Day, T. The
adaptive evolution of virulence: a review of theoretical predictions and empirical
tests. *Parasitology* **2016**, *143*, 915–930.

37. Gillum, A. M.; Tsay, E. Y.; Kirsch, D. R. Isolation of the *Candida albicans* gene for
orotidine-5'-phosphate decarboxylase by complementation of *S. cerevisiae ura3* and
E. coli pyrF mutations. *Mol Gen Genet* **1984**, *198*, 179–182.

38. Morschhäuser, J.; Ruhnke, M.; Michel, S.; Hacker, J. Identification of CARE-2-
negative Candida albicans isolates as Candida dubliniensis. *Mycoses* **1999**, *42*, 29–
32.

39. Jain, C.; Pastor, K.; Gonzalez, A. Y.; Lorenz, M. C.; Rao, R. P. The role of Candida
albicans AP-1 protein against host derived ROS in in vivo models of infection.
Virulence **2013**, *4*, 67–76.

40. Yapar, N. Epidemiology and risk factors for invasive candidiasis. *Ther Clin Risk*
Manag **2014**, *10*, 95–105.

41. Krcméry, V.; Barnes, A. J. Non-albicans Candida spp. causing fungaemia:
pathogenicity and antifungal resistance. *J. Hosp. Infect.* **2002**, *50*, 243–260.

42. Meisel, J. D.; Kim, D. H. Behavioral avoidance of pathogenic bacteria by
Caenorhabditis elegans. *Trends Immunol.* **2014**, *35*, 465–470.

43. Jain, C.; Yun, M.; Politz, S. M.; Rao, R. P. A pathogenesis assay using
Saccharomyces cerevisiae and Caenorhabditis elegans reveals novel roles for yeast
AP-1, Yap1, and host dual oxidase BLI-3 in fungal pathogenesis. *Eukaryotic Cell*
2009, *8*, 1218–1227.

44. Gravato-Nobre, M. J.; Nicholas, H. R.; Nijland, R.; O'Rourke, D.; Whittington, D. E.;
Yook, K. J.; Hodgkin, J. Multiple genes affect sensitivity of Caenorhabditis elegans to
the bacterial pathogen Microbacterium nematophilum. *Genetics* **2005**, *171*, 1033–
1045.

45. Garsin, D. A.; Villanueva, J. M.; Begun, J.; Kim, D. H.; Sifri, C. D.; Calderwood, S.
B.; Ruvkun, G.; Ausubel, F. M. Long-lived C. elegans daf-2 mutants are resistant to
bacterial pathogens. *Science* **2003**, *300*, 1921–1921.

46. Pfaller, M. A.; Diekema, D. J.; Gibbs, D. L.; Newell, V. A.; Ellis, D.; Tullio, V.;
Rodloff, A.; Fu, W.; Ling, T. A.; Global Antifungal Surveillance Group Results from
the ARTEMIS DISK Global Antifungal Surveillance Study, 1997 to 2007: a 10.5-year
analysis of susceptibilities of Candida Species to fluconazole and voriconazole as
determined by CLSI standardized disk diffusion. *J. Clin. Microbiol.* **2010**, *48*, 1366–
1377.

47. Vilela, M. M. S.; Kamei, K.; Sano, A.; Tanaka, R.; Uno, J.; Takahashi, I.; Ito, J.;
Yarita, K.; Miyaji, M. Pathogenicity and virulence of Candida dubliniensis:
comparison with C. albicans. *Med Mycol* **2002**, *40*, 249–257.

48. Silva, S.; Negri, M.; Henriques, M.; Oliveira, R.; Williams, D. W.; Azeredo, J.
Candida glabrata, Candida parapsilosis and Candida tropicalis: biology,
epidemiology, pathogenicity and antifungal resistance. *FEMS Microbiol. Rev.* **2012**,
36, 288–305.

Figure 1: Reduced nematode survival associated with *C. albicans* infection. Survival curves for *C. elegans* populations that are either uninfected (exposed to just an *E. coli* food source, black) or when infected with *C. albicans* SC5314 (red). Mean data values are plotted with SEM error bars, number of worms analyzed (n) for each treatment is indicated in the legend. **** indicates p < 0.0001, Log-rank (Mantel-Cox) test.

Figure 2. Live *C. albicans* reduces nematode fecundity. A) Box and whiskers plot of brood sizes for *C. elegans* (N2) exposed to *E. coli* food source alone (uninfected, black), heat-killed *C. albicans* (grey), or live *C. albicans* (red). Boxes indicate the 25-75th quartiles with median indicated. Error bars are the normalized range of the data and circles indicate outliers. **B)** Number of viable *C. elegans* progeny produced per day in uninfected, heat-killed, and live *C. albicans* treatments. Data represent the mean and SEM. **C)** The fraction of *C. elegans* progeny produced on Day4 or later during the 'Late Reproductive' window. Data represent the mean and SEM. **D)** Box and whiskers plot of average population size (representing the number of F1 and F2 progeny) produced from a single founder *C. elegans*. Boxes indicate the 25-75th quartiles with median indicated. Error bars are the normalized range of the data and circles indicate outliers. The number (n) of experimental samples analyzed is indicated for each treatment. Treatments that share letters are not significantly different, whereas treatments with differing letters are stastically significant, post-hoc Dunn's multiple comparison test.

Figure 3: *C. albicans* **severely impacts host fitness in immunocompromised nematodes.**
A) Box and whiskers plot of brood sizes for *C. elegans* (N2) exposure to *E. coli* food source alone
(uninfected, black), heat-killed *C. albicans* (grey), or live *C. albicans* (red). Boxes indicate the
25-75th quartiles with median indicated. Error bars are the normalized range of the data and
circles indicate outliers. **B)** The fraction of *C. elegans* progeny produced on Day 4 or later during
the 'Late Reproductive' window. Data represent the mean and SEM. **C)** Box and whiskers plot of
average population size (representing the number of F1 and F2 progeny) produced from a single
founder *C. elegans*. Boxes indicate the 25-75th quartiles with median indicated. Error bars are the
normalized range of the data and circles indicate outliers. The number (n) of experimental sam-
ples analyzed is indicated for each treatment. Stastistical significance is indicated when treat-
ments differ. Treatments that share letters are not significantly different, whereas treatments with
differing letters are stastically significant, post-hoc Dunn's multiple comparison tests.

Figure 4. Virulence of *Candida* species depends on host immune function.
Reductions in relative population size for healthy (N2, black bars) and immunocompromised
(*sek-1*, green bars) hosts exposed to *C. albicans*, *C. dublineinsis*, *C. tropicalis*, and *C. parapsilosis*. Dashed line indicates the relative population size of uninfected hosts.
Interactions between *Candida* species and host background was determined by two-way
ANOVA ($p < 0.001$). Treatments that share letters are not significantly different, whereas
treatments with differing letters are stastically significant, post-hoc Tukey's multiple
comparisons test. The number (n) of experimental samples analyzed is indicated for each
treatment in Table S1.