Toxoplasma does not secrete the GRA16 and GRA24 effectors beyond
the parasitophorous vacuole membrane of tissue cysts

Shruthi Krishnamurthy[1], Jeroen PJ Saeij[1*]

[1] Department of Pathology, Microbiology and Immunology, School of Veterinary Medicine,
University of California Davis, One Shields Ave., Davis, 95616, California, USA

* Correspondence:
Corresponding Author
jsaeij@ucdavis.edu

**Keywords: *Toxoplasma gondii*[1], secreted effectors[2], tissue cyst wall[3], tetracycline inducible
expression[4], GRA16[5], GRA24[6].**

Abstract

After invasion, *Toxoplasma* resides in a parasitophorous vacuole (PV) that is surrounded by the PV
membrane (PVM). Once inside the PV, tachyzoites secrete dense granule proteins (GRAs) of which
some, such as GRA16 and GRA24, are transported beyond the PVM likely *via* a putative translocon.
However, once tachyzoites convert into bradyzoites within cysts, it is not known if secreted GRAs
can traffic beyond the cyst wall membrane. We used the tetracycline inducible system to drive
expression of HA epitope tagged GRA16 and GRA24 after inducing stage conversion and show that
these proteins are not secreted beyond the cyst wall membrane.

1   Introduction

Toxoplasma gondii, which belongs to the phylum Apicomplexa, is an obligate intracellular
parasite that can cause disease in immuno-compromised patients and fetuses (Montoya and
Liesenfeld, 2004; Weiss and Dubey, 2009). It is the causative agent of toxoplasmosis, the 2nd most
common cause of food-borne illness in the USA (Jones and Dubey, 2012). Infectious tissue cysts are
present in brain and muscles of many warm-blooded chronically infected hosts (Kim and Weiss,
2004). Infected cats, which are the definitive hosts, shed infectious oocysts in their feces
contaminating food and water sources. Infection is initiated by ingestion of either tissue cysts
containing the bradyzoite life-cycle stage or oocysts (Dubey, 1998). Upon stage conversion into
tachyzoites and invasion of a host cell, *Toxoplasma* forms a parasitophorous vacuole (PV) that is
surrounded by the PV membrane (PVM)(Black and Boothroyd, 2000). During invasion, *Toxoplasma*
secretes effector proteins from rhoptries (ROPs), which mediate invasion, inhibition of host
restriction factors, and modulation of host signaling pathways (Dubremetz, 2007; Boothroyd and
Dubremetz, 2008). Once inside the PV, proteins from the dense granule secretory organelles (GRAs)
are secreted onto, and beyond the PVM into the host cell cytoplasm (Hakimi and Bougdour, 2015).
Three putative translocon proteins: Myc-regulation 1 (MYR1) along with MYR2 and MYR3
determine transport of GRA16 and GRA24 across the PVM into the host cell cytoplasm after which
they traffic to the host cell nucleus (Franco et al., 2016; Marino et al., 2018). In addition to these
putative translocon proteins, an aspartyl protease, ASP5 cleaves many secreted GRA proteins at a
characteristic RRLxx motif also known as the *Toxoplasma* export element (TEXEL) motif which is
important for their localization and function (Coffey et al., 2015; Hammoudi et al., 2015). Most of

these effectors that have been characterized are from the non-orally infectious tachyzoite stage. It is
unclear if bradyzoites within tissue cysts, akin to tachyzoites within the PV, can secrete GRAs
beyond the PVM as the cyst wall is built on the inside of the PVM and presents a potential barrier for
GRA secretion into the host cell.

GRA16, GRA24, GRA28 and IST (*T. gondii* inhibitor of STAT1 transcriptional activity) are
secreted by tachyzoites beyond the PVM and traffic to the host nucleus (Bougdour et al., 2013; Braun
et al., 2013; Gay et al., 2016; Nadipuram et al., 2016; Olias et al., 2016) where they modulate host
signaling pathways important for parasite fitness. In this brief report we use the tetracycline inducible
system to induce the expression of epitope tagged GRA16 and GRA24 after *in vitro* stage
conversion. We observed that ATc induced *GRA16*-HA and *GRA24*-HA do not traffic to host cell
nucleus. Instead, they accumulate within the *in vitro* tissue cysts.

2   Materials and Methods

2.1   Host cells and parasite strain

Human foreskin fibroblasts (HFFs) were used as host cells and were cultured under standard
conditions using Dulbecco Modified Eagle Medium (DMEM) with 10% fetal bovine serum (FBS)
(Rosowski et al., 2011). We chose GT1 parasites expressing Tet-R (Etheridge et al., 2014a) , a type I
strain that is capable of forming cysts during *in vitro* stage conversion (Lindsay et al., 1991; Fux et
al., 2007) induced by pH 8-8.2+low CO_2 (Skariah et al., 2010).

2.2   Plasmid construction

Using Gibson assembly (Gibson et al., 2009), we constructed Tet-On plasmids with a phleomycin
resistance cassette to express GRA16-HA and GRA24-HA. The pTeton vector backbone was
amplified with the following primers which was used to construct both the pTetONGRA16-HA
SRS22A3'UTR as well as the pTetONGRA24-HA SRS22A3UTR constructs:
vector TetON for-GCATCCACTAGTGCTCTTCAAGGTTTTACATCCGTTGCCT
Vector TetON rev-AATTGCGCCATTTTGACGGTGACGAAGCCACCTGAGGAAGAC
The following primers were used to amplify the pieces for pTetONGRA16-HA SRS22A3'UTR Gibson
assembly:
Vector GRA16 for-ACCGTCAAAATGGCGCAATTATGTATCGAAACCACTCAGGGATAC
SRS22UTRGRA16 rev-AATGACAGGTTCAAGCATAATCGGGAACGTCGTATG
GRA16HA SRS22A for-TTATGCTTGAACCTGTCATTTACCTCCAGTAAACATG
SRS22Avector rev-TGAAGAGCACTAGTGGATGCGTTCTAGTGCTGTACGGAAAAGCAAC
The following primers were used to amplify the pieces for pTetONGRA24-HA SRS22A3'UTR Gibson
assembly:
vectorGRA24 for -ACCGTCAAAATGGCGCAATTATGCTCCAGATGGCACGATATACCG
SRS22AUTRGRA24HA rev-AATGACAGGTTTAAGCATAATCGGGAACGTCGTATG
GRA24HASRS22A For-TTATGCTTAAACCTGTCATTTACCTCCAGTAAACATG

2.3   Parasite transfection and selection

The pTetOn vectors containing GRA16-HA and GRA24-HA were linearized using AseI restriction
enzyme. The linearized plasmids (50µg) were electroporated into $5x10^7$ GT1tetR parasites using the
protocol described in (Gold et al., 2015). After lysis of parasites from host cells, they were selected
twice with 50µg/ml phleomycin (Ble) and maintained in 5µg/ml Ble (Krishnamurthy et al., 2016)
until they were cloned out by limiting dilution.

3   Results

To test if GRAs are secreted beyond the tissue cyst wall membrane, we utilized the Tet-inducible
system (Etheridge et al., 2014a) to express an HA-tagged copy of GRA16 or GRA24 under the Tet
operator (Tet-O) in parasites expressing the tetracycline repressor (Tet-R). We could not just use
GRA16-HA or *GRA24*-HA expressed from the endogenous promoter because if we see these proteins
in the host nucleus we would not know if they were secreted beyond the PVM before the cyst wall
was made (as tachyzoites) or after the cyst wall was made (as bradyzoites). In the absence of
anhydrotetracycline (ATc), the Tet-R binds to Tet-O and represses the transcription of either *GRA16*-
HA or *GRA24*-HA under the RPS13 promoter (Etheridge et al., 2014b; Wang et al., 2016). ATc
binds TetR and relieves repression of transcription which allows for the expression of HA-tagged
GRA16 or GRA24. Since ATc is smaller than the size-exclusion limit of the cyst wall (Lemgruber et
al., 2011), we decided to use this system to answer our question.

To check if our constructs were able to stably express functional GRA16 and GRA24, we first
transfected them into the RH parasite strain and observed nuclear localization of these proteins (data
not shown). After transfection of the GT1 Tet-R expressing strain with the Tet-inducible GRA16-HA
or GRA24-HA construct and subsequent selection with phleomycin for stable integration, we show
by IFA that in tachyzoites GRA16-HA and GRA24-HA are only expressed in the presence of ATc
(fig. 1a).
A single parasite clone that expressed GRA16-HA or GRA24-HA only in the presence of
ATc was chosen for induction of stage differentiation *in vitro* in human foreskin fibroblast (HFFs).
The media was switched from DMEM with 10% FBS to tricine-buffered RPMI media with pH 8.0
and put in an incubator with low CO_2 after 24 hours of infection (MOI=0.1) to induce tachyzoite to
bradyzoite stage conversion. Five days post-switching, 2μM of ATc was added to the cultures to
induce the expression of GRA16-HA and GRA24-HA since we observed that at least 50% of the
parasites had converted to cysts by staining the cyst wall with DBA-lectin (Boothroyd et al., 1997)
(data not shown). The parasites were fixed 48 hours following addition of ATc to allow for sufficient
expression of GRA16-HA and GRA24-HA. We performed an indirect immunofluorescence assay
(IFA) to determine the localization of GRA16 and GRA24 using anti-HA antibody as well as DBA-
lectin to detect the cyst wall. We show that in host cells containing tissue cysts, GRA16-HA and
GRA24-HA were not detected in the host cell nucleus or beyond the tissue cyst wall membrane and
that instead they accumulated underneath the cyst wall. Almost 100% of vacuoles we observed were
DBA positive. We decided to observe HFFs only infected with one parasite and therefore containing
only one cyst as differences in the timing of conversion could affect the localization of GRA16 and
GRA24. Out of 142 (59 for GRA16-HA and 83 for GRA24-HA from four biological replicates)
images of singly infected host cells containing DBA positive cysts, both GRA16 and GRA24 were
expressed exclusively beneath the cyst wall only in the presence of ATc (Figs 1b and 2a). We
observed GRA24 localized to the host cell nucleus only in multiple infected cells containing
tachyzoites, along with *in vitro* cysts (Fig. 2b). Thus, our results show that GRA16 and GRA24 are
not secreted beyond the tissue cyst membrane into the host cell.

4   Discussion

We show here that GT1 parasites are able to form DBA lectin positive *in vitro* cysts. We also
show for the first time that ATc is able to cross the cyst wall *in vitro*. Even though bradyzoites within

tissue cysts are not as metabolically active compared to tachyzoites, it is becoming clear that they are
also not in a dormant state (Sinai et al., 2016). However, we observed that bradyzoites do not secrete
GRA16 and GRA24 beyond the *in vitro* cyst wall membrane. These proteins accumulated within the
cyst wall suggesting that their role in the host cell nucleus is not required at this stage. Possibly
bradyzoites require these proteins during natural oral infections after excystation from tissue cysts to
establish infection in gut epithelial cells of the host. Not secreting parasite proteins beyond the cyst
wall might help *Toxoplasma* to remain invisible and undetected by the host immune response during
the chronic phase of infection. Another possibility may be that the translocon proteins MYR1/2/3 or
ASP5 are not sufficiently expressed at this stage to effectively mediate transport of secreted GRAs
beyond the cyst wall membrane. Even though all the MYRs and ASP5 are expressed in tachyzoites,
sporozoites and bradyzoites, their expression is significantly lower in bradyzoites (Marino et al.,
2018). However, even if MYR1-3 and ASP5 are expressed, our data indicate that the cyst wall seems
to act as a barrier as ATc- induced GRA16 and GRA24 accumulated beneath the wall (Fig. 1 and 2).
141

5   Abbreviations

Tet-on: tetracycline inducible system; TetR: tetracycline repressor protein; ATc: anhydrotetracycline;
HA: hemagglutinin epitope tag; YFP: yellow fluorescent protein; Ble: phleomycin; HFF: human
foreskin fibroblast; DBA lectin: dolichos biflorus agglutinin; IFA: indirect immunofluorescence;
GRA: dense granule proteins; ROP: rhoptry proteins; IST: *T. gondii* inhibitor of STAT1
transcriptional activity; PV: parasitophorous vacuole; PVM: PV membrane.

6   Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

7   Author Contributions

SK generated all the data. SK and JS wrote and edited the manuscript.

8   Funding

This research was funded by NIH-2R01AI080621-06A1.

9   Acknowledgments

We thank all the members of the Saeij lab for their feedback on this manuscript. We also thank Dr.
Ronald D. Etheridge for proving us with GT1-TetR parasites.

10   References

Black, M.W., and Boothroyd, J.C. (2000). Lytic Cycle of *Toxoplasma gondii Microbiol Mol Biol Rev*
64(3)**,** 607–623.

Boothroyd, J.C., Black, M., Bonnefoy, S., Hehl, A., Knoll, L.N., Manger, I.D., et al. (1997). Genetic
and biochemical analysis of development in *Toxoplasma gondii*. *Phil.Trans. R. Soc. Lond. B*
353**,** 1347-1354.

Boothroyd, J.C., and Dubremetz, J.F. (2008). Kiss and spit: the dual roles of *Toxoplasma* rhoptries.
Nat Rev Microbiol 6(1)**,** 79-88. doi: 10.1038/nrmicro1800.

Bougdour, A., Durandau, E., Brenier-Pinchart, M.P., Ortet, P., Barakat, M., Kieffer, S., et al. (2013).
Host cell subversion by Toxoplasma GRA16, an exported dense granule protein that targets
the host cell nucleus and alters gene expression. *Cell Host Microbe* 13(4)**,** 489-500. doi:
10.1016/j.chom.2013.03.002.

Braun, L., Brenier-Pinchart, M.P., Yogavel, M., Curt-Varesano, A., Curt-Bertini, R.L., Hussain, T.,
et al. (2013). A Toxoplasma dense granule protein, GRA24, modulates the early immune
response to infection by promoting a direct and sustained host p38 MAPK activation. *J Exp*
Med 210(10)**,** 2071-2086. doi: 10.1084/jem.20130103.

Coffey, M.J., Sleebs, B.E., Uboldi, A.D., Garnham, A., Franco, M., Marino, N.D., et al. (2015). An
aspartyl protease defines a novel pathway for export of Toxoplasma proteins into the host
cell. *Elife* 4. doi: 10.7554/eLife.10809.

Dubey, J.P. (1998). Advances in the life cycle of *Toxoplasma gondii*. *Int J Parasitol* 28**,** 1019-1024.

Dubremetz, J.F. (2007). Rhoptries are major players in *Toxoplasma gondii* invasion and host cell
interaction. *Cell Microbiol* 9(4)**,** 841-848. doi: 10.1111/j.1462-5822.2007.00909.x.

Etheridge, R.D., Alaganan, A., Tang, K., Lou, H.J., Turk, B.E., and Sibley, L.D. (2014a). The
Toxoplasma pseudokinase ROP5 forms complexes with ROP18 and ROP17 kinases that
synergize to control acute virulence in mice. *Cell Host Microbe* 15(5)**,** 537-550. doi:
10.1016/j.chom.2014.04.002.

Franco, M., Panas, M.W., Marino, N.D., Lee, M.C., Buchholz, K.R., Kelly, F.D., et al. (2016). A
Novel Secreted Protein, MYR1, Is Central to Toxoplasma's Manipulation of Host Cells. *MBio*
7(1)**,** e02231-02215. doi: 10.1128/mBio.02231-15.

Fux, B., Nawas, J., Khan, A., Gill, D.B., Su, C., and Sibley, L.D. (2007). Toxoplasma gondii strains
defective in oral transmission are also defective in developmental stage differentiation. *Infect*
Immun 75(5)**,** 2580-2590. doi: 10.1128/IAI.00085-07.

Gay, G., Braun, L., Brenier-Pinchart, M.P., Vollaire, J., Josserand, V., Bertini, R.L., et al. (2016).
Toxoplasma gondii TgIST co-opts host chromatin repressors dampening STAT1-dependent
gene regulation and IFN-gamma-mediated host defenses. *J Exp Med* 213(9)**,** 1779-1798. doi:
10.1084/jem.20160340.

Gibson, D.G., Young, L., Chuang, R.Y., Venter, J.C., Hutchison, C.A., 3rd, and Smith, H.O. (2009).
Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Methods* 6(5)**,**
343-345. doi: 10.1038/nmeth.1318.

Gold, D.A., Kaplan, A.D., Lis, A., Bett, G.C., Rosowski, E.E., Cirelli, K.M., et al. (2015). The
Toxoplasma Dense Granule Proteins GRA17 and GRA23 Mediate the Movement of Small
Molecules between the Host and the Parasitophorous Vacuole. *Cell Host Microbe* 17(5)**,** 642-
652. doi: 10.1016/j.chom.2015.04.003.

Hakimi, M.A., and Bougdour, A. (2015). *Toxoplasma*'s ways of manipulating the host transcriptome
via secreted effectors. *Curr Opin Microbiol* 26**,** 24-31. doi: 10.1016/j.mib.2015.04.003.

Hammoudi, P.M., Jacot, D., Mueller, C., Di Cristina, M., Dogga, S.K., Marq, J.B., et al. (2015).
Fundamental Roles of the Golgi-Associated Toxoplasma Aspartyl Protease, ASP5, at the
Host-Parasite Interface. *PLoS Pathog* 11(10)**,** e1005211. doi: 10.1371/journal.ppat.1005211.

Jones, J.L., and Dubey, J.P. (2012). Foodborne toxoplasmosis. *Clin Infect Dis* 55(6)**,** 845-851. doi:
10.1093/cid/cis508.

Kim, K., and Weiss, L.M. (2004). *Toxoplasma gondii*: the model Apicomplexan,. *Int J Parasitol*
34(3). doi: 10.1016/j.ijpara.2003.12.009.

Krishnamurthy, S., Deng, B., Del Rio, R., Buchholz, K.R., Treeck, M., Urban, S., et al. (2016). Not a
Simple Tether: Binding of Toxoplasma gondii AMA1 to RON2 during Invasion Protects
AMA1 from Rhomboid-Mediated Cleavage and Leads to Dephosphorylation of Its Cytosolic
Tail. *MBio* 7(5). doi: 10.1128/mBio.00754-16.

Lemgruber, L., Lupetti, P., Martins-Duarte, E.S., De Souza, W., and Vommaro, R.C. (2011). The
organization of the wall filaments and characterization of the matrix structures of Toxoplasma
gondii cyst form. *Cell Microbiol* 13(12)**,** 1920-1932. doi: 10.1111/j.1462-5822.2011.01681.x.

Lindsay, D.S., Dubey, J.P., Blagburn, B.L., and Toivio-Kinnucan, M. (1991). Examination of Tissue
Cyst Formation by *Toxoplasma gondii* in Cell Cultures UsingBradyzoites, Tachyzoites, and
Sporozoites. *J Parasitol* 77(1)**,** 126-132.

Marino, N.D., Panas, M.W., Franco, M., Theisen, T.C., Naor, A., Rastogi, S., et al. (2018).
Identification of a novel protein complex essential for effector translocation across the
parasitophorous vacuole membrane of Toxoplasma gondii. *PLoS Pathog* 14(1)**,** e1006828.
doi: 10.1371/journal.ppat.1006828.

Montoya, J.G., and Liesenfeld, O. (2004). Toxoplasmosis. *The Lancet* 363(9425)**,** 1965-1976. doi:
10.1016/s0140-6736(04)16412-x.

Nadipuram, S.M., Kim, E.W., Vashisht, A.A., Lin, A.H., Bell, H.N., Coppens, I., et al. (2016). In
Vivo Biotinylation of the Toxoplasma Parasitophorous Vacuole Reveals Novel Dense
Granule Proteins Important for Parasite Growth and Pathogenesis. *MBio* 7(4). doi:
10.1128/mBio.00808-16.

Olias, P., Etheridge, R.D., Zhang, Y., Holtzman, M.J., and Sibley, L.D. (2016). Toxoplasma Effector
Recruits the Mi-2/NuRD Complex to Repress STAT1 Transcription and Block IFN-gamma-
Dependent Gene Expression. *Cell Host Microbe* 20(1)**,** 72-82. doi:
10.1016/j.chom.2016.06.006.

Rosowski, E.E., Lu, D., Julien, L., Rodda, L., Gaiser, R.A., Jensen, K.D., et al. (2011). Strain-
specific activation of the NF-kappaB pathway by GRA15, a novel Toxoplasma gondii dense
granule protein. *J Exp Med* 208(1)**,** 195-212. doi: 10.1084/jem.20100717.

Sinai, A.P., Watts, E.A., Dhara, A., Murphy, R.D., Gentry, M.S., and Patwardhan, A. (2016).
Reexamining Chronic Toxoplasma gondii Infection: Surprising Activity for a "Dormant"
Parasite. *Curr Clin Microbiol Rep* 3(4)**,** 175-185. doi: 10.1007/s40588-016-0045-3.

Skariah, S., McIntyre, M.K., and Mordue, D.G. (2010). Toxoplasma gondii: determinants of
tachyzoite to bradyzoite conversion. *Parasitol Res* 107(2)**,** 253-260. doi: 10.1007/s00436-
010-1899-6.

Wang, J.L., Huang, S.Y., Behnke, M.S., Chen, K., Shen, B., and Zhu, X.Q. (2016). The Past, Present,
and Future of Genetic Manipulation in Toxoplasma gondii. *Trends Parasitol* 32(7)**,** 542-553.
doi: 10.1016/j.pt.2016.04.013.

Weiss, L.M., and Dubey, J.P. (2009). Toxoplasmosis: A history of clinical observations. *Int J*
Parasitol 39(8)**,** 895-901. doi: 10.1016/j.ijpara.2009.02.004.

11   Figure Legends

Figure 1a. **GT1- TetR parasites express GRA16-HA and GRA24-HA only in the presence of**
ATc. Coverslips with a monolayer of HFFs were infected in the presence or absence of 2μM ATC
with GT1-TetR parasites or single clones of parasites expressing GRA16-HA and GRA24-HA under
the RPS13 promoter downstream of a Tet-O7 operator. The coverslips were fixed 16 hours post
infection with 3% formaldehyde and processed for IFA using rabbit anti-HA primary (Roche)
antibody followed by goat anti-rabbit Alexa-555 secondary antibody and Hoechst (invitrogen). Tet-R
is YFP tagged and GT1 tet-R parasites do not express any proteins that are HA-epitope tagged (first
panel). In the absence of ATc, Tet-R represses the expression of GRA16-HA (second panel) and
GRA24-HA (fourth panel). Repression by Tet-R is relieved only in the presence of 2μM ATc (third
and fifth panel) allowing for the proper localization of GRA16-HA and GRA24-HA to the parasite
PVM and host cell nucleus (yellow arrows). Images are scaled to 10μm.

Figure 1b. ***In vitro* DBA positive cysts do not secrete GRA16-HA beyond the cyst wall**
membrane. HFFs were infected with a single clone of GT1-tetR parasites that expressed GRA16-
HA only in the presence of ATc. After 24 hours of infection, stage differentiation was induced as
indicated in the text. After 5 days, 2 μM ATc was added and 2 days later the coverslips were fixed
with 100% cold methanol (also eliminates YFP signal from TetR) and processed for IFA. The cyst
wall was stained with DBA-FITC along with HA and Hoechst. In the absence of ATc, GRA16-HA
was not detected in parasites (first panel). GRA16-HA accumulated beneath the cyst wall only in the
presence of ATc (second to fourth panels). Images are scaled to 10μm.

Figure 2. **GRA24-HA accumulates underneath the DBA positive cyst wall**. (a) Same IFA protocol
was followed as described for figure 2. In the absence of ATc, there was no expression of GRA24-
HA (first panel). In the presence of ATc, GRA24-HA accumulates beneath the cyst wall (second to
fourth panel). (b) Multiple infected HFFs with tachyzoites and four *in vitro* cysts of parasites that
express GRA24-HA showing nuclear localization. Yellow arrows indicate GRA24-HA expressing *in*
vitro cysts. DBA-FITC staining is faint since 3% formaldehyde was used to fix parasites. Images are
scaled to 50μm.

Anti-HA

Anti-HA
TetR-YFP
Hoechst

| DBA-FITC | Anti-HA | Hoechst | Merge |