Dispensing a synthetic green leaf volatile to two plant species in a common garden differentially alters physiological responses and herbivory

Grace E. Freundlich[1], Maria Shields[1], Christopher J. Frost[1,2*]

[1] Department of Biology, University of Louisville, Louisville, KY, USA 40292
[2] Current Affiliation: BIO5 Institute, University of Arizona, Tucson, AZ, USA 85721-0240

*Corresponding author:

Christopher J. Frost
BIO5 Institute
University of Arizona
Tucson, AZ 85721-0240
Email: jasmonate@gmail.com

Abstract

Herbivore-induced plant volatile (HIPV)-mediated eavesdropping by plants is a well-documented, inducible phenomenon that has practical agronomic applications for enhancing plant defense and pest management. However, as with any inducible phenomenon, responding to volatile cues may incur physiological and ecological costs that limit plant productivity. In a common garden experiment, we tested the hypothesis that exposure to a single HIPV would decrease herbivore damage at the cost of reduced plant growth and reproduction. Lima bean (*Phaseolus lunatus*) and pepper (*Capsicum annuum*) plants were exposed to a persistent, low-dose (~10ng/hour) of the green leaf volatile *cis*-3-hexenyl acetate (*z3*HAC), which is an HIPV and damage-associated volatile. *z*3HAC-treated pepper plants were shorter, had less aboveground and belowground biomass, and produced fewer flowers and fruits relative to controls while *z*3HAC-treated lima bean plants were taller and produced more leaves and flowers than did controls. Natural herbivory was reduced in *z3*HAC-exposed lima bean plants, but not in pepper. Cyanogenic potential, a putative direct defense mechanism in lima bean, was lower in young *z3*HAC-exposed leaves, suggesting a growth-defense tradeoff from *z3*HAC exposure alone. Plant species-specific responses to an identical volatile cue have important implications for agronomic costs and benefits of volatile-mediated inter-plant communication under field conditions.

Key Words. *Phaseolus lunatus, Capsicum annuum* var. Cayenne, Fabaceae, Solanaceae, common garden, field study, *cis-3*-hexenyl acetate

Introduction

Production and utilization of airborne chemical cues is prevalent within the plant kingdom. Plants depend on airborne chemical signaling for pollination [1], indirect defense [2], protection from pathogens [3], and herbivore resistance [4]. Volatile communication is also pivotal for plant-plant signaling, and selection for such signaling depends on honest cues that reliably confer ecologically relevant information. For example, herbivory is a fundamental ecological interaction that impacts plant fitness, and many plants increase the production and emission of volatile compounds in response to herbivore damage [5]. Such herbivore-induced plant volatiles (HIPVs) are potentially reliable cues around which plant-plant eavesdropping could be evolutionarily adaptive [6]. Undamaged plants (or parts of the same plant [7,8]) eavesdropping on HIPVs from a plant experiencing herbivory may directly trigger stress responses [9-11], or alternatively prime responses for future potential herbivory [4,7].

HIPV-mediated eavesdropping appears to be a common phenomenon. For example, HIPVs prime or induce corn [12,13], tomato [14], poplar [4,7], blueberry [15] and lima bean [16,17] against herbivory. HIPVs can be diverse and taxa-specific [18,19], but are often comprised of monoterpenes, sesquiterpenes, benzenoids and green leaf volatiles GLVs [20,21]. In contrast to volatile terpenes and benzenoids [18,22], GLVs are immediately released into the airspace whenever leaves are mechanically damaged [23] serving as early indicators of wounding and herbivory. GLV exposure alters gene expression profiles related to specialized metabolite production and accumulated secondary metabolite precursors in preparation for inducing resistance [24]. For example, the GLV *cis*-3-hexenyl acetate (*z*3HAC) induces transcriptional changes in poplar [4] and maize [21] that prime oxylipin signaling and induced resistance. Like other GLVs, *z*3HAC can be emitted from wounded leaves alone, but may also represent a

reliable cue because it is typically released from herbivore-damaged leaves in a variety of species

[23], including tomato [14], maize [21], Arabidopsis [25], lima bean [8,17], pepper [26,27], and

poplar [4,28].

Costs incurred by plants responding to airborne HIPV cues alone are largely unknown.

Plant defense theory posits that induced resistance by plants against herbivores is a cost-savings

strategy to restrict the deployment of costly specialized defensive metabolites until necessary

[29,30]. However, inducible resistance generates a period of vulnerability between the time of

attack and the upregulation of resistance [31]. Perception of early reliable cues may overcome

such a vulnerability by allowing a plant to anticipate a probable attack and prime defenses before

herbivory occurs. Since HIPV-mediated priming is an inducible phenomenon, theory predicts

that responding to reliable cues alone should incur costs that outweigh and select against

maintaining a "primed state" [32,33]. In other words, perception of a priming stimulus, such as an

HIPV, may induce physiological changes that incur costs that are less expensive than induced

resistance itself. Previous work with non-volatile priming agents β-amino butyric acid (BABA)

[34] and snail mucus [35] both support this prediction. Similarly, costs associated with volatile

perception alone that initiate priming should be less severe than costs of induced resistance to

actual herbivory [36]. Yet there is currently limited experimental evidence of such costs with

respect to anti-herbivore volatile cues. For example, bacterial-derived volatiles 3-pentanol and

2-butanone increased fresh fruit weight in field-grown *Cucumis sativa* [37], wild tobacco

(*Nicotiana attenuata*) exposed to airspace of experimentally clipped sagebrush produce more

seeds (i.e., higher presumptive fitness) relative to control plants [38]. These results suggest that

ecological costs of exposure to volatile cues may be context dependent, but comparative

cost/benefit tradeoffs for perception of HIPVs alone among sympatric field-grown plants is

currently lacking.

Here, we report a common garden field experiment with lima bean (*Phaseolus lunatus*)

and chili pepper (*Capsicum annuum*) testing the hypothesis that field plants subject to a

persistent dose of an ostensibly reliable volatile cue incur consistent costs reflected in reduced

growth and reproduction. We treated individuals of both species to repeated low-dose

applications of *z3*HAC and measured their growth, reproduction, and herbivore damage

throughout the growing season. We predicted that exposure to *z*3HAC—regardless of plant

species identity—would reduce growth and reproductive output, while also reducing natural

herbivory.

Materials and Methods

Study Site and Plants

A common garden experiment was established on a 54m^2 plot within Blackacre

Conservancy's community garden in Louisville, Kentucky (38°11'33.8"N 85°31'28.3"W;

Supplemental Fig. 1). The field site was enclosed in a mesh fence to exclude mammalian

herbivores. *Phaseolus lunatus,* Fabaceae, variety Fordham Hook 242 ('lima bean') and the

Capsicum annuum, Solanaceae, variety Cayenne pepper, Joe Red Long ('pepper') were chosen

as phylogenetically distinct model plants with previously established defense profiles [39,40].

Seeds were purchased from the Louisville Seed Company (Louisville, KY, USA), and

germinated in Metromix 510© in May 2016 in the Biology Department's greenhouse. After

reaching ~20cm in height, 132 lima bean plants were transplanted to the field May 30, 2016, at 4

weeks old and 98 pepper plants at 8 weeks old were transplanted to the field June 28, 2016.

While both species were started at the same time, peppers were placed in the field later than the

lime beans because they needed additional maturation time before transferring to the field.

Within the field site, plants were planted in alternating rows of twos of lima bean and pepper.

Distance limitations exist regarding plant volatile perception [41] and consequent herbivore

resistance [7,42]. Previous studies with sagebrush [43] and lima bean [44] indicate that volatile

cues are effective over relatively short distances of less than 100cm. Therefore, all plants in our

experiment were spaced one meter apart from one another in all directions to reduce the risk of

interplant communication and cue crossover.

Volatile exposure manipulations

Plants were acclimated to the field for one week after planting before volatile treatments began.

To simulate a naturally occurring low dose [45,46], plants were exposed to lanolin infused with

the equivalent of 10ng/hr of *z*3HAC, a concentration 25% of that which previously primed

poplar [4] and maize [21]. A treatment vial contained 50mg of a 30ng/ul *z*3HAC/lanolin, while a

control vial contained 50mg of lanolin. Each glass vial had a 9mm aperture and was maintained

at -80°C until use. Each week, both the *z*3HAC-infused lanolin vials and lanolin-only controls

were placed at the bottom of their respective plants. Each vial was inverted and supported with a

wire stand and each vial was wrapped in aluminum foil to reduce photodegradation [47]

(Supplemental Fig. 2). These vials were left in the field and replaced every seven days for the

duration of the field season. Plants were randomly assigned to either *z*3HAC treatment (lima

bean n=63; pepper n=35) or lanolin control (lima bean n=72; pepper n=43). The unit of

replication was an individual plant and each plant received its own vial. Random assignment of

treatments was made using blocks of 4 adjacent plants; block was included as a random factor in

statistical models, and was not a significant factor in any of the models.

Growth, biomass, and reproduction measurements

We measured height and total leaf counts routinely on the experimental plants. Leaves

were only counted if they were wider than 2cm across for both species while height

measurements for both species were recorded from the base of the plant to the uppermost

branching point. For lima bean, height was determined by measuring the longest runner within

the bush, while pepper plants were measured from the base of the main stalk to the highest

branching point. Along with height, the total number of leaves per plant was measured

throughout the field season. A complete biomass harvest was conducted on pepper for leaves,

roots, and stems at the end of the field season. All leaves and fruits were separated into paper

bags before individual plants were extracted from of the ground. After removal, roots and stems

were separated, roots were washed with water to remove dirt, and placed into separate paper

bags. All materials were dried at 60°C for 24 hours and then weighed. A biomass harvest for

lima bean was not performed because an *Epliachna varivestis* (Mexican Bean Beetle) outbreak

late in the season removed much of the leaf tissue before we could determine reliable biomass

measurements.

We measured total flower and fruit production in both species. Flowers were recorded if

they were true flowers with fully mature pistils and stamen. If a flower was not fully mature, it

was recorded as a flower bud. Fruits were recorded as soon as fruit development was observed

with either initial pod or exocarp development. Throughout the field season, fruit and flower

counts per plant were recorded along with the number of mature and immature fruits.

From the fruits harvested from the final biomass harvest, ~10 randomly selected, mature

fruits from each pepper plant were chosen for seed count analysis (188 fruits from z3HAC-

treated plants and 210 fruits from controls). Dried fruits were dissected with a scalpel and all

seeds were isolated and counted.

Herbivory

Leaf chewing damage was assessed for both pepper and lima bean as percent leaf area

removed (LAR) using a visual estimation technique [48,49] with the following damage categories:

0%, 0-5%, 5-15%, 15-30%, 30-50%, 50-70%, 70-90%, and >90%. For each damage assessment,

every leaf on a plant was categorized into one of the damage categories, and an overall percent

damage was determined as a weighted average of all leaves. Plants were also routinely

monitored for the presence of naturally occurring chewing and piercing/sucking herbivores. In

particular, we observed an ephemeral, natural occurrence of the black bean aphid (*Aphis faba*),

and recorded its presence/absence on lima bean plants in the field.

Leaf collections and cyanide measurements

Since cyanogenic potential (CP) is an inducible herbivore defense in lima bean [50,51], we

used CP as a metric for induced responses in the presence of *z3*HAC. We collected source and

sink leaves [52,53] on July 10, 2016, which was approximately 6 weeks into the field season. We

developed a novel protocol in the lab for microscale colorimetric CP quantification by modifying

an existing macro-scale protocol [54,55]. Briefly, 5mg of lyophilized tissue was mixed with

200uL of citrate buffer (0.1 M, pH 5.5-6.5) in a 2mL centrifuge tube, into which a 200uL glass

vial (Agilent #5183-2090) containing 100uL 1.0M NaOH was also placed and the centrifuge

tube caped completely. After 15 hours, a 50uL aliquot of the 1.0M NaOH in the inner glass vial

was diluted to 0.1M, and a 30uL aliquot was neutralized with 30uL 0.5M acetic acid in a 96-well

reaction plate. Then, 75 µL of Reagent A (5 mg/ml succinimide [VWR, AAA13503] and 0.5

mg/ml n-chlorosuccinimide [VWR, AAA10310]) and then 30 µL of Reagent B (30 mg/ml

barbituric acid [VWR, BT134930] in 30% pyridine [Sigma, 270970]) was added to each well.

After 8 min, absorbance at 580nm was measured on a plate reader (Molecular Devices

SpectraMax M2). Quantification of CP was made against a standard curve of NaCN (VWR,

BT212960).

Statistical analyses

All statistical analyses were performed in R (version 3.4.2) with the lme4 and multcomp

packages. Growth data, such as plant height, leaf area removed, and flower counts, were

analyzed using repeated measures ANOVA with the aov function with a Gaussian distribution.

For repeated measures analyses, we treated date as a within-subjects effect and treatment as a

between-subjects effect for all analyses. Differences between treatments at each individual time

point, as well as all biomass data, were analyzed using one-way ANOVA (glmer function)

followed by a Tukey's post hoc comparison. Cyanogenesis data were log transformed to satisfy

assumptions of normality before being analyzed by a glm model followed by a Tukey's post hoc

comparison.

Results

z3HAC differentially effects growth of lima bean and pepper plants

Treatment with *z3*HAC differentially affected the growth of lima bean and pepper plants.

On average, *z3*HAC-treated lima bean grew 11% taller compared to control plants throughout the

field season (Fig.1A; $F_{2,927}$=9.688, *P*=0.002) and produced 17% more leaves overall than did

controls (Fig.1B; $F_{1,571}$=4.339, *P*=0.038). In contrast, *z3*HAC-treated pepper plants were 12%

shorter relative to controls (Fig.1C; $F_{1,48}$=6.168, *P*=0.017) and produced 23% fewer leaves over

the field season (Fig.1D; $F_{1,237}$=21.58, *P*<0.001). Consistent with height and leaf counts, *z3*HAC

treatment reduced overall biomass of pepper plants by 24% on average (Fig.2). When we

destructively harvested all pepper plant biomass at the end of the season, *z3*HAC-treated pepper

plants had lower leaf, stem, and root dry biomass by 21%, 31%, 29%, respectively (Fig.2A-C)

(Z=3.379, *P*<0.001; Z=-2.035, *P*=0.042; Z=-2.379, *P*=0.017). Despite these *z3*HAC-mediated

effects on biomass exposure, the aboveground-to-belowground biomass ratio was similar

regardless of treatment (Fig.2D; Z=0.31, *P*=0.757). That is, pepper plants treated with *z3*HAC

were smaller relative to control plants.

z3HAC reduces reproductive output in pepper plants

*z3*HAC treatment also differentially affected reproductive output between the two

species, and lowered fruit output in pepper. Flower production was 30% higher in lima bean

plants exposed to *z3*HAC (Fig.3A; $F_{1,576}$=15.044, *P*<0.001), while *z3*HAC-treated peppers

produced 37% fewer flowers relative to control plants at the end of the field season (Fig.3B;

$F_{1,43}$=14.48, *P*<0.001). *z3*HAC-treated pepper plants also produced 23% fewer fruits overall

relative to controls (Fig.4A; Z=-2.035, *P*=0.042), and the fruits that were produced by *z3*HAC-

treated plants had lower wet and dry masses (Fig.4B-C; Z=-2.88, *P*=0.004; Z=-2.439, *P*=0.015),

and 10% lower total seed counts (Fig.4D; Z=3.524, *P*<0.001) and total seed masses (Fig.4E;

Z=3.334, $P<0.001$), relative to controls. That said, the ratio of seed mass to fruit mass was

similar between z3HAC-treated and control plants (Fig.4F; Z=0.588, $P=0.807$), as was mass of

an individual pepper seed (Supplemental Fig.3). There was no apparent difference in lima bean

pod production (Supplemental Fig.3), though an unexpected field-wide premature pod drop

independent of treatment prevented us from fully determining lima bean pod/seed production

with confidence.

z3HAC exposure reduces herbivory on lima bean

z3HAC exposure reduced natural herbivory in lima bean but not pepper plants. Chewing

herbivory on pepper plants was low overall and statistically higher in z3HAC-treated plants only

during the first sampling date (Fig.5A, $F_{1,193}=5.627$, $P=0.019$).   In contrast, chewing damage to

lima bean leaves increased as the field season progressed, with z3HAC-treated plants having

overall 26% less chewing damage than did control plants (Fig.5B; $F_{1,539}=21.745$, $P<0.001$). In

addition to chewing herbivory, black bean aphids (*Aphis faba*) colonized 87% of the z3HAC-

treated lima bean plants, compared with only 21% of control plants (Fig.5C; $\chi^2 = 50.11$, df=1,

$P<0.001$). *A.faba* colonized early in the season and was only observed June 15-31 (Julian dates

166-181) because a heavy rainfall event reduced their population to undetectable levels.

Piercing/sucking herbivores were rare for the remainder of the experiment.

Cyanogenic potential in lima bean is decreased by z3HAC exposure

Exposure to *z3*HAC reduced cyanogenic potential in lima bean plants. *z3*HAC exposure

alone reduced cyanide concentration by 50% in sink leaves (Figure 6A; Z= -0.3257, $P=0.003$),

but not significantly in source leaves (Fig. 6B; $Z=-0.14387$, $P=0.146$). Baseline cyanogenic

potential was ~2x greater in sink leaves compared to source leaves (Figure 6).

Discussion

We show that a persistent, low-dose application of $z3$HAC differentially affects growth

and reproduction of two plant species grown in the same field. Based on previous work on plant

and sensory perception of volatiles [7,21], we hypothesized $z3$HAC application would decrease

growth and reproductive fitness in both plant species. The rationale for this hypothesis was a

central assumption of induced resistance theory that ecological costs modulate the deployment

particular defensive phenotypes until necessary [30,31,56-60]. Volatile-mediated priming, even if

regulated by a different mechanism from resistance [61], is an inducible phenomenon that

theoretically should incur such fitness costs [36]. Yet, our results clearly indicate that pepper and

lima bean had divergent fitness outcomes when subjected to a single GLV under identical field

conditions. Whereas $z3$HAC-treated pepper plants had reduced growth (Fig. 1) and no effect on

herbivore resistance (Fig. 5A) relative to controls, $z3$HAC-treated lima bean plants grew more

and produced more flowers (Fig. 1,3), and suffered less chewing herbivory (Fig. 5B) compared

to controls. This result—that some plants experience costs while others have minimal or even

positive effects when exposed to the same HIPV—has important implications for how volatile

cues may structure interspecific competition and ecological communities. HIPVs alone may be

sufficient to result in differential fitness effects among species. Moreover, the positive effects of

$z3$HAC on lima bean growth and flowering opens agronomic opportunities to exploit plant

volatiles to enhance both growth and pest resistance of important crop plants.

What might affect the response of plants to volatile exposure? One possibility is that the

signal integrity of HIPVs varies among plant species, which influences plant sensory perception

and the outcome of defense priming. That is, z3HAC provides different information to different

plants. Previous work on the role of HIPVs in plant anti-herbivore resistance focused on

priming-mediated defense with consistent results in wheat [62,63], corn [12,45], lima bean [8,11,64],

tomato [65], blueberry [15], sagebrush [43], Arabidopsis [66] and poplar [4]. In contrast, we

specifically focused on indicators of plant fitness in lima bean and pepper in a common garden

experiment. z3HAC treatment alone increased growth and flowering in lima bean, while

reducing growth and reproductive output in pepper (Fig. 3 and 5). This season-long evidence

illustrates that informational integrity on an HIPV varies between these two plant species. Such

divergent fitness effects from exposure to a single ubiquitous herbivore-associated cue

underscore the potential for functional similarity in the mechanisms by which plants modulate

responses to herbivory and volatile indicators of herbivory.

Flower and fruit production is a key component of plant fitness potential. We show that

z3HAC treatment alone differentially affected flower production in lima bean and pepper (Fig.

3). Insect herbivory can increase or decrease floral production depending on the system and

environmental conditions [67-69]. Whereas increased flower production is a strategy assumed to

ameliorate fitness losses in the presence of an environmental stress [56,70], decreased flower

production may be related to costs of chemically mediated defense [71]. Previous work with lima

bean has demonstrated stress-mediated compensation [72]. Our result that z3HAC alone was

sufficient to trigger increased flowering is consistent with this observation, suggesting that

z3HAC alone may stimulate a long-term stress response similar to herbivory. Therefore, even

though the mechanisms underlying *z*3HAC-mediated effects on flower and fruit production are

not yet known, they may be similar to those induced by herbivory [67].

Resource allocation between different tissues is pivotal for growth, reproduction, and

defense, and can be influenced by environmental stress. For example, direct herbivory alters

resource allocation between aboveground tissue and belowground tissue [49,73,74], as does

application of the anti-herbivore phytohormone jasmonic acid (JA) [75,76]. Volatile cues can also

affect biomass allocation. For example, barley exposed to volatiles from unwounded neighboring

plants of different cultivars increases root and leaf biomass [77], while exposure to volatiles

decreases aboveground biomass in other systems [78,79]. In our case, volatile treatment reduced

overall aboveground and belowground biomass in pepper, but did not appear to alter overall

biomass allocation patterns. Simply put, *z*3HAC-treated pepper plants were smaller overall, and

therefore produced fewer seeds.

Differential investment between growth and defense is vital for maximizing limited

resources and, ultimately, fitness. Inducible defenses in plants against herbivores and pathogens

modulate such growth/defense tradeoffs in a number of plant species [31]. In some cases,

exposure to VOCs alone is sufficient to affect such tradeoffs. For example, sagebrush exposed to

VOCs from damaged conspecifics had decreased growth [80], while HIPV-exposed tobacco had

increased herbivore resistance but decreased seed set [38]. In our experiment, *z*3HAC exposure

decreased cyanogenic potential in sink leaves (Fig. 6), which were 20-30% higher than in source

leaves. These results suggest that persistent exposure to *z*3HAC led to allocation shifts from

cyanogenic potential (a putative defense) towards growth (Fig. 1 and 3), as well as supporting

ontogeny-mediated cyanogenic potential [52,81].. That said, *z*3HAC-treated plants also

experienced less natural damage primarily on mature source leaves than did controls (Fig. 5,

personal observations), suggesting that the herbivory was deterred by defense mechanism other

than cyanogenic potential in source leaves.

Volatile cues may impact ecological communities in both expected and pleiotropic ways.

HIPVs are well-established mediators of multi-trophic antagonistic and mutualistic interactions

[82-84], and manipulations of chemical signals and volatile blends have been used for biological

control in a wide range of systems [85,86]. For example, HIPV-infused sticky traps in a grape

(*Vitis vinifera*) orchard differentially attracted lacewings, hoverflies, and parasitoids [87].

Exogenous GLV manipulation using "dispensers" under field conditions altered the arthropod

community composition in maize [47]. In our study, *A.faba* were clearly and unexpectedly

attracted to *z*3HAC-exposed plants (Fig. 5). Under glasshouse conditions, *A.faba* were repelled

by *z*3HAC alone [88], which suggests that the cue that mediated attraction was not our treatment

alone. It is tempting to speculate that aphid attraction combined with reduced chewing herbivory

in lima bean may be reflective of *z*3HAC effects on JA and salicylic acid (SA) signaling, which

would be consistent with a JA-SA tradeoff [89,90]. Ultimately, however, the utility of GLVs (or

other VOCs) in field applications will depend on understanding community-level effects of

volatile exposure.

As a caveat, volatile identity, concentration, and duration may affect the reliability of a

cue and therefore the costs associated with eavesdropping. Plants experiencing insect herbivory

frequently generate species-specific blends of volatile compounds [91,92], which can influence

fitness in neighboring plants [80,93,94]. Plant-derived compounds associated with herbivory

include GLVs [4,12,78], shikimate derivatives [13], and terpenes [95]. However, individual

compounds within a blend can affect plant defense and priming as much as the blend itself. We

used *z*3HAC in this study because it is released herbivore-damaged leaves [21], which ostensibly

allows *z*3HAC to confer reliable ecological information. That said, *z*3HAC can also be released

from wounded leaves without herbivores [91] [23]. However, plants detect and respond

specifically to *z*3HAC [4], and the costs associated with that response in the context of pest

resistance were the focus of this investigation. Additionally, concentration of a cue may

influence plant resistance [66,78]. For example, a repeated, low-dose exposure to a GLV blend

enhances plant resistance compared to a single application [46] while *z3*HAC emissions can be as

high as 66 ng/cm^3 after herbivory [20]. For these reasons, we chose to use a low-dose exposure to

*z*3HAC (25% of the concentration that primed poplar [4] and maize [21]), and still observed

divergent fitness effects between the two plant species (Figs.1, 3, and 5).

In summary, our key finding is that persistent application of a low dose of a single

volatile compound *z*3HAC, a common HIPV and GLV, in field conditions leads to divergent

growth and reproductive fitness effects between two plant species. This result underscores the

variable nature of volatile-mediated eavesdropping, and that plants may have evolved species-

specific mechanisms for responding to volatile cues. Given natural variation among species,

future work must assess costs with other species as well as within accessions and landraces of

our model plants over multiple years. Ultimately, the adaptive significance of eavesdropping for

enhancing plant immunity will depend on plant life history, physiology, and other ecological

factors to determine whether a plant will benefit from eavesdropping VOCs or not, and therefore

what impact volatile-mediated eavesdropping might have on plants and their insect pests.

Supplemental Material. Supplemental Tables describing the outputs of statistical analyses are

listed as tabs in a single spreadsheet ("Supp_Tables"), with tab names corresponding to each

table. Supplemental Figure 1 ("Supp_Fig1"). Aerial view of the field site within the Blackacre

Community Garden. The research area is outlined and the original picture was acquired from

Google Earth. Supplemental Figure 2 ("Supp_Fig2"). Volatile administration setup. Each 2mL

glass vial was inverted and supported by wire stands to prevent rain water accumulation.

Additionally, each vial was wrapped in aluminum foil to prevent photodegradation [47].

Supplemental Figure 3 ("Supp_Fig3"). Calculated individual seed mass for *C.annuum* (A) and

total pod production for *P.lunatus* (B). Box plots represent the raw data ranging from the upper

to the lower quartiles and the median. White boxes represent control plants; gray boxes

represent plants treated with *z*3HAC. Error bars represent the 5% and 95% of the data, and

individual dots are observations that fell outside of those parameters. *P*-values represent Tukey's

HSD comparisons.

Funding. This work was supported by the National Science Foundation grants IOS-1656625 and

IOS-2101059 to C.F.

Acknowledgments. We are grateful to Andrea Clavijo-McCormick for the invitation to submit

our work to this special issue. We thank Allie Peot and Abhinav Maurya for field assistance, and

to A.Peot for assistance processing the plant materials in the laboratory. Comments from Heidi

Appel, Amy Austin, and anonymous reviewers greatly improved the manuscript. We are

grateful to A. Dale Josey and Susan Ballerstedt for permission and logistical support to work in

the community gardens at the Blackacre Conservancy.

References

1. Muhlemann, J.K.; Klempien, A.; Dudareva, N. Floral volatiles: from biosynthesis to function. *Plant Cell and Environment* **2014**, *37*, 1936-1949, doi:10.1111/pce.12314.

2. Mumm, R.; Dicke, M. Variation in natural plant products and the attraction of bodyguards involved in indirect plant defense. *Canadian Journal of Zoology-Revue Canadienne De Zoologie* **2010**, *88*, 628-667, doi:10.1139/z10-032.

3. Farag, M.A.; Zhang, H.M.; Ryu, C.M. Dynamic Chemical Communication between Plants and Bacteria through Airborne Signals: Induced Resistance by Bacterial Volatiles. *Journal of Chemical Ecology* **2013**, *39*, 1007-1018, doi:10.1007/s10886-013-0317-9.

4. Frost, C.J.; Mescher, M.C.; Dervinis, C.; Davis, J.M.; Carlson, J.E.; De Moraes, C.M. Priming defense genes and metabolites in hybrid poplar by the green leaf volatile cis-3-hexenyl acetate. *New Phytologist* **2008**, *180*, 722-733, doi:10.1111/j.1469-8137.2008.02599.x.

5. Turlings, T.C.J.; Loughrin, J.H.; McCall, P.J.; Rose, U.S.R.; Lewis, W.J.; Tumlinson, J.H. How caterpillar-damaged plants protect themselves by attracting parasitic wasps. *Proceedings of the National Academy of Sciences of the United States of America* **1995**, *92*, 4169-4174, doi:10.1073/pnas.92.10.4169.

6. Rodriguez-Saona, C.R.; Frost, C.J. New evidence for a multi-functional role of herbivore-induced plant volatiles in defense against herbivores. *Plant signaling & behavior* **2010**, *5*, 58-60.

7. Frost, C.J.; Appel, M.; Carlson, J.E.; De Moraes, C.M.; Mescher, M.C.; Schultz, J.C. Within-plant signalling via volatiles overcomes vascular constraints on systemic signalling and primes responses against herbivores. *Ecology Letters* **2007**, *10*, 490-498, doi:10.1111/j.1461-0248.2007.01043.x.

8. Heil, M.; Silva Bueno, J.C. Within-plant signaling by volatiles leads to induction and priming of an indirect plant defense in nature. *Proceedings of the National Academy of Sciences of the United States of America* **2007**, *104*, 5467-5472, doi:10.1073/pnas.0610266104.

9. Heil, M.; Karban, R. Explaining evolution of plant communication by airborne signals. *Trends in Ecology & Evolution* **2010**, *25*, 137-144, doi:10.1016/j.tree.2009.09.010.

10. Pearse, I.S.; Hughes, K.; Shiojiri, K.; Ishizaki, S.; Karban, R. Interplant volatile signaling in willows: revisiting the original talking trees. *Oecologia* **2013**, *172*, 869-875, doi:10.1007/s00442-013-2610-2.

11. Arimura, G.; Ozawa, R.; Nishioka, T.; Boland, W.; Koch, T.; Kuhnemann, F.; Takabayashi, J. Herbivore-induced volatiles induce the emission of ethylene in neighboring lima bean plants. *Plant Journal* **2002**, *29*, 87-98, doi:10.1046/j.1365-313x.2002.01198.x.

12. Farag, M.A.; Fokar, M.; Zhang, H.A.; Allen, R.D.; Pare, P.W. (Z)-3-Hexenol induces
defense genes and downstream metabolites in maize. *Planta* **2005**, *220*, 900-909,
doi:10.1007/s00425-004-1404-5.

13. Erb, M.; Veyrat, N.; Robert, C.A.M.; Xu, H.; Frey, M.; Ton, J.; Turlings, T.C.J. *Indole is*
an essential herbivore-induced volatile priming signal in maize; *Nature communications*,
2015; pp 1-10.

14. Farag, M.A.; Paré, P.W. C6-Green leaf volatiles trigger local and systemic VOC
emissions in tomato. *Phytochemistry* **2002**, *61*, 545-554, doi:10.1016/s0031-
9422(02)00240-6.

15. Rodriguez-Saona, C.R.; Rodriguez-Saona, L.E.; Frost, C.J. Herbivore-induced volatiles
in the perennial shrub, *Vaccinium corymbosum*, and their role in inter-branch signaling.
Journal of Chemical Ecology **2009**, *35*, 163-175, doi:10.1007/s10886-008-9579-z.

16. Sarai Giron-Calva, P.; Molina-Torres, J.; Heil, M. Volatile dose and exposure time
impact perception in neighboring plants. *Journal of Chemical Ecology* **2012**, *38*, 226-
228, doi:10.1007/s10886-012-0072-3.

17. Arimura, G.I.; Kopke, S.; Kunert, M.; Volpe, V.; David, A.; Brand, P.; Dabrowska, P.;
Maffei, M.E.; Boland, W. Effects of feeding *Spodoptera littoralis* on lima bean leaves:
IV. Diurnal and nocturnal damage differentially initiate plant volatile emission. *Plant*
Physiology **2008**, *146*, 965-973, doi:10.1104/pp.107.111088.

18. Arimura, G.; Matsui, K.; Takabayashi, J. Chemical and Molecular Ecology of Herbivore-
Induced Plant Volatiles: Proximate Factors and Their Ultimate Functions. *Plant and Cell*
Physiology **2009**, *50*, 911-923, doi:10.1093/pcp/pcp030.

19. Copolovici, L.; Kannaste, A.; Remmel, T.; Vislap, V.; Niinemets, U. Volatile Emissions
from *Alnus glutionosa* Induced by Herbivory are Quantitatively Related to the Extent of
Damage. *Journal of Chemical Ecology* **2011**, *37*, 18-28, doi:10.1007/s10886-010-9897-9.

20. Boggia, L.; Sgorbini, B.; Bertea, C.M.; Cagliero, C.; Bicchi, C.; Maffei, M.E.; Rubiolo,
P. Direct Contact-Sorptive Tape Extraction coupled with Gas Chromatography-Mass
Spectrometry to reveal volatile topographical dynamics of lima bean (*Phaseolus lunatus*
L.) upon herbivory by *Spodoptera littoralis* Boisd. *BMC Plant Biology* **2015**, *15*, 1-13,
doi:10.1186/s12870-015-0487-4.

21. Engelberth, J.; Alborn, H.T.; Schmelz, E.A.; Tumlinson, J.H. Airborne signals prime
plants against insect herbivore attack. *Proceedings of the National Academy of Sciences*
of the United States of America **2004**, *101*, 1781-1785, doi:10.1073/pnas.0308037100.

22. Paschold, A.; Halitschke, R.; Baldwin, I.T. Using 'mute' plants to translate volatile
signals. *Plant Journal* **2006**, *45*, 275-291, doi:10.1111/j.1365-313X.2005.02623.x.

23. Matsui, K.; Sugimoto, K.; Mano, J.i.; Ozawa, R.; Takabayashi, J. Differential
Metabolisms of Green Leaf Volatiles in Injured and Intact Parts of a Wounded Leaf Meet

Distinct Ecophysiological Requirements. *Plos One* **2012**, *7*,
doi:10.1371/journal.pone.0036433.

24.   Pastor, V.; Balmer, A.; Gamir, J.; Flors, V.; Mauch-Mani, B. Preparing to fight back:
generation and storage of priming compounds. *Frontiers in Plant Science* **2014**, *5*,
doi:10.3389/fpls.2014.00295.

25.   D'Auria, J.C.; Pichersky, E.; Schaub, A.; Hansel, A.; Gershenzon, J. Characterization of a
BAHD acyltransferase responsible for producing the green leaf volatile (Z)-3-hexen-1-yl
acetate in *Arabidopsis thaliana*. *Plant Journal* **2007**, *49*, 194-207, doi:10.1111/j.1365-
313X.2006.02946.x.

26.   Addesso, K.M.; McAuslane, H.J.; Alborn, H.T. Attraction of pepper weevil to volatiles
from damaged pepper plants. *Entomologia Experimentalis Et Applicata* **2011**, *138*, 1-11,
doi:10.1111/j.1570-7458.2010.01070.x.

27.   Cardoza, Y.J.; Tumlinson, J.H. Compatible and incompatible Xanthomonas infections
differentially affect herbivore-induced volatile emission by pepper plants. *Journal of*
Chemical Ecology **2006**, *32*, 1755-1768, doi:10.1007/s10886-006-9107-y.

28.   McCormick, A.C.; Irmisch, S.; Reinecke, A.; Boeckler, G.A.; Veit, D.; Reichelt, M.;
Hansson, B.S.; Gershenzon, J.; Kollner, T.G.; Unsicker, S.B. Herbivore-induced volatile
emission in black poplar: regulation and role in attracting herbivore enemies. *Plant Cell*
and Environment **2014**, *37*, 1909-1923, doi:10.1111/pce.12287.

29.   Accamando, A.K.; Cronin, J.T. Costs and Benefits of Jasmonic Acid Induced Responses
in Soybean. *Environmental Entomology* **2012**, *41*, 551-561, doi:10.1603/en11277.

30.   Cipollini, D.; Purrington, C.B.; Bergelson, J. Costs of induced responses in plants. *Basic*
and Applied Ecology **2003**, *4*, 79-89, doi:10.1078/1439-1791-00134.

31.   Cipollini, D.; Heil, a.M. Costs and benefits of induced resistance to herbivores and
pathogens in plants. *CAB International* **2010**, *5*, 1-25.

32.   Douma, J.C.; Vermeulen, P.J.; Poelman, E.H.; Dicke, M.; Anten, N.P.R. When does it
pay off to prime for defense? A modeling analysis. *New Phytologist* **2017**, *216*, 782-797,
doi:10.1111/nph.14771.

33.   Frost, C.J.; Mescher, M.C.; Carlson, J.E.; De Moraes, C.M. Plant defense priming against
herbivores: Getting ready for a different battle. *Plant Physiology* **2008**, *146*, 818-824,
doi:10.1104/pp.107.113027.

34.   van Hulten, M.; Pelser, M.; van Loon, L.C.; Pieterse, C.M.J.; Ton, J. Costs and benefits
of priming for defense in Arabidopsis. *Proceedings of the National Academy of Sciences*
of the United States of America **2006**, *103*, 5602-5607, doi:10.1073/pnas.0510213103.

35. Orrock, J., Connolly, B., Choi, W., Guiden, P., Swanson, S., & Gilroy, S. . Plants eavesdrop on cues produced by snails and induce costly defenses that affect insect herbivores. *Oecologia* **2018**, *186*, 703-710.

36. Martinez-Medina, A.; Flors, V.; Heil, M.; Mauch-Mani, B.; Pieterse, C.M.J.; Pozo, M.J.; Ton, J.; van Dam, N.M.; Conrath, U. Recognizing Plant Defense Priming. *Trends in Plant Science* **2016**, *21*, 818-822, doi:10.1016/j.tplants.2016.07.009.

37. Song, G.C.; Ryu, C.M. Two volatile organic compounds trigger plant self-defense against a bacterial pathogen and a sucking insect in cucumber under open field conditions. *International Journal of Molecular Sciences* **2013**, *14*, 9803-9819, doi:10.3390/ijms14059803.

38. Karban, R.; Maron, J. The fitness consequences of interspecific eavesdropping between plants. *Ecology* **2002**, *83*, 1209-1213.

39. Ballhorn, D.J.; Schiwy, S.; Jensen, M.; Heil, M. Quantitative Variability of Direct Chemical Defense in Primary and Secondary Leaves of Lima Bean (*Phaseolus lunatus*) and Consequences for a Natural Herbivore. *Journal of Chemical Ecology* **2008**, *34*, 1298-1301, doi:10.1007/s10886-008-9540-1.

40. Zachariah, T.J.; Safeer, A.L.; Jayarajan, K.; Leela, N.K.; Vipin, T.M.; Saji, K.V.; Shiva, K.N.; Parthasarathy, V.A.; Mammootty, K.P. Correlation of metabolites in the leaf and berries of selected black pepper varieties. *Scientia Horticulturae* **2010**, *123*, 418-422, doi:10.1016/j.scienta.2009.09.017.

41. Frost, C.; Mescher, M.; Carlson, J.; De Moraes, C. Why do distance limitations exist on plant-plant signaling via airborne volatiles? *Plant Signaling & Behavior* **2008**, *3*, 466-468.

42. Dolch, R.; Tscharntke, T. Defoliation of alders (*Alnus glutinosa*) affects herbivory by leaf beetles on undamaged neighbours. *Oecologia* **2000**, *125*, 504-511, doi:10.1007/s004420000482.

43. Karban, R.; Shiojiri, K.; Huntzinger, M.; McCall, A.C. Damage-induced resistance in sagebrush: Volatiles are key to intra- and interplant communication. *Ecology* **2006**, *87*, 922-930.

44. Heil, M.; Adame-Alvarez, R.M. Short signalling distances make plant communication a soliloquy. *Biology Letters* **2010**, *6*, 843-845, doi:10.1098/rsbl.2010.0440.

45. Engelberth, J.; Seidl-Adams, I.; Schultz, J.C.; Tumlinson, J.H. Insect elicitors and exposure to green leafy volatiles differentially upregulate major octadecanoids and transcripts of 12-oxo phytodienoic acid reductases in *Zea mays*. *Molecular Plant-Microbe Interactions* **2007**, *20*, 707-716, doi:10.1094/mpmi-20-6-0707.

46.   Shiojiri, K.; Ozawa, R.; Matsui, K.; Sabelis, M.W.; Takabayashi, J. Intermittent exposure
to traces of green leaf volatiles triggers a plant response. *Scientific Reports* **2012**, *2*,
doi:10.1038/srep00378.

47.   von Merey, G.; Veyrat, N.; Mahuku, G.; Valdez, R.L.; Turlings, T.C.J.; D'Alessandro, M.
Dispensing synthetic green leaf volatiles in maize fields increases the release of
sesquiterpenes by the plants, but has little effect on the attraction of pest and beneficial
insects. *Phytochemistry* **2011**, *72*, 1838-1847, doi:10.1016/j.phytochem.2011.04.022.

48.   Frost, C.J.; Hunter, M.D. Insect canopy herbivory and frass deposition affect soil nutrient
dynamics and export in oak mesocosms. *Ecology* **2004**, *85*, 3335-3347, doi:10.1890/04-
0003.

49.   Frost, C.J.; Hunter, M.D. Herbivore-induced shifts in carbon and nitrogen allocation in
red oak seedlings. *New Phytologist* **2008**, *178*, 835-845, doi:10.1111/j.1469-
8137.2008.02420.x.

50.   Gleadow, R.M.; Moller, B.L. Cyanogenic Glycosides: Synthesis, Physiology, and
Phenotypic Plasticity. *Annual Review of Plant Biology, Vol 65* **2014**, *65*, 155-185,
doi:10.1146/annurev-arplant-050213-040027.

51.   Ballhorn, D.J.; Lieberei, R.; Ganzhorn, J.U. Plant cyanogenesis of Phaseolus lunatus and
its relevance for herbivore-plant interaction: The importance of quantitative data. *Journal*
of Chemical Ecology **2005**, *31*, 1445-1473, doi:10.1007/s10886-005-5791-2.

52.   Ochoa-Lopez, S.; Villamil, N.; Zedillo-Avelleyra, P.; Boege, K. Plant defence as a
complex and changing phenotype throughout ontogeny. *Annals of Botany* **2015**, *116*,
797-806, doi:10.1093/aob/mcv113.

53.   Schultz, J.C.; Appel, H.M.; Ferrieri, A.P.; Arnold, T.M. Flexible resource allocation
during plant defense responses. *Frontiers in Plant Science* **2013**, *4*,
doi:10.3389/fpls.2013.00324.

54.   Brinker, A.M.; Seigler, D. Methods for the detection and quantitative determination of
cyanide in plant materials. *Phytochemical Bulletin* **1989**, *21*, 24-31.

55.   Gleadow, R.; Bjarnholt, N.; Jørgensen, K.; Fox, J. Cyanogenic Glycosides. In *Soil*
Allelochemicals, Narwal, S.S., Ed. Studium Press LLC: 2011; Vol. 1, pp. 283-310.

56.   Agrawal, A.A. Induced responses to herbivory in wild radish: Effects on several
herbivores and plant fitness. *Ecology* **1999**, *80*, 1713-1723, doi:10.1890/0012-
9658(1999)080[1713:irthiw]2.0.co;2.

57.   Baldwin, I.T. Jasmonate-induced responses are costly but benefit plants under attack in
native populations. *Proceedings of the National Academy of Sciences of the United States*
of America **1998**, *95*, 8113-8118, doi:10.1073/pnas.95.14.8113.

58.   Didiano, T.J.; Turley, N.E.; Everwand, G.; Schaefer, H.; Crawley, M.J.; Johnson, M.T.J.
Experimental test of plant defence evolution in four species using long-term rabbit
exclosures. *Journal of Ecology* **2014**, *102*, 584-594, doi:10.1111/1365-2745.12227.

59.   Koricheva, J. Meta-analysis of sources of variation in fitness costs of plant antiherbivore
defenses. *Ecology* **2002**, *83*, 176-190, doi:10.2307/2680130.

60.   Mauricio, R. Costs of resistance to natural enemies in field populations of the annual
plant *Arabidopsis thaliana*. *American Naturalist* **1998**, *151*, 20-28, doi:10.1086/286099.

61.   Hilker, M.; Schwachtje, J.; Baier, M.; Balazadeh, S.; Baurle, I.; Geiselhardt, S.; Hincha,
D.K.; Kunze, R.; Mueller-Roeber, B.; Rillig, M.G., et al. Priming and memory of stress
responses in organisms lacking a nervous system. *Biological Reviews* **2016**, *91*, 1118-
1133, doi:10.1111/brv.12215.

62.   Ameye, M.; Audenaert, K.; De Zutter, N.; Steppe, K.; Van Meulebroek, L.; Vanhaecke,
L.; De Vleesschauwer, D.; Haesaert, G.; Smagghe, G. Priming of Wheat with the Green
Leaf Volatile Z-3-Hexenyl Acetate Enhances Defense against *Fusarium graminearum*
But Boosts Deoxynivalenol Production. *Plant Physiology* **2015**, *167*, 1671-1684,
doi:10.1104/pp.15.00107.

63.   Walters, D.R.; Paterson, L.; Walsh, D.J.; Havis, N.D. Priming for plant defense in barley
provides benefits only under high disease pressure. *Physiological and Molecular Plant*
Pathology **2008**, *73*, 95-100, doi:10.1016/j.pmpp.2009.03.002.

64.   Choh, Y.; Takabayashi, J. Herbivore-induced extrafloral nectar production in lima bean
plants enhanced by previous exposure to volatiles from infested conspecifics. *Journal of*
Chemical Ecology **2006**, *32*, 2073-2077.

65.   Acevedo, F.E.; Peiffer, M.; Tan, C.W.; Stanley, B.A.; Stanley, A.; Wang, J.; Jones, A.G.;
Hoover, K.; Rosa, C.; Luthe, D., et al. Fall Armyworm-Associated Gut Bacteria
Modulate Plant Defense Responses. *Molecular Plant-Microbe Interactions* **2017**, *30*,
127-137, doi:10.1094/mpmi-11-16-0240-r.

66.   Bissmeyer, S.; Freundlich, G.; Frost, C. The influence of dose of a plant-derived volatile
cue on *Arabidopsis thaliana* resistance against insect herbivores. *Kentucky Journal of*
Undergraduate Research **2018**, *2*, 84-95.

67.   Lucas-Barbosa, D. Integrating Studies on Plant-Pollinator and Plant-Herbivore
Interactions. *Trends in Plant Science* **2016**, *21*, 125-133,
doi:10.1016/j.tplants.2015.10.013.

68.   Agrawal, A.A.; Strauss, S.Y.; Stout, M.J. Costs of induced responses and tolerance to
herbivory in male and female fitness components of wild radish. *Evolution* **1999**, *53*,
1093-1104, doi:10.1111/j.1558-5646.1999.tb04524.x.

69.   Pashalidou, F.G.; Lucas-Barbosa, D.; van Loon, J.J.A.; Dicke, M.; Fatouros, N.E.
599   Phenotypic plasticity of plant response to herbivore eggs: effects on resistance to
600   caterpillars and plant development. *Ecology* **2013**, *94*, 702-713, doi:10.1890/12-1561.1.

601   70.   Agrawal, A.A. Overcompensation of plants in response to herbivory and the by-product
602   benefits of mutualism. *Trends in Plant Science* **2000**, *5*, 309-313, doi:10.1016/s1360-
603   1385(00)01679-4.

604   71.   Heath, J.J.; Kessler, A.; Woebbe, E.; Cipollini, D.; Stireman, J.O. Exploring plant defense
605   theory in tall goldenrod, *Solidago altissima*. *New Phytologist* **2014**, *202*, 1357-1370,
606   doi:10.1111/nph.12755.

607   72.   Cuny, M.A.; Gendry, J.; Hernández-Cumplido, J.; Benrey, B. Changes in plant growth
608   and seed production in wild lima bean in response to herbivory are attenuated by
609   parasitoids. *Oecologia* **2018**, *187*, 447-457.

610   73.   Machado, R.A.R.; Ferrieri, A.P.; Robert, C.A.M.; Glauser, G.; Kallenbach, M.; Baldwin,
611   I.T.; Erb, M. Leaf-herbivore attack reduces carbon reserves and regrowth from the roots
612   via jasmonate and auxin signaling. *New Phytologist* **2013**, *200*, 1234-1246,
613   doi:10.1111/nph.12438.

614   74.   Eichenberg, D.; Purschke, O.; Ristok, C.; Wessjohann, L.; Bruelheide, H. Trade-offs
615   between physical and chemical carbon-based leaf defence: of intraspecific variation and
616   trait evolution. *Journal of Ecology* **2015**, *103*, 1667-1679, doi:10.1111/1365-2745.12475.

617   75.   Gomez, S.; Ferrieri, R.A.; Schueller, M.; Orians, C.M. Methyl jasmonate elicits rapid
618   changes in carbon and nitrogen dynamics in tomato. *New Phytologist* **2010**, *188*, 835-
619   844, doi:10.1111/j.1469-8137.2010.03414.x.

620   76.   Schweiger, R.; Heise, A.M.; Persicke, M.; Muller, C. Interactions between the jasmonic
621   and salicylic acid pathway modulate the plant metabolome and affect herbivores of
622   different feeding types. *Plant Cell and Environment* **2014**, *37*, 1574-1585,
623   doi:10.1111/pce.12257.

624   77.   Ninkovic, V. Volatile communication between barley plants affects biomass allocation. *J
625   Exp Bot* **2003**, *54*, 1931-1939.

626   78.   Lu, H.B.; Xu, S.Y.; Zhang, W.J.; Xu, C.M.; Li, B.X.; Zhang, D.X.; Mu, W.; Liu, F.
627   Nematicidal Activity of trans-2-Hexenal against Southern Root-Knot Nematode
628   (*Meloidogyne incognita*) on Tomato Plants. *Journal of Agricultural and Food Chemistry*
629   **2017**, *65*, 544-550, doi:10.1021/acs.jafc.6b04091.

630   79.   Cipollini, D. Constitutive expression of methyl jasmonate-inducible responses delays
631   reproduction and constrains fitness responses to nutrients in *Arabidopsis thaliana*.
632   *Evolutionary Ecology* **2010**, *24*, 59-68, doi:10.1007/s10682-008-9290-0.

80. Karban, R. Tradeoff between resistance induced by volatile communication and over-
topping vertical growth. *Plant Signaling & Behavior* **2017**, *12*,
doi:10.1080/15592324.2017.1309491.

81. Ballhorn, D.J.; Heil, M.; Lieberei, R. Phenotypic plasticity of cyanogenesis in lima bean
Phaseolus lunatus - Activity and activation of beta-glucosidase. *Journal of Chemical
Ecology* **2006**, *32*, 261-275, doi:10.1007/s10886-005-9001-z.

82. Kessler, A.; Baldwin, I.T. Defensive function of herbivore-induced plant volatile
emissions in nature. *Science* **2001**, *291*, 2141-2144, doi:10.1126/science.291.5511.2141.

83. Heil, M. Indirect defence via tritrophic interactions. *New Phytologist* **2008**, *178*, 41-61,
doi:10.1111/j.1469-8137.2007.02330.x.

84. Peñaflor, M.F.G.V.; Gonçalves, F.G.; Colepicolo, C.; Sanches, P.A.; Bento, J.M.S.
Effects of single and multiple herbivory by host and non-host caterpillars on the
attractiveness of herbivore-induced volatiles of sugarcane to the generalist parasitoid
Cotesia flavipes. *Entomologia Experimentalis et Applicata* **2017**, *165*, 83-93,
doi:10.1111/eea.12623.

85. Stenberg, J.A.; Heil, M.; Ahman, I.; Bjorkman, C. Optimizing crops for biocontrol of
pests and disease. *Trends in Plant Science* **2015**, *20*, 698-712,
doi:10.1016/j.tplants.2015.08.007.

86. Peñaflor, M.F.G.V.; Bento, J.M.S. Herbivore-induced plant volatiles to enhance
biological control in agriculture. *Neotropical Entomology* **2013**, *42*, 331-343,
doi:10.1007/s13744-013-0147-z.

87. Lucchi, A.; Loni, A.; Gandini, L.M.; Scaramozzino, P.; Ioriatti, C.; Ricciardi, R.; Shearer,
P.W. Using herbivore-induced plant volatiles to attract lacewings, hoverflies and
parasitoid wasps in vineyards: achievements and constraints. *Bulletin of Insectology*
2017, *70*, 273-282.

88. Webster, B.; Bruce, T.; Dufour, S.; Birkemeyer, C.; Birkett, M.; Hardie, J.; Pickett, J.
Identification of volatile compounds used in host location by the black bean aphid, *Aphis
fabae*. *Journal of Chemical Ecology* **2008**, *34*, 1153-1161, doi:10.1007/s10886-008-
9510-7.

89. Huot, B.; Yao, J.; Montgomery, B.L.; He, S.Y. Growth-Defense Tradeoffs in Plants: A
Balancing Act to Optimize Fitness. *Molecular Plant* **2014**, *7*, 1267-1287,
doi:10.1093/mp/ssu049.

90. Wei, J.; van Loon, J.J.A.; Gols, R.; Menzel, T.R.; Li, N.; Kang, L.; Dicke, M. Reciprocal
crosstalk between jasmonate and salicylate defence-signalling pathways modulates plant
volatile emission and herbivore host-selection behaviour. *Journal of Experimental Botany*
2014, *65*, 3289-3298, doi:10.1093/jxb/eru181.

91.   Ameye, M.; Allmann, S.; Verwaeren, J.; Smagghe, G.; Haesaert, G.; Schuurink, R.C.;
670   Audenaert, K. Green leaf volatile production by plants: a meta-analysis. *New Phytologist*
671   **2017**, 10.1111/nph.14671, 1-18, doi:10.1111/nph.14671.

672   92.   Holopainen, J.K.; Gershenzon, J. Multiple stress factors and the emission of plant VOCs.
673   *Trends in Plant Science* **2010**, *15*, 176-184, doi:10.1016/j.tplants.2010.01.006.

674   93.   Caparrotta, S.; Boni, S.; Taiti, C.; Palm, E.; Mancuso, S.; Pandolfi, C. Induction of
675   priming by salt stress in neighboring plants. *Environmental and Experimental Botany*
676   **2018**, *147*, 261-270, doi:10.1016/j.envexpbot.2017.12.017.

677   94.   Kessler, A.; Halitschke, R.; Diezel, C.; Baldwin, I.T. Priming of plant defense responses
678   in nature by airborne signaling between *Artemisia tridentata* and *Nicotiana attenuata*.
679   *Oecologia* **2006**, *148*, 280-292, doi:10.1007/s00442-006-0365-8.

680   95.   Arimura, G.; Muroi, A.; Nishihara, M. Plant-plant-plant communications, mediated by
681   (E)-beta-ocimene emitted from transgenic tobacco plants, prime indirect defense
682   responses of lima beans. *Journal of Plant Interactions* **2012**, *7*, 193-196,
683   doi:10.1080/17429145.2011.650714.

Figure Captions

Fig. 1. **Height measurements and leaf counts for *Phaseolus lunatus* (lima bean) and *Capsicum annuum* (pepper) grown in a common garden field experiment.** Height for (A) lima bean was measured from the base of the longest runner to the uppermost branching point; (C) pepper height was measured from the base of the main stalk to the uppermost branching point. Leaf counts for (B) lima bean and (D) pepper included all mature leaves on each plant. Open circles represent control plants (receiving lanolin-filled vials); filled squares represent plants receiving a persistent application of vials containing 10ng/hr *cis*-3-hexenyl acetate (*z*3HAC) dissolved in lanolin. Dropdown lines indicate the initial application of *z*3HAC treatment: lima bean and pepper plants were first exposed on June 10, 2016 (Julian date 161) and July 11, 2016 (Julian date 192), respectively. Points represent averages +/- SE. Repeated measures ANOVAs (aov in R) were followed by one-way ANOVAs at each time-point. Asterisks (*) represent $P<0.05$ between treatment and control at each time point. See Supplemental Table 1 for complete statistics.

Fig. 2. **Biomass measurements of field-grown *Capsicum annuum* (pepper) plants.** (A) Leaf, (B) stem, (C) root biomass, and (D) the aboveground:belowground biomass ratio in *C.annuum* plants were determined at the end of the field season following destructive harvest. Bars represent means +/- S.E.M.. *P*-values represent Tukey's HSD comparisons. See Supplemental Table 2 for complete statistics.

Fig. 3. **Total flower production in (A) *Phaseolus lunatus* (lima bean) and (B) *Capsicum annuum* (pepper).** Open circles represent control plants (receiving lanolin-only vials); filled squares represent plants receiving a persistent application of vials containing 10ng/hr *cis*-3-hexenyl acetate (*z*3HAC) dissolved in lanolin. Dropdown lines indicate the initial application of *z*3HAC treatment: lima bean and pepper plants were first exposed on June 10, 2016 (Julian date 161) and July 11, 2016 (Julian date 192), respectively. Points represent averages +/- SE. Asterisks (*) represent $P<0.05$ between treatment and control at each time point. See Supplemental Table 3 for complete statistics.

Fig. 4. **Fruit and seed production in *Capsicum annuum* (pepper) plants grown in a common garden experiment treated with a persistent application of the green leaf volatile *cis*-3-hexenyl acetate (*z*3HAC).** The (A) total number of fruits were counted in the field, and (B) wet and (C) dry masses fruit masses were determined in the lab. (D) The total number of seeds per fruit and (E) the estimated mass per seed were determined from a subset of the fruits produced. (F) The ratio of seed mass to fruit mass was calculated to assess the efficiency of seed production. Bars represent means +/- S.E.M.. White boxes represent control plants; gray boxes represent plants treated with *z*3HAC. *P*-values represent Tukey's HSD comparisons. See Supplemental Table 4 for complete statistics.

Fig. 5. **Herbivore damage on *Capsicum annuum* (pepper) and *Phaseolus lunatus* (lima bean) plants in a common garden field experiment.** Chewing damage on (A) pepper and (B) lima bean plants was determined using a visual estimation technique (see Methods). Open circles represent control plants (receiving lanolin-filled vials); filled squares represent plants receiving a persistent application 10ng/hr *cis*-3-hexenyl acetate (*z*3HAC) dissolved in lanolin.

Dropdown lines indicate the initial application of $z3$HAC treatment. Points represent averages
+/- SE. Repeated measures ANOVAs (aov in R) were followed by one-way ANOVAs at each
time-point. Asterisks (*) represent $P<0.05$ between treatment and control at each time point. See
Supplemental Table 5 for complete statistics. (C) *Aphis faba* colonization on lima bean plants.
Bars represent the percentage of plants in each group where *A.faba* were observed ($\chi^2 = 50.11$,
df=1, $P<0.001$).
738
739 **Fig. 6. Cyanide induction in *Phaseolus lunatus* (Lima bean) as affected by *z3*HAC exposure**.
740 Cyanide concentration was determined by trapping volatile HCN in 1.0M NaOH followed by
741 colorometric quantification. Sink leaves (A) and source leaves (B) were exposed long-term
742 exposure of 10ng/hour $z3$HAC or left as a control. Points represent averages +/- SE and *
743 indicate a p-value <0.05.

756   **Figure 1**

768   **Figure 2**

773   **Figure 3**

 Figure 4

796   **Figure 5**

799 **Figure 6**

818