Single-cell chromatin accessibility in glioblastoma delineates cancer stem cell heterogeneity predictive of survival

Authors: P. Guilhamon[1,2], M.M. Kushida[2], A. Nikolic[3,4,5], D. Singhal[3,4,5], G. MacLeod[6], S.A. Madani Tonekaboni[1,7], F.M.G. Cavalli[2], C. Arlidge[1], N. Rajakulendran[6], N. Rastegar[2], X. Hao[3,8,13], R. Hassam[3,8,13], L.J. Smith[18], H. Whetstone[2], F.J. Coutinho[2], B. Nadorp[1], K.I. Ellestad[3,4], H.A. Luchman[3,8,13], J.A. Chan[3,4,16], M.S. Shoichet[18], M.D. Taylor[2,9], B. Haibe-Kains[1,7,10,11,12], S. Weiss[3,8,13,14], S. Angers[6,17], M. Gallo[3,4,5,14], P.B. Dirks[2,9,15,*], M. Lupien[1,7,11,*]

Affiliations:
[1]Princess Margaret Cancer Centre, University Health Network, Toronto, ON M5G 1L7, Canada.
[2]Developmental and Stem Cell Biology Program and Arthur and Sonia Labatt Brain tumor Research Centre, The Hospital for Sick Children, Toronto, ON M5G 0A4, Canada.
[3]Clark Smith Brain Tumour Centre, Arnie Charbonneau Cancer Institute, University of Calgary, Calgary, Alberta T2N 4N1, Canada.
[4]Alberta Children's Hospital Research Institute, University of Calgary, Calgary, Alberta T2N 4N1, Canada.
[5]Department of Biochemistry and Molecular Biology, Cumming School of Medicine, University of Calgary, Calgary, Alberta T2N 4N1, Canada.
[6]Leslie Dan Faculty of Pharmacy, University of Toronto, Toronto, ON M5S 3M2, Canada.
[7]Department of Medical Biophysics, University of Toronto, Toronto, ON M5S 1A8, Canada.
[8]Hotchkiss Brain Institute, University of Calgary, Calgary, AB T2N 4N1, Canada.
[9]Division of Neurosurgery, University of Toronto, Toronto, ON M5S 1A8, Canada.
[10]Department of Computer Science, University of Toronto, Toronto, ON M5S 1A8, Canada.
[11]Ontario Institute for Cancer Research, Toronto, ON M5G 0A3, Canada.
[12]Vector Institute, Toronto, ON M5G 0A3, Canada.
[13]Department of Cell Biology & Anatomy, University of Calgary, Calgary, AB T2N 4N1, Canada.
[14]Department of Physiology & Pharmacology, University of Calgary, Calgary, AB T2N 4N1, Canada.
[15]Departments of Molecular Genetics and Surgery, University of Toronto, Toronto, ON M5S 1A8, Canada.
[16]Department of Pathology and Laboratory Medicine, University of Calgary, Calgary, AB, Canada.
[17]Department of Biochemistry, Faculty of Medicine, University of Toronto, Toronto, ON, Canada.
[18]Institute of Biomaterials and Biomedical Engineering, University of Toronto, Toronto, ON, Canada.

*Correspondence to: Mathieu Lupien (mathieu.lupien@uhnresearch.ca) and Peter B Dirks (peter.dirks@sickkids.ca)

ABSTRACT

Chromatin accessibility discriminates stem from mature cell populations, enabling the identification of primitive stem-like cells in primary tumors, such as Glioblastoma (GBM) where self-renewing cells driving cancer progression and recurrence are prime targets for therapeutic intervention. We show, using single-cell chromatin accessibility, that primary GBMs harbor a heterogeneous self-renewing population whose diversity is captured in patient-derived glioblastoma stem cells (GSCs). In depth characterization of chromatin accessibility in GSCs identifies three GSC states: Reactive, Constructive, and Invasive, each governed by uniquely essential transcription factors and present within GBMs in varying proportions. Orthotopic xenografts reveal that GSC states associate with survival, and identify an invasive GSC signature predictive of low patient survival. Our chromatin-driven characterization of GSC states improves prognostic precision and identifies dependencies to guide combination therapies.

INTRODUCTION

Glioblastoma (GBM) is a lethal form of brain cancer with standard surgery and radiation giving a median survival of only 12.6 months[1]. The addition of temozolomide chemotherapy provides only an additional 2.5 months in the small subset of responsive patients[2]. Despite extensive characterization and stratification of the bulk primary tumors, no targeted therapies have been successfully developed[1,3]. GBM tumors are rooted in self-renewing tumor-initiating cells commonly referred to as glioblastoma stem cells (GSCs)[4] that drive disease progression *in vivo[5,6]* and display resistance to chemo- and radiotherapy leading to disease recurrence[7]. The promise of therapeutically targeting self-renewing tumor-initiating cancer cells depends on

our capacity to capture the full range of heterogeneity within this population from individual tumors. Intratumoral heterogeneity within primary GBM has recently been documented through single cell RNA-seq experiments and revealed a continuum between four cellular states[8]: neural-progenitor-like (NPC), oligodendrocyte-progenitor-like (OPC), astrocyte-like (AC), and mesenchymal-like (MES)[8]. A subsequent study[9] using single-cell gene-centric enrichment analysis placed GBM cells along a single axis of variation from proneural to mesenchymal transcriptional profiles, with cells expressing stem-associated genes lying at the extremes of this axis. Hence, primary GBM consists of distinct states, across which stem-like cells appear to be found. Whether these stem-like cells found across GBM states represent functionally distinct GSC populations with tumor-initiating properties and unique dependencies remains to be established to guide therapeutic progress. To address this issue, we combined single-cell technologies to define GSC composition in primary GBM with functional assays to reveal the unique dependencies across GSCs, reflective of invasive, constructive and reactive states that relate to patient outcome.

RESULTS

Chromatin accessibility readily discriminates stem from mature cell populations[10], which can be resolved at the single cell level through single-cell ATAC-seq[11,12] taking into account non-gene centric features, such as accessibility of noncoding elements and total amount of accessible DNA sequences. Applying single-cell chromatin accessibility profiling (scATAC-seq) across four *IDH* wild-type primary adult GBM tumors (3797 cells) revealed 7-9 accessibility modules in each tumor based on unsupervised clustering (Fig. 1A). We assigned

cells to each of the four scRNA-seq-derived cellular states[8] based on individual cells' chromatin

accessibility enrichment scores for the promoter regions of each state's signature genes. The

MES state reported from scRNA-seq[8] dominates the identity of two or more modules reported

from chromatin accessibility in every tumor (Fig. 1B-C). In contrast, the NPC and OPC states are

mixed within the same module defined based on chromatin accessibility, dominating over the

other states typically in at least two modules. Cells assigned to the AC state did not preferentially

cluster within a single module reported from chromatin accessibility (Fig. 1B-C). Collectively, our

results suggest that chromatin accessibility reflects a greater stratification of the MES state,

detects similarities between the OPC and NPC states and heterogeneity within the AC state.

To identify putative cancer stem cells within each primary tumor, we next focused on the

level of chromatin accessibility at promoters of 19 transcription factors previously associated

with self-renewal and tumor-propagating capacity in GBM[13] (Fig. 1D). Individual cells scoring

as putative cancer stem cells were not restricted to a unique module defined by chromatin

accessibility but were distributed across a subset of modules, suggesting heterogeneity across

cancer stem cells in primary GBM, in agreement with reports relying on single GBM cell

labelling[6,10,12–20] assessing the heterogeneity of self-renewing tumor-initiating cells.

Putative cancer stem cells identified in primary GBM through scATAC-seq were found in

modules ascribed to every one of the four cellular states defined by gene expression[8],

predominantly within NPC and OPC containing modules and a smaller fraction (<10%) in MES-

specific modules across all four tumors (Fig. 1E). This suggests that the core transcriptional unit

of cancer stem cells in primary GBM[13] is not restricted to a unique population defined by its

global transcriptional or chromatin accessibility profile with the resolution achieved with current

single-cell technologies.

To further probe the heterogeneity in chromatin accessibility within the GBM cancer

stem cell pool, we derived GSC populations from 27 adult *IDH* wild-type GBM tumors[21] and

profiled their chromatin accessibility by bulk ATAC-seq (Fig. 2A). Each patient-derived GSC

showed a similar enrichment for accessible chromatin regions in promoters and 5'UTRs, and

depletion in introns and distal intergenic regions (Fig. 2B). Collectively, we uncovered 92% of

the total predicted regions of accessible chromatin (255,891 regions) within GSCs based on a

saturation analysis using a self-starting nonlinear regression model across the 27 samples (Fig.

2C). We next assessed the similarity between these GSCs and the putative cancer stem cells

found by scATAC-seq in the four primary GBMs. GSCs were identified within each tumor by

calculating the enrichment of accessible chromatin regions shared by a majority of GSCs

(>14/27) in each tumor cell (Fig. 2D). Comparing the distribution of GSCs across the 7-9 modules

defined by scATAC-seq to that of the 19 transcription factor-derived cancer stem cell signature

demonstrates concordance between the two signatures (Fig. 2E). Moreover, the enrichment z-

scores for both cancer stem signatures (ie stem transcription factors signature and GSC

chromatin accessibility signature) are significantly correlated across cells in all four tumors (p ≤

$1.6e^{-5}$) (Fig. 2F). Additionally, an average of 91.2% (85.1-100%) of the cells identified by either

signature display the hallmark GBM copy number changes at chromosomes 7 and 10, confirming

their neoplastic status (Fig. S1A-D). Collectively, these results demonstrate that the patient-

derived GSC populations reflect the chromatin identity of putative cancer stem cells found in

primary brain tumors, highlighting the value of these GSCs as models to deepen our

understanding of individual cells within primary GBM with features found in self-renewing

tumor-initiating cells. Accordingly, spectral clustering of the 27 patient-derived GSC ATAC

profiles identifies three distinct states of self-renewing tumor-initiating cells (Fig. 3A).

Expression profiling of these 27 GSCs by RNA-seq reveals GSCs significantly enriched for the

signatures of each of the three TCGA GBM subtypes (proneural, classical, and

mesenchymal)[22] (Fig. 3B, top panel). However, the assignment of the proneural, classical and

mesenchymal subtypes across GSCs did not match the three clusters identified from ATAC-seq

(Fig. 3B, top panel). Conversely, clustering the GSCs by gene expression, independently of their

chromatin accessibility, did largely recapitulate the GSC states defined by chromatin

accessibility (Fig. 3B, bottom panel). This suggests that the mismatch between the GSC cluster

from chromatin accessibility profiles and TCGA expression subtypes is not mainly due to

differences between chromatin accessibility and gene expression. A more likely possibility is

that given the TCGA subtypes were determined from bulk GBM, they may not fully capture the

nature of rarer populations found within a tumor, such as the cancer stem cell populations.

Gene set enrichment analysis with GSEA[23] and g:profiler[24] using genes exclusively

enriched for both expression and promoter chromatin accessibility in each subtype reported

significantly enriched terms defining the largest GSC state as a Reactive state, with terms

related to immune cells and response (Fig. 3C, top panel). A second GSC state was enriched for

Constructive gene sets involved in brain, neuron, and glial cell development (Fig. 3C, middle

panel). The third and smallest GSC state presented an Invasive state characterized by terms

relating to the extracellular matrix and angiogenesis (Fig. 3C, bottom panel). We next mapped

copy number alterations (CNAs) across the 27 GSCs by applying the molecular neuropathology

classifier tool[25] to DNA methylation data from the GSCs (Fig. 3D). While common GBM CNAs,

including EGFR gains and CDKN2A/B loss, were observed, the CNA-based classification of GSCs

failed to match the three chromatin accessibility-derived states, suggesting the three GSC

states are not genetically defined (Fig. 3D). Further comparison of the accessible chromatin in

each GSC state reveals that only a small subset of accessible chromatin regions drives the three

GSC states (Fig. 3E and Fig. S1E-G). Our ability to discriminate GSC state-specific regions of

accessible chromatin is reflective of the comprehensiveness of our cohort to saturate the

detection of accessible regions to 93%, 88%, and 71% across the Reactive (n=13), Constructive

(n=9), and Invasive (n=5) state GSCs, respectively (Fig. 3F).

Considering that regions of accessible chromatin serve as binding sites for transcription

factors engaging in gene expression regulation, we next tested for DNA recognition motif family

enrichment across regions exclusively accessible in Reactive, Constructive or Invasive GSC states

(Fig. 4A and Fig. S2A). The most enriched DNA recognition motif families in each state were

either depleted or showed low-level enrichment in the other states. Specifically, the DNA

recognition motifs for the interferon-regulatory factor (IRF) and Cys2-His2 zinc finger (C2H2 ZF)

transcription factor families were enriched in the Reactive state (Fig. 4A, top panel). Regulatory

factor X (RFX) and basic helix-loop-helix (bHLH) DNA recognition motifs were enriched in the

Constructive state (Fig. 4A, middle panel), while the Forkhead motif family was enriched in the

Invasive state (Fig. 4A, bottom panel). Genome-wide CRISPR/Cas9 essentiality screens (Fig.

S2B) in three Reactive, two Constructive and one Invasive GSC[26] revealed the preferential

requirement for expressed transcription factors (Fig. S2C-E) recognizing the enriched DNA

recognition motif in a state-specific manner (Fig. 4B). Specifically, the SP1 regulatory network

is preferentially essential in the Reactive state GSCs (Fig 4B, top panel), ASCL1, OLIG2, AHR,

and NPAS3 are uniquely essential in the Constructive state GSCs (Fig. 4B, middle panel) and

FOXD1 in essential only in the Reactive state GSC (Fig. 4B, bottom panel). Notably, SP1 itself is

exclusively essential in only one Reactive GSC (G564). However, of the 36 transcription factors

from the Reactive-enriched families (IRF and C2H2 ZF) that were essential in at least one

Reactive GSC and not in any of the Constructive or Invasive GSCs, 13 are directly regulated by

SP1 (Fig. S2C), thus suggesting that the SP1 regulatory network as a whole, rather than SP1 on

its own, is key in the Reactive GSC state. Notably, all six transcription factors display significantly

higher expression in GBM compared to normal brain[27] (Fig. S2F), further supporting their

function as key regulators of tumor initiation and development. An additional gene set

enrichment analysis combining genes exclusively essential to each state with the putative

targets of the key transcription factors outlined above identifies additional enriched terms

supporting the identities of the three GSC states as Reactive, Constructive, and Invasive (Fig.

S2G).

Previous work suggests that GBM tumors harbor a heterogeneous population of

GSCs[14,15,28]. We therefore quantified the presence of Reactive, Constructive and Invasive

cancer stem cells in our four primary GBM based on their scATAC-seq profiles. The Constructive

state was dominant in every primary tumor, ranging from 9-21% of all cells captured by scATAC-

seq (Fig. 4C). The Reactive and Invasive states accounted for only 0-9% of all cells, in varying

proportions from one tumor to another (Fig. 4C). Collectively, our results further support the

heterogeneous nature of cancer stem cells that populate primary tumors.

While various classifications of GBMs and/or their constitutive bulk and stem tumor cells

have been reported, with some associating with patient survival[8,22,28–31], no molecular

signature in GBM has so far been reported that can significantly stratify the poorer prognosis

IDH wild-type patients by survival. We performed intracranial xenografts of 37 *IDH* wild-type

GSC populations and classified the transplanted cells by their GSC state to perform a differential

survival analysis (Fig. 4D). The overall survival times of the transplanted mice grouped by GSC

state were significantly different (LogRank test p = 0.041), with the Invasive state GSCs leading

to the worst prognosis. Relative to the mice injected with Reactive state GSCs, mice with

Constructive state GSCs and Invasive state GSCs had hazard ratios of 1.3 (95% CI: 0.57-2.97) and

3.5 (95% CI: 1.2-10.49), respectively. Next, we investigated the prognostic value of the GSC

states using the TCGA GBM cohort (*IDH* wild-type, n = 144). When classified by dominant GSC

state, TCGA tumors display the same trend as the xenografts with Invasive state-dominated

tumors showing the lowest survival (Fig. S3A). However, with only two tumors classified as

Invasive-dominant, the difference in survival between the three patient groups was not

statistically significant (p = 0.3) (Fig. S3A). We proceeded to rank the TCGA tumors solely by their

concordance to Invasive GSCs and classified the patient tumors into Invasive-low (z-score < 1),

Invasive-mid (z-score 1-1.65), and Invasive-high (z-score ≥ 1.65) groups (Fig. 4E). With this

stratification method, median patient survival per group not only decreased with increasing

Invasive GSC score, but we also identified an Invasive-high subset of tumors with significantly

lower survival (p = 0.019, HR = 2.8, 95% CI: 1.3-5.81)) (Fig. 4F). Invasion assays using

representative Invasive state GSCs additionally highlight the invasive properties of these

populations both *in vitro* and *in vivo* (Fig. S4). These results show that cancer stem cell states

defined based on the chromatin accessibility in GSCs can identify transcriptional programs

associated with poor prognosis and can serve as a signature to identify high-risk patients in *IDH*

wild-type GBM.

DISCUSSION

Defining the nature of self-renewing tumor-initiating cells in primary GBM is required to

identify vulnerabilities for therapeutic intervention. Quantifying their heterogeneity within

tumors can guide treatment strategies and assist in predicting the course of disease progression.

Here we show that chromatin accessibility assays capture a heterogeneity across self-renewing

tumor-initiating cells in primary GBM that extends beyond their genetic diversity, and underlies

the heterogeneity in bulk progeny[15]. This heterogeneity aligns with diversity in the three-

dimensional genome organization of GSCs[32] and agrees with how the three-dimensional

genome organization instructs cis-regulatory plexuses underlying gene regulation[33–37]. We

further reveal a specific cancer stem state that is significantly predictive of patient survival and

can be used as a signature to identify high-risk patients. Our results also highlight dependencies

unique to each cancer stem state. Specifically, the Reactive GSC state-associated transcription

factor SP1 and its regulatory partners are involved in cellular differentiation and growth,

apoptosis, response to DNA damage, chromatin remodelling[38], stimulation of *TERT*

expression in cancer stem cells[39], and increased stemness and invasion in GBM[40]. In

contrast, the Constructive GSC state rely on transcription factors including OLIG2, a known GSC

marker[41], AHR involved in tumor microenvironment responses and metabolic adaptation[42],

NPAS3, a regulator of Notch signaling and neurogenesis[43] and ASCL1, a critical regulator of

GSC differentiation and marker of sensitivity to Notch inhibition in GSCs[19,44]. Finally, the

Invasive GSC state relies on FOXD1, a pluripotency regulator and determinant of tumorigenicity

in GSCs regulating expression of the aldehyde dehydrogenase ALDH1A3, a functional marker

for invasive GSCs[45,46]. Collectively, our results support developing combination therapy using

targeting agents against each GSC state, such as Notch inhibitors[19] and small molecule

inhibitors of ALDH[45], to eradicate self-renewing tumor-initiating cells with the hope to cure

GBM patients.

METHODS

Patient samples and cell culture

All tissue samples were obtained following informed consent from patients, and all

experimental procedures were performed in accordance with the Research Ethics Board at The

Hospital for Sick Children (Toronto, Canada), the University of Calgary Ethics Review Board, and

the Health Research Ethics Board of Alberta - Cancer Committee (HREBA). Approval to

pathological data was obtained from the respective institutional review boards. Patient tumor

tissue samples were dissociated in artificial cerebrospinal fluid followed by treatment with

enzyme cocktail at 37°C. Patient tumor-derived GSCs were grown as adherent monolayer

cultures in serum-free medium as previously described[21]. Briefly, cells were grown adherently

on culture plates coated with poly-L-ornithine and laminin. Serum-free NS cell self-renewal

media (NS media) consisted of Neurocult NS-A Basal media, supplemented with 2 mmol/L L-

glutamine, N2 and B27 supplements, 75 µg/mL bovine serum albumin, 10 ng/mL recombinant

human EGF (rhEGF), 10 ng/mL basic fibroblast growth factor (bFGF), and 2 µg/mL heparin. A

subset (22/37) of the GSCs used for orthotopic xenografts were grown as non-adherent spheres

prior to single-cell dissociation and injection into the mice. Briefly, serum-free medium (SFM)

was used to initiate GSC cultures. Non-adherent spheres formed after 7-21 days in culture and

were expanded, then cryopreserved in 10% dimethyl sulfoxide (DMSO; Sigma-Aldrich) in SFM

until used in experiments.

ATAC-seq

ATAC-seq was used to profile the accessible chromatin landscape of 27 patient tumor-

derived GSCs. 50,000 cells were processed from each sample as previously described[47,48]. The

resulting libraries were sequenced with 50 bp single-end reads which were mapped to hg19.

Reads were filtered to remove duplicates, unmapped or poor quality (Q <30) reads,

mitochondrial reads, chrY reads, and those overlapping the ENCODE blacklist. Following

alignment, accessible chromatin regions/peaks were called using MACS2. Default parameters

were used except for the following: --keep-dup all -B --nomodel --SPMR -q 0.05 --slocal 6250 --

llocal 6250. The signal intensity was calculated as the fold enrichment of the signal per million

reads in a sample over a modelled local background using the bdgcmp function in MACS2.

Spectral clustering implemented in the SNFtool package[49] was run on the SNF fused similarity

matrix to obtain the groups corresponding to k=2 to 12. Enrichment for genomic features was

calculated using CEAS[50].

A given chromatin region was considered exclusive to one of the clusters if it was called

as a peak in any of the cluster's samples using a q-value filter of 0.05 and was not called as a peak

in any of the other samples using a q-value filter of 0.2, in order to ensure stringency of

exclusivity.

The ATAC-seq saturation analysis was performed by randomizing the order of samples,

and successively calculating the number of additional peaks discovered with the addition of each

new sample. This process was repeated 10,000 times and averaged. A self-starting non-linear

regression model was then fitted to the data to estimate the level of saturation reached.

For the xenograft survival analysis, 11/37 GSCs used overlap with the cohort of 27

described above. The other 26/37 GSCs were profiled by ATAC-seq independently following the

same protocol described above and assigned to a GSC state through unsupervised hierarchical

clustering with the original cohort of 27 GSCs.

<u>Single cell ATAC-seq</u>

The four tumors used were G4218 (primary GBM, IDH wt, Male, 64yrs), G4250 (primary

GBM, IDH wt, Male, 73yrs), G4275 (primary GBM, IDH wt, Female, 52yrs), G4349 (primary GBM,

IDH wt, Male, 62 yrs). Fragments of tumor were received fresh from the operating room, and

blunt dissected into individual fragments of approximately 0.3-0.7 cm3. Each fragment was

placed in 1 mL of freezing media (400 µL of NeuroCult NS-A Basal medium with proliferation

supplement (StemCell Technologies; #05751) containing 20 µg/mL rhEGF (Peprotech, AF-100-

15), 10 µg/mL bFGF (StemCell Technologies, #78003), and 2 µg/mL heparin (StemCell

Technologies, #07980); 500 µL of 25% BSA (Millipore-Sigma; A9647) in DMEM, and 100 µL

DMSO (Millipore-Sigma; D2650) in a 2 mL cryotube, and placed at -80 C in a CoolCell for at least

24 hours. Samples were then stored at -80C until use. Cryopreserved primary GBM samples were

washed at 1000 RPM for 5 minutes in PBS to remove DMSO, and then transferred to 1.5 mL

tubes. Samples were resuspended in cold ATAC resuspension buffer (10 mM Tris-HCl pH 7.4, 10

mM NaCl, 3 mM MgCl2, 0.1% NP-40, 0.1% Tween-20, 0.01% Digitonin, 1% BSA in PBS) on ice

and dissociated using a wide-bore P1000 pipette tip and vortexing, followed by 10 minutes of

incubation on ice. Cells were spun down at 500x g for 5 minutes at 4 C, washed in the ATAC

resuspension buffer, spun down again, and resuspended in ATAC-Tween wash buffer (10 mM

Tris-HCl pH 7.4, 10 mM NaCl, 3 mM MgCl2, 0.1% Tween-20, 1% BSA in PBS), then passed

through a cell strainer top FACS tube (Falcon; #38030) to remove debris. Nuclei quality and

quantity was evaluated using trypan blue on an Invitrogen Countess II device in duplicate, and a

subset of nuclei was spun down in a fresh tube and resuspended in 10X sample dilution buffer.

Nuclei were then used for single cell ATAC-seq library construction using the Chromium Single

Cell ATAC Solution v1.0 kit (10X Genomics) on a Chromium controller. Completed libraries were

further quality checked for fragment size and distribution using an Agilent TapeStation prior to

sequencing. Single-cell ATAC-seq samples were sequenced on a NextSeq 500 (Illumina)

instrument with 50 bp paired-end reads at the Centre for Health Genomics and Informatics

(CHGI) at the University of Calgary.

The raw sequencing data was demultiplexed using cellranger-atac mkfastq (Cell Ranger

ATAC, version 1.0.0, 10x Genomics). Single cell ATAC-seq reads were aligned to the hg19

reference genome (hg19, version 1.1.0, 10x Genomics) and quantified using cellranger-atac

count function with default parameters (Cell Ranger ATAC, version 1.1.0, 10x Genomics). The

resulting data were analysed using the chromVAR[51] and Signac[52] R packages (v1.4.1). The

number of accessibility modules in each sample was determined using the ElbowPlot method

implemented in Signac. Similarity between individual cells and GSC states was assessed using

the deviation scores calculated by chromVAR within the single cell data for significantly

differentially accessible sets of peaks (Fold Change Signal difference >2 and Wilcoxon test q-

value <=0.05) between the states as determined by bulk ATAC-seq. Similarity between

individual cells and the expression-derived cellular states was assessed using the deviation

scores calculated by chromVAR within the single cell data for promoter regions of the signature

genes of each of the cellular states[8]. A 2-fold cut-off was used to determine dominance of a

UMAP module by an individual or group of cellular states. Similarity between individual cells and

the GBM cancer stem cell signatures was assessed using the deviation scores calculated by

chromVAR within the single cell data for promoter regions of the 19 transcription factors

identified as markers of cancer stem cells in GBM[13].

Copy number variants in single cells were determined using CONICSmat[53] with default

parameters using the gene activity matrix generated by Signac as input. We focused on chr7

gains and chr10 losses as they are hallmark chromosomal changes in GBM and found the

following fractions of cells carrying these CNVs, on average across the four tumors: 76% of all

cells, 88% of cells allocated to scRNAseq cellular states[8], 95% of cancer stem cells based on

the 19 gene signature, 91% of GSCs based on shared accessible regions between 14/27 GSC

populations, 94% of GSCs identified based on the state-specific signatures.

DNA Methylation arrays

Bisulfite conversion of DNA for methylation profiling was performed using the EZ DNA

Methylation kit (Zymo Research) on 500 ng genomic DNA from all 27 samples. Conversion

efficiency was quantitatively assessed by quantitative PCR (qPCR). The Illumina Infinium

MethylationEPIC BeadChips were processed as per manufacturer's recommendations. The R

package ChAMP v2.6.4[54] was used to process and analyse the data. For the copy number

analysis, the raw IDAT files were uploaded to the MNP tool[25], which directly compares the

copy number profile estimated from the probe intensities on the methylation array to the

distribution observed across thousands of brain tumors in its database.

<u>RNA-seq</u>

RNA was extracted from GSCs using the Qiagen RNeasy Plus kit. RNA sample quality

was measured by Qubit (Life Technologies) for concentration and by Agilent Bioanalyzer for

RNA integrity. All samples had RIN above 9. Libraries were prepared using the TruSeq Stranded

mRNA kit (Illumina). Two hundred nanograms from each sample were purified for polyA tail

containing mRNA molecules using poly-T oligo attached magnetic beads, then fragmented

post-purification. The cleaved RNA fragments were copied into first strand cDNA using reverse

transcriptase and random primers. This is followed by second strand cDNA synthesis using

RNase H and DNA Polymerase I. A single "A" base was added and adapter ligated followed by

purification and enrichment with PCR to create cDNA libraries. Final cDNA libraries were verified

by the Agilent Bioanalyzer for size and concentration quantified by qPCR. All libraries were

pooled to a final concentration of 1.8nM, clustered and sequenced on the Illumina NextSeq500

as a pair-end 75 cycle sequencing run using v2 reagents to achieve a minimum of ~40 million

reads per sample. Reads were aligned to hg19 using the STAR aligner v2.4.2a [55] and transcripts

were quantified using RSEM v1.2.21[56] or vst transformed using DESeq2[57].

<u>Motif Enrichment</u>

Regions exclusively accessible in one of the GSC states and not the others were used as

input sequences for the motif enrichment, while the full ATAC-seq catalogue served as the

background set when running HOMER v4.7 to detect enrichments of transcription factor binding

motifs. Enriched motifs were then grouped into families based on similarities in DNA-binding

domains using the CIS-BP database[58]. Each family was assigned the fold-enrichment value of

the most enriched motif within the family.

The transcription factors whose motifs were found enriched in Reactive-exclusive

accessible regions, were run together through GSEA[23], and the gene set corresponding to

genes potentially regulated by SP1 was identified as significantly enriched (GSEA gene set

GGGCGGR_SP1_Q6). The expression levels of key transcription factors in tumor and normal

samples were analysed and displayed using GEPIA[27].

<u>Gene essentiality screen</u>

Illumina sequencing reads from genome-wide TKOv1 CRISPR screens in patient-derived

GSCs[26] were mapped using MAGECK[59] and analysed using the BAGEL algorithm with

version 2 reference core essential genes/non-essential genes[60,61]. Resultant raw Bayes Factor

(BF) statistics were used to determine essentiality of transcription factor genes using a minimum

BF of 3 and a 5% FDR cut-off. For visualisation purposes only, the essentiality scores were scaled

and the individual GSC essentiality thresholds subtracted from each score to obtain a common

threshold at 0 across GSCs.

Orthotopic xenografts

All animal procedures were performed according to and approved by the Animal Care Committee of the Hospital for Sick Children or the University of Calgary. All attempts are made to minimize the handling time during surgery and treatment so as not to unduly stress the animals. Animals are observed daily after surgery to ensure there are no unexpected complications. For intracranial xenografts, 100,000 GSC cells were stereotactically injected into the frontal cortex of 6-8 weeks old female NOD/SCID or C17/SCID mice. Mice were monitored and euthanized once neurological symptoms were observed or at the experimental endpoint of 12 months.

Invasion assay

Hydrogels were synthesized as previously described[62], with the following modifications: 1% w/v hyaluronan-methyl furan, 2.3mM MMP cleavable crosslinker, and 400µM fibronectin-derived peptide with sequence Mal-SKAGPHSRNGRGDSPG. Cells were plated on hydrogels at a density of 3500 cells/hydrogel and allowed to adhere for 24 h. 48 h after seeding, fresh media was added to each well. Cells were fixed with 4% PFA 4-5 days after seeding. Cells were counterstained with Hoechst (1/500) to label nuclei, Alexa FluorTM 488 Phalloidin (1/40) to label F-actin, and 15 µm FluosphereTM red beads (1/15) to label the surface of the hydrogel. Hydrogels were imaged using confocal microscopy at 10x magnification, taking images every 20 µm on the z-axis. Imaris Bitplane 8.3.1 software was used to prepare images and to analyze the positions of each cell and surface bead label. A custom Matlab script was used to calculate the

percent invasion, defined as the percentage of total cells located below a 75 µm threshold from

the surface of the hydrogel. Statistics were calculated using Graphpad Prism 7.04 software. Plots

are shown as mean with standard deviation, and statistics shown are from one-way ANOVA with

multiple comparisons with Dunnet's test correction, comparing all patient cell lines (dark grey)

to the hf6562 healthy control. Data were displayed with p values represented as * $p \leq 0.05$, ** p

≤ 0.01, *** $p \leq 0.001$, **** $p \leq 0.0001$.

<u>Immunohistochemistry</u>

Tissue samples were formalin fixed and paraffin embedded. Serial sections

deparaffinized, rehydrated through an alcohol gradient to water and antigen retrieval in citrate

buffer pH 6.0 was used for the human nucleolin antibody at 5.0 g/mL (ab13541) (Abcam,

Canbridge, MA). Endogenous peroxide activity and nonspecific binding was blocked with

3%(v/v) peroxide and 2% (v/v) normal horse serum. Primary antibody and anti-mouse

ImmPRESS-HRP secondary antibody were incubated for 1 hr and visualized using DAB (3,3'-

diaminobenzidine) (Vectorlabs, Burlingame, CA). Normal horse serum or monoclonal IgM was

used in control sections.

<u>Survival analysis</u>

Survival analysis on xenografts and TCGA data was performed using R packages

survival[63] and survminer[64]. The LogRank test was used in every analysis. See ATAC-seq

section for details on how each GSC used in the orthotopic xenografts was assigned to a GSC

state. TCGA samples were assigned to individual GSC states in the following way. 1) Using the

unsupervised clustering of RNA-seq data presented in Fig. 3B, the 23/27 GSCs that displayed

matched GSC state assignments by RNA-seq and ATAC-seq were used in this analysis. 2) Genes

preferentially enriched in each GSC state were determined using DEseq2[57] (q ≤ 0.05 and fold-

change ≥ 2). 3) The mean log2(FPKM+1) value for each of these genes over all GSCs in each state

was calculated to obtain a single representative value for each gene in each of the three GSC

states. 4) the concordance index was then calculated between each TCGA sample and each GSC

state and individual TCGA samples were assigned to the GSC state with the highest score.

Similarly, to assign TCGA samples to the three Invasive groups (Invasive-low, -mid, and -high),

the concordance to Invasive GSCs as calculated above was used. The z-score for each sample

was then used to classify each TCGA sample into the three subgroups of Invasive-low (Invasive

z-score < 1), Invasive-mid (Invasive z-score 1-1.65), and Invasive-high (Invasive z-score ≥ 1.65).

When changing the Invasive z-score thresholds for grouping the TCGA samples, the most

Invasive-high subgroup remains associated with the lowest survival (Fig. S3B-C).

Data and materials availability

The GSCs are available upon reasonable request from PBD and SW. The GSC ATAC-seq

and DNA methylation data have been deposited at GEO. The scATAC-seq data has been

deposited at GEO. RNA-seq data are available at EGA.

Code availability

All data analysis was performed using established methods implemented in published

software or R packages. Software and package versions and parameters are detailed in the

Methods section. All scripts used for the analysis are available upon request.

REFERENCES
1.   Carlsson SK, Brothers SP, Wahlestedt C (2014) Emerging treatment strategies for
glioblastoma multiforme. EMBO Mol Med 6: 1359–1370.
2.   Stupp R, Mason WP, van den Bent MJ, Weller M, Fisher B, Taphoorn MJB, Belanger K,
Brandes AA, Marosi C, Bogdahn U, Curschmann J, Janzer RC, Ludwin SK, Gorlia T, Allgeier
A, Lacombe D, Cairncross JG, Eisenhauer E, Mirimanoff RO, et al. (2005) Radiotherapy
plus concomitant and adjuvant temozolomide for glioblastoma. N Engl J Med 352: 987–
996.
3.   von Neubeck C, Seidlitz A, Kitzler HH, Beuthien-Baumann B, Krause M (2015)
Glioblastoma multiforme: emerging treatments and stratification markers beyond new
drugs. Br J Radiol 88: 20150354.
4.   Venere M, Fine HA, Dirks PB, Rich JN (2011) Cancer stem cells in gliomas: Identifying and
understanding the apex cell in cancer's hierarchy. Glia 59: 1148–1154.
5.   Chen J, Li Y, Yu T-S, McKay RM, Burns DK, Kernie SG, Parada LF (2012) A restricted cell
population propagates glioblastoma growth after chemotherapy. Nature 488: 522–526.
6.   Gallo M, Coutinho FJ, Vanner RJ, Gayden T, Mack SC, Murison A, Remke M, Li R,
Takayama N, Desai K, Lee L, Lan X, Park NI, Barsyte-Lovejoy D, Smil D, Sturm D, Kushida
MM, Head R, Cusimano MD, et al. (2015) MLL5 Orchestrates a Cancer Self-Renewal State
by Repressing the Histone Variant H3.3 and Globally Reorganizing Chromatin. Cancer Cell
28: 715–729.
7.   Bao S, Wu Q, McLendon RE, Hao Y, Shi Q, Hjelmeland AB, Dewhirst MW, Bigner DD, Rich
JN (2006) Glioma stem cells promote radioresistance by preferential activation of the
DNA damage response. Nature 444: 756–760.
8.   Neftel C, Laffy J, Filbin MG, Hara T, Shore ME, Rahme GJ, Richman AR, Silverbush D,
Shaw ML, Hebert CM, Dewitt J, Gritsch S, Perez EM, Nicolas Gonzalez Castro L, Lan X,
Druck N, Rodman C, Dionne D, Kaplan A, et al. (2019) An Integrative Model of Cellular
States, Plasticity, and Genetics for Glioblastoma. Cell 0. Available:
http://www.cell.com/article/S0092867419306877/abstract. Accessed 19 July 2019.
9.   Wang L, Babikir H, Muller S, Yagnik G, Shamardani K, Catalan F, Kohanbash G, Alvarado
B, Di Lullo E, Kriegstein A, Shah S, Wadhwa H, Chang SM, Philips JJ, Aghi MK, Diaz AA
(2019) The phenotypes of proliferating glioblastoma cells reside on a single axis of
variation. Cancer Discov. Available: http://dx.doi.org/10.1158/2159-8290.CD-19-0329.
10.   Stergachis AB, Neph S, Reynolds A, Humbert R, Miller B, Paige SL, Vernot B, Cheng JB,
Thurman RE, Sandstrom R, Haugen E, Heimfeld S, Murry CE, Akey JM,
Stamatoyannopoulos JA (2013) Developmental fate and cellular maturity encoded in

human regulatory DNA landscapes. Cell 154: 888–903.

11. Buenrostro JD, Corces MR, Lareau CA, Wu B, Schep AN, Aryee MJ, Majeti R, Chang HY,
Greenleaf WJ (2018) Integrated Single-Cell Analysis Maps the Continuous Regulatory
Landscape of Human Hematopoietic Differentiation. Cell 173: 1535–1548.e16.

12. Corces MR, Buenrostro JD, Wu B, Greenside PG, Chan SM, Koenig JL, Snyder MP,
Pritchard JK, Kundaje A, Greenleaf WJ, Majeti R, Chang HY (2016) Lineage-specific and
single-cell chromatin accessibility charts human hematopoiesis and leukemia evolution.
Nat Genet 48: 1193–1203.

13. Suvà ML, Rheinbay E, Gillespie SM, Patel AP, Wakimoto H, Rabkin SD, Riggi N, Chi AS,
Cahill DP, Nahed BV, Curry WT, Martuza RL, Rivera MN, Rossetti N, Kasif S, Beik S, Kadri
S, Tirosh I, Wortman I, et al. (2014) Reconstructing and reprogramming the tumor-
propagating potential of glioblastoma stem-like cells. Cell 157: 580–594.

14. Lan X, Jörg DJ, Cavalli FMG, Richards LM, Nguyen LV, Vanner RJ, Guilhamon P, Lee L,
Kushida MM, Pellacani D, Park NI, Coutinho FJ, Whetstone H, Selvadurai HJ, Che C, Luu B,
Carles A, Moksa M, Rastegar N, et al. (2017) Fate mapping of human glioblastoma reveals
an invariant stem cell hierarchy. Nature 549: 227–232.

15. Meyer M, Reimand J, Lan X, Head R, Zhu X, Kushida M, Bayani J, Pressey JC, Lionel AC,
Clarke ID, Cusimano M, Squire JA, Scherer SW, Bernstein M, Woodin MA, Bader GD, Dirks
PB (2015) Single cell-derived clonal analysis of human glioblastoma links functional and
genomic heterogeneity. Proc Natl Acad Sci U S A. Available:
http://dx.doi.org/10.1073/pnas.1320611111.

16. Orzan F, De Bacco F, Crisafulli G, Pellegatta S, Mussolin B, Siravegna G, D'Ambrosio A,
Comoglio PM, Finocchiaro G, Boccaccio C (2017) Genetic Evolution of Glioblastoma Stem-
Like Cells from Primary to Recurrent Tumor. Stem Cells. Available:
http://dx.doi.org/10.1002/stem.2703.

17. Rheinbay E, Suvà ML, Gillespie SM, Wakimoto H, Patel AP, Shahid M, Oksuz O, Rabkin
SD, Martuza RL, Rivera MN, Louis DN, Kasif S, Chi AS, Bernstein BE (2013) An aberrant
transcription factor network essential for Wnt signaling and stem cell maintenance in
glioblastoma. Cell Rep 3: 1567–1579.

18. Miller TE, Liau BB, Wallace LC, Morton AR, Xie Q, Dixit D, Factor DC, Kim LJY, Morrow JJ,
Wu Q, Mack SC, Hubert CG, Gillespie SM, Flavahan WA, Hoffmann T, Thummalapalli R,
Hemann MT, Paddison PJ, Horbinski CM, et al. (2017) Transcription elongation factors
represent in vivo cancer dependencies in glioblastoma. Nature 547: 355–359.

19. Park NI, Guilhamon P, Desai K, McAdam RF, Langille E, O'Connor M, Lan X, Whetstone H,
Coutinho FJ, Vanner RJ, Ling E, Prinos P, Lee L, Selvadurai H, Atwal G, Kushida M, Clarke
ID, Voisin V, Cusimano MD, et al. (2017) ASCL1 Reorganizes Chromatin to Direct Neuronal
Fate and Suppress Tumorigenicity of Glioblastoma Stem Cells. Cell Stem Cell. Available:
http://dx.doi.org/10.1016/j.stem.2017.06.004.

20. Liau BB, Sievers C, Donohue LK, Gillespie SM, Flavahan WA, Miller TE, Venteicher AS,
Hebert CH, Carey CD, Rodig SJ, Shareef SJ, Najm FJ, van Galen P, Wakimoto H, Cahill DP,
Rich JN, Aster JC, Suvà ML, Patel AP, et al. (2017) Adaptive Chromatin Remodeling Drives
Glioblastoma Stem Cell Plasticity and Drug Tolerance. Cell Stem Cell 20: 233–246.e7.

21. Pollard SM, Yoshikawa K, Clarke ID, Danovi D, Stricker S, Russell R, Bayani J, Head R, Lee
M, Bernstein M, Squire JA, Smith A, Dirks P (2009) Glioma stem cell lines expanded in

adherent culture have tumor-specific phenotypes and are suitable for chemical and
genetic screens. Cell Stem Cell 4: 568–580.

22. Wang Q, Hu B, Hu X, Kim H, Squatrito M, Scarpace L, deCarvalho AC, Lyu S, Li P, Li Y,
Barthel F, Cho HJ, Lin Y-H, Satani N, Martinez-Ledesma E, Zheng S, Chang E, Sauvé C-
EG, Olar A, et al. (2017) Tumor Evolution of Glioma-Intrinsic Gene Expression Subtypes
Associates with Immunological Changes in the Microenvironment. Cancer Cell 32: 42–
56.e6.

23. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, Paulovich A,
Pomeroy SL, Golub TR, Lander ES, Mesirov JP (2005) Gene set enrichment analysis: a
knowledge-based approach for interpreting genome-wide expression profiles. Proc Natl
Acad Sci U S A 102: 15545–15550.

24. Reimand J, Arak T, Adler P, Kolberg L, Reisberg S, Peterson H, Vilo J (2016) g:Profiler-a
web server for functional interpretation of gene lists (2016 update). Nucleic Acids Res 44:
W83–W89.

25. Capper D, Jones DTW, Sill M, Hovestadt V, Schrimpf D, Sturm D, Koelsche C, Sahm F,
Chavez L, Reuss DE, Kratz A, Wefers AK, Huang K, Pajtler KW, Schweizer L, Stichel D,
Olar A, Engel NW, Lindenberg K, et al. (2018) DNA methylation-based classification of
central nervous system tumours. Nature 555: 469–474.

26. MacLeod G, Bozek DA, Rajakulendran N, Monteiro V, Ahmadi M, Steinhart Z, Kushida
MM, Yu H, Coutinho FJ, Cavalli FMG, Restall I, Hao X, Hart T, Luchman HA, Weiss S, Dirks
PB, Angers S (2019) Genome-Wide CRISPR-Cas9 Screens Expose Genetic Vulnerabilities
and Mechanisms of Temozolomide Sensitivity in Glioblastoma Stem Cells. Cell Rep 27:
971–986.e9.

27. Tang Z, Li C, Kang B, Gao G, Li C, Zhang Z (2017) GEPIA: a web server for cancer and
normal gene expression profiling and interactive analyses. Nucleic Acids Res. Available:
http://dx.doi.org/10.1093/nar/gkx247.

28. Patel AP, Tirosh I, Trombetta JJ, Shalek AK, Gillespie SM, Wakimoto H, Cahill DP, Nahed
BV, Curry WT, Martuza RL, Louis DN, Rozenblatt-Rosen O, Suvà ML, Regev A, Bernstein
BE (2014) Single-cell RNA-seq highlights intratumoral heterogeneity in primary
glioblastoma. Science 344: 1396–1401.

29. Verhaak RGW, Hoadley KA, Purdom E, Wang V, Qi Y, Wilkerson MD, Miller CR, Ding L,
Golub T, Mesirov JP, Alexe G, Lawrence M, O'Kelly M, Tamayo P, Weir BA, Gabriel S,
Winckler W, Gupta S, Jakkula L, et al. (2010) Integrated genomic analysis identifies
clinically relevant subtypes of glioblastoma characterized by abnormalities in PDGFRA,
IDH1, EGFR, and NF1. Cancer Cell 17: 98–110.

30. Yin W, Tang G, Zhou Q, Cao Y, Li H, Fu X, Wu Z, Jiang X (2019) Expression Profile Analysis
Identifies a Novel Five-Gene Signature to Improve Prognosis Prediction of Glioblastoma.
Front Genet 10: 419.

31. Zuo S, Zhang X, Wang L (2019) A RNA sequencing-based six-gene signature for survival
prediction in patients with glioblastoma. Sci Rep 9: 2615.

32. Johnston M, Nikolic A, Ninkovic N, Guilhamon P, Cavalli F, Seaman S, Zemp F, Lee J,
Abdelkareem A, Ellestad K, Murison A, Kushida M, Coutinho F, Ma Y, Mungall A, Moore R,
Marra M, Taylor M, Dirks P, et al. (2019) High-resolution structural genomics reveals new
therapeutic vulnerabilities in glioblastoma. Genome Res. Available:

http://dx.doi.org/10.1101/gr.246520.118.

33.   Bailey SD, Desai K, Kron KJ, Mazrooei P, Sinnott-Armstrong NA, Treloar AE, Dowar M,
      Thu KL, Cescon DW, Silvester J, Yang SYC, Wu X, Pezo RC, Haibe-Kains B, Mak TW,
      Bedard PL, Pugh TJ, Sallari RC, Lupien M (2016) Noncoding somatic and inherited single-
      nucleotide variants converge to promote ESR1 expression in breast cancer. Nat Genet 48:
      1260–1266.

34.   Sallari RC, Sinnott-Armstrong NA, French JD, Kron KJ, Ho J, Moore JH, Stambolic V,
      Edwards SL, Lupien M, Kellis M (2017) Convergence of dispersed regulatory mutations
      predicts driver genes in prostate cancer. bioRxiv: 097451. Available:
      https://www.biorxiv.org/content/10.1101/097451v2. Accessed 20 November 2019.

35.   Kim K, Jang K, Yang W, Choi E-Y, Park S-M, Bae M, Kim Y-J, Choi JK (2016) Chromatin
      structure-based prediction of recurrent noncoding mutations in cancer. Nat Genet 48:
      1321–1326.

36.   Schmitt AD, Hu M, Ren B (2016) Genome-wide mapping and analysis of chromosome
      architecture. Nat Rev Mol Cell Biol 17: 743–755.

37.   Zheng H, Xie W (2019) The role of 3D genome organization in development and cell
      differentiation. Nat Rev Mol Cell Biol 20: 535–550.

38.   O'Connor L, Gilmour J, Bonifer C (2016) The Role of the Ubiquitously Expressed
      Transcription Factor Sp1 in Tissue-specific Transcriptional Regulation and in Disease. Yale
      J Biol Med 89: 513–525.

39.   Liu T, Yuan X, Xu D (2016) Cancer-Specific Telomerase Reverse Transcriptase (TERT)
      Promoter Mutations: Biological and Clinical Implications. Genes 7. Available:
      http://dx.doi.org/10.3390/genes7070038.

40.   Lee WS, Kwon J, Yun DH, Lee YN, Woo EY, Park M-J, Lee J-S, Han Y-H, Bae IH (2014)
      Specificity protein 1 expression contributes to Bcl-w-induced aggressiveness in
      glioblastoma multiforme. Mol Cells 37: 17–23.

41.   Trépant A-L, Bouchart C, Rorive S, Sauvage S, Decaestecker C, Demetter P, Salmon I
      (2015) Identification of OLIG2 as the most specific glioblastoma stem cell marker starting
      from comparative analysis of data from similar DNA chip microarray platforms. Tumour
      Biol 36: 1943–1953.

42.   Gabriely G, Wheeler MA, Takenaka MC, Quintana FJ (2017) Role of AHR and HIF-1α in
      Glioblastoma Metabolism. Trends Endocrinol Metab 28: 428–436.

43.   Michaelson JJ, Shin M-K, Koh J-Y, Brueggeman L, Zhang A, Katzman A, McDaniel L, Fang
      M, Pufall M, Pieper AA (2017) Neuronal PAS Domain Proteins 1 and 3 Are Master
      Regulators of Neuropsychiatric Risk Genes. Biol Psychiatry 82: 213–223.

44.   Rajakulendran N, Rowland KJ, Selvadurai HJ, Ahmadi M, Park NI, Naumenko S, Dolma S,
      Ward RJ, So M, Lee L, MacLeod G, Pasiliao C, Brandon C, Clarke ID, Cusimano MD,
      Bernstein M, Batada N, Angers S, Dirks PB (2019) Wnt and Notch signaling govern self-
      renewal and differentiation in a subset of human glioblastoma stem cells. Genes Dev 33:
      498–510.

45.   Cheng P, Wang J, Waghmare I, Sartini S, Coviello V, Zhang Z, Kim S-H, Mohyeldin A,
      Pavlyukov MS, Minata M, Valentim CLL, Chhipa RR, Bhat KPL, Dasgupta B, La Motta C,
      Kango-Singh M, Nakano I (2016) FOXD1-ALDH1A3 Signaling Is a Determinant for the
      Self-Renewal and Tumorigenicity of Mesenchymal Glioma Stem Cells. Cancer Res 76:

626   7219–7230.

627   46.   Koga M, Matsuda M, Kawamura T, Sogo T, Shigeno A, Nishida E, Ebisuya M (2014) Foxd1
628   is a mediator and indicator of the cell reprogramming process. Nat Commun 5: 3197.

629   47.   Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ (2013) Transposition of native
630   chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-binding
631   proteins and nucleosome position. Nat Methods 10: 1213–1218.

632   48.   Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA, Vesuna S,
633   Satpathy AT, Rubin AJ, Montine KS, Wu B, Kathiria A, Cho SW, Mumbach MR, Carter AC,
634   Kasowski M, Orloff LA, Risca VI, Kundaje A, Khavari PA, et al. (2017) An improved ATAC-
635   seq protocol reduces background and enables interrogation of frozen tissues. Nat
636   Methods 14: 959–962.

637   49.   Wang B, Mezlini AM, Demir F, Fiume M, Tu Z, Brudno M, Haibe-Kains B, Goldenberg A
638   (2014) Similarity network fusion for aggregating data types on a genomic scale. Nat
639   Methods 11: 333–337.

640   50.   Shin H, Liu T, Manrai AK, Liu XS (2009) CEAS: cis-regulatory element annotation system.
641   Bioinformatics 25: 2605–2606.

642   51.   Schep AN, Wu B, Buenrostro JD, Greenleaf WJ (2017) chromVAR: inferring transcription-
643   factor-associated accessibility from single-cell epigenomic data. Nat Methods 14: 975–
644   978.

645   52.   Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM 3rd, Hao Y,
646   Stoeckius M, Smibert P, Satija R (2019) Comprehensive Integration of Single-Cell Data.
647   Cell 177: 1888–1902.e21.

648   53.   Müller S, Cho A, Liu SJ, Lim DA, Diaz A (2018) CONICS integrates scRNA-seq with DNA
649   sequencing to map gene expression to tumor sub-clones. Bioinformatics 34: 3217–3219.

650   54.   Morris TJ, Butcher LM, Feber A, Teschendorff AE, Chakravarthy AR, Wojdacz TK, Beck S
651   (2014) ChAMP: 450k Chip Analysis Methylation Pipeline. Bioinformatics 30: 428–430.

652   55.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M,
653   Gingeras TR (2013) STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29: 15–21.

654   56.   Li B, Dewey CN (2011) RSEM: accurate transcript quantification from RNA-Seq data with
655   or without a reference genome. BMC Bioinformatics 12: 323.

656   57.   Anders S, Huber W (2010) Differential expression analysis for sequence count data.
657   Genome Biol 11: R106.

658   58.   Weirauch MT, Yang A, Albu M, Cote AG, Montenegro-Montero A, Drewe P, Najafabadi
659   HS, Lambert SA, Mann I, Cook K, Zheng H, Goity A, van Bakel H, Lozano J-C, Galli M,
660   Lewsey MG, Huang E, Mukherjee T, Chen X, et al. (2014) Determination and inference of
661   eukaryotic transcription factor sequence specificity. Cell 158: 1431–1443.

662   59.   Li W, Xu H, Xiao T, Cong L, Love MI, Zhang F, Irizarry RA, Liu JS, Brown M, Liu XS (2014)
663   MAGeCK enables robust identification of essential genes from genome-scale
664   CRISPR/Cas9 knockout screens. Genome Biol 15: 554.

665   60.   Hart T, Tong AHY, Chan K, Van Leeuwen J, Seetharaman A, Aregger M, Chandrashekhar
666   M, Hustedt N, Seth S, Noonan A, Habsid A, Sizova O, Nedyalkova L, Climie R,
667   Tworzyanski L, Lawson K, Sartori MA, Alibeh S, Tieu D, et al. (2017) Evaluation and Design
668   of Genome-Wide CRISPR/SpCas9 Knockout Screens. G3 7: 2719–2727.

61.   Hart T, Chandrashekhar M, Aregger M, Steinhart Z, Brown KR, MacLeod G, Mis M,

Zimmermann M, Fradet-Turcotte A, Sun S, Mero P, Dirks P, Sidhu S, Roth FP, Rissland
OS, Durocher D, Angers S, Moffat J (2015) High-Resolution CRISPR Screens Reveal
Fitness Genes and Genotype-Specific Cancer Liabilities. Cell 163: 1515–1526.
62. Tam RY, Yockell-Lelièvre J, Smith LJ, Julian LM, Baker AEG, Choey C, Hasim MS,
Dimitroulakos J, Stanford WL, Shoichet MS (2019) Rationally Designed 3D Hydrogels
Model Invasive Lung Diseases Enabling High-Content Drug Screening. Adv Mater 31:
e1806214.
63. Therneau T (2015) A Package for Survival Analysis in S. version 2.38. Available:
https://CRAN.R-project.org/package=survival.
64. Drawing Survival Curves using "ggplot2" [R package survminer version 0.4.6] (n.d.).
Available: https://cran.r-project.org/web/packages/survminer/index.html. Accessed 16
October 2019.
682

ACKNOWLEDGMENTS

We would like to thank the Princess Margaret Genomics Centre for their assistance in this study

and 10x Genomics, especially Adam Jerauld, Ariel Royall and John Chevillet for training and

support. We also thank Aude Gerbaud and Stacey Krunholtz for their help with figure formatting

and David Vetrie for his thoughtful input on the manuscript. This work was supported by the

Princess Margaret Cancer Foundation, the Canadian Epigenetics, Environment, and Health

Research Consortium (CEEHRC) initiative from the Canadian Institute for Health Research

(CIHR) (TGH-158221 to ML, SA and PBD) and SU2C Canada Cancer Stem Cell Dream Team

Research Funding (SU2C-AACR-DT-19-15) provided by the Government of Canada through

Genome Canada and the Canadian Institute of Health Research, with supplemental support

from the Ontario Institute for Cancer Research, through funding provided by the Government

of Ontario. Stand Up To Cancer Canada is a Canadian Registered Charity (Reg. # 80550 6730

RR0001). Research Funding is administered by the American Association for Cancer Research

International - Canada, the Scientific Partner of SU2C Canada. This study was also conducted

with the support of the Ontario Institute for Cancer Research through funding provided by the

Government of Ontario for the Brain Cancer Translational Research Initiative. PG was supported

by a CIHR Fellowship (MFE 338954). Funding to HAL and SW was from CIHR. MG holds a Canada

Research Chair in Brain Cancer Epigenomics and is supported by CIHR, NSERC and the Alliance

for Cancer Gene Therapy. PBD is also supported by CIHR, OICR, the Terry Fox Research Institute,

and the Hospital for Sick Children Foundation, Jessica's Footprint, Hopeful Minds, and the

Bresler Family. PBD holds a Garron Chair in Childhood Cancer Research at the Hospital for Sick

Children. SA is supported by the CIHR, the Terry Fox Institute and the Canadian Cancer Society.

ML holds an Investigator Award from the Ontario Institute for Cancer Research and a Canadian

Institutes of Health Research (CIHR) New Investigator Award.

AUTHOR CONTRIBUTIONS

ML, PG, and PBD conceptualized and designed the study assisted by FJC. PG conducted the

genomics experiments, designed and/or implemented most of the computational and statistical

approaches, and made the figures. MMK and RH performed all the tissue culture, under the

supervision of PBD, HAL and SW. AN, KE, and DS generated the scATAC-seq data from samples

provided by JAC, under the supervision of MG. BN performed the alignment of the scATAC-seq

data. NR, XH and RH performed the xenografts, under the supervision of PBD, HAL, and SW. CA

generated some of the ATAC-seq data used in the xenograft classification. FMGC ran the

spectral clustering, under the supervision of MDT. GM, NR and SA contributed the essentiality

screen data. HW conducted the endpoint xenograft staining. LJS performed the invasion assay

under the supervision of MSS. SAMT contributed to the computational analysis design, under

supervision of BHK and ML. The manuscript was written by PG, PBD and ML with input from all

other authors.

COMPETING INTERESTS

The authors declare no competing interests.

FIGURES

Fig. 1. The diverse GBM cancer stem cell pool. (A) UMAP representation of chromatin

accessibility across four primary GBM. **(B)** UMAPs with tumor cells assigned to cellular states.

(C) UMAP modules are grouped by dominant cellular state. **(D)** UMAPs with cancer stem cells

highlighted based on the enrichment of GBM cancer stem transcription factor promoters. **(E)**

Distribution of cancer stem cells across the modules dominated by each cellular state.

Fig. 2. GSCs recapitulate the GBM cancer stem cell population. (A) Schematic representation

of the GSC derivation process, from patient tumor to GSC-enriched population. **(B)** Genomic

feature enrichment of accessible chromatin peaks. **(C)** Saturation curve for the 27 GSCs. **(D)**

UMAPs with GSCs highlighted based on the enrichment of shared accessible regions across

GSCs. **(E)** Proportion of UMAP modules assigned to cancer stem cells and GSCs. **(F)** Correlation

of z-scores for each signature for each cell in each primary GBM.

Fig. 3. Three GSC states driven by chromatin accessibility. (A) Spectral clustering of ATAC-

seq signal across peaks in 27 GSCs. **(B)** Top: enrichment of TCGA subtypes across GSCs and

comparison to GSC states; Bottom: spectral clustering of gene expression across GSCs and

comparison to chromatin-derived GSC states. **(C)** Gene set enrichment analysis in each GSC

state. **(D)** CNAs across GSCs identified from DNA methylation array data cluster GSCs into four

subgroups. **(E)** Number and percentage of peaks unique and shared in each GSC state. **(F)**

Saturation analysis of each individual state.

Fig. 4. Functional diversity between GSC states drives survival in GBM. (A) Motif family

enrichment in each cluster; log2(Fold Enrichment) > 0.5 threshold selected based on the

distribution of values in each cluster (Fig. S1A). **(B)** Z-score distribution of key essential genes in

each cluster. Red line corresponds to the empirically determined threshold for essentiality in

each tested line, scaled and adjusted to zero. Boxplot whiskers in this case extend to data

extremes. Side barplots show the total count of the key cluster-specific regulators found

essential in each subtype. **(C)** UMAPs with GSCs from each state highlighted based on the

enrichment of the top differentially accessible regions in each GSC state. **(D)** Kaplan-Meier plot

for orthotopic xenografts grouped by GSC state. The dotted lines indicate median survival. The

pairwise p-values are also significant for Invasive vs Reactive (p=0.02) and Invasive vs

Constructive (p=0.045) but not for Reactive vs Constructive (p=0.45). **(E)** TCGA samples ordered

by increasing concordance with Invasive GSCs and grouped into three subgroups: <1, 1-

1.65,>1.65. **(F)** Kaplan-Meier plot for TCGA samples grouped by concordance with Invasive

GSCs. The dotted line indicates median survival. When considering pairwise comparisons, only

the Invasive-high and Invasive-low subgroups were significantly different (p=0.0043). Further

subgrouping of the TCGA samples into smaller intervals of concordance z-score yielded no

benefit, preserving the Invasive-high subgroup as the only one with significantly poorer

prognosis (Fig. S3D-E).

SUPPLEMENTARY DATA

Fig. S1. (A-D) UMAPs showing cells confirmed to be GBM tumor cells, with gains of chr7 or

losses of chr10. **(E-G)** Overlaid ATAC signal tracks for the 27 GSCs at the 92 most differentially

accessible regions between GSC states as determined by pairwise wilcoxon test and a median

signal fold change equal or greater than 2. (E) Most accessible in Reactive state GSCs. (F) Most

accessible in Constructive state GSCs. (G) Most accessible in Invasive state GSCs.

Fig. S2. (A) The log2(fold enrichment) for all motif families were ordered and plotted. A cut-off

threshold to select families for follow-up was selected based on the steep inflection of the curve

above 0.5. **(B)** Schematic of drop-out essentiality screen using GSCs stably expressing Cas9 and

gRNA libraries. **(C-E)** The expression values determined by RNA-seq for all transcription factors

whose motif is enriched in each GSC state and exclusively essential in at least one GSC of each

state is plotted. Bold colors highlight factors essential in all tested GSCs in that state. In the case

of the Reactive state, blue boxplots correspond to SP1 and members of its regulatory network.

(F) Expression levels of key transcription factors in tumor and normal samples, analysed and

displayed using GEPIA[27]. **(G)** Gene enrichment analysis of subtype-specific essential genes

(Essentiality), upregulated and differentially accessible genes (ATAC and RNA combined), and

putative targets of key transcription factors.

Fig. S3. (A) Kaplan-Meier plot for TCGA samples grouped by GSC state. The dotted lines indicate

median survival. **(B)** TCGA samples ordered by increasing concordance with Invasive GSCs and

grouped into 8 subgroups. **(C)** Kaplan-Meier plot for TCGA samples grouped by concordance

with Invasive GSCs. The dotted line indicates median survival.

Fig. S4. (A) Left: Top and side views of confocal z-stack images of Invasive GSCs G705, G837 and

healthy control hf6562 cultured on hyaluronan-based hydrogels. Right: percent invasion,

defined as the percentage of total cells located below a 75 μm threshold from the surface of the

hydrogel. Plots are shown as mean with standard deviation, and statistics shown are from one-

way ANOVA with multiple comparisons with Dunnet's test correction, comparing all patient cell

lines (dark grey) to the hf6562 healthy control. Data were displayed with p values represented

as * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$, **** $p \leq 0.0001$. **(B)** Human-specific staining of mouse

brains injected with GSCs from G411 (upper panel) and G837 (lower panel) with higher

magnification (black inset) at the mass boundaries.

A

B

C

1: Reactive

2: Constructive

3: Invasive

D

E

F