Phosphorylation of the Type VI secretion system sheath controls *Francisella* virulence

Jason Ziveri[1], Cerina Chhuon[2], Anne Jamet[1], Guénolé Prigent[1], Fabiola Tros[1], Monique Barel[1], Claire Lays[4], Thomas Henry[4], Nicholas H Keep[3], Ida Chiara Guerrera[2]*, Alain Charbit[1]*

[1] Université Paris Descartes, Sorbonne Paris Cité, Bâtiment Leriche, Paris, INSERM U1151 - CNRS UMR 8253, Institut Necker-Enfants Malades. Team 11: Pathogenesis of Systemic Infections, Paris, France

[2] Plateforme protéomique 3P5-Necker, Université Paris Descartes - Structure Fédérative de Recherche Necker, INSERM US24/CNRS UMS3633, Paris 75014, France

[3] Crystallography, Institute for Structural and Molecular Biology, Department of Biological Sciences Birkbeck, University of London, United Kingdom

[4] CIRI, Centre International de Recherche en Infectiologie, Université Lyon, Inserm, U1111, University Claude Bernard Lyon 1, CNRS, UMR5308, École Normale Supérieure de Lyon, Labex Ecofect, Eco-evolutionary dynamics of infectious diseases, F-69007, LYON, France

Running title: Phosphorylation of IglB is critical for *Francisella* sheath assembly

Keywords: Type 6 secretion system, Phosphoproteome, *Francisella tularensis*, IglB.

Bâtiment Leriche. 14 Rue Maria Helena Vieira Da Silva CS 61431 - 75993 PARIS – FRANCE

*Corresponding authors: Ida Chiara Guerrera, Alain Charbit.

e-mail: alain.charbit@inserm.fr; chiara.guerrera@inserm.fr

Tel: 33 1 – 72 60 65 11 — Fax: 33 1 - 72 60 65 13

Abstract

Many Gram-negative bacteria use type VI secretion systems (T6SS) to translocate effector proteins into eukaryotic or prokaryotic cells. *Francisella* possesses a non-canonical T6SS encoded on the *Francisella* pathogenicity island (FPI) that is essential for efficient phagosomal escape and access to the cytosol of infected host cells. Using a global and site-specific phosphoproteomic analysis of *Francisella*, we identified a unique phosphorylation site on tyrosine 139 of IglB, the TssC homologue and a key component of the T6SS contractile sheath. Phosphorylated IglB was exclusively found in the non-oligomerized form of the protein, suggesting that sheath assembly requires IglB dephosphorylation. All the different amino acid substitutions at site 139 of IglB all impaired sheath assembly. The data presented suggest that the phosphorylation status of IglB plays a critical role in T6SS activity and, hence, *Francisella* pathogenicity. We propose that this post-translational modification of the sheath may constitute a previously unrecognized mechanism to modulate the dynamics of assembly-disassembly of the T6SS.

Author summary

Francisella possesses a non-canonical T6SS that is essential for efficient phagosomal escape and access to the cytosol of infected host cells. Using a global and site-specific phosphoproteomic analysis of *Francisella* we identified here a unique phosphorylation site on IglB, the TssC homologue and a key component of the T6SS contractile sheath. Our data suggest that the phosphorylation status of IglB plays a critical role in the assembly of a functional T6SS. This post-translational modification of the sheath may constitute a previously unrecognized mechanism to modulate the dynamics of assembly-disassembly of the T6SS.

Introduction

Francisella tularensis is the causative agent of the zoonotic disease tularemia (1-4). This

facultative intracellular pathogen is able to infect a variety of different cell types but, *in vivo*, is

thought to replicate and disseminate mainly in phagocytes (5). The four major subspecies

(subsp) of *F. tularensis* currently listed are the subsps: *tularensis*, *holarctica, mediasiatica*

and *novicida* (the latter is also called *F. novicida*). These subspecies differ in their virulence

and geographical origin (6) but all cause a fulminant disease in mice that is similar to

tularemia in humans (7). Although *F. novicida* is rarely pathogenic in humans*,* its genome

shares a high degree of nucleotide sequence conservation with the human pathogenic

species and is thus widely used as a model to study highly virulent subspecies (8-10). The

efficiency of *Francisella* systemic dissemination is tightly linked to its capacity to multiply in

the cytosolic compartment of infected macrophages. The transcriptional and post-

transcriptional regulatory processes, controlling phagosomal escape and cytosolic bacterial

multiplication, have been already well-characterized (11-17). In contrast, the contribution of

post-translational protein modifications (PTMs) has remained largely underestimated. For

example, protein glycosylation has been only very recently identified in *Francisella* (18, 19)

and the importance of protein phosphorylation in *Francisella* pathogenesis is still largely

unknown (20), although it constitutes one of the most prevalent PTM involved in controlling

protein activity.

Two major systems of protein phosphorylation exist in Gram-negative bacteria: i) two-

component systems (TCSs), composed of a membrane-bound sensor kinase that auto-

phosphorylates at a conserved histidine residue and a cognate response regulator that is

phosphorylated at a conserved aspartate residue; and ii) serine/threonine or tyrosine protein

kinases (Ser/Thr/Tyr-K) and their associated phosphatases (21-23 and references therein).

In addition, many Gram-negative bacteria also possess a phospho-relay system, designated

phosphoenol pyruvate:carbohydrate phosphotransferase system (or PTS), that is more

specifically devoted to sugar transport (24, 25 and references therein). *Francisella* genomes,

do not encode any PTS (26) but encode elements of two-component regulatory systems

(TCS). Whereas most *Francisella* strains possess only incomplete TCS pairs (27), *F.*

novicida has two complete and two incomplete TCS pairs (28, 29)*.*

Our careful inspection of the *Francisella* genomes did not identify any gene encoding for a

putative serine/threonine-protein kinase or tyrosine kinase. In spite of this lack of canonical

kinases, we hypothesized that such post-translational protein modifications might occur in

Francisella and contribute to pathogenesis. By using state-of-the art mass spectrometry

analyses, we identified and quantified 82 peptides bearing phosphorylated residues in the

proteome of *F. novicida.* One peptide, corresponding to a phosphorylation of IglB on tyrosine

139 (Tyr139), particularly attracted our attention. Indeed, IglB is analogous to the TssC

protein in canonical T6SS and oligomerizes with IglA (TssB homologue) to form the

contractile sheath (30) of its type six secretion apparatus (T6SS).

T6SS have been described as contractile syringes used by bacterial species either to kill

neighboring bacteria competing for the same niche or to target eukaryotic cells upon

infection. Bioinformatic analyses have divided bacterial T6SS in three phylogenetically

distinct subtypes (designated T6SSi-iii) (31). Whereas most of the well-characterized T6SS

belong to the T6SSi (including in *P. areruginosa* and *V. cholerae*), *Francisella* is currently the

only bacterium to possess a T6SSii family member (32). *Francisella* T6SS has been shown,

thus far, to be exclusively involved in the intracellular life cycle of the pathogen. Phagosomal

escape occurs as early as 30 minutes post-infection (33) and critically depends on the T6SS

that is encoded by a 33 kb locus named *Francisella* pathogenicity island (FPI) (34-36, 37,

38). The FPI (**Supplementary Fig. 1**) also encodes a number of proteins of unknown

functions, some of which (such as IglF,PdpC or PdpD) have been proposed to be T6SS

effectors(9). Remarkably, the FPI does not encode any homologue of the unfoldase ClpV,

present in all the canonical T6SS, involved in the recycling of the contracted sheaths.

However, it has been recently shown that, in *F. novicida*, the general chaperone ClpB (39)

could fulfill ClpV function(8) by specifically colocalizing with the contracted sheath and

promoting its disassembly. Of note, the FPI is duplicated in all the subspecies of *F. tularensis*

except in *F. novicida*, in which it is present in a single copy.

Our current knowledge of the structure and functional assembly of the *Francisella* T6SSii

are mainly based on recent cryo-electron microscopy data (30) and structural homologies

with other members of the T6SSi. It has been proposed that the tube comprised of IglC

subunits (Hcp homologue) that form hexameric rings, fits within the cavity of the IglA/IglB

sheath (TssB/C in *Pseudomonas*, VipA/B in *Vibrio*). Contraction of the sheath results in the

ejection of the Hcp tube, together with effector proteins located within the tube and on top of

the tip of the tube (40). By analogy with other T6SS structural data, it has been suggested

but not formally proven that, in *Francisella*, the sheath assembles on a preassembled IglC

tube (30). This assumption awaits additional structural information on the different

conformations of the *Francisella* T6SS (from fully elongated to fully contracted stages).

The data presented here reveal that the phosphorylation status of a single tyrosine

residue of IglB is essential for the proper activity of the T6SS and, hence, *Francisella*

pathogenicity. We propose that, upon infection, this post-translational mechanism may

participate in the control of T6SS dynamics, for a rapid and reversible response to its

different intracellular environments.

Results

The phosphoproteome of *Francisella*

We carried out a global and site-specific phosphoproteomic analysis of *F. novicida* (strain U112), based on phosphopeptide enrichment and high-resolution LC-MS/MS determination of whole cell lysates (see Methods). This analysis detected 82 phosphopeptides (68 with high probability for site attribution). The majority of the phosphosites (53 out of 82) were found on serine residues (S), while similar lower number of sites, 13 and 16, were found on threonine (T) and tyrosine (Y), respectively (**Fig 1A and S1 and Tables**).

Fig1. The phosphoproteome of *F. novicida*

(**A**) Distribution of phosphosites according to the modified amino acid (tyrosine,Y; threonine, T and serine, S). (**B**) Logos of phosphopeptide amino acid sequences; (**C**) Histograms of the most represented classes of protein. The values correspond to the number of proteins bearing phosphosites in each category.

In order to highlight possible consensus sequences, we aligned the sequences of the phosphoserine / threonine / tyrosine peptides. Although the number of peptides is perhaps too low to be conclusive, we could observe the recurrence of some amino acids in specific positions, like lysine (K) in position n-1 of phosphothreonines (pT) or apolar amino acids valine and phenylalanine (V, F) in n-1 of phosphotyrosines (pY) (**Fig 1B**). These phosphosites corresponded to 49 proteins, of which 13 presented multiple phosphorylation sites. The 49 proteins belonged to various functional categories and the most represented classes were carbohydrate metabolism, energy production and conversion and translation (**Fig 1C**).

Remarkably, the two proteins that constitute the T6SS sheath(30, 41, 42) (IglA and IglB,

S1 Fig) appeared to bear a unique phosphorylation site (at Ser173 for IglA, and Tyr139 for

IglB). This somewhat unexpected observation prompted us to determine whether the

phosphorylated forms of the two proteins were incorporated into the sheath upon its

assembly. For this, we first monitored sheath formation using the procedure previously

described in (30). Briefly, lysates of wild-type *F. novicida,* grown in the presence of 5% KCl in

order to trigger sheath formation, were layered onto 15%-55% sucrose gradients (**Fig 2A**)

and the different fractions were tested by western blotting with anti-Igl (A,B or C) antibodies.

IglA and IglB were detected in all the fraction of the gradient, although with variable

intensities (**Fig 2B**). In contrast, IglC was only detected in the upper fractions of the gradient

(in the 15% and in part of the 35% fractions). The fractions sedimenting in equilibrium at 55%

sucrose and below (55% Optiprep, grey arrow, **Fig 2A**) were further dialyzed and used for

negative staining TEM imaging. Confirming earlier reports, we observed rod-shaped particles

of variable length in this fraction by electron microscopy, corresponding to assembled

sheath-like structures (**Fig 2A**). We then compared by mass spectrometry the fractions at the

bottom of the gradient (sucrose 55%-Optiprep 55% density zone) that contained these

sheath-like particles, to the fractions at the top of the gradient (15-35% sucrose density

zone), presumably containing non-polymerized forms of the IglA and IglB proteins (hereafter

called "heavy fraction", H or "light fraction", L, respectively). The mass spectrometry analysis

of the H and L fractions identified over 1,145 proteins (**S3 Table**). Of note, proteins

exclusively identified in the H fraction included several Type IV pili assembly proteins (fimT,

PilE, PilF) that might interact with the sheath. IglA and IglB were detected in both H and L

fractions. In order to estimate the proportion of the Igl proteins in the two fractions, we

calculated the ratio of the intensity of the protein, proportional to the amount of protein in the

fraction. Fully supporting the western blotting analyses, IglA and B were more abundant in

the L fraction than in the H fraction and only 2% of IglC was detected in the H fraction (**Fig

2C**).

As illustrated in **Fig 2D**, the peptide bearing the phosphorylated tyrosine residue (pY139)

of IglB was exclusively detected in the L fraction of the gradient, suggesting that only the

soluble forms of IglB are phosphorylated (**S3 Table**). Unmodified peptides containing Tyr139

were however detected in both H and L fractions, indicating that soluble IglB include both

phosphorylated and unphosphorylated forms of the protein. The phosphopeptide, bearing the

phosphorylated serine residue of IglA, identified in the whole cell lysate, was not detected in

either of the two samples analyzed.   This lack of detection could be due either to an

insufficient amounts (below the threshold of detection), to its localization in other fractions of

the gradient that were not tested or to the fact that KCl treatment might have triggered its

dephosphorylation.

Fig 2. Importance of IglB amino acid Y139 in sheath formation

(**A**) Sucrose gradient. Lysates of bacteria grown in Schaedler K3 medium in the presence of

KCl were laid on top of a discontinuous sucrose gradient 15%-55%/Optiprep 55%. Left panel,

composition of the gradient; Right panel, transmission electron microscopy (TEM) of the

"heavy" fraction (grey arrow, topping the Optiprep cushion). The assembled sheath-like

structure, sedimenting at equilibrium to below 55% sucrose, showed rod-shaped particles of

variable length (100 - 600 nm). (**B**) Western blotting analysis of the different fractions, using

anti-Igl (A, B or C) antibodies. (**C**) Mass spectrometry analysis of the H and L fractions. Ratio

of the intensities of the proteins IglA, B and C detected in the H and the L fractions (H / L %).

(**D**) Proteome and phosphoproteome of cellular fractions. Intensities (in log2) of the non

phosphorylated (left) and phosphorylated (right) Tyr139-containing peptide in the "heavy" (H)

and "light" (L) fractions. The signal of the phosphopeptide was only detectable after

phosphoenrichment. nd, not detectable.

We decided to focus here on IglB and evaluate the importance of the phosphorylation on

Tyr139 on T6SS functionality and *Francisella* pathogenicity.

Comparative sequence analyses on 33 IglB proteins from 19 genomes of the *Francisella* genus were performed to evaluate the conservation of residue Tyr139 (**S2 Fig**). All, but 5 genomes, encode two IglB proteins (due to the duplication of the FPI in these genomes). Sequences were highly conserved between 31 IglB proteins sharing 93 to 100% amino acid identity. In contrast, the proteins encoded by *FTN_0043,* encoded in another genomic island termed the *F. novicida* island or FNI(42)) and *F7308_1917* (Fsp_TX077308) exhibit only 48% amino acid identity with the 31 other IglB proteins whereas they both share 85% identity. Excluding these 2 outliers, among 506 sites, 458 were without polymorphism (91%), with Tyr139 conserved in all IglB sequences (**S2 Fig**). Of note, since IglB is among the most conserved components of T6SS(30, 41) (**S1 Fig**) we were able to retrieve orthologues of IglB encoded outside of *Francisella* genus. Out of 535 non-redundant IglB orthologues (also named VipB, TssC or EvpB/VC_A0108), we found 90 sequences with a tyrosine in the vicinity of Tyr139 site outside of *Francisella* genus. In 82 sequences, the predicted location of the tyrosine residue was similar to that of Tyr139 (in a turn region between two helices) (**S2 Fig**).

3D analysis

Inspection of the 3.7 Å EM reconstruction of the *F. novicida* sheath[32] (PDB 3j9o) does not show any atoms close enough to form a hydrogen bond with the hydroxyl of Tyr139. However, the carboxylates of Asp114 from IglB, and Asp63 of IglA in the next dimer, are around 5 Å away. At the resolution of the reconstruction, side chain positions are not clearly defined and so these may approach close enough to Tyr139 OH to form hydrogen bond interactions stabilizing the assembly.

There is space for a phosphate group on this atom as the OH is pointing into a pocket in the structure (**Fig 3**). This was modeled by adding a phosphate group (using ccp4mg to position a phosphate and then editing the PDB file to generate a PTR residue) onto the hydroxyl and showing there were no steric clashes. However, the negatively charged

phosphate would induce charge repulsion with the nearby Asps from the next dimer most

likely preventing assembly of the strands when Tyr139 is phosphorylated.

Hence, the charge repulsion provoked by addition of the phosphogroup (when Y139 is

phosphorylated) is likely to weaken the stability of the assembled sheath.

Fig 3. Predicted role of Y139 of IglB in sheath assembly

(**A**) Overview of the sheath that consists of 12 helical strands based on EM reconstruction at

3.7 Å. Each unit consists of a IglA/IglB dimer. The same colors were used in the further

pictures. The chains near Tyr139 of IglB consist of 2 IglA/IglB dimers from the same strand

(pink/grey and coral/sky blue). The N-terminal tail of an IglA (blue) from an adjacent strand

wraps round to close to Y139 of this strand. (**B**) A closer view of just the selected chains.

(**C**) Close up of the region of Tyr139. The hydroxyl is pointing into a pocket and not

obviously making a Hydrogen bond. However, the carboxylates of Asp114 from IglB and

Asp63 of IglA in the next dimer are around 5 Å away. At the resolution of the reconstruction

side chain density is not well defined and these could in fact be close enough to H bond.

(**D**) Surface of the assembly clipped to show Tyr139. There is enough space for a Phospho

group on the Tyr. The two Asps will however be close to the phosphogroup and so there is

likely to be charge repulsion impairing the sheath assembly or weakening the sheath stability

when Y139 is phosphorylated.

Role of IglB phosphorylation on IglA/IglB interaction and sheath formation

We evaluated the importance of the phosphorylation site in IglB on T6SS assembly and

bacterial virulence by constructing two IglB variants in which Tyr139 was substituted either

by an alanine (Ala) or by the non-phosphorylatable aromatic analogue phenylalanine (Phe).

These IglB mutated proteins (designated Y139A and Y139F, respectively) were expressed *in*

trans in a Δ*iglB* mutant of *F. novicida* carrying a chromosomal deletion of the entire *iglB*

gene, (see Methods). A *F. novicida* mutant with a deletion of the entire *FPI*, unable to escape

from phagosomes and hence to grow in macrophages (43), was used as a negative control.

The mutations Y139A and Y139F had no effect on bacterial growth in broth (**S3 Fig**). The

impact of Y139A and Y139F substitutions was first tested by Western blotting on whole cell

lysates. As expected (**Fig 4A**), IglB was detected in wild-type and complemented strains but

not in the ΔiglB and ΔFPI strains. IglA was detected in all the strains, except the ΔFPI mutant,

but in clearly lower amounts in the ΔiglB strain. This observation is in agreement with earlier

observations (44), suggesting that the presence of IglB increases the stability of the IglA

protein.

A classical measure of basal T6SS function is the export of an Hcp-related protein.

Therefore, we examined IglB-dependent IglC export in the different *F. novicida* mutant

strains by Western-blot on bacteria grown in the presence -or absence- of 5% KCl. In

agreement with previous reports, IglC was detected in the culture supernatant in the

presence, but not in the absence, of KCl, in the wild-type and $iglB_{WT}$ complemented strains.

In contrast, deletion of *iglB* as well as the two single amino acid substitutions in IglB

abolished the secretion of IglC into the culture medium (**Fig 4B**), revealing that a wild-type

form of IglB is required for IglC secretion. KCl treatment did not modify the amounts of IglA,

IglB and IglC proteins detected by western blotting on whole cells (bacterial pellet fractions).

To test if any of the two substitutions at Tyr139 would impair IglA/IglB heterodimer

formation, we performed co-immunoprecipitation assays (**Fig 4C**), using either anti-IglA to

precipitate the complex, followed by western blotting with anti-IglB (upper panel) or anti-IglB,

followed by western blotting with anti-IglA (lower panel). In both assays, the two IglB mutant

alleles (Y139A and Y139F) were able to interact with IglA like wild-type IglB.

Since the presence of sheath-like structures correlated with the presence of non-

phosphorylated forms of IglB (**Fig 2**), we anticipated that the non-phosphorylatable Y139F

mutant might permanently produce oligomerized sheath-like structures. To test this

possibility, we performed sucrose gradient fractionation followed by western blotting (as

described previously) on the Y139F mutant strain. In contradiction of the above hypothesis,

the proteins IglA and IglB were detected in the L fractions of the gradient but not in the H

fraction corresponding to the sheath (**Fig 3D**) and no sheath-like structure could be

visualized by TEM in the H fraction of the Y139F mutant. Mass spectrometry analysis of the

fractions confirmed that only a very low portion of the IglA, B and C is present in the H

fractions compared to the L (**Fig 4E**). Since the only difference between the non-

phosphorylated form of Tyr139 and its non-phosphorylatable Phe analogue is the presence

of an additional hydroxyl moiety, it is likely that this hydroxyl contributes to sheath

formation/stability.

Fig 4. Immunodetection

(**A**) Western Blots (WB) of whole cell protein extracts. Upper gel: WB with anti-IglB; lower

gel: WB with anti-IglA. (**B**) Western Blots (WB) with anti-IglC on bacteria / culture filtrates.

Bacteria were grown in Schaedler-K3 supplemented or not with 5% KCl until late log phase

and then harvested by centrifugation. Culture supernatants were collected after filtration on

0.2 µM Millipore filters and concentrated using Amicon 3 kDa membranes. The equivalent of

200 µg of total protein were loaded onto each well. Upper gel, bacterial cultures grown

without KCl; lower gel, bacterial cultures supplemented with 5% KCl. (**C**) Co-

immunoprecipitations (Co-IP). Upper gel: IP-anti-IglA, followed by WB with anti-IglB; lower

gel: IP-anti-IglB, followed by WB with anti-IglA. (**D**) WBs of sucrose gradient fractions of the

Y139F mutant strain. Upper gel, WB with anti-IglB; Middle gel, WB with anti-IglA; lower gel,

WB with anti-IglC. (**E**) Mass spectrometry analysis of the H and L fractions of the Y139F

mutant strain. Ratio of the intensities of the proteins IglA, B and C detected in the H and the

L fractions (H / L %).

Critical role of IglB residue Y139 in _Francisella_ pathogenesis

We next evaluated the impact of the Y139A and Y139F mutations on bacterial virulence by

studying their behavior in macrophages and in the mouse model. The ability of wild-type _F._

novicida (WT), Δ*iglB* and complemented strains (CpWT, Y139A and Y139F) to survive and

multiply in murine macrophage-like J774.1 cells was monitored over a 24 h-period (**Fig 5A**).

Confirming earlier reports(45), intracellular multiplication of the Δ*iglB* mutant was essentially

abolished and comparable to that of the Δ*FPI* mutant. The single amino acid substitution of

Tyr139 by Ala (Y139A) was also sufficient to abolish intracellular bacterial multiplication. A

severe intracellular growth defect was also recorded when Tyr139 was substituted by Phe

(Y139F), until 10 h after infection (almost fifty-fold less bacterial counts). However, at 24 h,

the Y139F mutant strain multiplied and showed a ten-fold reduction of bacterial counts

compared to wild-type, suggesting that a late phagosomal escape had occurred in the Y139F

mutant, therefore promoting some cytosolic bacteria multiplication. As expected, functional

complementation (*i.e.,* introduction of a plasmid-born wild-type *iglB* allele into the Δ*iglB*

mutant strain (CpWT) restored wild-type growth.

We next tested the intracellular behavior of the Δ*iglB* mutant and complemented strains in

primary bone marrow-derived macrophages (BMMs). Growth of the Δ*iglB* deletion mutant as

well as that of the two Y139A or Y139F was identical to that of the Δ*FPI* mutant (**Fig 5B**) at

all time points tested. The fact that no multiplication of Y139F was recorded at 24 h is most

likely due to the higher bactericidal activity of primary macrophages compared to the J774-1

cell line. Functional complementation restored normal intracellular bacterial replication (in

CpWT).

The impact of the mutations in *iglB* on bacterial virulence was evaluated in an *in vivo*

competition assay between wild-type (WT) and Δ*iglB* mutant bacteria. For this, we monitored

the bacterial burden in spleen and livers 2.5 days after infection of 7-week-old female BALB/c

mice by the intra-peritoneal route (**Fig 5C**). The competition index recorded for the *iglB*

mutants (Y139A and Y139F) was very low (≤10^{-6} in both target organs for Y139F; ≤10^{-5} in

livers and ≤10^{-6} in spleens for Y139A), demonstrating the importance of residue Tyr139 in

Francisella virulence in the mouse model. As a control, we also performed an *in vivo*

competition assay between wild-type and the Δ*iglB* complemented strain (WT and CpWT).

The competition index recorded close to 1 for CpWT (≤10^{-1} in both target organs), confirmed

almost complete functional complementation.

To determine whether this lack of intracellular multiplication, associated with reduced

virulence, was due to impaired access to the host cytosol or aborted cytosolic multiplication,

we compared the ability of the Δ*iglB* mutants to escape from the phagosomal compartment to

that of the wild-type strain. For this, we first monitored co-localization of intracellular bacteria

with LAMP-1, in J774-1 macrophages (**Fig 5D**), using specific antibodies and automated

quantification (**Fig 5E**). The frequency of bacteria co-localizing with LAMP-1 at all 3 time

points was elevated (between 60% and 90%), with Δ*iglB* and the two complemented strains

expressing mutated *iglB* alleles (designated Y139A and Y139F; respectively). In contrast, co-

localization of the wild-type strain with LAMP-1 was below 20% after 1 h and remained in the

same range throughout the infection. These results strongly suggest that the Δ*iglB* mutant as

well as the two complemented strains expressing mutated *iglB* alleles, are still trapped in the

phagosomal compartment after 10 h, whereas the wild-type strain has already escaped into

the cytosol after 1 h.

The ability of the Δ*iglB* mutants to induce phagosomal membrane rupture was further

tested by using a CCF4 assay in BMMs, essentially as previously described(42, 46). Briefly,

bacteria were added to the macrophages at multiplicity of infection (MOI) of 100. After 1 h of

infection, cells were loaded for 1 h with CCF4. *F. novicida* naturally secretes β-lactamase

that is able to cleave the cytosolic CCF4 substrate. Once CCF4 is lysed, its emission

spectrum changes from 535 nm (green) to 450 nm (blue). Wild-type *F. novicida* and CpWT

strains showed similar amount of cleaved CCF4 at the three time-points tested (1 h, 3 h and

6 h) whereas there was no (or marginal) loss of FRET signal with Δ*iglB* (**Fig 5F**), Y139A,

Y139F, and in the two negative control strains Δ*FPI* and Δ*bla,* carrying a deletion of the

functional copy of the β-lactamase-encoding gene. These data confirmed that the two *iglB*

mutant strains (Y139A, Y139F) remained confined into intact phagosomes at least up to 6 h

after infection.

Fig 5. IglB mutants show impaired intracellular survival and attenuated virulence

(**A, B**) Intracellular bacterial multiplication of wild-type *F. novicida* (WT, Grey squares), isogenic Δ*iglB* mutant (Δ*iglB*, blue triangles) and complemented Δ*iglB* strain (Cp-WT, grey circles; Y139A, blue square; Y139F, red squares) and the Δ*FPI* control (black lines), was monitored during 24 h in J774A.1 macrophage cells (**A**) and bone marrow-derived macrophages (BMDM) from BALB/c mice (**B**). In J774.1, the values recorded for Y139F, at 24 h, were significantly different from those of Δ*FPI*, Δ*iglB* or Y139A. In both cell types, the values recorded for WT or CpWT, at 10 h and 24 h, were significantly different from those of Δ*FPI*, Δ*iglB,* Y139A or Y139F. **p<0.001 (determined by student's t-test). (**C**) Groups of five BALB/c mice were infected intraperitoneally with 100 CFU of wild-type *F. novicida* and 100 CFU of Δ*iglB* mutant strain or complemented strain (Cp-WT, Y139A, Y139F). Bacterial burden was quantified in liver (L) and spleen (S) of mice. The data represent the competitive index (CI) value (in ordinate) for CFU of mutant/wild-type of each mouse, after 48 h infection, divided by CFU of mutant/wild-type in the inoculum. Bars represent the geometric mean CI value. (**D, E**) Confocal microscopy of intracellular bacteria with J774-1 infected with wild-type *F. novicida* (WT), Δ*iglB* or complemented strains and their co-localization with the phagosomal marker LAMP1 observed at 1 h, 4 h and 10 h. J774.1 were stained for *F. novicida* (green), LAMP-1 (red) and host DNA (blue, DAPI stained) (**D**). Analysis was performed with ImageJ software (**E**). (**F**) CCF4 assay. Infection of $ASC^{-/-}$ BMDM were performed at MOI of 100 and the cytosol of BMDMs was loaded with CCF4, as described previously(46). *F. novicida* naturally expresses a β-lactamase able to cleave the CCF4. Bacterial escape into the host cytosol is thus associated with a shift of the CCF4 probe fluorescence emission from 535 nm to 450 nm. The number of cells demonstrating CCF4 cleavage, determined at different time points (1 h, 3 h and 6 h) by flow cytometry, reached above 50% at 6 h for both WT and CpWT complemented strains whereas it was below 1% for the other mutant strains tested. *p<0.01 (determined by student's t-test).

Cytosolic escape of wild-type *F. novicida* is associated with the release of DNA that is
sensed by the host macrophage and triggers innate immune responses, namely secretion of
type I interferon (IFN), and activation of the AIM2 inflammasome (47-49). In contrast,
mutants that remain trapped in the phagosomal compartment generally fail to induce type I
IFN secretion (42). To check whether in spite of their phagosomal confinement, the *iglB*
mutants could still promote some cytosolic immune response, we quantified Type I IFN
secretion triggered upon infection with the *iglB* mutant strains, in supernatants of BMMs
infected for different time points (1 h, 3 h and 6 h; **Fig 6A**) by an ISRE-luciferase bioassay
(50). IFN secretion reached between 25 and 30 U/mL after 6 h, for both WT and CpWT
complemented strains whereas it remained below 5 U/mL for the other mutant strains tested,
further demonstrating that the *iglB* mutants remained trapped in phagosomes during the time
course of these assays.

Finally, cell death kinetics were determined to monitor possible cytotoxicity in infected
BMMs (measured by real time propidium iodide incorporation). Indeed, host cell death could
be an alternate explanation for the lack of bacterial intracellular multiplication. As expected,
delayed cell death (starting five hours after infection) was observed in BMM infected with
wild-type *F. novicida* (**Fig 6B**) as a result of normal intracellular bacterial multiplication and
activation of the AIM2 inflammasome. The Δ*iglB* complemented strain (CpWT) showed
similar cell death kinetics to the WT strain. In contrast, none of the other strains tested had
any substantial effect on BMM death, demonstrating that the intracellular multiplication
defects of the Y139A, Y139F mutant strains were not due to increased cytotoxicity and
further demonstrating that these mutants are likely to be stuck in intact phagosomes.

Fig 6. Type I IFN secretion and cell death assays

(**A**) Type I IFN secretion in the culture supernatant of BMDMs infected with the indicated
strains at an MOI of 1 was determined by the ISRE-luciferase bioassay and normalized using
rIFNβ. The supernatants of infections (50 µL) collected at different time points (1 h, 3 h and 6

h) were analyzed. Type I IFN secretion reached between 25 and 30 U/mL after 6 h, for both

WT and CpWT complemented strains whereas it remained below 5 U/mL for the other

mutant strains tested. (**B**) Cell death of BMDMs infected with the indicated strains at an MOI

of 1 was monitored in real time by monitoring propidium iodide fluorescence (expressed in

arbitrary units -AU) every 15 min. Mean and SD of triplicates from one experiment

representative of two independent experiments are shown in panels A, B. Unpaired t-tests

were performed, two-tailed P-values were $P \leq 0.001$.

These data demonstrated that residue Y139 of IglB is critical for functional T6SS

assembly, phagosomal escape and bacterial virulence and suggested that only the

unphosphorylated form of the protein was incorporated into sheath structures.

To verify that a residue possibly mimicking the phosphorylated state of Tyr139 would be

deleterious for functional T6SS assembly, we replaced Tyr139 by two phosphomimetics *i.e.*

the negatively charged amino acids Aspartate and Glutamate. We first evaluated the impact

of these amino acid substitutions (Y139D and Y139E) on intracellular multiplication in J774-1

cells (**S4 Fig**). The two mutants led to severe defects in intracellular multiplication and, at all

time points tested, the replication defect was comparable to that of the Y139A or Δ*FPI*

mutant. We next evaluated the capacity of the Y139D and Y139E mutants to form sheath-like

structure by using the sucrose gradient assay. In the two mutants, the IglB protein was

exclusively detected by Western blot in the L fractions of the sucrose gradient (**S4 Fig**),

strongly suggesting that the two amino acid substitutions at Tyr139 also prevented sheath-

like assembly.

Altogether, the data presented support the notion that dephosphorylated Tyr139 is

required for functional sheath assembly. Furthermore, they suggest that the free hydroxyl

group, present on the aromatic ring of the dephosphorylated form of Tyr139 (absent in the

Y139F mutant; **S4 Table**), plays an important role in the assembly of IglA/IglB oligomers.

Discussion

We show here that phosphorylation status of IglB critically influences T6SS functional

assembly, bacterial phagosomal escape and virulence. This post-translational modification of

the sheath could constitute a mechanism controlling the dynamics of T6SS assembly.

Phosphorylation of the sheath, a novel player in control of the T6SS

Cryo-electron microscopy analyses (30) have demonstrated that the FPI of *F. novicida*

encodes a genuine T6SS and proposed a model in which the sheath of the T6SS is

composed of discs of heterodimers of inter-digitated IglA/IglB proteins. The dynamics of

T6SS sheaths have also been studied by live imaging approaches (8, 51 and references

therein). However, the molecular mechanisms associated with T6SS sheath contraction

remain speculative. Interestingly, the recently published structure of the sheath of one of

T6SS of *P. aeruginosa* (TssB1C1 from H1-T6SS) (52) suggests that the sheath shifts from

an extended state to a contracted state using a spring-like motion that might be conserved in

other T6SS.

Formation of the *Francisella* T6SS is believed to occur in the phagosome before cytosolic

escape of the bacterium. Nevertheless, T6SS elements are also likely to be required during

the cytosolic stage of the infectious cycle. Indeed, most FPI genes are induced during late

intra-macrophage growth (53) and IglC has been shown to be required for intracellular

growth of *F. novicida* that are microinjected directly into the cytosol of HeLa cells (54). IglC-

export in *F. novicida* depends on the T6SS core components IglA and IglB (55). Of note,

Clemens *et al*. (30) reported that some split-GFP fusions with truncated IglA proteins,

although still able to form IglA/IglB-heterodimers, had lost their IglC secretory function, which

suggests that heterodimer formation is not sufficient to generate a functional sheath.

Mass spectrometry analyses revealed that IglB contained a phosphorylated residue

(Tyr139). Remarkably, this residue was found in a phosphorylated form exclusively in non-

oligomerized IglB proteins, suggesting that the phosphorylation status of IglB residue Tyr139

could be instrumental in the biogenesis of a functional T6SS. The substitution of Tyr139

either by an alanine or by the non-phosphorylatable aromatic analogue phenylalanine still

allowed the interaction of IglA and IglB in co-immunoprecipitation assays. However, both

substitutions prevented secretion of IglC in the culture supernatant, revealing impaired T6SS

expression in these mutants. Fully supporting the notion that phosphorylation of Tyr139

impairs sheath assembly, replacement of this residue by a phosphomimetic (Y139D or

Y139E) led to a severe intracellular multiplication defect and prevented sheath-like structure

formation.

Our 3D predictions suggest that the OH of tyrosine, that is absent in the aromatic ring of

phenylalanine (**Fig 3 and S4 Table**), might be important for sheath formation. Upon addition

of a phosphoryl moiety at this position, charge repulsion between the phosphate group and

nearby aspartate residues would lead to the interaction between IglA and IglB being

destabilized. Therefore, phosphorylation is unfavorable for the formation of stable hexameric

rings and sheath-like structures. The fact that a late phagosomal escape of the Y139F

mutant apparently occurred in J774-1 macrophages (translated by some bacterial

multiplication at 24 h) tends to suggest that a partially active T6SS could still assemble in the

Y139F mutant.

Possible regulatory mechanisms involved in tyrosine phosphorylation

Protein phosphorylation on tyrosine is catalyzed by autophosphorylation of ATP-dependent

protein-tyrosine kinases (56). Bacterial tyrosine kinases do not share sequence homology

with eukaryotic tyrosine kinases (also called Hanks-type kinases) and, hence, have been

named BY-kinases. To date, the best-characterized function of BY-kinases is related to

capsular polysaccharide synthesis (57). Two main classes of bacterial tyrosine phosphatases

catalyze the reverse reaction: conventional eukaryotic-like phosphatases and low molecular

weight acidic phosphatases. *M. tuberculosis* possesses only one tyrosine kinase (PtkA), but

encodes two tyrosine phosphatases (PtpA and PtpB); the latter being also active against

phospho-Ser and phospho-Thr residues). These two tyrosine phosphatases have been

shown to act on host tyrosine signaling pathways. *S. aureus* also encodes one PtpA

orthologue (designated PtpA, sharing 32% amino acid identity with PtpA of *M. tuberculosis*)

that is phosphorylated by the tyrosine kinase CapA1B2 and secreted by the bacterium (57),

suggesting that, like in *M. tuberculosis*, the *S. aureus* PtpA protein could act as a host cell

signaling molecule(58).

As mentioned earlier, the *Francisella* genomes do not encode a canonical tyrosine kinase.

However, we identified a unique gene encoding a highly conserved protein (97.5% amino

acid identity between the different *F. tularensis* subsps), that is predicted be a low molecular

weight phosphotyrosine protein phosphatase (FTN_1046 in *F. novicida*). This protein,

designated PtpA, is composed of 161 amino acids and shares 40.1% amino acid identity with

its orthologue in *Staphylococcus aureus* (SAAG_02400）, 29.2% with PtpA of *M.

tuberculosis* (TMBG_01746) and less than 28% amino acid identity with the tyrosine

phosphatase of *Escherichia coli* (Wzb and Etp(59); **S5 Fig**).

This prompted us to evaluate the impact of *ptpA* inactivation on *F. novicida* virulence,

using a *ptpA* transposon insertion mutant (from the *F. novicida* Tn insertion library (60). We

evaluated the capacity of the *ptpA* mutant to survive and multiply in J774.1 cells, over a 24 h-

period. We assumed that if PtpA served to dephosphorylate IglB, its inactivation would lead

to a constitutive production of sheath-like structures in this mutant that might affect normal

intracellular bacterial multiplication. The *ptpA* mutant multiplied in these cells like the wild-

type strain (**S5 Fig**). We also evaluated the functionality of the T6SS in this mutant by

monitoring IglC secretion in the supernatant of KCl-induced bacterial cultures and on the

formation of sheath-like structures by sucrose fractionation and western blotting. Confirming

the kinetics of intracellular multiplication, iglC secretion and T6SS assembly were similar in

the *ptpA* mutant to that of the wild-type strain. Altogether, these data strongly indicate that, in

Francisella, the protein PtpA does not play a role in T6SS assembly. At this stage, it cannot

be excluded that this putative phosphatase could play a similar function to its secreted

orthologues in *S. aureus* or *M. tuberculosis* that act on host proteins.

Thanks to the recent improvements in phosphoproteomic approaches, novel families of

protein kinases with a Ser/Thr/Tyr kinase activity have been identified (23). It is likely that

Francisella expresses non-canonical Ser/Thr/Tyr kinase-like activities that still await

authentication and characterisation.

Structural and functional implications of IglB phosphorylation

Tyrosine phosphorylation can regulate protein activity in multiple ways, including causing

electrostatic effects and allosteric transitions, but the most important function of

phosphotyrosine is to serve as a docking site for specific interaction with a target protein that

contains a phosphotyrosine binding domain. The Src homology 2 (SH2) domains represent

the largest class of known phosphotyrosine recognition domains in eukaryotic proteins (61)

and tyrosine phosphorylation serves to recruit SH2-containing targets.

Remarkably, no specific phosphotyrosine binding proteins have been found in bacteria.

Since the phosphate on Tyr is linked to the O^4 position of the phenolic ring, and therefore lies

much further away from the peptide backbone than the phosphate on the β-OH groups of

Ser and Thr, the effects of Tyr phosphorylation in bacteria may be exerted primarily through

allosteric/electrostatic effects. Looking at the full helical reconstruction, one can see that

Y139 is close to both D63 of IglA and D114 of IglB. These are from two different symmetry-

related chains (**Fig 3**). Putting a phosphate on Y139 is likely to repel the aspartate residues

and to disassemble the structure. Although, Aspartates and a Glutamate are not a likely

environment for a Phosphate, there is a quite large pocket which may be able to

accommodate the phosphate.

Since both Y139A and Y139F substitutions in IglB impaired secretion of IglC in the

supernatant of KCl-stimulated bacterial cultures (**Fig 6B**) and prevented normal phagosomal

escape in BMDM macrophages (**Figs 4 and 5**), it is clear that Tyr139 of IglB plays a critical

role in the functional assembly of the T6SS. Moreover, the fact that Y139F substitution also

prevented the formation of a sheath-like structure in KCl-stimulated bacteria, further indicates

that the presence of the non-phosphorylated OH moiety on the C4 of the phenol ring is also

important for proper assembly of the T6SS. The absence of phosphorylated IglB peptide (carrying Y139p) in sheath like structures (**Fig 2C** and **S2 Table**), suggests that phosphorylation of IglB is not favorable for sheath assembly (**Fig 7**). Hence, it is tempting to propose that modulating the phosphorylation status of IglB could represent a simple and rapid mean for *Francisella* to regulate T6SS assembly/disassembly process (**Fig 7**).

Fig 7. Proposed model of phosphorylation-dependent TSS6 assembly-disassembly

(**Left**) Upon bacterial entry into the phagosomal compartment, phosphorylation of IglB may decrease and non-phosphorylated IglB can form stable heterodimers with IglA. Sheath assembly initiates at the bacterial pole. (**Right**) Within 30 min, T6SS sheath contraction triggers the secretion of effectors and the rupture of the phagosomal membrane. Once in the cytosol, the T6SS sheath disassembles, thanks to the concerted action of the ClpB unfoldase and of the phosphorylation of IglB residue Y139.

Upon entry into the host cell phagosome, induced expression of IglB generated by induction of the FPI (dephosphorylated form due to a reduced phosphorylation and/or increased dephosphorylation) would promote T6SS assembly. Once in the cytosol, phosphorylation of IglB (due to a reduced dephosphorylation and/or and increased dephosphorylation) would then favor disassembly the contracted T6SS in concert with the general stress chaperone ClpB (39) and allow active cytosolic bacterial multiplication. In the Tyr139 mutant, bacteria remain trapped in the phagosome because of their inability to promote efficient sheath assembly.

The post-translational phosphorylation of protein is a rapid and reversible mechanism. Hence, yet unidentified phosphorylation-dependent mechanisms controlling the dynamics of T6SS assembly might exist in other pathogens and should be carefully re-examined.

Methods

Ethics Statement. All experimental procedures involving animals were conducted in accordance with guidelines established by the French and European regulations for the care and use of laboratory animals (Decree 87–848, 2001–464, 2001–486 and 2001–131 and European Directive 2010/63/UE) and approved by the INSERM Ethics Committee (Authorization Number: 75-906).

Strains and culture conditions

All strains used in this study are derived from *F. tularensis* subsp. *novicida* U112 (*F. novicida*) as described in **Supplementary Table 5**. Strains were grown at 37°C on pre-made chocolate agar PolyViteX plates (BioMerieux), Schaedler K3 or Chemically Defined Medium (CDM). The CDM used for *F. novicida* corresponds to standard CDM (62) without threonine and valine (63). For growth condition determination, bacterial strains were inoculated in the appropriate medium at an initial OD_{600} of 0.05 from an overnight culture in Schaedler K3 medium.

Construction of a ∆*iglB* deletion mutant

We inactivated the gene *iglB* in *F. novicida* (*FTN_1323*) by allelic replacement resulting in the deletion of the entire gene (4 first codons and 7 last codons were conserved). We constructed a recombinant PCR product containing the upstream region of the gene *iglB* (*iglB*-UP), a kanamycin resistance cassette (*nptII* gene fused with *pGro* promoter) and the downstream region of the gene *iglB* (*iglB*-DN) by overlap PCR. Primers *iglB* upstream FW (p1) and *iglB* upstream (spl_K7) RV (p2) amplified the 689 bp region upstream of position +3 of the *iglB* coding sequence (*iglB*-UP), primers *pGro* FW (p3) and *nptII* RV (p4) amplified the 1091 bp kanamycin resistance cassette (*nptII* gene fused with *pGro* promoter); and primers

iglB downstream (spl_K7) FW (p5) and *iglB* downstream RV (p6) amplified the 622 bp region

downstream of the position +1520 of the *iglB* gene coding sequence (*iglB*-DN). Primers p2

and p5 have an overlapping sequence of 12 nucleotides with primers p3 and p4 respectively

resulting in fusion of *iglB*-UP and *iglB*-DN with the cassette after cross-over PCR. All single

fragment PCR reactions were realized using Phusion High-Fidelity DNA Polymerase

(ThermoScientific) and PCR products were purified using NucleoSpin® Gel and PCR Clean-

up kit (Macherey-Nagel). Overlap PCRs were carried out using 100 ng of each purified PCR

products and the resulting fragment of interest was purified from agarose gel. This fragment

was then directly used to transform wild type *F. novicida* by chemical transformation (64).

Recombinant bacteria were isolated on Chocolate agar plates containing kanamycin (10 µg

mL^{-1}). The mutant strains were checked for loss of the wild type gene by PCR product direct

sequencing (GATC-biotech) using appropriate primers.

Functional complementation

The Δ*iglB* mutant strain was transformed with the recombinant pKK214-derived plasmids

(pKK-*iglBY139A$_{cp}$*, pKK-*iglBY139F$_{cp}$*, pKK-*iglBY139D$_{cp}$* and pKK-*iglBY139E$_{cp}$*) described

below (**Supplementary Table 5**). Primers pGro(65) FW and pGro RV amplified the 328 bp of

the *pGro* promoter and primers *iglB* FW / *iglB*[PstI] RV amplified the 1,534 bp *iglB* gene from

U112. PCR products were purified and SmaI (*pGro* promoter) or PstI (*iglB*) digested in

presence of FastAP Thermosensitive Alkaline Phosphatase (ThermoScientific) to avoid self-

ligation. Mixtures of *pGro* promoter and fragments of the genes of interest were then

incubated with T4 DNA Ligase (New England Biolabs) to allow blunt end ligation and

fragments were then cloned in pKK214 vector after SmaI/PstI double digest and transformed

in *E. coli* TOP10. Recombinant plasmid pKK-*iglB$_{cp}$* was purified and directly used for

chemical transformation in *F. novicida* Δ*iglB* (64). Recombinant colonies were selected on

Chocolate agar plates containing tetracycline (5 µg mL^{-1}) and kanamycin (10 µg mL^{-1}).

For site-directed mutagenesis of *iglB* gene, we used plasmid pKK-*iglB_cp.* The recombinant

plasmids pKK-*iglBY139A_cp*, pKK-*iglBY139F_cp*, pKK-*iglBY139D_cp* and pKK-*iglBY139E_cp* were

constructed using the primer pairs iglB (Y/A, Y/F,Y/D, or Y/E) FW and iglB RV

(**Supplementary Table 6**). The three first bases of primers iglB FW were modified to change

Tyr139 to Ala, Phe, Asp or Glu

PCR products were purified, phosphorylated by the T4 Polynucleotide Kinase and then

incubated with T4 DNA Ligase (New England Biolabs) and transformed in *E. coli* TOP10.

Recombinant plasmids were purified and directly used for chemical transformation in *F.*

novicida Δ*iglB*, as described previously (64). Recombinant colonies were selected on

Chocolate agar plates containing tetracycline (5 μg mL^{-1}) and kanamycin (10 μg mL^{-1}).

Immunodetection

Antibodies to IglA, IglB and IglC were obtained through the NIH Biodefense and Emerging

Infections (BEI) Research Resources Repository, NIAID, NIH.

Immunoblotting analysis. Protein lysates for immunoblotting were prepared by using

Laemmli sample buffer. Protein lysates corresponding to equal OD$_{600}$ were loaded on 12%

Bis/Tris gels (Invitrogen), and run in TGS buffer. Primary antibodies (anti-IglA or anti-IglB)

were used at final dilution of 1:2,000. Secondary horseradish peroxidase (HRP)-conjugated

goat anti-mouse antibody (Santa Cruz Biotechnology, CA, USA) or (HRP)-conjugated goat

anti-rabbit antibody and the enhanced Chemiluminescence system (ECL) (Amersham

Biosciences, Uppsala, Sweden) were used as previously described (66).

Co-immunoprecipitations. Wild-type *F. novicida* was grown to late exponential phase in

Schaedler K3; collected by centrifugation and lysed with lysozyme and 1% TritonX-100

detergent in 20 mM Tris HCl, (pH 7.8) with 1 mM EDTA, protease inhibitor cocktail and

benzonase nuclease. The lysate was centrifuged at 15,000 g for 15 min at 4°C to pellet

bacterial debris. Monoclonal anti-IglB antibody was incubated 40 min with Dynabeads protein

G (Invitrogen) and the complex was incubated at room temperature with the bacterial lysate

for 1 hour. After washing, the complex was loaded on 12% Bis/Tris gels (Invitrogen), and run

in TGS buffer.

Purification of T6SS

T6SS were prepared essentially as described in (30). Briefly, wild-type *F. novicida* was

grown to late exponential phase in Schaedler K3 containing 5% KCl; pelleted by

centrifugation and lysed with lysozyme and 1% TritonX-100 detergent in 20 mM Tris HCl, (pH

7.8) with 1 mM EDTA, protease inhibitor cocktail and benzonase nuclease. The lysate was

centrifuged 3 times at 15,000 g for 30 min at 4°C to pellet bacterial debris, and the

supernatant was layered onto a 10%–55% sucrose gradient overlying a 55% Optiprep

cushion and centrifuged at 100,000 g for 18 hr. Fractions were collected and examined by

TEM negative staining using 2% uranyl acetate. The sheath-like structures sedimented to

just below the 55% sucrose/Optiprep interface.

Phosphoproteomic analyses

Reagents and chemicals. For protein digestion, dithiothreitol, iodoacetamide and

ammonium bicarbonate were purchased from Sigma-Aldrich (St Louis, MO, USA). For

phosphopeptide enrichment and LC-MS/MS analysis, trifluoroacetic acid (TFA), formic acid,

acetonitrile and HPLC-grade water were purchased from Fisher Scientific (Pittsburgh, PA,

USA) at the highest purity grade.

Protein digestion. For proteomic analysis, *F. novicida* was analysed in three independent

biological replicates. Protein concentration was determined by DC assay (Biorad, CA, USA)

according to the manufacturer's instructions. An estimated 1.2 mg of proteins for each

biological replicates were digested following a FASP protocol (67) slightly modified. Briefly,

proteins were reduced using 100 mM dithiothreitol in 50 mM ammonium bicarbonate for 1h at

60°C. Proteins were then split into four samples of 300 µg and applied on 30 kDa MWCO

centrifugal filter units (Microcon, Millipore, Germany, Cat No MRCF0R030). Samples were

mixed with 200 µL of 8M urea in 50 mM ammonium bicarbonate (UA buffer) and centrifuged

for 20 min at 15,000 x g. Filters were washed with 200 µL of UA buffer. Proteins were

alkylated for 30min by incubation in the dark at room temperature with 100 µL of 50 mM

iodoacetamide in UA buffer. Filters were then washed twice with 100 µL of UA buffer (15,000

x g for 15 min) followed by two washes with 100 µL of 50 mM ammonium bicarbonate

(15,000 x g for 10 min). Finally, sequencing grade modified trypsin (Promega, WI, USA) was

added to digest the proteins in 1:50 ratio for 16 h at 37°C. Peptides were collected by

centrifugation at 15,000 x g for 10min followed by one wash with 50mM ammonium

bicarbonate and vacuum dried.

Phosphopeptides enrichment by titanium dioxide (TiO$_2$) and phosphopeptides

purification by graphite carbon (GC). Phosphopeptide enrichment was carried out using a

Titansphere TiO$_2$ Spin tip (3 mg/200 µL, Titansphere PHOS-TiO, GL Sciences Inc, Japan) an

estimated 1.2 mg of digested proteins for each biological replicate. Briefly, the TiO$_2$ Spin tips

were conditioned with 20 µL of solution A (80% acetonitrile, 0,1% TFA), centrifuged at 3,000

x g for 2min and equilibrated with 20µL of solution B (75% acetonitrile, 0,075% TFA, 25%

lactic acid) followed by centrifugation at 3,000 x g for 2 min. Peptides were resuspended in

10 µL of 2% TFA, mixed with 100 µL of solution B and centrifuged at 1,000 x g for 10min.

Sample was applied back to the TiO$_2$ Spin tips two more times in order to increase the

adsorption of the phosphopeptides to the TiO$_2$. Spin tips were washed with, sequentially, 20

µL of solution B and two times with 20 µL of solution A. Phosphopeptides were eluted by the

sequential addition of 50 µL of 5% NH$_4$OH and 50 µL of 5% pyrrolidine. Centrifugation was

carried out at 1,000 x g for 5 min.

Phosphopeptides were further purified using GC Spin tips (GL-Tip, Titansphere, GL Sciences

Inc, Japan). Briefly, the GC Spin tips were conditioned with 20 µL of solution A, centrifuged

at 3,000 x g for 2 min and equilibrated with 20 µL of solution C (0,1% TFA in HPLC-grade

water) followed by centrifugation at 3,000 x g for 2 min. Eluted phosphopeptides from the

TiO$_2$ Spin tips were added to the GC Spin tips and centrifuged at 1,000 x g for 5 min. GC

Spin tips were washed with 20 µL of solution C. Phosphopeptides were eluted with 70 µL of

solution A (1,000 x g for 3 min) and vacuum dried.

nanoLC-MS/MS protein identification and quantification. Samples were resuspended in

12 µL of 0.1% TFA in HPLC-grade water. For each run, 5 µL was injected in a nanoRSLC-Q

Exactive PLUS (RSLC Ultimate 3000, Thermo Scientific, MA, USA). Phosphopeptides were

loaded onto a µ-precolumn (Acclaim PepMap 100 C18, cartridge, 300 µm i.d.×5 mm, 5 µm,

Thermo Scientific, MA, USA) and were separated on a 50 cm reversed-phase liquid

chromatographic column (0.075 mm ID, Acclaim PepMap 100, C18, 2 µm, Thermo Scientific,

MA, USA). Chromatography solvents were (A) 0.1% formic acid in water, and (B) 80%

acetonitrile, 0.08% formic acid. Phosphopeptides were eluted from the column with the

following gradient 5% to 40% B (180 min), 40% to 80% (6 min). At 181 min, the gradient

returned to 5% to re-equilibrate the column for 20 minutes before the next injection. Two

blanks were run between each sample to prevent sample carryover. Phosphopeptides

eluting from the column were analyzed by data dependent MS/MS, using top-8 acquisition

method. Phosphopeptides were fragmented using higher-energy collisional dissociation

(HCD). Briefly, the instrument settings were as follows: resolution was set to 70,000 for MS

scans and 17,500 for the data dependent MS/MS scans in order to increase speed. The MS

AGC target was set to 1×10^6 counts with maximum injection time set to 250 ms, while

MS/MS AGC target was set to 2×10^5 with maximum injection time set to 250 ms. The MS

scan range was from 400 to 1.800 m/z. MS and MS/MS scans were recorded in profile

mode. Dynamic exclusion was set to 30 seconds.

Data Processing Following nanoLC-MS/MS acquisition. The MS files were processed

with the MaxQuant software version 1.5.3.30 and searched with Andromeda search engine

against the UniProtKB/Swiss-Prot *F. tularensis* subsp. *novicida* database (release 28-04-

2014, 1719 entries). To search parent mass and fragment ions, we set an initial mass

deviation of 4.5 ppm and 0.5 Da respectively. The minimum peptide length was set to 7

amino acids and strict specificity for trypsin cleavage was required, allowing up to two missed

cleavage sites. Carbamidomethylation (Cys) was set as fixed modification, whereas oxidation (Met), N-term acetylation and phosphorylation (Ser, Thr, Tyr) were set as variable modifications. The match between runs option was enabled with a match time window of 0.7 min and an alignment time window of 20 min. The false discovery rates (FDRs) at the protein and peptide level were set to 1%. Scores were calculated in MaxQuant as described previously(68). The reverse and common contaminants hits were removed from MaxQuant output.

The phosphopeptides output table and the corresponding logarithmic intensities were used for phosphopeptide analysis. The phosphopeptide table was expanded to separate individual phosphosites, and we kept all sites identified at least once in the three independent replicates of the analysis of *F. novicida*.

All data analysis of mass spectrometry data was performed in Perseus 1.6.0.7. For functional class annotation, we also integrated COG and EggNog databases in Perseus 1.6.0.7, freely available at www.perseus-framework.org. Data are available via ProteomeXchange with identifier PXD009225.

Cell cultures and cell infection experiments

J774A.1 (ATCC® TIB-67™) cells were propagated in Dulbecco's Modified Eagle's Medium (DMEM, PAA), containing 10% fetal bovine serum (FBS, PAA) unless otherwise stated. The day before infection, approximately 2×10^5 eukaryotic cells per well were seeded in 12-well cell tissue plates and bacterial strains were grown overnight in 5 mL of Schaedler K3 at 37°C. Infections were realized at a multiplicity of infection (MOI) of 100 and incubated for 1 h at 37°C in culture medium. After 3 washes with cellular culture medium, plates were incubated for 4, 10 and 24 h in fresh medium supplemented with gentamycin (10 µg mL^{-1}). At each kinetic point, cells were washed 3 times with culture medium and lysed by addition of 1 mL of distilled water for 10 min at 4°C. The titre of viable bacteria was determined by

781 spreading preparations on chocolate plates. Each experiment was conducted at least twice

782 in triplicates.

Confocal experiments

J774.1 macrophage cells were infected (MOI of 1,000) with wild-type *F. novicida* U112, the

isogenic Δ*iglB* mutant, the Δ*iglB* complemented either with wild-type *iglB* (CpWT) Y139A,

Y139F mutants or an isogenic Δ*FPI* strain, in standard DMEM (DMEM-glucose) for 30 min at

37°C. Cells were then washed three times with PBS and maintained in fresh DMEM

supplemented with gentamycin (10 μg mL^{-1}) until the end of the experiment. Three kinetic

points (*i.e.* 1 h, 4 h and 10 h) were sampled. For each time point, cells were washed with 1X

PBS, fixed 15 min with 4% paraformaldheyde, and incubated 10 min in 50 mM NH$_4$Cl in 1X

PBS to quench free aldehydes. Cells were then blocked and permeabilised with PBS

containing 0.1% saponin and 5% goat serum for 10 min at room temperature. Cells were

then incubated for 30 min with anti-*F. novicida* mouse monoclonal antibody (1/500e final

dilution, Creative Diagnostics) and anti-LAMP1 rabbit polyclonal antibody (1/100e, ABCAM).

After washing, cells were incubated for 30 min with Alexa488-conjugated goat anti mouse

and Alexa546 conjugated donkey anti rabbit secondary antibodies (1/400e, AbCam). After

washing, DAPI was added (1/1,000) for 1 min and glass coverslips were mounted in Mowiol

(Cityfluor Ltd.). Cells were examined using an X63 oil-immersion objective on a Zeiss

Apotome 2 microscope. Co-localisation tests were quantified by using Image J software; and

mean numbers were calculated on more than 500 cells for each condition. Confocal

microscopy analyses were performed at the Cell Imaging Facility (Faculté de Médecine

Necker Enfants-Malades).

BMDMs Infections

Infection of WT or *ASC*$^{-/-}$ BMDMs was performed as described previously (42). Briefly,

BMDMs were differentiated in DMEM (Invitrogen) with 10% v/v FCS (Thermo Fisher

Scientific), 15% MCSF (L929 cell supernatant), 10 mM HEPES (Invitrogen), and non-

essential amino acids (Invitrogen). One day before infection, macrophages were seeded into

12- 48- or 96-well plates at a density of 2 x10^5, 1.5 x10^5 or 5 x10^4 cells per well, respectively

and incubated at 37°C, 5% CO_2. The overnight culture of bacteria was added to the

macrophages at multiplicity of infection (MOI) of 100. The plates were centrifuged for 15 min

at 500 g to ensure comparable adhesion of the bacteria to the cells and placed at 37°C for

1h. After 3 washes with PBS, fresh medium with 5 μg mL^{-1} gentamycin (Invitrogen) was

added to kill extracellular bacteria and plates were incubated for the desired time.

Phagosomal rupture assay

Quantification of vacuolar escape using the β-lactamase/CCF4 assay (Life technologies) was

performed as previously described(49). *ASC*$^{-/-}$ BMDMs seeded onto non-treated plates were

infected as described above for 2 h, washed and incubated in CCF4 for 1 h at room

temperature in the presence of 2.5 mM probenicid (Sigma). Propidium iodide negative cells

were considered for the quantification of cells containing cytosolic *F. novicida* using

excitation at 405 nm and detection at 450 nm (cleaved CCF4) or 510 nm (intact CCF4).

Type I IFN secretion

Type I IFN secretion was determined in *ASC*$^{-/-}$ BMDMs by an ISRE-luciferase bioassay(50).

L929 ISRE-luciferase cells were plated at 5x10^4 cells per well in a 96 wells-plate and

incubated 24 h at 37°C with 5% CO_2. 50 μl of supernatants from infected BMDMs were

added for 4 h onto the ISRE-luciferase cells. Luciferase luminescence was detected using

Bright Glo Assay (Promega) following the manufacturer's instructions. Mouse rINF-β was

used as standard to determine the concentration of type I IFN in each sample.

Cell death assays

834 After infection 1h at MOI=100, WT BMMs were washed 3 times and incubated with

835 Propidium Iodide (PI; 5 μg mL^{-1}) in CO_2-independent medium (ThermoFisher Scientific). PI

836 incorporation was determined by measuring the fluorescence emission at 635 nm every 15

837 min on a microplate fluorescent reader (Tecan).

839 **Mouse infection**

840 Wild-type *F. novicida*, the Δ*iglB* mutant and the Δ*iglB* mutant complemented either with wild-

841 type *iglB* (CpWT) or with Y139A and Y139F mutants, were grown in Schaedler K3 to

842 exponential growth phase and diluted to the appropriate concentrations. 6 to 8-week-old

843 female BALB/c mice (Janvier, Le Genest St Isle, France) were intraperitoneally (i.p.)

844 inoculated with 200 μL of bacterial suspension. The actual number of viable bacteria in the

845 inoculum was determined by plating dilutions of the bacterial suspension on chocolate plates.

846 For competitive infections, wild-type *F. novicida* and mutant bacteria were mixed in 1:1 ratio

847 and a total of 100 bacteria were used for infection of each of five mice. After two days, mice

848 were sacrificed. Homogenized spleen and liver tissue from the five mice in one experiment

849 were mixed, diluted and spread on to chocolate agar plates. Kanamycin selection to

850 distinguish wild-type and mutant bacteria was performed. Competitive index (CI) [(mutant

851 output/WT output)/(mutant input/WT input)]. Statistical analysis for CI experiments was as

852 described in (69). Macrophage experiments were analyzed by using the Student's unpaired t

853 test.

855 **Sequence analyses**

856 To retrieve IglB homologous proteins, we used the Hidden Markov Model (HMM) profile

857 "T6SSii_iglB.hmm" from reference (32). We scanned a dataset of 2,462 predicted proteomes

858 of complete bacterial genomes retrieved from GenBank Refseq (last accessed September

859 2016)(70) using *hmmsearch* program of HMMER v.3.1b2 (gathering threshold = 25).

Our dataset included 19 genomes of the *Francisella* genus. All IglB sequences encoded by

Francisella genomes were aligned with MUSCLE v.3.8.31(71). Due to differences in the

annotation of the first methionine codon, we used the sequence of FTN_1323 as a reference

for the multiple sequence alignment.

Redundancy of the sequence set retrieved from *hmmsearch* was reduced using cd-hit-

v4.6.7 (72) with a 90% identity threshold. The longest representative sequence of each

cluster was then aligned with MUSCLE v.3.8.31 (71). Outliers that were very divergent in

sequence length were removed. PROMALS3D and ESPript3.0 (73) servers were used to

visualize multiple sequence and structure alignment using 3J9O.B PDB structure reference

file(30). To determine the conservation of the phosphotyrosine site identified in *Francisella*

genus (Tyr139 or Y139) we focused on the 12 amino acids surrounding the *Francisella*

Tyr139 residue in the 535 aligned sequences.

To determine the conservation of the phosphotyrosine site identified in *Francisella* genus

(Tyr139) we focused on the 12 amino acids surrounding the *Francisella* Y139 residue in the

535 aligned sequences. Out of the 535 aligned sequences, we found 90 sequences with a

tyrosine residue in the vicinity of *Francisella* Tyr139 residue. For illustrative purpose, we

selected 12 representative sequences out of 90 as shown in **Fig. 2D**.

Data Availability

The mass spectrometry proteomics data have been deposited in the ProteomeXchange

Consortium via the PRIDE (74) partner repository with the dataset identifier PXD009225.

Acknowledgements

We thank Dr A. Sjostedt for providing the *Francisella* strain U112 and its *ptpA* transposon insertion derivative (*FTN_1046*) and Alain Schmitt (Cochin Institute Electron Microscopy Facility) for excellent technical support.

References

1.   Sjostedt A. Tularemia: history, epidemiology, pathogen physiology, and clinical manifestations. Ann N Y Acad Sci. 2007;1105:1-29.

2.   Sjostedt A, editor. Francisella tularensis and tularemia: Fontiers Media SA; 2011.

3.   Luque-Larena JJ, Mougeot F, Arroyo B, Vidal MD, Rodriguez-Pastor R, Escudero R, et al. Irruptive mammal host populations shape tularemia epidemiology. PLoS Pathog. 2017;13(11):e1006622.

4.   Maurin M, Gyuranecz M. Tularaemia: clinical aspects in Europe. Lancet Infect Dis. 2016;16(1):113-24.

5.   Santic M, Molmeret M, Klose KE, Abu Kwaik Y. Francisella tularensis travels a novel, twisted road within macrophages. Trends Microbiol. 2006;14(1):37-44.

6.   McLendon MK, Apicella MA, Allen LA. Francisella tularensis: taxonomy, genetics, and Immunopathogenesis of a potential agent of biowarfare. Annu Rev Microbiol. 2006;60:167-85.

7.   Kingry LC, Petersen JM. Comparative review of Francisella tularensis and Francisella novicida. Front Cell Infect Microbiol. 2014;4:35.

8.   Brodmann M, Dreier RF, Broz P, Basler M. Francisella requires dynamic type VI secretion system and ClpB to deliver effectors for phagosomal escape. Nat Commun. 2017;8:15853.

9.   Eshraghi A, Kim J, Walls AC, Ledvina HE, Miller CN, Ramsey KM, et al. Secreted

Effectors Encoded within and outside of the Francisella Pathogenicity Island Promote

Intramacrophage Growth. Cell Host Microbe. 2016;20(5):573-83.

10.   Lagrange B, Benaoudia S, Wallet P, Magnotti F, Provost A, Michal F, et al. Human

caspase-4 detects tetra-acylated LPS and cytosolic Francisella and functions differently from

murine caspase-11. Nat Commun. 2018;9(1):242.

11.   Charity JC, Blalock LT, Costante-Hamm MM, Kasper DL, Dove SL. Small molecule

control of virulence gene expression in Francisella tularensis. PLoS Pathog.

2009;5(10):e1000641.

12.   Charity JC, Costante-Hamm MM, Balon EL, Boyd DH, Rubin EJ, Dove SL. Twin RNA

polymerase-associated proteins control virulence gene expression in Francisella tularensis.

PLoS Pathog. 2007;3(6):e84.

13.   Cuthbert BJ, Ross W, Rohlfing AE, Dove SL, Gourse RL, Brennan RG, et al.

Dissection of the molecular circuitry controlling virulence in Francisella tularensis. Genes

Dev. 2017;31(15):1549-60.

14.   Meibom KL, Forslund AL, Kuoppa K, Alkhuder K, Dubail I, Dupuis M, et al. Hfq, a

novel pleiotropic regulator of virulence-associated genes in Francisella tularensis. Infect

Immun. 2009;77(5):1866-80.

15.   Meibom KL, Barel M, Charbit A. Loops and networks in control of Francisella

tularensis virulence. Future Microbiol. 2009;4:713-29.

16.   Celli J, Zahrt TC. Mechanisms of Francisella tularensis intracellular pathogenesis.

Cold Spring Harb Perspect Med. 2013;3(4):a010314.

17.   Ziveri J, Barel M, Charbit A. Importance of Metabolic Adaptations in Francisella

Pathogenesis. Front Cell Infect Microbiol. 2017;7:96.

18.   Balonova L, Mann BF, Cerveny L, Alley WR, Chovancova E, Forslund AL, et al.

Characterization of protein glycosylation in Francisella tularensis subsp. holarctica;

identification of a novel glycosylated lipoprotein required for virulence. Mol Cell Proteomics.

2012;11(7):M111.015016.

19.   Dankova V, Balonova L, Link M, Straskova A, Sheshko V, Stulik J. Inactivation of

Francisella tularensis Gene Encoding Putative ABC Transporter Has a Pleiotropic Effect

upon Production of Various Glycoconjugates. J Proteome Res. 2016;15(2):510-24.

20.   Barel M, Charbit A. Role of Glycosylation/Deglycolysation Processes in Francisella

tularensis Pathogenesis. Front Cell Infect Microbiol. 2017;7:71.

21.   Canova MJ, Molle V. Bacterial serine/threonine protein kinases in host-pathogen

interactions. J Biol Chem. 2014;289(14):9473-9.

22.   Shi L, Pigeonneau N, Ravikumar V, Dobrinic P, Macek B, Franjevic D, et al. Cross-

phosphorylation of bacterial serine/threonine and tyrosine protein kinases on key regulatory

residues. Front Microbiol. 2014;5:495.

23.   Nguyen HA, El Khoury T, Guiral S, Laaberki MH, Candusso MP, Galisson F, et al.

Expanding the Kinome World: A New Protein Kinase Family Widely Conserved in Bacteria. J

Mol Biol. 2017;429(20):3056-74.

24.   Deutscher J, Ake FM, Derkaoui M, Zebre AC, Cao TN, Bouraoui H, et al. The

bacterial phosphoenolpyruvate:carbohydrate phosphotransferase system: regulation by

protein phosphorylation and phosphorylation-dependent protein-protein interactions.

Microbiol Mol Biol Rev. 2014;78(2):231-56.

25.   Rodionova IA, Zhang Z, Mehla J, Goodacre N, Babu M, Emili A, et al. The

phosphocarrier protein HPr of the bacterial phosphotransferase system globally regulates

energy metabolism by directly interacting with multiple enzymes in Escherichia coli. J Biol

Chem. 2017;292(34):14250-7.

26.   Meibom KL, Charbit A. Francisella tularensis metabolism and its relation to virulence.

Front Microbiol. 2010;1:140.

27.   Dai S, Mohapatra NP, Schlesinger LS, Gunn JS. Regulation of Francisella tularensis

virulence. Frontiers in Microbiology. 2011;2.

28.   Alkhuder K, Meibom KL, Dubail I, Dupuis M, Charbit A. Identification of trkH,

Encoding a Potassium Uptake Protein Required for Francisella tularensis Systemic

Dissemination in Mice. PLoS One. 2010;5(1):e8966.

29. Bell BL, Mohapatra NP, Gunn JS. Regulation of virulence gene transcripts by the

Francisella novicida orphan response regulator PmrA: role of phosphorylation and evidence

of MglA/SspA interaction. Infect Immun. 2010;78(5):2189-98.

30. Clemens DL, Ge P, Lee BY, Horwitz MA, Zhou ZH. Atomic structure of T6SS reveals

interlaced array essential to function. Cell. 2015;160(5):940-51.

31. Russell AB, Wexler AG, Harding BN, Whitney JC, Bohn AJ, Goo YA, et al. A type VI

secretion-related pathway in Bacteroidetes mediates interbacterial antagonism. Cell Host

Microbe. 2014;16(2):227-36.

32. Abby SS, Cury J, Guglielmini J, Neron B, Touchon M, Rocha EP. Identification of

protein secretion systems in bacterial genomes. Sci Rep. 2016;6:23080.

33. Pizarro-Cerda J, Charbit A, Enninga J, Lafont F, Cossart P. Manipulation of host

membranes by the bacterial pathogens Listeria, Francisella, Shigella and Yersinia. Semin

Cell Dev Biol. 2016.

34. Lindgren H, Golovliov I, Baranov V, Ernst RK, Telepnev M, Sjostedt A. Factors

affecting the escape of Francisella tularensis from the phagolysosome. J Med Microbiol.

2004;53(Pt 10):953-8.

35. Nano FE, Zhang N, Cowley SC, Klose KE, Cheung KK, Roberts MJ, et al. A

Francisella tularensis pathogenicity island required for intramacrophage growth. J Bacteriol.

2004;186(19):6430-6.

36. Lauriano CM, Barker JR, Yoon SS, Nano FE, Arulanandam BP, Hassett DJ, et al.

MglA regulates transcription of virulence factors necessary for Francisella tularensis

intraamoebae and intramacrophage survival. Proc Natl Acad Sci U S A. 2004;101(12):4246-

9.

37. Nano FE, Schmerk C. The Francisella pathogenicity island. Ann N Y Acad Sci.

2007;1105:122-37.

38. Broms JE, Sjostedt A, Lavander M. The Role of the Francisella Tularensis

Pathogenicity Island in Type VI Secretion, Intracellular Survival, and Modulation of Host Cell

Signaling. Front Microbiol. 2010;1:136.

39. Meibom KL, Dubail I, Dupuis M, Barel M, Lenco J, Stulik J, et al. The heat-shock

protein ClpB of Francisella tularensis is involved in stress tolerance and is required for

multiplication in target organs of infected mice. Mol Microbiol. 2008;67(6):1384-401.

40. Vettiger A, Basler M. Type VI Secretion System Substrates Are Transferred and

Reused among Sister Cells. Cell. 2016;167(1):99-110 e12.

41. Bröms JE, Sjöstedt A, Lavander M. The Role of the Francisella Tularensis

Pathogenicity Island in Type VI Secretion, Intracellular Survival, and Modulation of Host Cell

Signaling. Front Microbiol. 2010;1:136.

42. Rigard M, Broms JE, Mosnier A, Hologne M, Martin A, Lindgren L, et al. Francisella

tularensis IglG Belongs to a Novel Family of PAAR-Like T6SS Proteins and Harbors a

Unique N-terminal Extension Required for Virulence. PLoS Pathog. 2016;12(9):e1005821.

43. Weiss DS, Brotcke A, Henry T, Margolis JJ, Chan K, Monack DM. In vivo negative

selection screen identifies genes required for Francisella virulence. Proc Natl Acad Sci U S

A. 2007;104(14):6037-42.

44. Broms JE, Lavander M, Sjostedt A. A conserved alpha-helix essential for a type VI

secretion-like system of Francisella tularensis. J Bacteriol. 2009;191(8):2431-46.

45. Spidlova P, Stulik J. Francisella tularensis type VI secretion system comes of age.

Virulence. 2017;8(6):628-31.

46. Ramond E, Gesbert G, Guerrera IC, Chhuon C, Dupuis M, Rigard M, et al.

Importance of host cell arginine uptake in Francisella phagosomal escape and ribosomal

protein amounts. Mol Cell Proteomics. 2015;14(4):870-81.

47. Henry T, Brotcke A, Weiss DS, Thompson LJ, Monack DM. Type I interferon signaling

is required for activation of the inflammasome during Francisella infection. J Exp Med.

2007;204(5):987-94.

48. Jones JW, Kayagaki N, Broz P, Henry T, Newton K, O'Rourke K, et al. Absent in

melanoma 2 is required for innate immune recognition of Francisella tularensis. Proc Natl

Acad Sci U S A. 2010;107(21):9771-6.

49.   Meunier E, Wallet P, Dreier RF, Costanzo S, Anton L, Ruhl S, et al. Guanylate-

binding proteins promote activation of the AIM2 inflammasome during infection with

Francisella novicida. Nat Immunol. 2015;16(5):476-84.

50.   Jiang Z, Georgel P, Du X, Shamel L, Sovath S, Mudd S, et al. CD14 is required for

MyD88-independent LPS signaling. Nat Immunol. 2005;6(6):565-70.

51.   Vettiger A, Winter J, Lin L, Basler M. The type VI secretion system sheath assembles

at the end distal from the membrane anchor. Nat Commun. 2017;8:16088.

52.   Salih O, He S, Planamente S, Stach L, MacDonald JT, Manoli E, et al. Atomic

Structure of Type VI Contractile Sheath from Pseudomonas aeruginosa. Structure.

2018;26(2):329-36 e3.

53.   Wehrly TD, Chong A, Virtaneva K, Sturdevant DE, Child R, Edwards JA, et al.

Intracellular biology and virulence determinants of Francisella tularensis revealed by

transcriptional profiling inside macrophages. Cell Microbiol. 2009;11(7):1128-50.

54.   Meyer L, Broms JE, Liu X, Rottenberg ME, Sjostedt A. Microinjection of Francisella

tularensis and Listeria monocytogenes reveals the importance of bacterial and host factors

for successful replication. Infect Immun. 2015;83(8):3233-42.

55.   Ludu JS, de Bruin OM, Duplantis BN, Schmerk CL, Chou AY, Elkins KL, et al. The

Francisella pathogenicity island protein PdpD is required for full virulence and associates

with homologues of the type VI secretion system. J Bacteriol. 2008;190(13):4584-95.

56.   Grangeasse C, Nessler S, Mijakovic I. Bacterial tyrosine kinases: evolution, biological

function and structural insights. Philos Trans R Soc Lond B Biol Sci. 2012;367(1602):2640-

55.

57.   Brelle S, Baronian G, Huc-Brandt S, Zaki LG, Cohen-Gonsaud M, Bischoff M, et al.

Phosphorylation-mediated regulation of the Staphylococcus aureus secreted tyrosine

phosphatase PtpA. Biochem Biophys Res Commun. 2016;469(3):619-25.

58.   Pereira SF, Goss L, Dworkin J. Eukaryote-like serine/threonine kinases and

phosphatases in bacteria. Microbiol Mol Biol Rev. 2011;75(1):192-212.

59.   Vincent C, Duclos B, Grangeasse C, Vaganay E, Riberty M, Cozzone AJ, et al.
Relationship between exopolysaccharide production and protein-tyrosine phosphorylation in
gram-negative bacteria. J Mol Biol. 2000;304(3):311-21.

60.   Gallagher LA, Ramage E, Jacobs MA, Kaul R, Brittnacher M, Manoil C. A
comprehensive transposon mutant library of Francisella novicida, a bioweapon surrogate.
Proc Natl Acad Sci U S A. 2007;104(3):1009-14.

61.   Hunter T. The genesis of tyrosine phosphorylation. Cold Spring Harb Perspect Biol.
2014;6(5):a020644.

62.   Chamberlain RE. Evaluation of Live Tularemia Vaccine Prepared in a Chemically
Defined Medium. Appl Microbiol. 1965;13:232-5.

63.   Gesbert G, Ramond E, Rigard M, Frapy E, Dupuis M, Dubail I, et al. Asparagine
assimilation is critical for intracellular replication and dissemination of Francisella. Cell
Microbiol. 2014;16(3):434-49.

64.   Gesbert G, Ramond E, Tros F, Dairou J, Frapy E, Barel M, et al. Importance of
branched-chain amino acid utilization in Francisella intracellular adaptation. Infect Immun.
2015;83(1):173-83.

65.   Klionsky DJ, Abdelmohsen K, Abe A, Abedin MJ, Abeliovich H, Acevedo Arozena A,
et al. Guidelines for the use and interpretation of assays for monitoring autophagy (3rd
edition). Autophagy. 2016;12(1):1-222.

66.   Ziveri J, Tros F, Guerrera IC, Chhuon C, Audry M, Dupuis M, et al. The metabolic
enzyme fructose-1,6-bisphosphate aldolase acts as a transcriptional regulator in pathogenic
Francisella. Nat Commun. 2017;8(1):853.

67.   Lipecka J, Chhuon C, Bourderioux M, Bessard MA, van Endert P, Edelman A, et al.
Sensitivity of mass spectrometry analysis depends on the shape of the filtration unit used for
filter aided sample preparation (FASP). Proteomics. 2016;16(13):1852-7.

68.   Cox J, Mann M. MaxQuant enables high peptide identification rates, individualized
p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat Biotechnol.
2008;26(12):1367-72.

69. Brotcke A, Weiss DS, Kim CC, Chain P, Malfatti S, Garcia E, et al. Identification of

MglA-regulated genes reveals novel virulence factors in Francisella tularensis. Infect Immun.

2006;74(12):6642-55.

70. Coordinators NR. Database Resources of the National Center for Biotechnology

Information. Nucleic acids research. 2017;45(D1):D12-D7.

71. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic acids research. 2004;32(5):1792-7.

72. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-

generation sequencing data. Bioinformatics. 2012;28(23):3150-2.

73. Robert X, Gouet P. Deciphering key features in protein structures with the new

ENDscript server. Nucleic acids research. 2014;42(Web Server issue):W320-4.

74. Vizcaino JA, Csordas A, Del-Toro N, Dianes JA, Griss J, Lavidas I, et al. 2016 update

of the PRIDE database and its related tools. Nucleic Acids Res. 2016;44(22):11033.

1089

1090

1091 ## Supporting Information Legends

1092

S1 Fig. Schematic representation of the FPI locus and overview of the T6SS

structure of _Francisella._ The _Francisella_ FPI locus representation (_anmK_ to _pdpA_ gene)

with nomenclature of the canonical T6SS and the _F. novicida_ T6SS. A color was assigned for

each gene function and applied on the overview of the T6SS structure.

Red/Green/Blue/Orange, structural protein; Grey, secreted protein; White, hypothetical

protein

1099

S2 Fig. Alignments of orthologous IglBs proteins. (**A**) Alignments of orthologous

IglBs proteins showing conservation of Y139 in _Francisellae_. The IglB sequences retrieved

from 19 _Francisella_ genomes were aligned using MUSCLE [71] and rendered using ESPript3.0

[73]. Secondary structure elements information is extracted from 3J9O PDB file [32] and is

displayed at the top of the alignment. Sequences are identified by corresponding species

name followed by strain name and locus tag. Species names are abbreviated as follow: Fn,

Francisella tularensis subsp. *novicida*; Ftm, *Francisella tularensis* subsp. *mediasiatica*; Fth,

Francisella tularensis subsp. *holarctica*; Ftt, *Francisella tularensis* subsp. *tularensis*; Ft,

Francisella tularensis; Fno, *Francisella noatunensis*; Fsp, *Francisella* sp. (**B**) Alignments of

orthologous IglBs in other genera showing conservation of a tyrosine residue. Sequences of

11 representative IglB orthologs with a tyrosine residue in the vicinity of Y139 were aligned

using MUSCLE [71] and rendered using ESPript3.0 [73]. Secondary structure elements

information is extracted from 3J9O PDB file [32] and is displayed at the top of the alignment.

Sequences are identified by corresponding Uniprot accession number. Species names are

abbreviated as follow: FRATN, *Francisella tularensis subsp. novicida*; SALTY, *Salmonella*

Typhimurium; YERPE, *Yersinia pestis; BURMA, Burkholderia mallei; XANC5, Xanthomonas*

campestris; RALP1, Ralstonia pickettii; PHOAA, Photorhabdus asymbiotica; ERWAC,

Erwinia amylovora; ENTCC, Enterobacter cloacae; BRAJP, Bradyrhizobium japonicum;

RHIFR, Rhizobium fredii; 9GAMM, Serratia nematodiphila.

1119

1120   **S3 Fig. Effect of IglB mutations on growth in broth.** Wild-type *F. novicida* (WT, Grey

1121   squares), isogenic Δ*iglB* mutant (Δ*iglB*, blue triangles), and complemented *iglB* strain (Cp-

1122   WT*,* grey circles; Y139A, purple square; Y139F, red square), were grown in Schaedler K3

1123   (left panel) or in Tryptic Soy broth (TSB) supplemented with 0,2% cysteine (right panel).

1124

S4 Fig. Impact of substitution of Tyr139 by phosphomimetics (Glu or Asp) on

Francisella **intracellular multiplication and T6SS assembly.** (**A**) J774.1

macrophage-like cells were infected in DMEM with 100 MOI of wild-type *F. novicida* (WT)

and mutated strains (Y139E*,* Y139D*,* Δ*FPI*) for 24 h. Results are shown as the average of

$\log_{10}$ cfu mL^{-1} ± standard deviation. Each experiment was performed in triplicate. **, $p<0.001$

(as determined by two-tailed unpaired Student's t-test). (**B**) WB with anti-IglB of sucrose

gradient fractions.

1132

S5 Fig. Impact of *ptpA* inactivation on *Francisella* intracellular multiplication

and T6SS assembly. (**A**) J774.1 macrophage-like cells were infected in DMEM with 100

MOI of wild-type *F. novicida* (WT), Δ*iglB* mutant, Δ*ptpA* mutant and Δ*FPI* mutant, for 24 h.

Results are shown as the average of log10 cfu mL^{-1} ± standard deviation. Each experiment

was performed in triplicate. **, $p<0.001$ (as determined by two-tailed unpaired Student's t-

test). (**B**) Alignments of orthologous PtpAs proteins from different species (FTN, *Francisella*

novicida; Wtb, *Mycobacterium tuberculosis*; *Sau*, *Staphylococcus aureus*). The PtpAs

sequences were aligned using CLUSTALW. (**C**) WBs on bacteria / culture filtrates. Bacteria

were grown in Schaedler-K3 supplemented with 5% KCl until late log phase and then

harvested by centrifugation. Culture supernatants were collected after filtration on 0.2 µM

Millipore filters and concentrated on Amicon 3 kDa. The equivalent of 200 µg of total protein

were loaded onto each well. Left gel, WB with anti-IglC for the culture pellets; right gel, WB

with anti-IglC for the culture supernatants. (**D**) WB with anti-IglB of sucrose gradient fractions.

1146

S1 Table. Detailed list of phosphosites identified in *F. novicida.* In this table, we

report: i) the average of the log2 (intensity) of the phosphosites identified from three

independent replicates; ii) the amino acid carrying the phosphorylation ("Amino Acid"); iii) its

position on the protein (Position); iv) the "localization probability" of the phosphosite (1 being

100%); v) the number of replicates the phosphosite has been identified in ("identified"); vi)

the "sequence window" of 15 amino acids before and after the amino acid phosphorylated;

vii) the probability and score of identification and localization (PEP, score, score for

localization); viii) the "charge" of the peptide; and ix) the annotation according to COG and

EggNOG.

1156

1157 **S2 Table 2**. **Table of intensities for IglB peptide carrying Y139 phosphorylation.**

1158

1159 **S3 Table. List of proteins identified in the "Heavy" (H) and "Light" (L) Fraction**

1160 **in WT and in the Y139F mutant.** WT_H, WT_L, Y139_H, Y139F_L report the log2

1161 (intensity) of the proteins identified in the fractions. Peptides and sequence coverage

1162 indicates the number of total peptides identified and the primary sequence coverage for that

1163 protein across samples. The Q value and the score refer to the quality of the identification.

1164

1165 **S4 Table. Impact of amino acid substitutions at site 139 of IglB on sheath**

1166 **assembly.**

1167

1168 **S5 Table. Strains and plasmids.**

1169

1170

1171

A — IFNβ Secretion (U/mL) versus Time (hours post-infection) for WT, Δbla, ΔFPI, ΔiglB, CpWT, Y139A, Y139F at 1, 3, 6 hours.

B — PI fluorescence (AU) versus Time (hours post-infection) for WT, CpWT, Y139F, Y139A, ΔiglB, ΔFPI.