A post-translational modification of the sheath modulates *Francisella* type VI secretion system assembly and function

Jason Ziveri[1], Cerina Chhuon[2], Anne Jamet[1], Guénolé Prigent[1], Héloïse Rytter[1], Fabiola Tros[1], Monique Barel[1], Mathieu Coureuil[1], Claire Lays[4], Thomas Henry[4], Nicholas H Keep[3], Ida Chiara Guerrera[2]*, Alain Charbit[1]*

[1] Université Paris Descartes, Sorbonne Paris Cité, Bâtiment Leriche, Paris, INSERM U1151 - CNRS UMR 8253, Institut Necker-Enfants Malades. Team 11: Pathogenesis of Systemic Infections, Paris, France

[2] Plateforme protéomique 3P5-Necker, Université Paris Descartes - Structure Fédérative de Recherche Necker, INSERM US24/CNRS UMS3633, Paris 75014, France

[3] Crystallography, Institute for Structural and Molecular Biology, Department of Biological Sciences Birkbeck, University of London, United Kingdom

[4] CIRI, Centre International de Recherche en Infectiologie, Université Lyon, Inserm, U1111, University Claude Bernard Lyon 1, CNRS, UMR5308, École Normale Supérieure de Lyon, Labex Ecofect, Eco-evolutionary dynamics of infectious diseases, F-69007, LYON, France

Running title: IglB phosphorylation and sheath assembly.

Keywords: Type 6 secretion system, Phosphoproteome, *Francisella tularensis*, IglB.

Bâtiment Leriche. 14 Rue Maria Helena Vieira Da Silva CS 61431 - 75993 PARIS – FRANCE

*Corresponding authors: Ida Chiara Guerrera, Alain Charbit.

e-mail: alain.charbit@inserm.fr; chiara.guerrera@inserm.fr

Tel: 33 1 – 72 60 65 11 — Fax: 33 1 - 72 60 65 13

Abstract

Francisella tularensis is a facultative intracellular pathogen that causes the zoonotic disease tularemia in human and animal hosts. This bacterium possesses a non-canonical type VI secretion systems (T6SS) required for phagosomal escape and access to its replicative niche in the cytosol of infected macrophages. KCl stimulation has been previously used to trigger assembly and secretion of the Francisella T6SS in culture. We found that the amounts of essentially all the TSS6 proteins remained unchanged upon KCl stimulation. We therefore hypothesized that a post-translational modification might be involved in T6SS assembly. A whole cell phosphoproteomic analysis allowed us to identify a unique phosphorylation site on IglB, the TssC homologue and key component of the T6SS sheath. Importantly, the phosphorylated form of IglB was not present in the contracted sheath and 3D modeling indicated that the charge repulsion provoked by addition of a phosphogroup on tyrosine 139 was likely to weaken the stability of the sheath structure. Substitutions of the phosphorylatable residue of IglB (tyrosine 139) with alanine or with phosphomimetics prevented T6SS formation and totally impaired phagosomal escape. In contrast, the substitution with the non-phosphorylatable aromatic analog phenylalanine impaired but did not prevent phagosomal escape and cytosolic bacterial multiplication in J774-1 macrophages. Altogether these data suggest that phosphorylation of the sheath participates to T6SS disassembly. Post-translational modifications of the sheath may represent a previously unrecognized mechanism to finely modulate the dynamics of T6SS assembly-disassembly.

Data are available via ProteomeXchange with identifier PXD012507.

Synopsis

Francisella possesses a non-canonical T6SS that is essential for efficient phagosomal escape and access to the cytosol of infected macrophages. KCl stimulation has been previously used to trigger assembly and secretion of the Francisella T6SS in culture. We found that KCl stimulation did not result in an increased production of TSS6 proteins. We therefore hypothesized that a post-translational modification might be involved in T6SS assembly. Using a global and site-specific phosphoproteomic analysis of *Francisella* we identified a unique phosphorylation site on IglB, the TssC homologue and a key component of the T6SS contractile sheath. We show that this site plays a critical role in T6SS biogenesis and propose that phosphorylation may represent a new mechanism affecting the dynamics of sheath formation.

Introduction

Francisella tularensis is the causative agent of the zoonotic disease tularemia (Luque-Larena, Mougeot et al., 2017, Maurin & Gyuranecz, 2016, Sjostedt, 2007, Sjostedt, 2011). This facultative intracellular pathogen is able to infect a variety of different cell types but, *in vivo*, is thought to replicate and disseminate mainly in phagocytes (Santic, Molmeret et al., 2006). Four major subspecies (subsp) of *F. tularensis* are currently listed that all cause a fulminant disease in mice that is similar to tularemia in human (Kingry & Petersen, 2014, McLendon, Apicella et al., 2006). Although the subsp *novicida* (also called *F. novicida) is* rarely pathogenic in human*,* its genome shares a high degree of nucleotide sequence conservation with the human pathogenic species and is thus widely used as a model organism (Brodmann, Dreier et al., 2017, Eshraghi, Kim et al., 2016, Lagrange, Benaoudia et al., 2018).

The transcriptional and post-transcriptional regulatory processes, controlling *Francisella* phagosomal escape and cytosolic multiplication, have been already well characterized (Celli & Zahrt, 2013, Charity, Blalock et al., 2009, Cuthbert, Ross et al., 2017, Meibom, Barel et al., 2009, Sjostedt, 2011, Ziveri, Barel et al., 2017). Notably, the crucial role of a 30-kb locus in *Francisella* virulence (**Figure supplement 1**), designated FPI for "Francisella pathogenicity island" (Broms, Sjostedt et al., 2010), has been extensively documented (Clemens, Lee et al., 2018). Of note, the FPI is duplicated three subspecies of *F. tularensis* (subsps *holarctica*, *mediasiatica*, and *tularensis*), but is present in a single copy in *F. novicida* and *F. philomiragia*

The FPI encodes a Type VI secretion sytem (T6SS) that is essential to promote bacterial phagosomal escape and access to the cytosolic replication niche (Lauriano, Barker et al., 2004, Meyer, Broms et al., 2015, Nano, Zhang et al., 2004). Whereas the transcriptional regulation of the FPI locus has been deeply characterized (Ramsey, Osborne et al., 2015), the molecular mechanisms triggering T6SS assembly and contraction remain largely

unknown. Bioinformatic analyses have divided bacterial T6SS in three phylogenetically distinct subtypes (designated T6SSi-iii) (Russell, Wexler et al., 2014). Whereas most of the well-characterized T6SS belong to the T6SSi (including in *P. areruginosa* and *V. cholerae*), *Francisella* is currently the only bacterium to possess a T6SSii (Abby, Cury et al., 2016) known to be exclusively involved in the intracellular life cycle of the pathogen. Phagosomal escape occurs as early as 30 minutes post-infection (Pizarro-Cerda, Charbit et al., 2016). The FPI encodes, in addition to the T6SS components, a number of proteins of unknown functions some of which (such as IglF,PdpC or PdpD) have been proposed to be T6SS effectors (Eshraghi et al., 2016). Very recently, Mougous and co-workers elucidated the function of one substrate of the T6SS (Ledvina, Kelly et al., 2018). This effector (named OpiA for outside pathogenicity island protein A) is a phosphatidylinositol (PI) 3-kinase that acts on the Francisella-containing phagosomal membrane to promote bacterial escape into the cytoplasm.

Our current knowledge of the structure and functional assembly of the *Francisella* T6SS are mainly based on recent cryo-electron microscopy data (Clemens, Ge et al., 2015, Nazarov, Schneider et al., 2018) and structural homologies with other members of the T6SSi (Clemens et al., 2015, Nazarov et al., 2018) and references therein). It has been proposed that the tube comprised of IglC subunits (Hcp homologue) that form hexameric rings, fits within the cavity of the IglA/IglB sheath (TssB/C in *Pseudomonas*, VipA/B in *Vibrio*). Contraction of the sheath results in the ejection of the Hcp tube, together with effector proteins located within the tube and on top of the tip of the tube (Vettiger & Basler, 2016). Formation of the contracted *Francisella* T6SS can be mimicked in vitro by high K^+ concentration (Clemens et al., 2015). Yet, the molecular relay for this environmental cue remains uncharacterized.

In order to understand the mechanisms underlying this KCl-induced T6SS production, we performed a global proteomic approach on KCl-induced and non-induced bacteria. This analysis revealed that the relative amounts of most FPI proteins were not increased upon KCl stimulation that suggested the possible implication of a post-translational control of T6SS polymerization. We hypothesized that protein phosphorylation might occur in *Francisella* and

contribute to the dynamics of T6SS assembly-disassembly. By using state-of-the art mass spectrometry analyses, we indeed identified more than one hundred peptides bearing serine/threonine/or tyrosine phosphorylated residues in the proteome of *F. novicida*. One peptide, corresponding to a phosphorylation of IglB on tyrosine 139 (Tyr139), particularly attracted our attention. IglB is analogous to the TssC protein in canonical T6SS and forms with IglA (TssB homologue) the T6SS contractile sheath (Clemens et al., 2015). The data presented here reveal that phosphorylation of IglB is essential for the proper activity of the T6SS and could contribute to the fine control of T6SS dynamics.

Results

We first monitored sheath formation upon KCl stimulation, using the procedure previously described (Clemens et al., 2015). Briefly, lysates of wild-type *F. novicida,* grown in the presence of 5% KCl, were layered onto 15%-55% sucrose gradients (**Figure 1A**). The fraction sedimenting in equilibrium at 55% sucrose and below (55% Optiprep, grey arrow) from KCl-induced cultures were further dialyzed and used for negative staining TEM imaging. As expected, rod-shaped particles were detected by transmission electron microscopy in this fraction that corresponded to contracted sheath-like structures. The different fractions of the gradient were next tested by western blotting with anti-IglA and anti-IglB antibodies. IglA and IglB were detected in all the fraction of the gradient in KCl-induced conditions (**Figure 1B**). In contrast, they were only detected in the fractions of the upper half of the gradient (in the 15% and in part of the 35% sucrose fractions) in the non KCl-induced conditions. We next decided to perform a whole cell proteome analysis of KCl-induced and non-induced bacterial cultures.

Whole cell proteome analyses

Whole cells extracts from KCl-induced and non-induced cultures were analyzed in order to identify proteins whose intensity was altered by the KCl treatment. We could quantify 1,395 proteins that represent 80% of the predicted *Francicella* proteome (**Figure 1C**). Upon KCl stimulation, 113 proteins were up-regulated after KCl stimulation and 97 down-regulated, confirming that KCl stimulation strongly alters the proteome of the cell (**Table supplement 1**). Remarkably, the amounts of most of the FPI-encoded proteins (17 out the 18 proteins encoded by the FPI were identified) did not change significantly upon KCl stimulation, including notably the sheath and tube proteins IglA, IglB, IglC (blue dots in **Figure 1C**).

The down-regulated proteins comprised seven proteins (FTN_0042 to FTN_0048) encoded by the "Francisella novicida island" (or FNI), a genomic island that shows some similarities with the FPI (Rigard, Broms et al., 2016), FTN_0042 and FTN_0043 corresponding to the orthologues of IglA and IglB, respectively). Oppositely, two sets of proteins emerged as up-regulated (**Figure supplement 2A**): proteins involved in Iron-sulfur (Fe/S) clusters biogenesis (FTN_0751 to FTN_0754, FTN_0850 to FTN_0853 and FTN1082) as well as most of the protein encoded by the *fig* (also designated *fsl*) operon (Kiss, Liu et al., 2008, Sullivan, Jeffery et al., 2006), involved in iron acquisition (FTN_1681, FTN_1682, FTN_1684 to FTN_1687) (**Figure supplement 2B**). On the basis of the known regulation of Fe/S clusters in other bacterial species (Mettert & Kiley, 2015, Roche, Aussel et al., 2013), these Fe/S cluster proteins could be regulated by FTN_0810, a predicted transcriptional regulator sharing modest sequence identity with the Fe/S regulator IscR.

The phosphoproteome of *Francisella*

The fact that the amounts of T6SS proteins remained unchanged upon KCl induction, led up to hypothesize that a post-translational mechanism might be responsible for KCl-dependent sheath polymerization. We decided to test if protein phosphorylation could be involved in the process and carried out a global and site-specific phosphoproteomic analysis of *F.*

novicida (strain U112), based on phosphopeptide enrichment and high-resolution LC-MS/MS

analysis of whole cell lysates from KCl-induced and non-induced cultures (**Figure 2**). Overall,

this analysis allowed the identification of 103 phosphopeptides, of which 78 were robustly

quantified. The majority of the phosphosites (47 out of 78) were found on serine residues (S),

while similar lower number of sites, 19 and 12, were found on threonine (T) and tyrosine (Y),

respectively (**Figure 2A, B; Tables supplement 1**, **2**). These phosphosites corresponded

to 59 proteins, of which 7 presented multiple phosphorylation sites. These proteins belonged

to various functional categories and the most represented classes were carbohydrate

metabolism, energy production and conversion and translation.

IgIB was the only protein of the T6SS to be phosphorylated (**Figure 2C**). Furthermore, IgIB

which constitutes with IgIA the T6SS sheath of *Francisella* (Broms et al., 2010, Clemens et

al., 2015, Rigard et al., 2016) bore a unique phosphorylation site on Tyr139 (**Table**

supplement 3). In order to determine if the phosphorylated forms of IgIB was incorporated

into the sheath upon its assembly, we further analyzed the fractions at the bottom of the

gradient (sucrose 55%-Optiprep 55% density zone), containing the sheath-like particles) to

the fractions at the top of the gradient (15-35% sucrose density zone, presumably containing

non-polymerized forms of the IgIA and IgIB proteins (hereafter called "heavy fraction", H or

"light fraction", L, respectively) (**Figure 2D**). The mass spectrometry analysis of the H and L

fractions identified over 1,145 proteins (**Table supplement 4**) and confirmed the efficiency of

the sucrose gradient separation. Unmodified peptides containing Tyr139 were detected in

both H and L fractions, indicating that soluble IgIB include both phosphorylated and non-

phosphorylated forms of the protein. Importantly, the peptide bearing the phosphorylated

tyrosine residue (pY139) of IgIB was only detected in the L fraction of the gradient (**Table**

supplement 4), suggesting that tyrosine phosphorylation might be unfavorable for sheath

polymerization.

Fully supporting this notion, our 3D predictions (**Figure 3**) indicate that, upon addition of a

phosphoryl moiety at this position, charge repulsion between the phosphate group and

nearby aspartate residues should be unfavorable for the formation of stable hexameric rings

and sheath contraction. Since the phosphate on Y139 is linked to the O^4 position of the

phenolic ring, the effects of Tyr phosphorylation may be exerted primarily through

allosteric/electrostatic effects. Looking at the full helical reconstruction, one can see that

Y139 is close to both D63 of IglA and D114 of IglB. These are from two different symmetry-

related chains (**Figure 3**). Putting a phosphate on Y139 is thus likely to repel the aspartate

residues and to disassemble the structure.

Sequence analyses

Comparative sequence analyses on 33 IglB proteins from 19 genomes of the *Francisella*

genus were performed to assess the conservation of residue Tyr139 (**Figure supplement 3**).

Sequences were highly conserved between 31 IglB proteins sharing 93 to 100% amino acid

identity. In contrast, the proteins encoded by *FTN_0043* in the FNI (Rigard et al., 2016) and

F7308_1917 (Fsp_TX077308) exhibited only 48% amino acid identity with the 31 other IglB

proteins whereas they both share 85% identity. Excluding these 2 outliers, among 506 sites,

458 were without polymorphism (91%), with Tyr139 conserved in all IglB sequences (**Figure**

supplement 3, upper part). Since IglB is among the most conserved components of T6SS

(Broms et al., 2010, Clemens et al., 2015), we were able to retrieve orthologues of IglB

encoded outside of *Francisella* genus. Out of 535 non-redundant IglB orthologues (also

named VipB, TssC or EvpB/VC_A0108), we found 90 sequences with a tyrosine in the

vicinity of Tyr139 site outside of *Francisella* genus. In 82 sequences, the predicted location of

the tyrosine residue was similar to that of Tyr139 (in a turn region between two helices)

(**Figure supplement, lower part**).

We hereafter thoroughly evaluated the importance of Y139 phosphorylation on T6SS

functionality and *Francisella* pathogenicity. For this, we constructed two IglB variants in which

Tyr139 was substituted either by an alanine (Ala) or by the non-phosphorylatable aromatic

amino acid analogue of tyrosine, phenylalanine (Phe). These IglB mutated proteins

(designated Y139A and Y139F, respectively) were expressed *in trans* in a Δ*iglB* mutant of *F.*

novicida carrying a chromosomal deletion of the entire *iglB* gene, (see Materials and

Methods). A *F. novicida* mutant with a deletion of the entire *FPI*, unable to escape from

phagosomes and hence to grow in macrophages (Weiss, Brotcke et al., 2007), was used as

a negative control.

A role of IglB phosphorylation in sheath formation

IglB was detected by Western blotting on whole cell lysates in wild-type and complemented

mutated strains (Y139A and Y139F) but not in the Δ*iglB* and Δ*FPI* strains (**Figure 4A**). IglA

was also detected in wild-type and complemented mutated strains (Y139A and Y139F), but

in lower amounts in the Δ*iglB* strain and not in the Δ*FPI* mutant. This observation is in

agreement with earlier observations (Broms, Lavander et al., 2009) that suggested that the

presence of IglB increases the stability of IglA.

A classical measure of basal T6SS function is the export of an Hcp-related protein.

Therefore, we examined IglB-dependent IglC export in the different *F. novicida* mutant

strains by Western-blot on bacteria grown in the presence -or absence- of 5% KCl. In

agreement with previous reports, IglC was detected in the culture supernatant in the

presence, but not in the absence, of KCl, in the wild-type and $iglB_{WT}$ complemented strains.

In contrast, deletion of *iglB* as well as the two single amino acid substitutions Y139A and

Y139F abolished the secretion of IglC into the culture medium (**Figure 4B**). KCl treatment did

not modify the amounts of IglA, IglB and IglC proteins detected by western blotting on whole

cells (bacterial pellet fractions).

To check the impact of the two amino acid substitutions on IglA/IglB heterodimer formation,

we performed co-immunoprecipitation assays (**Figure 4C**). We used either anti-IglA to

precipitate the complex, followed by western blotting with anti-IglB (upper panel) or anti-IglB,

followed by western blotting with anti-IglA (lower panel). In both conditions, the two IglB

mutants were still able to interact with IglA to form IglA/IglB heterodimers.

We next performed sucrose gradient fractionation (as described above), followed by western

blotting with anti-IglB on the Y139F and Y139A mutant strains. The Y139F IglB mutant

protein was detected in both the L and H fractions of the gradient (**Figure 4D**), indicating that

some sheath polymerization still occurred in the Y139F mutant. In contrast, the Y139A

mutant protein was detected in the upper fractions of the gradient but not in the lower

fractions (corresponding to the contracted sheath). Fully supporting the western-blot data,

proteomic analysis of the distribution of IglA and IglB in the Heavy (H) and light (L) sucrose

fraction revealed that their distribution was similar in the strains expressing either WT IglB

and Y139F IglB with, in both case, a slightly higher proportion of IglA and IglB in the L

fraction compared to the H fraction. In sharp contrast, with the Y139A mutant, IglB was

absent from the H fraction and IglA also almost exclusively found in the L fraction (**Figure**

4E).

Altogether, these data comforted the infection assays described above and suggested that

the non-phosphorylatable Y139F mutant could still assemble into not fully functional sheath-

like structures.

Critical role of residue Y139 in *Francisella* pathogenesis

The mutations Y139A and Y139F had no effect on bacterial growth in broth (**Figure**

supplement 4). The ability of wild-type *F. novicida* (WT), Δ*iglB* and complemented strains

(CpWT, Y139A and Y139F) to survive and multiply in murine macrophage-like J774.1 cells

was monitored over a 24 h-period (**Figure 5A**). Confirming earlier reports (Spidlova & Stulik,

2017), intracellular multiplication of the Δ*iglB* mutant was essentially abolished and

comparable to that of the Δ*FPI* mutant. Remarkably, the single amino acid substitution of

Tyr139 by Ala (Y139A) also abolished intracellular bacterial multiplication. In contrast, when

Tyr139 was substituted by Phe (Y139F), a severe intracellular growth defect was recorded

until 10 h after infection (almost fifty-fold less bacterial counts) but at 24 h, the Y139F mutant

strain had multiplied and showed only a ten-fold reduction of bacterial counts compared to

wild-type, suggesting that a late phagosomal escape had occurred therefore promoting cytosolic bacteria multiplication.

As expected, functional complementation (*i.e.,* introduction of a plasmid-born wild-type *iglB* allele into the Δ*iglB* mutant strain (CpWT) restored wild-type growth.

We compared the ability of the Δ*iglB* mutants to escape from the phagosomal compartment to that of the wild-type strain by monitoring co-localization of intracellular bacteria with LAMP-1 in J774-1 macrophages (**Figure 5B**), using specific antibodies and automated quantification. The frequency of bacteria co-localizing with LAMP-1 at all 3 time points was elevated (between 60% and 90%), with Δ*iglB* and the two complemented strains expressing mutated *iglB* alleles (designated Y139A and Y139F; respectively). In contrast, co-localization of the wild-type strain with LAMP-1 was below 20% after 1 h and remained in the same range throughout the infection, suggesting that the Δ*iglB* mutant as well as the two strains expressing mutated *iglB* alleles, are still trapped in the phagosomal compartment after 10 h, whereas the wild-type strain has already escaped into the cytosol after 1 h.

However, supporting the kinetics data, transmission electron microscopy confirmed that a clear cytosolic multiplication was visible at 24 h with the Y139F mutant but not with the Y139A mutant (**Figure 6A**). We therefore next used time-lapse video microscopy to visualize in real time multiplication of the Y139F mutant in J774-1 macrophages. For this, we inserted a GFP-encoding gene into plasmids pKK214::pGro*iglB* and pKK214::pGro*iglB*Y139F (yielding recombinant plasmids pKK214::pGro*iglB*-pGro*gfp* and pKK214::pGro*iglB*Y139F- pGro*gfp,* respectively). *F. novicida* Δ*iglB* was transformed with these GFP-expressing recombinant plasmids. We generated J774-1 cells with red nuclei by transduction with NucLight Red Lentivirus to facilitate cell recognition and counting (see Materials and Methods). J774-1 cells with red nuclei were infected by GFP-expressing bacteria at an MOI of 1,000. Infection was followed over a 24h-period, using a fully automated microscope Incucyte® 531 S3 (Essen BioScience). Images were taken every 20 minutes with the 20X objective during 24 h, starting 1 h after infection (**Figure 6B** snapshots; **Videos supplement 1, 2**). Under these conditions, an active intracellular multiplication was observed with both

strains. In the conditions used, we observed only a minor delay in bacterial multiplication was

observed with the strain complemented with the mutated *iglB* allele Y139F compared to the

strain complemented with the wild-type allele, In contrast to the kinetics and LAMP co-

localization data that suggested a retarded phagosomal escape of the Y139F mutant. Hence,

it is likely that the Y139F substitution allows the production of sufficient assembled T6SS to

promote bacterial phagosomal escape and cytosolic multiplication in these cells.

Finally, we replaced Tyr139 by two phosphomimetics *i.e.* the negatively charged amino acids

Aspartate and Glutamate and evaluated the impact of these amino acid substitutions (Y139D

and Y139E) on intracellular multiplication in J774-1 cells (**Figure supplement 6A**). The two

mutants led to severe defects in intracellular multiplication and, at all time points tested, the

replication defect was comparable to that of the Y139A or Δ*FPI* mutant. We next evaluated

the capacity of the Y139D and Y139E mutants to form sheath-like structure by using the

sucrose gradient assay. In the two mutants, the IglB protein was exclusively detected by

Western blot in the upper fractions of the sucrose gradient (**Figure supplement 6B**). These

data strongly suggest that residues possibly mimicking the phosphorylated state of Y139 also

prevented sheath-like assembly.

Of note, in primary bone marrow-derived macrophages (BMMs), growth of the Δ*iglB* deletion

mutant as well as that of the two Y139A or Y139F was similar to that of a Δ*FPI* mutant

(**Figure supplement 5A**) at all time points tested. Functional complementation restored

normal intracellular bacterial replication (in CpWT). The ability of the Δ*iglB* mutants to induce

phagosomal membrane rupture in BMMs was further tested by using a CCF4 assay,

essentially as previously described (Ramond, Gesbert et al., 2015, Rigard et al., 2016). Wild-

type *F. novicida* and CpWT strains showed similar amount of cleaved CCF4 at the three

time-points tested (1 h, 3 h and 6 h) whereas there was no (or marginal) loss of FRET signal

with Δ*iglB* (**Figure supplement 5B**), Y139A, Y139F, and the two negative control strains

(Δ*FPI* and Δ*bla*), indicating that the Y139A, Y139F mutant strains remained stuck in intact

phagosomes during the time-period tested in BMMS. Hence in primary macrophages, a

tyrosine residue is required for an optimal T6SS assembly likely due to their high bactericidal

activity. Moreover, in the mouse model of tularemia (see Materials and Methods for details),

the two mutants Y139A and Y139F also showed an almost complete loss of virulence (**Fig**

S5C) that confirmed the critical role of tyrosine residue 139 of IglB in Francisella

pathogenicity.

Searching for putative kinases

Our careful inspection of the *Francisella* genomes did not identify any gene encoding for a

putative serine/threonine- or tyrosine-kinase. We therefore searched for proteins containing

potential tyrosine kinase domains. Using the PFAM database (https://pfam.xfam.org/), we

could identify three candidates proteins (FTN_1082, FTN_0459 and FTN_1433). FTN_1082

is predicted to belong to the uncharacterized ACR,YdiU motif/UPF0061 family (PFAM

database PF02696). This protein family notably comprises the SELO proteins (for

selenoproteins) that have been proposed to constitute a novel ubiquitous kinase-like family of

proteins. Recent bioinformatic analyses have identified FTN_1082 as a potential SELO

family-member (Dudkiewicz, Szczepinska et al., 2012). FTN_0459, is predicted to encode a

UbiB homologue, The UbiB protein family is also a widespread family of kinase-like present

in the three kingdoms of life, often required for the biosynthesis of isoprenoid lipids. In

Escherichia coli, UbiB has been shown to be required for the aerobic biosynthesis of the

redox-active lipid ubiquinone, coenzyme Q (Aussel, Pierrel et al., 2014). It has been

assumed but not formally established that UbiB family members were protein kinases.

However, very recently it was demonstrated that a UbiB family member, the mitochondrial

protein ADCK3, adopted a PKL fold and bound adenine nucleotides in a divalent cation-

dependent manner strongly supporting the hypothesis that UbiB proteins are bona fide

kinases (Stefely & Pagliarini, 2017). Finally, FTN_1433, is a protein bearing a G-rich domain

identified on putative tyrosine kinase (PFAM database PF02706). This family notably

includes proteins involved in lipopolysaccharide biosynthesis (Wzz proteins).

We anticipated that if one of these genes was involved in IglB phosphorylation, its inactivation would impair phagosomal escape and hence cytosolic bacterial multiplication. We therefore monitored intracellular bacterial multiplication of the three corresponding transposon insertion mutants of *F. novicida* (Gallagher, Ramage et al., 2007) after 10 hours in J774-1 macrophages (**Figure supplement 7**). The three mutants showed wild-type intracellular multiplication in this assay, suggesting that either these proteins work synergistically or that there exists another yet unidentified enzyme with a moonlight tyrosine kinase activity.

Discussion

We show here that the unique phosphorylated residue of the sheath component IglB plays a critical role in T6SS functional assembly, bacterial phagosomal escape and virulence. This post-translational modification of IglB might contributes to the fine tuning of T6SS disassembly.

Pivotal role of residue 139 in sheath assembly

The assembly of the T6SS-H1 of *Pseudomonas aeruginosa* has been shown to be spatially regulated by a post-translational mechanism coined « the Threonine Protein Phosphorylation » (or TPP) Pathway. During this process, involved in the defense of bacteria to a neighboring attack, a membrane-bound threonine kinase phosphorylates a forkhead-associated domain-containing Protein (FhA or TagH), which promotes the assembly of an active conformation of the T6SS at the site of the attack (Cianfanelli, Monlezun et al., 2016, Mougous, Gifford et al., 2007). More recently, the offensive anti-bacterial T6SS of *Serratia marcescens* has also been shown be controlled by the opposing actions of the TPP pathway and TagF on assembly of the core machinery (Ostrowski, Cianfanelli et al., 2018). The TPP

represents the only currently identified post-translational mechanism regulating T6SS biogenesis.

In bacteria such as *Escherichia coli, Vibrio cholerae* or *Pseudomonas aeruginosa*, the sheath contracts spontaneously during the repeated rounds of ultraspeed purification required for their isolation from cell cultures (Nguyen, Douzi et al., 2018). In *Francisella*, T6SS contraction can be triggered in culture either by placement under coverslips or by KCl stimultation (Clemens et al., 2015). Our mass spectrometry analyses revealed that T6SS proteins were not upregulated upon KCl treatment and that IglB was the only T6SS protein to contain a phosphorylated residue (Tyr139). Importantly, the phosphorylated form of IglB was not associated with the contracted sheath (**Figure 2**) that suggested that the phosphorylation status of IglB residue Tyr139 could participate to T6SS dissociation.

Formation of the *Francisella* T6SS is believed to occur in the phagosome before cytosolic escape of the bacterium and the T6SS seems to be dispensable for cytosolic multiplication. Nevertheless, some T6SS elements are likely to be required during the cytosolic stage of the infectious cycle. Indeed, most FPI genes are induced during late intra-macrophage growth (Wehrly, Chong et al., 2009) and IglC has been shown to be required for intracellular growth of *F. novicida* that are microinjected directly into the cytosol of HeLa cells (Meyer et al., 2015). IglC secretion in *F. novicida* depends on the T6SS core components IglA and IglB (Ludu, de Bruin et al., 2008). Of note, truncated IglA proteins still able to form IglA/IglB-heterodimers have been reported to have lost their IglC secretory function (Clemens et al., 2015) that suggests that heterodimer formation is not sufficient to generate a functional sheath.

The drastic impact of both Y139A and Y139F substitutions on bacterial virulence (**Figure supplement 5**) demonstrated that tyrosine residue 139 of IglB plays a critical role in the functional assembly of the T6SS. Remarkably, the Y139F mutant, which corresponds to the substitution of a tyrosine residue by its non-phosphorylatable aromatic analog phenylalanine, could still assemble unstable sheath-like structure (**Figure 4D, E**) and multiply in J774-1 macrophages (**Figure 5A; Figure 6A, B; video supplement 2**) and. At this stage, the

inability of the Y139F mutant to form stable T6SS might be explained by the fact that it lacks

the OH moiety on the C4 of the phenol ring. Indeed, several reports support the notion that

the OH moiety of tyrosine residues may participate to protein functionality and stability. For

example it has been shown that intramolecular hydrogen may contribute significantly to

conformational protein stability (Bolen & Rose, 2008). Since the OH moiety is capable of

forming such intra- or intermolecular hydrogen bonds with the carboxy oxygen of adjacent

amino acid residues, when a Tyr-Phe substitution is made in a protein, it is likely to impact its

folding or stability (Pace, Horn et al., 2001). More recently, (Baroni, Pandini et al., 2012) also

demontrated that a single tyrosine Hydroxyl Group controled the specificity of *Plasmodium*

falciparum ferredoxin-NADP+ reductase. Although inspection of the 3.7 Å EM reconstruction

of the *F. novicida* contracted sheath (PDB 3j9o) does not show any atoms close enough to

form a hydrogen bond with the hydroxyl of Tyr139, the carboxylates of Asp114 from IglB, and

Asp63 of IglA in the next dimer, are around 5 Å away and these may approach close enough

to Tyr139 OH to form hydrogen bond interactions stabilizing the assembly.

3D modeling also indicated that there is space for a phosphate group on this atom as the OH

is pointing into a pocket in the structure (**Figure 3**). However, the negatively charged

phosphate would induce charge repulsion with the nearby Asps from the next dimer most

likely preventing assembly of the strands when Tyr139 is phosphorylated. Hence, the charge

repulsion provoked by addition of the phosphogroup (when Y139 is phosphorylated) is likely

to weaken the stability of the contracted sheath. Altogether these data suggest that the

phosphorylation status of IglB could modulate T6SS assembly/disassembly process.

Supporting the notion that phosphorylation of Tyr139 impairs sheath contraction,

replacement of this residue by a phosphomimetic (Y139D or Y139E) led to a severe

intracellular multiplication defect and prevented sheath-like structure formation.

IglB phosphorylation, a new partner of T6SS dissociation

Basler and co-workers have recently shown that *F. novicida* assembled its T6SS sheath on

the bacterial poles (Brodmann et al., 2017) and that the sheath cycled through assembly,

contraction and disassembly, similarly to what was previously described for other canonical

T6SSs. The FPI does not encode any homologue of the unfoldase ClpV, present in all the

canonical T6SS, but instead uses the general chaperone ClpB for the recycling of the

contracted sheaths (Brodmann et al., 2017). ClpB specifically localizes with the contracted

sheaths and is responsible for their disassembly within phagosomes of infected

macrophages. We found that the amounts of ClpB were not (or marginally) up-regulated

upon KCl induction (*i.e.* only 1.3-fold more ClpB in KCl-induced conditions compared to non-

induced conditions). ClpB was also mainly associated with the soluble fraction of the sucrose

gradient of KCl-induced bacteria (**Table supplement 1**). The kinase activity, leading to IglB

phosphorylation might assemble, like ClpB, at the pole of the bacterium only when the

sheath is contracted and assist sheath dissociation.

In spite of our repeated attempts, we were unable to identify a candidate tyrosine kinase.

This suggests that a yet unidentified enzyme with either a non-canonical tyrosine kinase

activity or a protein predicted to be involved in a different enzymatic reaction with a

« moonlight » tyrosine kinase activity, is responsible for IglB phosphorylation. If so, its

identification might prove extremely difficult and would require extensive future work. It is

also possible that a single kinase mutant might have no or very limited effect on phagosomal

disruption since sheath polymerization would occur and be sufficient to promote phagosomal

membrane disruption. Indeed, it has been shown (Meyer et al., 2015) that only a minor

fraction of functional T6SS is sufficient to promote phagosomal escape and, once in the

cytosol, FPI mutants can multiply like wild-type bacteria.

Our whole cell proteomic analysis revealed that the amounts of more than two hundred

proteins were altered upon KCl stimulation (**Figure 1**). Notably, the down-regulated proteins

comprized seven proteins (FTN_0042 - FTN_0048) out of the 13 proteins potentially

encoded by the *Francisella novicida* island (FNI, FTN_0042 - FTN_0054) (Rigard et al.,

2016). Although inactivation of the whole FNI locus had no effect on *F. novivcida* virulence

(Rigard et al., 2016), a possible contribution of the orthologues of IglA and IglB (FTN_0042

and FTN_0043) to the sheath assembly-disassembly process cannot be excluded. The

proteins involved in Iron-sulfur (Fe/S) clusters biogenesis or in iron acquisition, that were

upregulated upon KCL stimulation might also contribute to T6SS biogenesis.

In conclusion, we have identified in the present work a critical residue of IglB whose

phosphorylation may represent a new player involved in the fine-tuning of the T6SS

dynamics of *Francisella*. Such post-translational sheath modifications might exist in other

bacteria that need to be explored.

Materials and Methods

Ethics Statement. All experimental procedures involving animals were conducted in accordance with guidelines established by the French and European regulations for the care and use of laboratory animals (Decree 87–848, 2001–464, 2001–486 and 2001–131 and European Directive 2010/63/UE) and approved by the INSERM Ethics Committee (Authorization Number: 75-906).

Strains and culture conditions

All strains used in this study are derived from *F. tularensis* subsp. *novicida* U112 (*F. novicida*) as described in **Table supplement 5**. Strains were grown at 37°C on pre-made chocolate agar PolyViteX plates (BioMerieux), Schaedler K3 or Chemically Defined Medium (CDM). The CDM used for *F. novicida* corresponds to standard CDM (Chamberlain, 1965) without threonine and valine (Gesbert, Ramond et al., 2014). For growth condition determination, bacterial strains were inoculated in the appropriate medium at an initial OD_{600} of 0.05 from an overnight culture in Schaedler K3 medium.

Construction of a Δ*iglB* deletion mutant

We inactivated the gene *iglB* in *F. novicida* (*FTN_1323*) by allelic replacement, resulting in the deletion of the entire gene (4 first codons and 7 last codons were conserved). We constructed a recombinant PCR product containing the upstream region of the gene *iglB* (*iglB*-UP), a kanamycin resistance cassette (*nptII* gene fused with *pGro* promoter) and the downstream region of the gene *iglB* (*iglB*-DN) by overlap PCR. Primers *iglB* upstream FW (p1) and *iglB* upstream (spl_K7) RV (p2) amplified the 689 bp region upstream of position +3 of the *iglB* coding sequence (*iglB*-UP), primers *pGro* FW (p3) and *nptII* RV (p4) amplified the 1091 bp kanamycin resistance cassette (*nptII* gene fused with *pGro* promoter); and primers *iglB* downstream (spl_K7) FW (p5) and *iglB* downstream RV (p6) amplified the 622 bp region

downstream of the position +1520 of the *iglB* gene coding sequence (*iglB*-DN). Primers p2

and p5 have an overlapping sequence of 12 nucleotides with primers p3 and p4 respectively

resulting in fusion of *iglB*-UP and *iglB*-DN with the cassette after cross-over PCR (**Table S6**).

All single fragment PCR reactions were realized using Phusion High-Fidelity DNA

Polymerase (ThermoScientific) and PCR products were purified using NucleoSpin® Gel and

PCR Clean-up kit (Macherey-Nagel). Overlap PCRs were carried out using 100 ng of each

purified PCR products and the resulting fragment of interest was purified from agarose gel.

This fragment was then directly used to transform wild type *F. novicida* by chemical

transformation (Gesbert, Ramond et al., 2015). Recombinant bacteria were isolated on

Chocolate agar plates containing kanamycin (10 µg mL^{-1}). The mutant strains were checked

for loss of the wild type gene by PCR product direct sequencing (GATC-biotech) using

appropriate primers.

Functional complementation

The Δ*iglB* mutant strain was transformed with the recombinant pKK214-derived plasmids

(pKK-*iglBY139A$_{cp}$*, pKK-*iglBY139F$_{cp}$*, pKK-*iglBY139D$_{cp}$* and pKK-*iglBY139E$_{cp}$*) described

below (**Table supplement 5**). Primers pGroFW and pGro RV amplified the 328 bp of the

pGro promoter; primers *iglB* FW/ and *iglB*[PstI] RV amplified the 1,534 bp *iglB* gene from

U112. PCR products were purified and SmaI (*pGro* promoter) or PstI (*iglB*) digested in

presence of FastAP Thermosensitive Alkaline Phosphatase (ThermoScientific) to avoid self-

ligation. Mixtures of *pGro* promoter and fragments of the genes of interest were then

incubated with T4 DNA Ligase (New England Biolabs) to allow blunt end ligation and

fragments were then cloned in pKK214 vector after SmaI/PstI double digest and transformed

in *E. coli* TOP10. Recombinant plasmid pKK-*iglB$_{cp}$* was purified and directly used for

chemical transformation in *F. novicida* Δ*iglB* (Gesbert et al., 2015). Recombinant colonies

were selected on Chocolate agar plates containing tetracycline (5 µg mL^{-1}) and kanamycin

(10 µg mL^{-1}).

For site-directed mutagenesis of *iglB* gene, we used plasmid pKK-*iglB$_{cp.}$* The recombinant

plasmids pKK-*iglBY139A$_{cp}$*, pKK-*iglBY139F$_{cp}$*, pKK-*iglBY139D$_{cp}$* and pKK-*iglBY139E$_{cp}$* were

constructed using the primer pairs iglB (Y/A, Y/F,Y/D, or Y/E) FW and iglB RV (**Table**

supplement 6). The three first bases of primers iglB FW were modified to change Tyr139 to

Ala, Phe, Asp or Glu

PCR products were purified, phosphorylated by the T4 Polynucleotide Kinase and then

incubated with T4 DNA Ligase (New England Biolabs) and transformed in *E. coli* TOP10.

Recombinant plasmids were purified and directly used for chemical transformation in *F.*

novicida ∆iglB, as described previously (Gesbert et al., 2015). Recombinant colonies were

selected on Chocolate agar plates containing tetracycline (5 µg mL^{-1}) and kanamycin (10 µg

mL^{-1}).

Gfp-expressing *F. novicida*

The *∆iglB* mutant strain was transformed with the recombinant plasmids pKK-GFP-*iglBWT$_{cp}$*

and pKK-GFP-*iglBY139F$_{cp,}$* derived from pKK214-GFP (Abd, Johansson et al., 2003).

Primers pGro (KlionskyAbdelmohsen et al.)FW and *iglB*(Klionsky et al.) RV (**Table**

supplement 5) amplified the 1,870 bp of the *pGro-iglB* from the pKK-*iglBWT$_{cp}$* and pKK-

iglBY139F$_{cp}$. PCR products were purified and digested with SmaI and the *pGro-iglB WT* and

pGro-iglB Y139F fragments were then incubated with T4 DNA Ligase (New England Biolabs)

to allow blunt end ligation and fragments were then cloned in the unique SmaI site of

pKK214-(GFP) vector and transformed in *E. coli* TOP10. Recombinant plasmid pKK-(GFP)-

iglBWT$_{cp}$ and pKK-(GFP)-*iglBY139F$_{cp}$* were purified and directly used for chemical

transformation in *F. novicida ∆iglB* (Gesbert et al., 2015). Recombinant colonies were

selected on Chocolate agar plates containing tetracycline (5 µg mL^{-1}) and kanamycin (10 µg

mL^{-1}).

Immunodetection

Antibodies to IglA, IglB and IglC were obtained through the NIH Biodefense and Emerging Infections (BEI) Research Resources Repository, NIAID, NIH.

Immunoblotting analysis. Protein lysates for immunoblotting were prepared by using Laemmli sample buffer. Protein lysates corresponding to equal OD_{600} were loaded on 12% Bis/Tris gels (Invitrogen), and run in TGS buffer. Primary antibodies (anti-IglA or anti-IglB) were used at final dilution of 1:2,000. Secondary horseradish peroxidase (HRP)-conjugated goat anti-mouse antibody (Santa Cruz Biotechnology, CA, USA) or (HRP)-conjugated goat anti-rabbit antibody and the enhanced Chemiluminescence system (ECL) (Amersham Biosciences, Uppsala, Sweden) were used as previously described (Ziveri, Tros et al., 2017).

Co-immunoprecipitations. Wild-type *F. novicida* was grown to late exponential phase in Schaedler K3; collected by centrifugation and lysed with lysozyme and 1% TritonX-100 detergent in 20 mM Tris HCl, (pH 7.8) with 1 mM EDTA, protease inhibitor cocktail and benzonase nuclease. The lysate was centrifuged at 15,000 g for 15 min at 4°C to pellet bacterial debris. Monoclonal anti-IglB antibody was incubated 40 min with Dynabeads protein G (Invitrogen) and the complex was incubated at room temperature with the bacterial lysate for 1 hour. After washing, the complex was loaded on 12% Bis/Tris gels (Invitrogen), and run in TGS buffer.

T6SS purification

T6SS were prepared essentially as described in (Clemens et al., 2015). Briefly, wild-type *F. novicida* was grown to late exponential phase in Schaedler K3 containing 5% KCl; pelleted by centrifugation and lysed with lysozyme and 1% TritonX-100 detergent in 20 mM Tris HCl, (pH 7.8) with 1 mM EDTA, protease inhibitor cocktail and benzonase nuclease. The lysate was centrifuged 3 times at 15,000 g for 30 min at 4°C to pellet bacterial debris, and the supernatant was layered onto a 10%–55% sucrose gradient overlying a 55% Optiprep cushion and centrifuged at 100,000 g for 18 hr. Fractions were collected and examined by TEM negative staining using 2% uranyl acetate. The sheath-like structures sedimented to

just below the 55% sucrose/Optiprep interface.

Phosphoproteomic analyses

Reagents and chemicals. For protein digestion, dithiothreitol, iodoacetamide and

ammonium bicarbonate were purchased from Sigma-Aldrich (St Louis, MO, USA). For

phosphopeptide enrichment and LC-MS/MS analysis, trifluoroacetic acid (TFA), formic acid,

acetonitrile and HPLC-grade water were purchased from Fisher Scientific (Pittsburgh, PA,

USA) at the highest purity grade.

Protein digestion. For proteomic analysis, *F. novicida* was analysed in three independent

biological replicates. Protein concentration was determined by DC assay (Biorad, CA, USA)

according to the manufacturer's instructions. An estimated 1.2 mg of proteins for each

biological replicates were digested following a FASP protocol (Lipecka, Chhuon et al., 2016)

slightly modified. Briefly, proteins were reduced using 100 mM dithiothreitol in 50 mM

ammonium bicarbonate for 1h at 60°C. Proteins were then split into four samples of 300 µg

and applied on 30 kDa MWCO centrifugal filter units (Microcon, Millipore, Germany, Cat No

MRCF0R030). Samples were mixed with 200 µL of 8M urea in 50 mM ammonium

bicarbonate (UA buffer) and centrifuged for 20 min at 15,000 x g. Filters were washed with

200 µL of UA buffer. Proteins were alkylated for 30min by incubation in the dark at room

temperature with 100 µL of 50 mM iodoacetamide in UA buffer. Filters were then washed

twice with 100 µL of UA buffer (15,000 x g for 15 min) followed by two washes with 100 µL of

50 mM ammonium bicarbonate (15,000 x g for 10 min). Finally, sequencing grade modified

trypsin (Promega, WI, USA) was added to digest the proteins in 1:50 ratio for 16 h at 37°C.

Peptides were collected by centrifugation at 15,000 x g for 10min followed by one wash with

50mM ammonium bicarbonate and vacuum dried.

Phosphopeptides enrichment by titanium dioxide (TiO$_2$) and phosphopeptides

purification by graphite carbon (GC). Phosphopeptide enrichment was carried out using a

Titansphere TiO$_2$ Spin tip (3 mg/200 µL, Titansphere PHOS-TiO, GL Sciences Inc, Japan) an

estimated 1.2 mg of digested proteins for each biological replicate. Briefly, the TiO$_2$ Spin tips

were conditioned with 20 µL of solution A (80% acetonitrile, 0,1% TFA), centrifuged at 3,000

x g for 2min and equilibrated with 20µL of solution B (75% acetonitrile, 0,075% TFA, 25%

lactic acid) followed by centrifugation at 3,000 x g for 2 min. Peptides were resuspended in

10 µL of 2% TFA, mixed with 100 µL of solution B and centrifuged at 1,000 x g for 10min.

Sample was applied back to the TiO_2 Spin tips two more times in order to increase the

adsorption of the phosphopeptides to the TiO_2. Spin tips were washed with, sequentially, 20

µL of solution B and two times with 20 µL of solution A. Phosphopeptides were eluted by the

sequential addition of 50 µL of 5% NH_4OH and 50 µL of 5% pyrrolidine. Centrifugation was

carried out at 1,000 x g for 5 min.

Phosphopeptides were further purified using GC Spin tips (GL-Tip, Titansphere, GL Sciences

Inc, Japan). Briefly, the GC Spin tips were conditioned with 20 µL of solution A, centrifuged

at 3,000 x g for 2 min and equilibrated with 20 µL of solution C (0,1% TFA in HPLC-grade

water) followed by centrifugation at 3,000 x g for 2 min. Eluted phosphopeptides from the

TiO_2 Spin tips were added to the GC Spin tips and centrifuged at 1,000 x g for 5 min. GC

Spin tips were washed with 20 µL of solution C. Phosphopeptides were eluted with 70 µL of

solution A (1,000 x g for 3 min) and vacuum dried.

nanoLC-MS/MS protein identification and quantification. Samples were resuspended in

12 µL of 0.1% TFA in HPLC-grade water. For each run, 5 µL was injected in a nanoRSLC-Q

Exactive PLUS (RSLC Ultimate 3000, Thermo Scientific, MA, USA). Phosphopeptides were

loaded onto a µ-precolumn (Acclaim PepMap 100 C18, cartridge, 300 µm i.d.×5 mm, 5 µm,

Thermo Scientific, MA, USA) and were separated on a 50 cm reversed-phase liquid

chromatographic column (0.075 mm ID, Acclaim PepMap 100, C18, 2 µm, Thermo Scientific,

MA, USA). Chromatography solvents were (A) 0.1% formic acid in water, and (B) 80%

acetonitrile, 0.08% formic acid. Phosphopeptides were eluted from the column with the

following gradient 5% to 40% B (180 min), 40% to 80% (6 min). At 181 min, the gradient

returned to 5% to re-equilibrate the column for 20 minutes before the next injection. Two

blanks were run between each sample to prevent sample carryover. Phosphopeptides

eluting from the column were analyzed by data dependent MS/MS, using top-8 acquisition

method. Phosphopeptides were fragmented using higher-energy collisional dissociation

(HCD). Briefly, the instrument settings were as follows: resolution was set to 70,000 for MS

scans and 17,500 for the data dependent MS/MS scans in order to increase speed. The MS

AGC target was set to 1×10^6 counts with maximum injection time set to 250 ms, while

MS/MS AGC target was set to 2×10^5 with maximum injection time set to 250 ms. The MS

scan range was from 400 to 1.800 m/z. MS and MS/MS scans were recorded in profile

mode. Dynamic exclusion was set to 30 seconds.

Data Processing Following nanoLC-MS/MS acquisition. The MS files were processed

with the MaxQuant software version 1.5.3.30 and searched with Andromeda search engine

against the UniProtKB/Swiss-Prot *F. tularensis* subsp. *novicida* database (release 28-04-

2014, 1719 entries). To search parent mass and fragment ions, we set an initial mass

deviation of 4.5 ppm and 0.5 Da respectively. The minimum peptide length was set to 7

amino acids and strict specificity for trypsin cleavage was required, allowing up to two missed

cleavage sites. Carbamidomethylation (Cys) was set as fixed modification, whereas oxidation

(Met), N-term acetylation and phosphorylation (Ser, Thr, Tyr) were set as variable

modifications. The match between runs option was enabled with a match time window of 0.7

min and an alignment time window of 20 min. The false discovery rates (FDRs) at the protein

and peptide level were set to 1%. Scores were calculated in MaxQuant as described

previously(Cox & Mann, 2008). The reverse and common contaminants hits were removed

from MaxQuant output.

The phosphopeptides output table and the corresponding logarithmic intensities were used

for phosphopeptide analysis. The phosphopeptide table was expanded to separate individual

phosphosites, and we kept all sites identified at least once in the three independent

replicates of the analysis of *F. novicida*.

All data analysis of mass spectrometry data was performed in Perseus 1.6.0.7. For functional

class annotation, we also integrated COG and EggNog databases in Perseus 1.6.0.7, freely

available at www.perseus-framework.org. Data are available via ProteomeXchange with
identifier PXD009225.

Cell cultures and cell infection experiments

J774A.1 (ATCC® TIB-67™) cells were propagated in Dulbecco's Modified Eagle's Medium
(DMEM, PAA), containing 10% fetal bovine serum (FBS, PAA) unless otherwise stated. The
day before infection, approximately $2x10^5$ eukaryotic cells per well were seeded in 12-well
cell tissue plates and bacterial strains were grown overnight in 5 mL of Schaedler K3 at
37°C. Infections were realized at a multiplicity of infection (MOI) of 100 and incubated for 1 h
at 37°C in culture medium. After 3 washes with cellular culture medium, plates were
incubated for 4, 10 and 24 h in fresh medium supplemented with gentamycin (10 µg mL^{-1}). At
each kinetic point, cells were washed 3 times with culture medium and lysed by addition of 1
mL of distilled water for 10 min at 4°C. The titre of viable bacteria was determined by
spreading preparations on chocolate plates. Each experiment was conducted at least twice
in triplicates.

Time lapse microscopy

J774-1 cells were transduced with the IncuCyte® NucLight Red Lentivirus, following the
manufacturer's recommendations, to obtain red nuclear labelling of living cells due to the
expression of the red fluorescent protein mKate2 containing a nuclear localization signal.
One prior to infection, J774-1 cells were seeded in 12-well cell tissue plates. Cells were
infected (MOI of 1,000) with wild-type and mutant GFP-expressing cell. Synchronization of
bacterial entry was realized by a 5 minutes centrifugation at 1,000 rpm. Plates were then
incubated for 1 h at 37°C in culture medium. After 3 washes with cellular culture medium,
plates were incubated at 5% CO_2 and 37°C for 24 hours in fresh medium supplemented with
gentamycin (10 µg mL^{-1}). Bacterial multiplication was monitored in the fully automated
microscope Incucyte® S3 (Essen BioScience). Images were taken every 20 minutes with the

20X objective. Analysis and time-lapse videos (from which images were extracted) were

generated by using Incucyte® S3 software.

Confocal experiments

J774.1 macrophage cells were infected (MOI of 1,000) with wild-type *F. novicida* U112, the

isogenic Δ*iglB* mutant, the Δ*iglB* complemented either with wild-type *iglB* (CpWT) Y139A,

Y139F mutants or an isogenic Δ*FPI* strain, in standard DMEM (DMEM-glucose) for 30 min at

37°C. Cells were then washed three times with PBS and maintained in fresh DMEM

supplemented with gentamycin (10 μg mL^{-1}) until the end of the experiment. Three kinetic

points (*i.e.* 1 h, 4 h and 10 h) were sampled. For each time point, cells were washed with 1X

PBS, fixed 15 min with 4% paraformaldheyde, and incubated 10 min in 50 mM NH$_4$Cl in 1X

PBS to quench free aldehydes. Cells were then blocked and permeabilised with PBS

containing 0.1% saponin and 5% goat serum for 10 min at room temperature. Cells were

then incubated for 30 min with anti-*F. novicida* mouse monoclonal antibody (1/500e final

dilution, Creative Diagnostics) and anti-LAMP1 rabbit polyclonal antibody (1/100e, ABCAM).

After washing, cells were incubated for 30 min with Alexa488-conjugated goat anti mouse

and Alexa546 conjugated donkey anti rabbit secondary antibodies (1/400e, AbCam). After

washing, DAPI was added (1/1,000) for 1 min and glass coverslips were mounted in Mowiol

(Cityfluor Ltd.). Cells were examined using an X63 oil-immersion objective on a Zeiss

Apotome 2 microscope. Co-localisation tests were quantified by using Image J software; and

mean numbers were calculated on more than 500 cells for each condition. Confocal

microscopy analyses were performed at the Cell Imaging Facility (Faculté de Médecine

Necker Enfants-Malades).

BMDMs Infections

733 Infection of WT or *ASC*$^{-/-}$ BMDMs was performed as described previously (Rigard et al.,

734 2016). Briefly, BMDMs were differentiated in DMEM (Invitrogen) with 10% v/v FCS (Thermo

Fisher Scientific), 15% MCSF (L929 cell supernatant), 10 mM HEPES (Invitrogen), and non-essential amino acids (Invitrogen). One day before infection, macrophages were seeded into 12- 48- or 96-well plates at a density of 2 x10^5, 1.5 x10^5 or 5 x10^4 cells per well, respectively and incubated at 37°C, 5% CO_2. The overnight culture of bacteria was added to the macrophages at multiplicity of infection (MOI) of 100. The plates were centrifuged for 15 min at 500 g to ensure comparable adhesion of the bacteria to the cells and placed at 37°C for 1h. After 3 washes with PBS, fresh medium with 5 µg mL^{-1} gentamycin (Invitrogen) was added to kill extracellular bacteria and plates were incubated for the desired time.

Phagosomal rupture assay

Quantification of vacuolar escape using the β-lactamase/CCF4 assay (Life technologies) was performed as previously described(Meunier, Wallet et al.). *ASC$^{-/-}$* BMDMs seeded onto non-treated plates were infected as described above for 2 h, washed and incubated in CCF4 for 1 h at room temperature in the presence of 2.5 mM probenicid (Sigma). Propidium iodide negative cells were considered for the quantification of cells containing cytosolic *F. novicida* using excitation at 405 nm and detection at 450 nm (cleaved CCF4) or 510 nm (intact CCF4).

Mouse infection

We evaluated the impact of the mutations Y139A and Y139F on bacterial virulence in the mouse model of tularemia, using an *in vivo* competition assay (Gesbert et al., 2014). Wild-type *F. novicida*, the Δ*iglB* mutant and the Δ*iglB* mutant complemented either with wild-type *iglB* (CpWT) or with Y139A and Y139F mutants, were grown in Schaedler K3 to exponential growth phase and diluted to the appropriate concentrations. 6 to 8-week-old female BALB/c mice (Janvier, Le Genest St Isle, France) were intraperitoneally (i.p.) inoculated with 200 µL of bacterial suspension. The actual number of viable bacteria in the inoculum was determined by plating dilutions of the bacterial suspension on chocolate plates. For competitive infections, wild-type *F. novicida* and mutant bacteria were mixed in 1:1 ratio and a total of 100 bacteria were used for infection of each of five mice. After two days, mice were

sacrificed. Homogenized spleen and liver tissue from the five mice in one experiment were

mixed, diluted and spread on to chocolate agar plates. Kanamycin selection to distinguish

wild-type and mutant bacteria was performed. The competitive index (CI) was determined as

the ratio (mutant output/WT output) / (mutant input/WT input). Statistical analysis for CI

experiments was as described in (Brotcke, Weiss et al., 2006). Macrophage experiments

were analyzed by using the Student's unpaired t test.

Sequence analyses

To retrieve IglB homologous proteins, we used the Hidden Markov Model (HMM) profile

"T6SSii_iglB.hmm" from reference (Abby et al., 2016). We scanned a dataset of 2,462

predicted proteomes of complete bacterial genomes retrieved from GenBank Refseq (last

accessed September 2016) using *hmmsearch* program of HMMER v.3.1b2 (gathering

threshold = 25).

Our dataset included 19 genomes of the *Francisella* genus. All IglB sequences encoded by

Francisella genomes were aligned with MUSCLE v.3.8.31 (Edgar, 2004). Due to differences

in the annotation of the first methionine codon, we used the sequence of FTN_1323 as a

reference for the multiple sequence alignment.

Redundancy of the sequence set retrieved from *hmmsearch* was reduced using cd-hit-

v4.6.7 (Fu, Niu et al., 2012) with a 90% identity threshold. The longest representative

sequence of each cluster was then aligned with MUSCLE v.3.8.31 (Edgar, 2004). Outliers

that were very divergent in sequence length were removed. PROMALS3D and ESPript3.0

(Robert & Gouet, 2014) servers were used to visualize multiple sequence and structure

alignment using 3J9O.B PDB structure reference file (Clemens et al., 2015). To determine

the conservation of the phosphotyrosine site identified in *Francisella* genus (Tyr139 or Y139)

we focused on the 12 amino acids surrounding the *Francisella* Tyr139 residue in the 535

aligned sequences.

To determine the conservation of the phosphotyrosine site identified in *Francisella* genus

(Tyr139) we focused on the 12 amino acids surrouding the *Francisella* Y139 residue in the

535 aligned sequences. Out of the 535 aligned sequences, we found 90 sequences with a tyrosine residue in the vicinity of *Francisella* Tyr139 residue. For illustrative purpose, we selected 12 representative sequences out of 90 as shown in **Fig. S3**.

Data Availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (Perez-Riverol, Xu et al., 2016) partner repository with the dataset identifier PXD012507.

Acknowledgements

We thank Dr A. Sjostedt for providing the *Francisella* strain U112 and its *ptpA* transposon insertion derivative (*FTN_1046*) and Alain Schmitt (Cochin Institute Electron Microscopy Facility) for excellent technical support.

Funding

These studies were supported by INSERM, CNRS and Université Paris Descartes Paris Cité Sorbonne. Jason Ziveri was funded by a fellowship from the "Délégation Générale à l'Armement". Claire Lays was funded by a fellowship from the LABEX ECOFECT (ANR-11-LABX-0048) of Université de Lyon, within the program "Investissement d'Avenir" (ANR-11-IDEX-0007) operated by the French National Research Agency (ANR).

Authors contribution

J.Z. performed most of the in vitro experiments; F.T. performed murine in vivo experiments and some immunofluorescence experiments; I.C.G and C.C. performed the proteomic analyses and I.C.G analyzed and compiled the data; N. K. performed the 3D analysis; A.J. performed the sequence alignments and analyzed the data ; H.R and M.C. performed the time lapse microscopy analyses; G.P. performed some in vitro experiments; M.B. and T.H. analyzed the data ; A.C. designed the experiments and J.Z. and A.C. analyzed the data, and wrote the paper.

Conflict of interest

The authors declare that they have no conflict of interest.

References

Abby SS, Cury J, Guglielmini J, Neron B, Touchon M, Rocha EP (2016) Identification of protein secretion systems in bacterial genomes. Sci Rep 6: 23080

Abd H, Johansson T, Golovliov I, Sandstrom G, Forsman M (2003) Survival and growth of Francisella tularensis in Acanthamoeba castellanii. Appl Environ Microbiol 69: 600-6

Aussel L, Pierrel F, Loiseau L, Lombard M, Fontecave M, Barras F (2014) Biosynthesis and physiology of coenzyme Q in bacteria. Biochim Biophys Acta 1837: 1004-11

Baroni S, Pandini V, Vanoni MA, Aliverti A (2012) A single tyrosine hydroxyl group almost entirely controls the NADPH specificity of Plasmodium falciparum ferredoxin-NADP+ reductase. Biochemistry 51: 3819-26

Bolen DW, Rose GD (2008) Structure and energetics of the hydrogen-bonded backbone in

protein folding. Annu Rev Biochem 77: 339-62

Brodmann M, Dreier RF, Broz P, Basler M (2017) Francisella requires dynamic type VI

secretion system and ClpB to deliver effectors for phagosomal escape. Nat Commun 8:

15853

Broms JE, Lavander M, Sjostedt A (2009) A conserved alpha-helix essential for a type VI

secretion-like system of Francisella tularensis. J Bacteriol 191: 2431-46

Broms JE, Sjostedt A, Lavander M (2010) The Role of the Francisella Tularensis

Pathogenicity Island in Type VI Secretion, Intracellular Survival, and Modulation of Host Cell

Signaling. Front Microbiol 1: 136

Brotcke A, Weiss DS, Kim CC, Chain P, Malfatti S, Garcia E, Monack DM (2006)

Identification of MglA-regulated genes reveals novel virulence factors in Francisella

tularensis. Infect Immun 74: 6642-55

Celli J, Zahrt TC (2013) Mechanisms of Francisella tularensis intracellular pathogenesis.

Cold Spring Harb Perspect Med 3: a010314

Chamberlain RE (1965) Evaluation of Live Tularemia Vaccine Prepared in a Chemically

Defined Medium. Appl Microbiol 13: 232-5

Charity JC, Blalock LT, Costante-Hamm MM, Kasper DL, Dove SL (2009) Small molecule

control of virulence gene expression in Francisella tularensis. PLoS Pathog 5: e1000641

Cianfanelli FR, Monlezun L, Coulthurst SJ (2016) Aim, Load, Fire: The Type VI Secretion

System, a Bacterial Nanoweapon. Trends Microbiol 24: 51-62

Clemens DL, Ge P, Lee BY, Horwitz MA, Zhou ZH (2015) Atomic structure of T6SS reveals

interlaced array essential to function. Cell 160: 940-951

Clemens DL, Lee BY, Horwitz MA (2018) The Francisella Type VI Secretion System. Front

Cell Infect Microbiol 8: 121

Cox J, Mann M (2008) MaxQuant enables high peptide identification rates, individualized

p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat Biotechnol 26:

1367-72

Cuthbert BJ, Ross W, Rohlfing AE, Dove SL, Gourse RL, Brennan RG, Schumacher MA

(2017) Dissection of the molecular circuitry controlling virulence in Francisella tularensis.

Genes Dev 31: 1549-1560

Dudkiewicz M, Szczepinska T, Grynberg M, Pawlowski K (2012) A novel protein kinase-like

domain in a selenoprotein, widespread in the tree of life. PLoS One 7: e32138

Edgar RC (2004) MUSCLE: a multiple sequence alignment method with reduced time and

space complexity. BMC Bioinformatics 5: 113

Eshraghi A, Kim J, Walls AC, Ledvina HE, Miller CN, Ramsey KM, Whitney JC, Radey MC,

Peterson SB, Ruhland BR, Tran BQ, Goo YA, Goodlett DR, Dove SL, Celli J, Veesler D,

Mougous JD (2016) Secreted Effectors Encoded within and outside of the Francisella

Pathogenicity Island Promote Intramacrophage Growth. Cell Host Microbe 20: 573-583

Fu L, Niu B, Zhu Z, Wu S, Li W (2012) CD-HIT: accelerated for clustering the next-generation

sequencing data. Bioinformatics 28: 3150-2

Gallagher LA, Ramage E, Jacobs MA, Kaul R, Brittnacher M, Manoil C (2007) A

comprehensive transposon mutant library of Francisella novicida, a bioweapon surrogate.

Proc Natl Acad Sci U S A 104: 1009-14

Gesbert G, Ramond E, Rigard M, Frapy E, Dupuis M, Dubail I, Barel M, Henry T, Meibom K,

Charbit A (2014) Asparagine assimilation is critical for intracellular replication and

dissemination of Francisella. Cell Microbiol 16: 434-49

Gesbert G, Ramond E, Tros F, Dairou J, Frapy E, Barel M, Charbit A (2015) Importance of

branched-chain amino acid utilization in Francisella intracellular adaptation. Infect Immun 83:

173-83

Kingry LC, Petersen JM (2014) Comparative review of Francisella tularensis and Francisella

novicida. Front Cell Infect Microbiol 4: 35

Kiss K, Liu W, Huntley JF, Norgard MV, Hansen EJ (2008) Characterization of fig operon

mutants of Francisella novicida U112. FEMS Microbiol Lett 285: 270-7

Klionsky DJ, Abdelmohsen K, Abe A, Abedin MJ, Abeliovich H, Acevedo Arozena A, Adachi

H, Adams CM, Adams PD, Adeli K, Adhihetty PJ, Adler SG, Agam G, Agarwal R, Aghi MK,

Agnello M, Agostinis P, Aguilar PV, Aguirre-Ghiso J, Airoldi EM et al. (2016) Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). Autophagy 12: 1-222

Lagrange B, Benaoudia S, Wallet P, Magnotti F, Provost A, Michal F, Martin A, Di Lorenzo F, Py BF, Molinaro A, Henry T (2018) Human caspase-4 detects tetra-acylated LPS and cytosolic Francisella and functions differently from murine caspase-11. Nat Commun 9: 242

Lauriano CM, Barker JR, Yoon SS, Nano FE, Arulanandam BP, Hassett DJ, Klose KE (2004) MglA regulates transcription of virulence factors necessary for Francisella tularensis intraamoebae and intramacrophage survival. Proc Natl Acad Sci U S A 101: 4246-9

Ledvina HE, Kelly KA, Eshraghi A, Plemel RL, Peterson SB, Lee B, Steele S, Adler M, Kawula TH, Merz AJ, Skerrett SJ, Celli J, Mougous JD (2018) A Phosphatidylinositol 3-Kinase Effector Alters Phagosomal Maturation to Promote Intracellular Growth of Francisella. Cell Host Microbe 24: 285-295 e8

Lipecka J, Chhuon C, Bourderioux M, Bessard MA, van Endert P, Edelman A, Guerrera IC (2016) Sensitivity of mass spectrometry analysis depends on the shape of the filtration unit used for filter aided sample preparation (FASP). Proteomics 16: 1852-7

Ludu JS, de Bruin OM, Duplantis BN, Schmerk CL, Chou AY, Elkins KL, Nano FE (2008) The Francisella pathogenicity island protein PdpD is required for full virulence and associates with homologues of the type VI secretion system. J Bacteriol 190: 4584-95

Luque-Larena JJ, Mougeot F, Arroyo B, Vidal MD, Rodriguez-Pastor R, Escudero R, Anda P, Lambin X (2017) Irruptive mammal host populations shape tularemia epidemiology. PLoS Pathog 13: e1006622

Maurin M, Gyuranecz M (2016) Tularaemia: clinical aspects in Europe. Lancet Infect Dis 16: 113-124

McLendon MK, Apicella MA, Allen LA (2006) Francisella tularensis: taxonomy, genetics, and Immunopathogenesis of a potential agent of biowarfare. Annu Rev Microbiol 60: 167-85

Meibom KL, Barel M, Charbit A (2009) Loops and networks in control of Francisella tularensis virulence. Future Microbiol 4: 713-29

Mettert EL, Kiley PJ (2015) How Is Fe-S Cluster Formation Regulated? Annu Rev Microbiol

69: 505-26

Meunier E, Wallet P, Dreier RF, Costanzo S, Anton L, Ruhl S, Dussurgey S, Dick MS,

Kistner A, Rigard M, Degrandi D, Pfeffer K, Yamamoto M, Henry T, Broz P (2015)

Guanylate-binding proteins promote activation of the AIM2 inflammasome during infection

with Francisella novicida. Nat Immunol 16: 476-484

Meyer L, Broms JE, Liu X, Rottenberg ME, Sjostedt A (2015) Microinjection of Francisella

tularensis and Listeria monocytogenes reveals the importance of bacterial and host factors

for successful replication. Infect Immun 83: 3233-42

Mougous JD, Gifford CA, Ramsdell TL, Mekalanos JJ (2007) Threonine phosphorylation

post-translationally regulates protein secretion in Pseudomonas aeruginosa. Nat Cell Biol 9:

797-803

Nano FE, Zhang N, Cowley SC, Klose KE, Cheung KK, Roberts MJ, Ludu JS, Letendre GW,

Meierovics AI, Stephens G, Elkins KL (2004) A Francisella tularensis pathogenicity island

required for intramacrophage growth. J Bacteriol 186: 6430-6

Nazarov S, Schneider JP, Brackmann M, Goldie KN, Stahlberg H, Basler M (2018) Cryo-EM

reconstruction of Type VI secretion system baseplate and sheath distal end. EMBO J 37

Nguyen VS, Douzi B, Durand E, Roussel A, Cascales E, Cambillau C (2018) Towards a

complete structural deciphering of Type VI secretion system. Curr Opin Struct Biol 49: 77-84

Ostrowski A, Cianfanelli FR, Porter M, Mariano G, Peltier J, Wong JJ, Swedlow JR, Trost M,

Coulthurst SJ (2018) Killing with proficiency: Integrated post-translational regulation of an

offensive Type VI secretion system. PLoS Pathog 14: e1007230

Pace CN, Horn G, Hebert EJ, Bechert J, Shaw K, Urbanikova L, Scholtz JM, Sevcik J (2001)

Tyrosine hydrogen bonds make a large contribution to protein stability. J Mol Biol 312: 393-

404

Perez-Riverol Y, Xu QW, Wang R, Uszkoreit J, Griss J, Sanchez A, Reisinger F, Csordas A,

Ternent T, Del-Toro N, Dianes JA, Eisenacher M, Hermjakob H, Vizcaino JA (2016) PRIDE

Inspector Toolsuite: Moving Toward a Universal Visualization Tool for Proteomics Data

Standard Formats and Quality Assessment of ProteomeXchange Datasets. Mol Cell

Proteomics 15: 305-17

Pizarro-Cerda J, Charbit A, Enninga J, Lafont F, Cossart P (2016) Manipulation of host

membranes by the bacterial pathogens Listeria, Francisella, Shigella and Yersinia. Semin

Cell Dev Biol

Ramond E, Gesbert G, Guerrera IC, Chhuon C, Dupuis M, Rigard M, Henry T, Barel M,

Charbit A (2015) Importance of host cell arginine uptake in Francisella phagosomal escape

and ribosomal protein amounts. Mol Cell Proteomics 14: 870-81

Ramsey KM, Osborne ML, Vvedenskaya IO, Su C, Nickels BE, Dove SL (2015) Ubiquitous

promoter-localization of essential virulence regulators in Francisella tularensis. PLoS Pathog

11: e1004793

Rigard M, Broms JE, Mosnier A, Hologne M, Martin A, Lindgren L, Punginelli C, Lays C,

Walker O, Charbit A, Telouk P, Conlan W, Terradot L, Sjostedt A, Henry T (2016) Francisella

tularensis IglG Belongs to a Novel Family of PAAR-Like T6SS Proteins and Harbors a

Unique N-terminal Extension Required for Virulence. PLoS Pathog 12: e1005821

Robert X, Gouet P (2014) Deciphering key features in protein structures with the new

ENDscript server. Nucleic Acids Res 42: W320-4

Roche B, Aussel L, Ezraty B, Mandin P, Py B, Barras F (2013) Iron/sulfur proteins biogenesis

in prokaryotes: formation, regulation and diversity. Biochim Biophys Acta 1827: 455-69

Russell AB, Wexler AG, Harding BN, Whitney JC, Bohn AJ, Goo YA, Tran BQ, Barry NA,

Zheng H, Peterson SB, Chou S, Gonen T, Goodlett DR, Goodman AL, Mougous JD (2014) A

type VI secretion-related pathway in Bacteroidetes mediates interbacterial antagonism. Cell

Host Microbe 16: 227-236

Santic M, Molmeret M, Klose KE, Abu Kwaik Y (2006) Francisella tularensis travels a novel,

twisted road within macrophages. Trends Microbiol 14: 37-44

Sjostedt A (2007) Tularemia: history, epidemiology, pathogen physiology, and clinical

manifestations. Ann N Y Acad Sci 1105: 1-29

Sjostedt A (2011) Francisella tularensis and tularemia. In Fontiers Research Topic Ebooks,

Fontiers Media SA

Spidlova P, Stulik J (2017) Francisella tularensis type VI secretion system comes of age.

Virulence 8: 628-631

Stefely JA, Pagliarini DJ (2017) Biochemistry of Mitochondrial Coenzyme Q Biosynthesis.

Trends Biochem Sci 42: 824-843

Sullivan JT, Jeffery EF, Shannon JD, Ramakrishnan G (2006) Characterization of the

siderophore of Francisella tularensis and role of fslA in siderophore production. J Bacteriol

188: 3785-95

Vettiger A, Basler M (2016) Type VI Secretion System Substrates Are Transferred and

Reused among Sister Cells. Cell 167: 99-110 e12

Wehrly TD, Chong A, Virtaneva K, Sturdevant DE, Child R, Edwards JA, Brouwer D, Nair V,

Fischer ER, Wicke L, Curda AJ, Kupko JJ, 3rd, Martens C, Crane DD, Bosio CM, Porcella

SF, Celli J (2009) Intracellular biology and virulence determinants of Francisella tularensis

revealed by transcriptional profiling inside macrophages. Cell Microbiol 11: 1128-50

Weiss DS, Brotcke A, Henry T, Margolis JJ, Chan K, Monack DM (2007) In vivo negative

selection screen identifies genes required for Francisella virulence. Proc Natl Acad Sci U S A

104: 6037-6042

Ziveri J, Barel M, Charbit A (2017) Importance of Metabolic Adaptations in Francisella

Pathogenesis. Front Cell Infect Microbiol 7: 96

Ziveri J, Tros F, Guerrera IC, Chhuon C, Audry M, Dupuis M, Barel M, Korniotis S, Fillatreau

S, Gales L, Cahoreau E, Charbit A (2017) The metabolic enzyme fructose-1,6-bisphosphate

aldolase acts as a transcriptional regulator in pathogenic Francisella. Nat Commun 8: 853

1006

1007

1008

Figure legends

Figure 1. KCl stimulation of sheath formation

(**A**) Sucrose gradient. Lysates of bacteria grown in Schaedler K3 medium in the presence of KCl were laid on top of a discontinuous sucrose gradient 15%-55%/Optiprep 55%. Left panel, composition of the gradient; Right panel, transmission electron microscopy (TEM) of the "heavy" fraction (grey arrow, topping the Optiprep cushion). The assembled sheath-like structure, sedimenting at equilibrium to below 55% sucrose, showed rod-shaped particles of variable length (100 - 600 nm). (**B**) Western blotting analysis of the different fractions, using anti-Igl (A, B or C) antibodies. (**C**) The proteome of KCl-stimulated and non-stimulated *F. novicida.* Volcano plot representing the statistical comparison of the protein LFQ intensities of KCl- stimulation vs non-stimulated cells. Inner volcano was established using S0=0.5, FDR=0.01 (class B proteins) and the outer volcano using S0=0.5, FDR=0.001 (class A proteins). Proteins belonging to the FPI, the Fe uptale and Fe-S clusters are highlighted in color as indicated.

Figure 2. The phosphoproteome of *F. novicida*

(A) Distribution of phosphosites according to the modified amino acid (tyrosine,Y; threonine, T and serine, S). (B) Histograms of the most represented classes of protein. The values correspond to the number of proteins bearing phosphosites in each category. (C) Heat map of phosphoproteins altered upon KCl stimulation (Pink to right of the FTN numbers corresponds to proteins with multiple phosphorylation sites). (D) IglB phosphorylation distribution. Intensities (in log2) of the non phosphorylated (left) and phosphorylated (right) Tyr139-containing peptide in the "heavy" (H) and "light" (L) fractions. The signal of the phosphopeptide was only detectable after phosphoenrichment. nd, not detectable.

Figure 3. Predicted role of Y139 of IglB in sheath assembly

(**A**) Overview of the sheath that consists of 12 helical strands based on EM reconstruction at 3.7 Å. Each unit consists of a IglA/IglB dimer. The same colors were used in the further pictures. The chains near Tyr139 of IglB consist of 2 IglA/IglB dimers from the same strand (pink/grey and coral/sky blue). The N-terminal tail of an IglA (blue) from an adjacent strand wraps round to close to Y139 of this strand. (**B**) A closer view of just the selected chains. (**C**) Close up of the region of Tyr139. The hydroxyl is pointing into a pocket and not obviously making a Hydrogen bond. However, the carboxylates of Asp114 from IglB and Asp63 of IglA in the next dimer are around 5 Å away. At the resolution of the reconstruction side chain density is not well defined and these could in fact be close enough to H bond. (**D**) Surface of the assembly clipped to show Tyr139. There is enough space for a Phospho group on the Tyr. The two Asps will however be close to the phosphogroup and so there is likely to be charge repulsion impairing the sheath assembly or weakening the sheath stability when Y139 is phosphorylated.

Figure 4. Immunodetection of IglA, IglB and IglC in wild-type and IglB mutant strains

(**A**) Western Blots (WB) of whole cell protein extracts. Upper gel: WB with anti-IglB; lower gel: WB with anti-IglA. (**B**) Western Blots (WB) with anti-IglC on bacteria / culture filtrates. Bacteria were grown in Schaedler-K3 supplemented or not with 5% KCl until late log phase and then harvested by centrifugation. Culture supernatants were collected after filtration on 0.2 µM Millipore filters and concentrated using Amicon 3 kDa membranes. The equivalent of 200 µg of total protein were loaded onto each well. Upper gel, bacterial cultures grown without KCl; lower gel, bacterial cultures supplemented with 5% KCl. (**C**) Co-immunoprecipitations (Co-IP). Upper gel: IP-anti-IglA, followed by WB with anti-IglB; lower gel: IP-anti-IglB, followed by WB with anti-IglA. (**D**) WBs of sucrose gradient fractions of the Y139F and Y139A mutant strains. (**E**) Distribution of the intensities (trasformed in log 2) of

1062 the proteins IglA and IglB in the Heavy (H) and light (L) sucrose fraction, in cp WT (Δ*iglB*

1063 complemented with WT IglB), Y139F and Y139A (Δ*iglB* complemented with IglB Y139F and

1064 IglB Y139A, respectively).

1065

Figure 5. Y139Aand Y139F IglB mutants show impaired intracellular survival

and attenuated virulence

(**A**) Intracellular bacterial multiplication of wild-type *F. novicida* (WT, Grey squares), isogenic

Δ*iglB* mutant (Δ*iglB*, blue triangles) and complemented Δ*iglB* strain (Cp-WT, grey circles;

Y139A, blue square; Y139F, red squares) and the Δ*FPI* control (black lines), was monitored

during 24 h in J774A.1 macrophage cells. The values recorded for Y139F, at 24 h, were

significantly different from those of Δ*FPI*, Δ*iglB* or Y139A. In both cell types, the values

recorded for WT or CpWT, at 10 h and 24 h, were significantly different from those of Δ*FPI*,

Δ*iglB,* Y139A or Y139F. **p<0.001 (determined by student's t-test). (**B**) Confocal microscopy

of intracellular bacteria with J774-1 infected with wild-type *F. novicida* (WT), Δ*iglB* or

complemented strains and their co-localization with the phagosomal marker LAMP1

observed at 1 h, 4 h and 10 h. Upper par: J774.1 were stained for *F. novicida* (green), LAMP-

1 (red) and host DNA (blue, DAPI stained). Lower part: analysis was performed with ImageJ

software. *p<0.01 (determined by student's t-test).

1080

1081

Figure 6. Electron microscopy and time lapse video microscopy

(**A**) Transmission electron microscopy of J774-1 cells infected with Y139F, Y139A or WT *F.*

novicida at 24 h. Black arrowheads (in cells infected with Y139A) point to single bacteria. (**B**)

Time lapse video microscopy. J774-1 cells expressing a nuclear restricted Red Fluorescent

Protein were infected with GFP-expressing WT and Y139 IglB proteins. Representative

1087 images were taken at 4,h, 8 h and 20 h (see Video S1 and S2). Red: cell nuclei. Green: GFP

1088 expressing bacteria.

1089

Figure 7. Proposed model of phosphorylation-dependent TSS6 disassembly

Upon bacterial entry into the phagosomal compartment, non-phosphorylated IglB form stable

heterodimers with IglA and sheath polymerization initiates at the bacterial pole. Within 30 min,

T6SS sheath contraction triggers the secretion of effectors and the rupture of the

phagosomal membrane. Immediately after, the T6SS sheath disassembles, thanks to the

concerted action of the ClpB unfoldase and of the phosphorylation of IglB residue Y139.

1096

1097

Supplementary materials

 Figure supplement 1. Schematic representation of the FPI locus and overview

 of the T6SS structure of *Francisella*

 The *Francisella* FPI locus representation (*anmK* to *pdpA* gene) with nomenclature of the

 canonical T6SS and the *F. novicida* T6SS. A color was assigned for each gene function and

 applied on the overview of the T6SS structure. Red/Green/Blue/Orange, structural protein;

 Grey, secreted protein; White, hypothetical protein

 Figure supplement 2. Schematic representation of the genomic regions

 encoding proteins whose expression vary upon KCl stimulation.

 (**A**) Genomic organization of the FNI; (**B**). the Fe-S cluster- and iron uptake-associated loci.

 Genes colored in blue correspond to proteins present in decreased amounts; genes colored

 in rose, to proteins present in increased amounts, genes in white, unchanged.

 Figure supplement 3. Alignments of orthologous IglB proteins

  (**A**) Alignments of orthologous IglBs proteins showing conservation of Y139 in *Francisellae*.

 The IglB sequences retrieved from 19 *Francisella* genomes were aligned using MUSCLE

 (Edgar, 2004) and rendered using ESPript3.0 (Robert & Gouet, 2014). Secondary structure

 elements information is extracted from 3J9O PDB file (Clemens et al., 2015) and is displayed

 at the top of the alignment. Sequences are identified by corresponding species name

 followed by strain name and locus tag. Species names are abbreviated as follow: Fn,

 Francisella tularensis subsp. *novicida*; Ftm, *Francisella tularensis* subsp. *mediasiatica*; Fth,

 Francisella tularensis subsp. *holarctica*; Ftt, *Francisella tularensis* subsp. *tularensis*; Ft,

 Francisella tularensis; Fno, *Francisella noatunensis*; Fsp, *Francisella* sp. (**B**) Alignments of

 orthologous IglBs in other genera showing conservation of a tyrosine residue. Sequences of

11 representative IglB orthologs with a tyrosine residue in the vicinity of Y139 were aligned

using MUSCLE (Edgar, 2004) and rendered using ESPript3.0 (Robert & Gouet, 2014).

Secondary structure elements information is extracted from 3J9O PDB file (Clemens et al.,

2015) and is displayed at the top of the alignment. Sequences are identified by

corresponding Uniprot accession number. Species names are abbreviated as follow: FRATN,

Francisella tularensis subsp. novicida; SALTY, *Salmonella* Typhimurium; YERPE, *Yersinia*

pestis; BURMA, Burkholderia mallei; XANC5, Xanthomonas campestris; RALP1, Ralstonia

pickettii; PHOAA, Photorhabdus asymbiotica; ERWAC, Erwinia amylovora; ENTCC,

Enterobacter cloacae; BRAJP, Bradyrhizobium japonicum; RHIFR, Rhizobium fredii;

9GAMM, Serratia nematodiphila.

1134

1135 **Figure supplement 4. Effect of IglB mutations on growth in broth**

1136 Wild-type *F. novicida* (WT, Grey squares), isogenic Δ*iglB* mutant (Δ*iglB*, blue triangles), and

1137 complemented *iglB* strain (Cp-WT*,* grey circles; Y139A, purple square; Y139F, red

1138 square), were grown in Schaedler K3 (left panel) or in Tryptic Soy broth (TSB) supplemented

1139 with 0,2% cysteine (right panel).

1140

Figure supplement 5. Intracellular multiplication in bone marrow-derived

macrophages

(**A**) Intracellular bacterial multiplication of wild-type *F. novicida* (WT, Grey squares), isogenic Δ*iglB*

mutant (Δ*iglB*, blue triangles) and complemented Δ*iglB* strain in bone marrow-derived macrophages

(BMDM) from BALB/c mice. (**B**) CCF4 assay. Infection of $ASC^{-/-}$ BMDM were performed at MOI of 100

and the cytosol of BMDMs was loaded with CCF4, as described previously (Ramond et al., 2015). *F.*

novicida naturally expresses a β-lactamase able to cleave the CCF4. Bacterial escape into the host

cytosol is thus associated with a shift of the CCF4 probe fluorescence emission from 535 nm to 450

nm. The number of cells demonstrating CCF4 cleavage, determined at different time points (1 h, 3 h

and 6 h) by flow cytometry, reached above 50% at 6 h for both WT and CpWT complemented strains

whereas it was below 1% for the other mutant strains tested. (**C**) Groups of five female BALB/c

mice were infected intraperitoneally with a mixture of 100 CFU of wild-type *F. novicida* and

100 CFU of Δ*iglB* mutant strain or complemented strain (Cp-WT, Y139A, Y139F). Bacterial

burden was quantified in liver (L) and spleen (S) of mice. The data represent the competitive

index (CI) value (in ordinate) for CFU of mutant/wild-type of each mouse, after 48 h infection,

divided by CFU of mutant/wild-type in the inoculum. The CI recorded for the two *iglB* mutants

was very low in both spleen and livers 2.5 days after infection (close to 10^{-7}). As a control, we

also performed an *in vivo* competition assay between wild-type and the Δ*iglB* complemented

strain (WT and CpWT). The competition index recorded close to 1 for CpWT ($\leq 10^{-1}$ in both

target organs), confirmed almost complete functional complementation.

Bars represent the geometric mean CI value.

1162

Figure supplement 6. Substitution of Tyr139 by phosphomimetics (Glu or Asp)

(**A**) J774.1 macrophage-like cells were infected in DMEM with 100 MOI of wild-type *F.*

novicida (WT) and mutated strains (Y139E*,* Y139D*,* Δ*FPI*) for 24 h. Results are shown as the

average of $\log_{10}$ cfu mL^{-1} ± standard deviation. Each experiment was performed in triplicate.

, *p*<0.001 (as determined by two-tailed unpaired Student's *t*-test). (B**) WB with anti-IglB of

sucrose gradient fractions.

Figure supplement 7. Inactivation of genes encoding proteins bearing a

putative tyrosine kinase domain

J774.1 macrophage-like cells were infected in DMEM-High glucose with an MOI of 100 with

wild-type *F. novicida* (WT), three mutants corresponding to a transposon insertion into genes

FTN_1082, *FTN_1433*, *FTN_0459* (tnfn1_pw060323p01q162, tnfn1_pw060420p02q142,

tnfn1_pw060323p07q132, respectively from the BEI resources repository), here designated

Δ*1082*, Δ*1433*, Δ*0459,* for simplification), and a Δ*FPI* mutant (Δ*FPI*). Intracellular

multiplication was monitored at 10 h. Results are shown as the average of log10 cfu mL^{-1} ±

standard deviation. Each experiment was performed in triplicate.

1178

1179

1180 **Video supplement 1.** WT: J774.1 macrophage-like cells were infected in DMEM-High

1181 glucose at an MOI of 1,000 with wild-type *F. novicida* Δ*iglB* expressing pKK214::pGro*iglB*-

1182 pGro*gfp* (designated cpWT-GFP).

1183

1184 **Video supplement 2.** Y139F: J774.1 macrophage-like cells were infected in DMEM-High

1185 glucose at an MOI of 1,000 with wild-type *F. novicida* Δ*iglB* expressing

1186 pKK214::pGro*iglB*Y139F-pGro*gfp* (designated cpY139F-GFP)

1187

1188

Table supplement 1. Whole cell proteome of non-induced and KCl-induced *F.

novicida. In this table, we report the log2 (intensity) of the proteins identified in whole cell

fractions from non-induced and KCl-induced wild-type *F. novicida*. Peptides and sequence

coverage indicates the number of total peptides identified and the primary sequence

coverage for that protein across samples. The Q value and the score refer to the quality of

the identification.

Table supplement 2. Detailed list of phosphosites identified in *F. novicida*

In this table, we report: i) the average of the log2 (intensity) of the phosphosites identified

from three independent replicates; ii) the amino acid carrying the phosphorylation ("Amino

Acid"); iii) its position on the protein (Position); iv) the "localization probability" of the

phosphosite (1 being 100%); v) the number of replicates the phosphosite has been identified

in ("identified"); vi) the "sequence window" of 15 amino acids before and after the amino acid

phosphorylated; vii) the probability and score of identification and localization (PEP, score,

score for localization); viii) the "charge" of the peptide; and ix) the annotation according to

COG and EggNOG.

1204

1205 **Table supplement 3**. **Table of intensities for IglB peptide carrying Y139**

1206 **phosphorylation.**

1207

Table supplement 4. List of proteins identified in the "Heavy" (H) and "Light"

(L) Fraction in WT and in the Y139F mutant

WT_H, WT_L, Y139_H, Y139F_L report the log2 (intensity) of the proteins identified in the

fractions. Peptides and sequence coverage indicates the number of total peptides identified

and the primary sequence coverage for that protein across samples. The Q value and the

score refer to the quality of the identification.

1214

1215 **Table supplement 5. Strains and plasmids.**

1216

1217 **Table supplement 6. Primers.**

1218

1219

Figure 1. Ziveri et al.

A. **78 p-sites** (S, T, Y)

B. Classes of phosphorylated proteins

- Translation — 11
- Carbohydrate metabolism and transport — 7
- Coenzyme metabolism — 5
- Replication and repair — 4
- Energy production and conversion — 4
- PTM, chaperone — 4

C.

FTN_1323 iglB (Y139)
FTN_0159 rpsA (T232)
FTN_0883 pheS (S97)
FTN_0752 (S46)
FTN_0164 ftsZ (S376)
FTN_1590 oppD (S307)
FTN_0900 (S505, Y507,T510)
FTN_1576 tuf (S9)
FTN_1397 (Y628,Y629)
FTN_1359 recC (T33,S39)
FTN_1317 (S2)
FTN_1246 me (Y504)
FTN_1569 rplL (S34)
FTN_0761 (T134)
FTN_0352 tolQ (S234)
FTN_0610 erpA (S2)
FTN_0253 rpsH (S2)
FTN_1434 icd (S130)
FTN_1329 fbaA (T278)
FTN_0171 glsA (Y108)
FTN_0266 htpG (S336)
FTN_0491 (Y141)
FTN_1537 (T96,S97)
FTN_1329 fbaA (S50)
FTN_1237 ubiH (T128)
FTN_0706 (T158,T159,T160,T162)
FTN_0594 sucC (S221)
FTN_1109 (S66)
FTN_0756 fopA (S383)
FTN_0311 prmB (Y311)
FTN_1194 gcvH1 (S126)
FTN_1170 (S26)
FTN_1518 relA (S389)
FTN_1339 fabG (S41)
FTN_0781 talA (S248)
FTN_0347 fkpB (T153)
FTN_1332 gapA (T211)

noKCl_P_3, noKCl_P_4, noKCl_P_1, noKCl_P_2, KCl_P_3, KCl_P_2, KCl_P_4, KCl_P_1

D.

IglB phosphorylation

NLYDISSSDFFKK — H, L (log$_2$ intensity)

NLpYDISSSDFFKK — H, L (log$_2$ intensity)

Sucrose %
B
L — 15
35
55 — H
Optiprep 55

Figure 2. Ziveri et al.

Figure 3. Ziveri et al.

Figure 4. Ziveri et al.

A

B

WT ΔiglB Y139A Y139F

1h

4h

10h

Figure 5. Ziveri et al.

Figure 6. Ziveri et al.

Figure 7. Ziveri et al.